- ACTFR, (1994). *Burdekin River Irrigation Area Flora and Fauna Survey*. Australian Center for Tropical Freshwater Research, James Cook University, Townsville. Report No. 94/10.
- Arthington, A.H., J. Marshall, G. Rayment, H. Hunter and S. Bunn. (1997). Potential impacts of sugarcane production on the riparian and freshwater environment. In: "Intensive Sugar Cane Production: Meeting the Challenges Beyond 2000" (Eds B.A. Keating and J.R Wilson), CAB International, Wallingford, UK. pp. 403-421.
- ANZECC/ARMCANZ. 2000. National Water Quality Guidelines. Australia New Zealand Environment Conservation Council.
- Australian Nature Conservation Agency (1996). A directory of important wetlands in Australia: Second Edition. ANCA, Canberra.

Cech et al. (1985) Envirinmental Biol of Fishes 13(4): 297-307 (o2 - Gambusia affinis)

- Herbert and Wells (2001) J. Comparative Physiol B Biochemical Systemic and Environmental Physiol 171(7): 603-612 (gourami 02)
- Battle, J.M. & Mihuc, T.B. (2000). Decomposition dynamics of aquatic macrophytes in the lower Atchafalaya, a large floodplain river. Hydrobiologia, **418**: 123-136.

Bayley, P.B. (1995). Understanding Large river-floodplain ecosystems. Bioscience 45: 153-158.

Bayly I.A.E and Williams W.D., 1973. Inland waters and their ecology. Longman, Australia.

- Beumer, J.P. 1979. Reproductive cycles of two Australian freshwater fishes: the spangled perch, *Therapon unicolor* Gunther, 1959 and the East Queensland rainbowfish, *Nematocentris splendida* Peters, 1866.
- Beumer, J.P. 1980. Hydrology and fish diversity of a north Queensland tropical stream. Aust. J. Ecol. 5:159-186.
- Bishop, K.A. & Forbes, M.A. (1991). The freshwater fishes of northern Australia. pp. 79-107. In Monsoonal Australia: Landscape, ecology and man in the northern lowlands. (Eds.) C.D. Haynes, M.G. Ridpath & M.A.J. Williams. A.A. Balkema, Rotterdam.

- Bishop, K.A., Allen S.A., Pollard D.A. and Cook M.G., 1980. Ecological studies on the freshwater fishes of the Alligator Rivers Region, Northern Territory. Supervising Scientists for the Alligator Rivers Region Research Report 4 Vol. I: Outline of the study, summary conclusions and recommendations.
- Bishop, K.A., Pidgeon, R.W. & Walden, D.J. (1995). Studies on fish movement dynamics in a tropical floodplain river: Prerequisites for a procedure to monitor the impacts of mining. Aust. J. Ecol. 20:81-107.
- Bishop, K.A.; Allen, S.A.; Pollard, D.A. & Cook, M.G. (2001). Ecological studies on the freshwater fishes of the Alligator Rivers Region, Northern Territory: Autecology. Supervising Scientists Report 145, Supervising Scientists, Darwin.
- Blackman, J.G.; Perry, T.W.; Ford, G.I.; Craven, S.A.; Gardiner, S.J. & De Lai, R.J. (1999). Characteristics of important wetlands in Queensland. Environmental Protection Agency. Queensland.
- Bruton, M.N. (1985). The effects of suspensoids on fishes. PP.221-224. *In:* B.R. Davies & R.D.Walmsley (ed.) Southern Hemisphere limnology, Dr. W. Junk Publishers, Dordrecht.
- Bunn S E; Davies P M; Kellaway D M; Prosser I P. (1998). Influence of invasive macrophytes on channel morphology and hydrology in an open tropical lowland stream, and potential control by riparian shading. Freshwater Biology. 39(1). 171-178.
- Bunn S E; Davies P M; Mosisch T D. 1999. Ecosystem measures of river health and their response to riparian and catchment degradation. Freshwater Biology. 41(2). 333-345.
- Bunn S.E.; Boon P.I. (1993). What sources of organic carbon drive food webs in billabongs? A study based on stable isotope analysis. Oecologia (Heidelberg). 96(1). 85-94.
- Bunn Stuart E; Davies Peter M; Kellaway Dominica M. 1997 Contributions of sugar cane and invasive pasture grass to the aquatic food web of a tropical lowland stream. Marine & Freshwater Research. 48(2). 173-179.

Bunn, S.E. & Arthington, A.H. (2002). Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. Env. Mngt. 30: 492-507.

Butler, B. 2003.

- Cappo, M., Alongi, D.M., Williams, D. McB. & Duke, N. (1998). A review and synthesis of Australian fisheries habitat research. Vol. 2: Scoping Review. Australian Institute of Marine Science. 371 p.
- Cech, J.J. Jr., Massingill, M.J., Vondrachek, B. and Linden, A.L. (1985). Respiratory metabolism of Mosquito fish, *Gambusia affinis*: effects of temperature, dissolved oxygen and sex difference. Environmental Biology of fishes 13: 297-307
- Clapcott, J.E. & Bunn, S.E. (2003). Can C₄ plants contribute to aquatic food webs of subtropical streams? Freshwater Biology **48**: 1105-1116.
- Congdon, R.A., Pearson, R.G., Lukacs, G.P. and Nolen, J. (1994). Limnology and classification of tropical floodplain wetlands, with particular reference to the effects of irrigation drainage. Australian Center for Tropical Freshwater Research, James Cook University, Townsville. Report No. 94/02.
- Cordo, H. & Center, T.D. (2000) Watch out water-hyacinth! New jungle enemies are coming. Argicultural Research, March 2001.
- Cotterell, E. (1998). Fish passage in streams: Fisheries guidelines for design of stream crossings. Fish Habitat Guideline FHG 001, Fisheries Group, DPI Queensland.
- Davis, T.L.O., (1988). Temporal changes in the fish fauna entering a tidal swamp system in tropical Australia. Environmental Biology of Fishes. 21: p. 161-172.

- Dehadrai, P.V. & Tripathi, S.D. (1976) Environment and ecology of freshwater air-breathing teleosts. Pages 39-72. In Hughes, G.M. (Ed.) *Respiration of amphibious vertebrates*. Academic Press, London.
- Douglas, M.M. & O'Connor, R.A. (2003). Effects of exotic macrophyte, para grass (Urochloa mutica), on benthic and epiphytic macroinvertebrates of a tropical floodplain. Freshwater Biology, 48: 962-971.
- EPA. (1999). State of the Environment Queensland 1999. Queensland Environmental Protection Agency, Brisbane.
- Erskine, W.D. and Webb, A.A. (2003) Desnagging to resnagging: New directions in river rehabilitation in Southeastern Australia. River Res. Applic. **19**: 233-249.
- Finlayson, C.M.; Hall, R. & Bayliss, B. (1998). Regional review of wetland management issues: Wet-Dry tropics of northern Australia. LWRRDC National wetlands R&D Program, LWRRDC Occasional paper 03/97, Canberra.
- Gehrke, P.C. (1990). Spatial and temporal dispersion patterns of golden perch, *Macquaria ambigua*, larvae in an artificial floodplain environment. J. Fish Biol. **37**: 225-236.
- Gehrke, P.C. (1991). Avoidance of inundated floodplain habitat by larvae of golden perch (*Macquaria ambigua* Richardson): influence of water quality or food distribution? Aust. J. Mar. Freshwater Res. 42: 707-719.
- Geiger, S.P., Torres, J.J. and Crabtree, R.E. (2000). Air breathing and gill ventilation frequencies in juvenile tarpon, *Megalops atlanticus*: responses to changes in dissolved oxygen, temperature, hydrogen sulfide, and pH. Env. Biol. Fishes **59**: 181-190.
- Gopal, B. & Junk, W.J. (2000). Biodiversity in wetlands: An introduction. In "Biodiversity in wetlands: Assessment, Function and Conservation. (Eds Gopal, B.; Junk, W.J. & Davis, J.A.), Backhuys Publishing, Leiden. pp.

Gore, J.A. & Shields, F.D.Jr. (1995). Can large rivers be restored?. Bioscience 45: 142-152.

- Gutierrez, E.L., Ruiz, E.G., Uribe, E.G. & Martinez, J.M. (2001) Biomass and productivity of Water Hyacinth and their application in control programs. In "Biological and integrated control of Water Hyacinth: Eichhornia crassipes. Proceedings of the second meeting of the global working group for the biological and integrated control of Water Hyacinth, Beijing, China, 9-12 October 2000" (Eds. Julien, M.H., Hill, M.P., Center, T.D. & Jianquing, D.). Australian Centre for International Agricultural Research, Canberra. pp.109-120.
- Halliday, I., Ley, J., Tobin, A., Garrett, R., Gribble, N. and Mayer, D. (2001) The effects of net fishing: addressing biodiversity and bycatch issues in Queensland inshore waters.
- Harris, J.H. (1995) The use of fish in ecological assessments. Aust. J. Ecol. 20: 65-80.
- Hayes C.D., Ridpath M.G. and Williams M.A.J., (Eds.).1991. Monsoonal Australia: Landscape, ecology and man in the northern lowlands. A.A. Balkema, Rotterdam.
- Herbert, B., Peeters, J., Graham, P. & Hogan, A. (1995). Natural Resources Assessment Program Fish Fauna Survey Project NR 10, CYPLUS report, QDPI.
- Herbert, N. A. and Wells, R. M. G. (2001) The aerobic physiology of the air-breathing blue gourami, Trichogaster trichopterus, necessitates behavioural regulation of breath-hold limits during hypoxic stress and predatory challenge. Journal of Comparative Physiology - B, Biochemical, Systemic, & Environmental Physiology. 171: 603-612.
- Hillman, T. J. 1986, Billadongs. In De Dekker P. and Williams W. D. Limnology in Australia. CSIRO\W. Junk Publishing.Melborne.
- Hogan, A. & Graham, P. (1994). Tully-Murray floodplain fish distribution and fish habitat. Freshwater fisheries and aquaculture center, Department of Primary Industries. Walkamin, Queensland
- Hopley, D. (1970). The geomorphology of the Burdekin Delta North Queensland. Monograph Series No. 1, James Cook University, Townsville.

- Hortle, K.G. & R.G. Pearson. 1990. Fauna of the Annan River system, far north Queensland, with reference to the impact of tin mining. I. Fishes. Aust. J. Mar. Freshw. Res. 41: 677-694.
- Houston, W.A. & Deivenvoorden, L.J. (2002). Replacement of littoral native vegetation with the ponded pasture grass *Hymenachne amplexicaulis*: effects on plant, macroinvertebrate and fish biodiversity of backwaters in the Fitzroy River, Central Queensland, Australia. J. Mar. Freshwater Res. 53: 1235-44.
- Humphries P.; King Alison J; Koehn John D. 1999. Fish, flows and flood plains: Links between freshwater fishes and their environment in the Murray-Darling river system, Australia. Environmental Biology of Fishes. 56(1-2). 129-151.
- Hunsaker, C.T and Levine, D.A. (1995) Hierarchical approaches to the study of water quality in rivers. Biosciense **45**: 193-203.
- Jepsen, D.B. (1997). Fish species diversity in sand bank habitats of a neotropical river. Env. Biol. Fish. **49**: 449-460.
- Johnson, A.K.L., Ebert, S.P. & Murray, A.E. (1997). Spatial and temporal distribution of wetland and riparian zones and opportunities for their management in catchments adjacent to the Great Barrier Reef Marine Park. Proc. GBRMPA Cross-sectoral workshop on wetlands and water quality, Babinda, Queensland, 25-26 September.
- Johnson, B.L., Richardson, W.B. & Naimo, T.J. (1995). Past, Present and Future Concepts in Large River Ecology. Bioscience. **45**: 134-141.
- Julien M. H., Griffiths M. W. and Stanley J. N., 2001 Biological control of water hyacinth 2: The moths *Niphograpta albiguttalis* and *Xubida infusellus*: biologies, host ranges, and rearing, releasing and monitoring techniques for biological control of *Eichhornia crassipes*. ACIAR Monograph series, Canberra.
- Junk W.J. (ed.), (1997). The central Amazon floodplain: Ecology of a pulsing system. Springer, Berlin.

- Junk W.J., Bayley, P.B. & Sparks, R.E. (1989). The flood pulse concept in river-floodplain systems. Pp. 110-27. In *Proceedings of the International large river* syposium. Can. Spec. Publ. Fish. Aquat. Sci. 106.
- Kaenel, B.R.; Buehrer, H. & Uehlinger, U. (2000). Effects of aquatic plant management on stream metabolism and oxygen balance in streams. Freshwater Biol. 45: 85-95.
- Kennard, M.J. (1995). Factors influencing freshwater fish assemblages in floodplain lagoons of the Normanby River, Cape York Peninsula: a large tropical Australian river. Masters of Philosophy Thesis, Griffith University, Brisbane. 225 p.
- Kerr, J. (1994) Black snow and liquid gold: a history of the Burdekin Shire. Burdekin Shire Publication. Ayr, Queensland, Australia. 320pp.
- Lokkers, C., Perry, T., James, D. and Perna, C. (2000). Property management assessment, *Wongaloo* grazing property. Earthworks Environmental Services report No. 98c14.2. Townsville, Queensland, Australia.
- Lowe-McConnell R.H., (1979). Ecological aspects of seasonality in fishes of tropical waters. Symp. zool. Soc. Lond. 44: 219-241.
- Lowe-McConnell R.H., 1987. Ecological studies in tropical fish communities. Cambridge University Press. Cambridge.
- Lukacs, G.P. (1995) Wetlands of the lower Burdekin region, North Queensland. In *Wetland* research in the wet-dry tropics of Australia, (ed.) C.M. Finlayson. Supervising Scientists Report 101, pp. 86-92.
- Lukacs, G.P. and Finlayson, C.M. (in press). Status of wetlands in Northern Australia. In Proceedings of the Australian Water Association "Sustaining our aquatic environments conference". Townsville, November, 2001.

- Lynch Romeny J; Bunn Stuart E ; Catterall Carla P. 2002. Adult aquatic insects: Potential contributors to riparian food webs in Australia's wet-dry tropics. Austral Ecology. 27. 515-526
- Mackey, A.P. (1991) Aspects of limnology of Yeppen Yeppen Lagoon, Central Queensland. Aust. J. Mar. Freshwater Res., 42:309-325.
- Masifwa, W.F., Twongo, T. & Denny, P. (2001). The impact of water hyacinth, *Eichhornia crassipes* (Mart) Solms of the abundance and diversity of aquatic macroinvertebrates along the shores of northern Lake Victoria, Uganda. Hydrobiologia, **452**: 79-88.
- Matthews, W.J. (1998) Patterns in freshwater fish ecology. Chapman & Hall, New York.
- McCormick, P.V. & Laing, J.A. (2003). Effects of increased phosphorus loading on dissolved oxygen in a subtropical wetland, the Florida Everglades. Wetlands Ecology and Management. **11:** 199-215.
- Munshi, J.S.D. (1976). Respiratory organs of air-breathing fishes. Pages 73-104. In Hughes, G.M. (Ed.) *Respiration of amphibious vertebrates* Academic press. London.
- Nicholls, A.O. (1980) Vegetation. In Burdekin Project Ecological Sutdy Eds., Fleming, P.M., Gunn, R.H., Reece, A.M. and McAlpine, J.R., Australian Government Publishing Service, Canberra.
- Nilsson, C. & Svedmark, M. (2002). Basic principles and ecological consequences of changing water regimes: riparian plant communities. Env. Mangt. **30**: 468-480.
- Norris, R.H. and Norris, K.H. (1995) The need for biological assessment of water quality: Australian perspective. Australian Journal of Ecology. **20**: 1-6.
- North Burdekin Water Board. (2000) Thirty-fifth annual report 1999/2000. North Burdekin Water Board, Ayr, Queensland, Australia.
- North Burdekin Water Board. (2001) Thirty-sixth annual report 2000/2001. North Burdekin Water Board, Ayr, Queensland, Australia.

- North Burdekin Water Board. (2002) Thirty-seventh annual report 2001/2002. North Burdekin Water Board, Ayr, Queensland, Australia.
- Ogutu-Ohwayo, R., Hecky, R.E. Cohen, A.S. & Kaufman, L. (1997). Human impacts of the African Great Lakes. Env. Biol. Fish. **50**: 117-131.
- Pearson, R.G., (2003). Effects of Cane-Field drainage of the ecology of tropical waterways. Australian Center for Tropical Freshwater Research, James Cook University, Townsville. Report No. 03/04
- Perna C. 1996. Seasonal dynamics of coastal stream fish communities in the Townsville region., Unpublished Honours Thesis, James Cook University, Townsville. 83 p.
- Perna C. and West A. 1998. Water Management for sustainable wet tropical floodplains. Dept. of Natural Resources, Sth. Johnstone, QLD.
- Puckridge, J.T.; Sheldon, F.; Walker, K.F. & Boulton, A.J. (1998). Flow variablility and the ecology of large rivers. J. Mar. Freshwater Res. **49**: 55-72.
- Pusey B J ; Arthington A H; Bird J R; Close P G. (2001).Reproduction in three species of rainbowfish (Melanotaeniidae) from rainforest streams in northern Queensland, Australia. Ecology of Freshwater Fish. 10(2). 75-87.
- Pusey B J; Arthington A H; Read M G. (2000) (b). The dry-season diet of freshwater fishes in monosoonal tropical rivers of Cape York Peninsula, Australia. Ecology of Freshwater Fish. 9(3). 177-190.
- Pusey B J; Kennard M J; Arthington A H. (2000) (a). Discharge variability and the development of predictive models relating stream fish assemblage structure to habitat in northeastern Australia. Ecology of Freshwater Fish. 9(1-2). 30-50.

- Pusey Bradley J ; Arthington Angela H; Read Martin G. 1998. Freshwater fishes of the Burdekin River, Australia: Biogeography, history and spatial variation in community structure. Environmental Biology of Fishes. 53(3). 303-318.
- Pusey Bradley J ; Kennard Mark J ; Arthur James M; Arthington Angela H . 1998. Quantitative sampling of stream fish assemblages: Single-vs multiple-pass electrofishing. *Australian* Journal of Ecology. 23(4). 365-374.
- Pusey Bradley J ; Kennard Mark J. 1996. Species richness and geographical variation in assemblage structure of the freshwater fish fauna of the wet tropics region of northern Queensland. Marine & Freshwater Research. 47(3). 563-573.
- Pusey Bradley J ; Read Martin G; Arthington Angela H. 1995 (b). The feeding ecology of freshwater fishes in two rivers of the Australian wet tropics. Environmental Biology of Fishes. 43(1). 85-103.
- Pusey Bradley J; Arthington Angela H ; Read Martin G. 1995. (a) Species richness and spatial variation in fish assemblage structure in two rivers of the Wet Tropics of northern Queensland, Australia. Environmental Biology of Fishes. 42(2). 181-199.
- Pusey, B.J. & Arthington, A.H. (2003). Importance of the riparian zone to the conservation and management of freshwater fishes: a review. J. Mar. Freshwater Res.
- Rayment, G.E. (2002). Land use and surface water quality in sugar catchments. CRC for Sustainable Sugar Production, Water Issue. Indooroopilly.
- Roggeri H. 1995. Tropical freshwater wetlands: A guide to current knowledge and sustainable management. Klewer Academic, Dordrecht.
- Russell, D.J. Hales, P.W. (1993). Stream habitat and fisheries resources of the Johnstone River Catchment, Northern Fisheries Centre, Queensland Department of Primary Industries, Cairns.

- Russell, D.J.; Hales, P.W. & Helmke, S.A. (1996). Stream habitat and fish resources in the Russell and Mulgrave Rivers catchment, Information Series QI96008, Northern Fisheries Centre, Queensland Department of Primary Industries, Cairns.
- Scheffer, M., Szaba, S., Grangnani, A., van Nes, E.H., Rinald, S., Kautsky, N., Norberg, J., Roijackers, R.M.M., and Franken, R.J.M. (2003). Floating plant dominance as a stable state. Proc. Nat. Acad. Sci. USA. 100(7): 4040-4045.
- Schlosser, I.J. (1987). The role of predation in age- and size-related habitat use by stream fishes. Ecology. **68:** 651-659.
- South Burdekin Water Board. (2000) Thirty-fifth annual report 1999/2000. South Burdekin Water Board, Home Hill, Queensland, Australia.
- South Burdekin Water Board. (2000) Thirty-sixth annual report 2000/2001. South Burdekin Water Board, Home Hill, Queensland, Australia.
- South Burdekin Water Board. (2000) Thirty-seventh annual report 2001/2002. South Burdekin Water Board, Home Hill, Queensland, Australia.
- Sparks, R.E. (1995). Need for ecosystem management of large rivers and their floodplains. Bioscience. **45**: 168-182.
- Storrs, M.J. & Finlayson, C.M. (1998). Overview of the conservation status of wetlands of the Northern Territory. Supervising Scientists Report 116, Supervising Scientists, Canberra.
- Swales Stephen [a]; Storey Andrew W; Bakowa Kayemen A [a]. Temporal and spatial variations in fish catches in the Fly River system in Papua New Guinea and the possible effects of the Ok Tedi copper mine. Environmental Biology of Fishes. 57(1). Jan., 2000. 75-95.
- Swales Stephen [a]; Storey Andrew W; Roderick Ian D; Figa Boga S. Fishes of floodplain habitats of the Fly River system, Papua New Guinea, and changes associated with El Nino droughts and algal blooms. Environmental Biology of Fishes. 54(4). April, 1999. 389-404.

- Tait, J. & Perna, C. (2001) Fish Habitat Management Challenges on an Intensively Developed Tropical Floodplain. RipRap 19. Land and Water Australia, Canberra
- Tait, J. (1994). Lowland habitat mapping and management recommendations: Tully-Murray catchments. Final Report. (An ICM initiative of the Queensland Department of Primary Industries), Cardwell Shire Catchment Coordinating Committee, Cardwell.
- Tallqvist, Minna; Sandberg-Kilpi, Eva & Bonsdorff, Erik (1999) Juvenile flounder, *Platichthys flesus* (L.), under hypoxia: effects on tolerance, ventilation rate and predation efficiency. Journal of Experimental Marine Biology and Ecology 242: 75 93.
- Taylor, J.C. & Miller, J.M. (2001) Physiological performance of juvenile southern flounder, *Paralichthys lethostigma* (Jordan and Gilbert, 1884), in chronic and episodic hypoxia. Journal of Experimental Marine Biology and Ecology 258: 195-214.
- Tejerina-Garro, F.L. & De Merona, B. (2001). Spatial variability of biotic and abiotic factors of the aquatic habitat in French Guiana. Regul. Rivers: Res. & Mngt. **17**: 157-169.
- Tejerina-Garro, F.L.; Fortin, R. & Rodriquez, M.A. (1998). Fish community structure in relation to environmental variation in floodplain lakes of the Araguaia River, Amazon Basin. Env. Biol. Fish. 51: 399-410.
- Thorpe, J.H. & Delong, M.D. (1994). The riverine productivity model: an heuristic view of carbon sources and organic processing in large river ecosystems. Oikos **70**: 307-309.
- Townsend, C.R. (1989). The patch dynamics concept of stream community ecology. J. Nth. Am. Benthol. Soc. 8: 36-50.
- Vannote, R.L., Minshall, G.W., Cummins, K.W., Sedell, J.R. & Cushing, C.E. (1980). The river continuum concept. Can. J. Fish. Aquat. Sci. 37; 130-137.
- Walker Keith F; Puckridge Jim T; Blanch Stuart J. 1997. Irrigation development on Cooper Creek, central Australia-prospects for a regulated economy in a boom-and-bust ecology. Aquatic Conservation. 7(1). 63-73.

- Webb, A.C. (2003). The ecology of invasions of non-indigenous fishes in Northern Queensland. PhD Thesis, James Cook University, Townsville, Australia
- Welcomme, R.L. (1979). Fisheries ecology of floodplain rivers. Longman, London.
- Wetzel, R.G. (1983). Limnology: Second Edition.. CBS college Publishing, New York. 767 p.
- Wijeyaratne, M.J.S. & Perera, W.M.D.S.K. (2000). Studies on the feasibility of using indigenous fishes for controlling aquatic macrophytes in Sri Lanka. J. Aquaculture in the Tropics. 15(3):252-260.
- Willoughby, N., Watson, I.G., Lauer, S. & Grant, I.F. (1993). An investigation into the effect of water hyacinth in the biodiversity and abundance of fish and invertebrates in Lake Victoria, Uganda. NRI Project Report 10066 Ao32g, Natural Resources Institute, Chathan, U.K.
- Winemiller, K.O. & Leslie, M.A. (1992). Fish assemblages across a complex, tropical freshwater/marine ecotone. Env. Biol. Fish. **34**: 29-50.
- Wood, P.J. & Armitage, P.D. (1997). Biological effects of fine sediment in the lotic environment. Env. Mngt. 21: 203-217
- Zalewski, M. and Welcomme, R. (2001) Restoration of sustainability of physically degraded fish habitats: The model of Intermediate Restoration. Ecohydrology & Hydrobiology. **1(3)**: 279-282.