

Title: Are myocardial infarction-associated single nucleotide polymorphisms associated with ischemic stroke?

Authors:

Yu-Ching Cheng, PhD¹; Christopher D. Anderson, MD^{2,3,4}; Silvia Bione, PhD⁵; Keith Keene, PhD^{6,7}; Jane M. Maguire PhD, RN^{8,9}; Michael Nalls, PhD¹⁰; Asif Rasheed, MBBS¹¹; Marion Zeginigg, MSc¹²; John Attia, PhD, MD, FRCPC, FRACP⁸; Ross Baker, FRACP, FRCPA^{13,14}; Simona Barlera, MSc¹⁵; Alessandro Biffi, MD^{2,3,4}; Ebony Bookman, PhD¹⁶; Thomas G. Brott, MD¹⁷; Robert D. Brown Jr., MD¹⁸; Fang Chen, PhD⁷; Wei-Min Chen, PhD^{7,19}; Emilio Ciusani, PhD²⁰; John W. Cole, MD^{21,22}; Lynelle Cortellini, MSc^{2,3,4}; John Danesh, FRCP²³; Kimberly Doheny, PhD²⁴; Luigi Ferrucci, MD, PhD²⁵; Maria Grazia Franzosi, PhD¹⁵; Philippe Frossard, PhD, DSc¹¹; Karen L. Furie, MD, MPH³; Jonathan Golledge, MChir, FRACS²⁶; Graeme J. Hankey, MD, FRACP, FRCP, FRCPE^{27,28}; Dena Hernandez, MS¹⁰; Elizabeth G. Holliday, PhD⁸; Fang-Chi Hsu, PhD^{29,30}; Jim Jannes, PhD, BMBS BMBS FRACP³¹; Ayeesha Kamal, MD, FAHA, ABVN³²; Muhammad Saleem Khan, MSc³³; Steven J. Kittner, MD^{21,22}; Simon A. Koblar, PhD, BMBS FRACP³¹; Martin Lewis, PhD³¹; Lisa Lincz, PhD⁸; Antonella Lisa, PhD⁵; Mar Matarin, PhD³⁴; Pablo Moscato, PhD³⁵; Josyf C. Mychaleckyj, DPhil^{7,19}; Eugenio A. Parati, MD²⁰; Silvia Parolo⁵; Elizabeth Pugh, PhD²⁴; Natalia S. Rost, MD^{2,3,4}; Michael Schallert MSc¹²; Helena Schmidt, MD, PhD¹²; Rodney J. Scott, PhD, PD, FRCPath, FHGSA⁸; Jonathan W. Sturm, PhD, MD³⁶; Sunaina Yadav, MSc³³; Moazzam Zaidi, MBBS¹¹; GARNET Collaborative Research Group³⁷; GENEVA Consortium³⁸; Giorgio B. Boncoraglio, MD²⁰; Christopher Royce Levi, MD, FRACP, RACP⁸; James F. Meschia, MD¹⁷; Jonathan Rosand, MD, MSc^{2,3,4}; Michele Sale, PhD^{6,7,39}; Danish Saleheen, MBBS, PhD^{11,23,40}; Reinhold Schmidt, MD¹²; Pankaj Sharma,

MD PhD FRCP³³; Bradford Worrall, MD, MSc^{19,41}; Braxton D. Mitchell, PhD¹; on behalf of the International Stroke Genetics Consortium

Institutions:

¹ Department of Medicine, University of Maryland, Baltimore, MD USA

² Center for Human Genetic Research, Massachusetts General Hospital, Boston, MA, USA

³ Department of Neurology, Massachusetts General Hospital, Boston, MA, USA

⁴ Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA

⁵ Institute of Molecular Genetics, National Research Council, Pavia, Italy

⁶ Department of Medicine, University of Virginia, VA, USA

⁷ Center for Public Health Genomics, University of Virginia, VA, USA

⁸ Hunter Medical Research Institute (HMRI), University of Newcastle, Australia

⁹ School of Nursing and Midwifery, University of Newcastle, Australia

¹⁰ Laboratory of Neurogenetics, National Institute of Aging, National Institute of Health, Bethesda, MD, USA

¹¹ Center for Non-Communicable Diseases, Karachi, Pakistan

¹² Department of Neurology and Institute of Molecular Biology and Biochemistry, Medical University Graz, Austria

¹³ Centre for Thrombosis and Haemophilia, Murdoch University, Australia

¹⁴ Department of Haematology, Royal Perth Hospital, Australia

¹⁵ Istituto di Ricerche Farmacologiche Mario Negri, Milano, Italy

¹⁶ National Human Genome Research Institute, National Institutes of Health, Rockville, MD, USA

¹⁷ Department of Neurology, Mayo Clinic, Jacksonville, FL, USA

- ¹⁸ Department of Neurology, Mayo Clinic, Rochester, MN, USA
- ¹⁹ Department of Public Health Sciences, University of Virginia, VA, USA
- ²⁰ Fondazione IRCCS Istituto Neurologico "Carlo Besta", Milano, Italy
- ²¹ Department of Neurology, Veterans Affairs Medical Center, Baltimore, MD, USA
- ²² Department of Neurology, University of Maryland School of Medicine, Baltimore, MD, USA
- ²³ Department of Public Health and Primary Care, University of Cambridge, UK
- ²⁴ Center for Inherited Disease Research, Johns Hopkins University School of Medicine,
Baltimore, MD, USA
- ²⁵ Longitudinal Studies Section, Clinical Research Branch, National Institute of Aging, National
Institute of Health, Bethesda, MD, USA
- ²⁶ Vascular Biology Unit, School of Medicine, James Cook University, Australia
- ²⁷ Department of Neurology, Royal Perth Hospital, Australia
- ²⁸ School of Medicine and Pharmacology, University of Western Australia, Australia
- ²⁹ Department of Biostatistical Sciences, Wake Forest School of Medicine, NC, USA
- ³⁰ Center for Cancer Genomics, Wake Forest School of Medicine, NC, USA
- ³¹ Stroke Research Programme, Schools of Medicine & Molecular Biomedical Science,
University of Adelaide, Australia
- ³² Section of Neurology, Aga Khan University, Pakistan
- ³³ Imperial College Cerebrovascular Research Unit (ICCRU), Imperial College London, UK
- ³⁴ Department of Clinical and Experimental Epilepsy, UCL Institute of Neurology, London, UK
- ³⁵ Centre for Bioinformatics, Biomarker Discovery and Information-based Medicine,
The University of Newcastle, Australia
- ³⁶ Neurosciences Department, Gosford Hospital, University of Newcastle, Australia

³⁷ The Genomics and Randomized Trials Network

³⁸ Gene-Environment Association Studies

³⁹ Department of Biochemistry and Molecular Genetics, University of Virginia, VA, USA

⁴⁰ University of Pennsylvania, USA

⁴¹ Department of Neurology, University of Virginia, VA, USA

Corresponding author:

Braxton D. Mitchell, PhD

Department of Medicine, University of Maryland School of Medicine

660 W. Redwood St., Howard Hall Room 492, Baltimore, MD 21201, USA

Phone: 410-706-0161

Fax: 410-706-1622

Email: bmitchel@medicine.umaryland.edu

Running Title: MI-associated SNPs and Ischemic Stroke

List of Tables:

Table 1: Previous published SNPs that were associated with MI or CAD

Table 2: Association results of the 11 MI/CAD-risk loci and risk scores with ischemic stroke

Table 3: Association results of the 11 MI/CAD-risk loci and risk scores with ischemic stroke, by TOAST subtypes

Subject Codes: [8] Epidemiology, [55] Genetics of stroke

Key words: Cerebral infarct, Genetics, Ischemia

Word counts: 6,116

Abstract

Background and Purpose — Ischemic stroke (IS) shares many common risk factors with coronary artery disease (CAD). We hypothesized that genetic variants associated with myocardial infarction (MI) or CAD may be similarly involved in the etiology of IS. To test this hypothesis, we evaluated whether single nucleotide polymorphisms (SNPs) at 11 different loci recently associated with MI or CAD through genome-wide association studies were associated with IS.

Methods — Meta-analyses of the associations between the 11 MI-associated SNPs and IS were performed using 6,865 cases and 11,395 controls recruited from 9 studies. SNPs were either genotyped directly or imputed; in a few cases a surrogate SNP in high linkage disequilibrium was chosen. Logistic regression was performed within each study to obtain study-specific betas and standard errors. Meta-analysis was conducted using an inverse variance weighted approach assuming a random effect model.

Results — Despite having power to detect odds ratio of 1.09-1.14 for overall IS and 1.20-1.32 for major stroke subtypes, none of the SNPs were significantly associated with overall IS and/or stroke subtypes after adjusting for multiple comparisons.

Conclusions — Our results suggest that the major common loci associated with MI risk do not have effects of similar magnitude on overall IS, but do not preclude moderate associations restricted to specific IS subtypes. Disparate mechanisms may be critical in the development of acute ischemic coronary and cerebrovascular events.

Introduction

Ischemic strokes (IS) comprise 87% of all strokes¹. Nearly 800,000 individuals in the US experience a stroke each year, making stroke one of the leading causes of serious, long-term disability in developed countries^{1,2}. The tendency of stroke to aggregate within families implies a genetic etiology, although few genetic variants have as yet been unequivocally linked to the common forms of IS.

Some of the difficulty in identifying stroke susceptibility genes may arise from heterogeneity within the stroke phenotype, making it imperative to carry out genetic studies within specific subtypes of IS. IS syndromes can, in fact, be recognized by experienced clinicians with a high degree of inter-rater agreement³. The major categories, which are based upon the presumed stroke mechanism, include atherosclerotic (i.e. large artery) stroke, cardioembolic stroke, and small vessel (lacunar) stroke. Given disparate mechanisms, it is likely that different risk alleles would contribute to these different stroke subtypes. Previous studies showing that a family history of stroke is more common in younger compared to older age of onset strokes⁴⁻⁶ raise the additional possibility that the genetic contribution to stroke may differ by age of onset.

IS shares many risk factors with myocardial infarction (MI) and coronary artery disease (CAD), including smoking and high blood pressure. The possibility that common susceptibility genes might predispose to both IS and MI/CAD prompted us to examine whether genetic variants previously associated with MI/CAD might also be associated with IS. To address this hypothesis, we evaluated whether DNA sequence variants at 11 different loci recently associated with MI or CAD through genome-wide association (GWA) studies were also associated with IS. Our analysis included 6,865 IS cases from 9 different studies, allowing us to test whether the

genetic variants were associated not only with overall IS, but also with the major IS subtypes and with age of stroke onset.

Materials and Methods

Study Populations

This analysis includes 6,865 IS cases and 11,395 controls from 9 different studies: Australian Stroke Genetics Collaborative (ASGC), Bio-Repository of DNA in Stroke (BRAINS), Besta Cerebrovascular Diseases Registry (CEDIR), Genes Affecting Stroke Risk and Outcomes Study (GASROS), Genetics of Early Onset Stroke (GEOS), Graz Stroke Study (GRAZ), Ischemic Stroke Genetics Study and Siblings with Ischemic Stroke Study (ISGS/SWISS), Risk Assessment of Cerebrovascular Events (RACE) and the Vitamin Intervention for Stroke Prevention (VISP) study. All stroke cases, except those from VISP study, were adjudicated for ischemic stroke subtype using the original TOAST system³, which assigns each case to one of the following categories: cardioembolic, large artery, small vessel, other known causes and undetermined causes. Controls free of stroke were selected based on study-specific criteria. For this report, cases and controls with a known history of MI were excluded from the analyses. Details of the study designs for each participating study are included in the Supplemental Method (<http://stroke.ahajournals.org>).

All participating studies were conducted with the consent of study subjects and were approved by the Institutional Review Board in each institution.

SNP selection

SNPs shown previously to be associated with MI or CAD in Caucasians were identified through a review of genome-wide association studies published before May 2010. MI/CAD-associated SNPs reported in these previously published GWA studies were selected for inclusion in this study only if their associations with MI/CAD reached genome-wide significance ($P < 5 \times 10^{-8}$) and were replicated in at least one large independent Caucasian cohort. Based on these criteria, we identified 11 loci from nine chromosomes that were consistently associated with MI/CAD (see Table 1)⁷⁻¹³.

Genotyping and Imputation

The 11 MI-associated SNPs selected for inclusion in this study were genotyped by existing commercial genome-wide SNP platforms, KASPar SNP array (KBioscience, Herts, UK) and/or by Taqman assays (Applied Biosystems, Foster City, USA). For SNPs that were not directly genotyped, genotypes were obtained either via imputation or the targeted SNP was replaced by a surrogate SNP that was in high linkage disequilibrium ($LD r^2 > 0.8$) with the target SNP. A detailed description on the genotyping and imputation methods for each participating site is provided in Supplemental Table S1.

Statistical Analysis

Site-specific logistic regression analysis was performed to test for associations of each individual SNP with IS under an additive model, and then site-specific beta coefficients (β) and standard errors (S.E.) were pooled in meta-analyses. A genotype risk score was also computed for each individual by summing the number of MI/CAD-risk alleles across each of the 11 loci (score 1). Each SNP was given equal weights rather than weights based on their effect sizes (β)

for MI/CAD in the risk score calculation given that we do not know the effects of these SNPs on stroke and the assumption that they would have similar effects on stroke as they do on MI seems unjustified. Additionally, we computed a genotype risk score that summed the number of MI/CAD-risk alleles across 10 loci, excluding SNP rs4977574 on chromosome 9, because SNPs near this locus have previously been reported to be associated with large artery stroke (score 2)¹⁴. Each genotype risk score was used as a continuous independent variable in the logistic regression model to obtain the joint additive effects of all MI/CAD-associated SNPs on the risk of IS. The β of the genotype risk score represents the increase in log odds of stroke associated with having one additional MI/CAD-risk allele. The GRAZ study was not included in the score analyses because three of the 11 SNPs were not genotyped directly and could not be imputed with acceptable quality scores (rs6725887, rs9818870 and rs9982601). Overall, about 35% of the individuals in whom one or more SNPs were not genotyped or imputed were removed from the genotype risk score analysis. For ease of result interpretation, all odds ratios were designated with the previously published non-risk allele as the reference so that the odds ratio reflects the effect of the MI-associated risk allele. Association analyses were adjusted for study-specific covariates (e.g. age, sex and population structure) (see Supplemental Table S2).

Meta-analysis was performed using a variance-weighted approach assuming a random effect model (DerSimonian & Laird method)¹⁵, which takes heterogeneity of the genetic effects across studies into account. Between-study heterogeneity was estimated using Cochran's Q statistic, which is the summation of the squared weighted difference between the study-specific effect and the overall effect size (estimated by the fixed effect), as well as I² statistics, which represent the percent of total variation across studies due to heterogeneity after accounting for variability due to chance. All meta-analyses were performed using the *metan* module

implemented in STATA 10 (StataCorp, College Station, TX). Analyses were carried out for total IS, by stroke subtype and by age of stroke onset (≤ 50 yrs, 50-70 yrs, and ≥ 70 yrs).

With our total sample size (6,865 cases), we had 80% power to detect odds ratios of 1.09 ~ 1.14 for overall IS and odds ratios of 1.20-1.32 for major stroke subtypes (~1,000 cases) for variants with allele frequency 10-50% (Bonferroni-corrected $\alpha = 0.0045 = 0.05/11$ SNPs). In this report, nominal p-values were provided for all the analyses.

Results

Study population characteristics

This study included 6,865 cases and 11,395 controls recruited from 9 studies across North America, Europe, Australia and South Asia. Stroke cases ranged in age between 15 to 101 years. Among cases with subtype information available, the three major stroke subtypes, cardioembolic, large artery (atherosclerotic) and small vessel (lacunar), accounted for ~ 20.9% (n = 1,216), 19.6% (n = 1,137) and 18.0% (n = 1,043) of the cases, respectively; the remaining cases were attributed to either “other known causes” (5.6%; n = 324) or “undetermined causes” (35.9%; n = 2,084). The majority of the study participants were of European ancestry with the exception of those enrolled in the RACE study (1890 cases and 4625 controls), which has enrolled participants of South Asian origin. Detailed characteristics of the participating studies are provided in Supplemental Table S3.

Associations of MI/CAD SNPs and overall IS risk

In the total sample, the meta-analysis odds ratios for all 11 SNPs ranged from 0.92 to 1.11 (Table 2). Only two SNPs were nominally associated with IS: rs11206510 at *PCSK9* (OR of

0.92; 95% CI: 0.86-0.99; $P=0.03$; allele T) and rs3184504 at *SH2B3* (OR of 1.11; 95% CI: 1.01-1.21; $P=0.03$; allele T). This association would not have withstood a conservative Bonferroni correction ($\alpha = 0.05/11$, or 0.0045). Although there was no significant evidence for heterogeneity of effect size across studies for rs11206510, the direction of effect was not consistent across studies with ORs ranging from 0.80 (RACE study) to 1.12 (CEDIR study) (Cochran's Q statistic = 8.1, $P = 0.4$) (Figure S1). rs3184504, on the other hand, showed significance evidence for heterogeneity (Cochran's Q statistic = 20.9, $P = 0.01$) with one study (BRAINS) showing opposite effect as compared with other studies. We also examined the joint effect of these MI/CAD-associated SNPs by testing for association with the genotype scores and found no significant effect for either genotype score 1 or score 2 (OR = 1.02, $P = 0.2$ for both scores). Removing the RACE study, the only study that had participants of South Asian origin, did not change results significantly although the association with rs11206510 and rs3184504 became less significant (OR=0.94, $P=0.10$ and 1.10, $P=0.06$, respectively) (Supplementary Table S4).

We further stratified the associations between MI/CAD-associated SNPs and overall IS according to age of stroke onset: early (≤ 50 yrs), intermediate ($50 \text{ yrs} < \text{age} < 70 \text{ yrs}$) and late onset ($\text{age} \geq 70 \text{ yrs}$). None of the associations were statistically significant in the age-stratified analyses, either (Supplemental Table S5).

Associations with major ischemic stroke subtypes

We next examined the effects of MI/CAD-associated SNPs on TOAST-defined subtypes (Table 3). Nominally significant associations ($0.03 < p < 0.05$) were observed for four of the different TOAST subtypes. The MI/CAD-associated risk allele at rs9818870 in *MRAS* was

associated with the cardioembolic stroke subtype (OR= 1.17, 95% C.I.: 1.02-1.35; P=0.03); the MI-associated risk allele at rs17465637 in *MIA3* was associated with the small vessel stroke subtype (OR=1.12, 95% C.I.: 1.00-1.26; P=0.05); and the MI-associated risk allele at rs3184504 in *SH2B3* was associated with strokes due to other known causes (OR=1.33, 95% C.I.: 1.02-1.72; P=0.04). In addition, rs11206510 was associated with strokes due to other undetermined causes, although the allele conferring protection against MI was associated with stroke risk (OR = 0.86, 95% C.I.: 0.76-0.99; P =0.03). The ASGC study had a much higher OR and a wide confidence interval for rs11206510 association, but this study contributed only 11 cases of undetermined causes in the analysis. Excluding the ASGC study from the meta-analysis resulted in an OR of 0.86 (95% CI: 0.76-0.97, P=0.01) for the association between rs11206510 and undetermined causes of stroke. The *ANRIL* rs4977574 SNP was very modestly (and not significantly) associated with the large artery stroke subtype (OR=1.09, 95% C.I.: 0.99-1.20; P=0.09), nor was it associated with any other stroke subtype. None of these results withstand correction for multiple testing. Forest plots showing the most strongly associated SNP with each subtype are in Figure S2.

Discussion

It is of interest to consider whether common loci influence susceptibility to MI/CAD and stroke to the extent that such loci could elucidate common genetic and/or biologic pathways. Given that the 11SNPs tested in this study were associated with MI/CAD in previous GWA studies, they likely are among the common SNPs having the largest effect sizes. The risk alleles at these loci were associated with increases of 12-20% in MI risk, with the exception of rs4977574 in *ANRIL*, for which the risk allele was associated with a 28% increase in Europeans

^{7, 8, 13}. In contrast, none of the MI/CAD risk alleles at these loci were significantly associated with risk of overall IS, with the estimated odds ratios ranging from 0.92 to 1.11 (i.e., 8% reduction to 11% increase in odds of stroke per allele).

It is possible that failure to detect associations of these SNPs with the combined IS endpoint might be partially attributable to the heterogeneity of the IS phenotype. The large size of our sample provided us with the opportunity to evaluate the association of these SNPs to subtypes of ischemic stroke. Similar to the combined IS endpoint, we did not find any significant associations of MI/CAD SNPs (or the joint genotyping risk scores) with any of the stroke subtypes. Given that this study included more than 6,800 total cases and more than 1,000 cases for the major stroke subtypes, we have sufficient power to detect even small effects on overall stroke although more modest power to detect associations with stroke subtype. Therefore, our findings suggest that these MI/CAD-associated SNPs do not have effects of similar magnitude on overall stroke as they do on MI/CAD. If an association does exist, it is likely to be very modest for overall IS or restricted to a single stroke subtype.

Although the SNP associations with stroke subtypes did not reach statistical significance, it is interesting that the strongest associations we observed in the subtype analysis were genes involved in cell regulation or signaling. For example, the strongest association with cardioembolic stroke was the MI-risk allele at rs9818870 in *MRAS* (OR= 1.17, P=0.03). *MRAS*, an intracellular signal transducer, is highly expressed in the cardiovascular system, and may be involved in cell adhesion ^{7, 8, 16}. The strongest association with small vessel stroke was the MI-risk allele at rs17465637 in *MIA3* (OR=1.12, P=0.05), which encodes melanoma inhibitory activity 3 and is associated with cell adhesion and migration in inflammatory pathways¹⁷. It should also be noted that rs4977574 near *ANRIL* was very modestly associated with the large

artery subtype (OR=1.09, P=0.09). Although the association did not achieve statistical significance in our study, other SNPs in the *ANRIL* region on chromosome 9p21 have previously been associated with large artery stroke, with the strongest signal found with SNP rs1537378 (OR=1.21, 95% CI: 1.09-1.35, P=0.0005)¹⁴. Interestingly, rs1537378 is not in high LD with our target SNP rs4977574 ($r^2=0.5$), suggesting these 2 SNPs may represent different association signals or each of them may be tagging the causal variant imperfectly (assuming the association is indeed true). *ANRIL* encodes an antisense noncoding RNA that may disrupt other genes in the region, including *CDKN2A* and *CDKN2B*, two genes that play a key role in regulating cell proliferation, senescence, and apoptosis. We also observed the MI-associated risk allele at rs3184504 in *SH2B3* to be nominally significantly associated with strokes due to other known causes (OR=1.33, 95% C.I.: 1.02-1.72; P=0.04). It has been hypothesized that this SNP could reduce the anti-inflammatory activity of *SH2B3*, which is expressed in the vasculature, thereby leading to the development of plaque development and/or progression in coronary arteries¹³.

In contrast, SNPs whose effects on MI are likely to be mediated through lipid metabolism, e.g., rs1122608 at *SMARCA4/LDLR* (19p13), rs646776 at the *CELSR2/PSRC1/SORT1* locus (1p13) and rs11206510 at *PCSK9* (1p32), have minimal effects on IS or any of its subtypes, and in some cases, even showed opposite effects on stroke. For example, the MI/CAD-associated risk allele of rs11206510 at *PCSK9*, a gene involved in cholesterol homeostasis, was modestly associated with decreased risk of overall stroke (OR=0.92, P=0.03) or strokes due to undetermined causes (OR=0.86, P=0.03). Our analyses suggest that genes involved in lipid metabolism do not predispose individuals to increased stroke risk as they do for MI although replications will be needed to confirm these associations.

Taken together, our results suggest that the major common loci associated with MI/CAD risk do not have large effects on stroke. However, although our analyses did not reveal any significant associations between MI/CAD SNPs and stroke, the chromosomal regions harboring these loci might still contain stroke-related SNPs that are not in high LD with the MI-associated SNPs, as the example of chromosome 9p region¹⁴. A limitation of our meta-analysis is that all included studies are case-control-based and some have included recurrent stroke, thus allowing for the possibility of selection toward milder strokes (due to survival bias). It is possible that these genetic MI/CAD variants could be associated with more severe forms of stroke, but longitudinal studies that follow healthy cohorts for stroke occurrence prospectively would be needed to address this issue. Finally, our analyses were based on 11 SNPs unequivocally associated with MI and/or CAD identified from the first wave of meta-analyses for these traits. A second wave of even larger meta-analyses of these traits is in progress that will likely generate additional associations. We can expect most of these newly associated SNPs to have even smaller effect sizes on MI and CAD. In fact, in September 2011, 17 additional CAD-associated SNPs were identified by even larger meta-analyses, and the reported effect sizes for these newly identified SNPs, as expected, were smaller (ranging from 1.06-1.17) than the original 11 SNPs used in the present analysis¹⁸⁻²¹. But this does not necessarily mean that these SNPs will have smaller effect sizes on stroke. Indeed, it is very possible that such SNPs may be in gene pathways that are more relevant to stroke than are the pathways associated with the currently known MI SNPs. Thus, it will be important to monitor new SNPs identified, even those with small effect sizes, as they may generate new biological insights into the etiology of stroke.

Summary

Common variants previously associated with MI/CAD risk do not have effects of similar magnitude on overall IS, but do not preclude moderate associations restricted to specific IS subtypes.

Acknowledgements

Australian population control data was derived from the Hunter Community Study. ASGC study would like to thank the men and women of the Hunter region who participated in this study. Genotyping for GEOS and VISP studies were provided by the Johns Hopkins University Center for Inherited Disease Research (CIDR) and data cleaning were assisted by GENEVA coordinating center under National Institutes of Health Genes, Environment and Health Initiative (GEI). RACE study would like to acknowledge the contributions made by the RACE investigators.

Sources of Funding

ASGC: We are grateful to the University of Newcastle for funding. The ASGC was funded by grants from the Australian National and Medical Health Research Council (NHMRC Project Grant ID: 569257) and the Australian National Heart Foundation (NHF Project Grant ID: G 04S 1623). EGH is supported by an NHMRC Training (Postdoctoral) Fellowship.

BRAINS: BRAINS is supported by a Dept of Health (UK) senior fellowship to PS and also by a grant from the Henry Smith Charity and the British Council (UKIERI).

CEDIR: The Besta Cerebrovascular Diseases Registry (CEDIR) was supported by the Italian Ministry of Health, years 2007–2010 (Annual Research Funding; Grant Numbers: RC 2007/LR6, RC 2008/LR6; RC 2009/LR8; RC 2010/LR8). This study was supported also by Cariplo Foundation (Grant n. 2010-0253). Silvia Parolo is supported by a PhD fellowship of the University of Pavia (PhD program in Genetic and Biomolecular Sciences). The PROCARDIS study was supported by the European Community Sixth Framework Program (LSHM-CT-2007-037273).

GASROS: GASROS was supported by the American Heart Association/Bugher Foundation Centers for Stroke Prevention Research (0775010N); NIH-NINDS grant numbers K23NS064052 (N.S.R.), R01NS063925 (J.R.) and P50NS051343 (J.R. and K.L.F.); the Deane Institute for Integrative Study of Atrial Fibrillation and Stroke; and the Keane Stroke Genetics Research Fund. The MIGen study was funded by the U.S. National Institutes of Health and National Heart, Lung, and Blood Institute's STAMPEED genomics research program (R01 HL087676) and a grant from the National Center for Research Resources. Drs. Biffi, Rost and Anderson were supported in part by the American Heart Association/Bugher Foundation Centers for Stroke Prevention Research.

GEOS: The GEOS study was supported by the National Institutes of Health Genes, Environment and Health Initiative (GEI) Grant U01 HG004436, as part of the GENEVA consortium under GEI. Genotyping services were provided by the Johns Hopkins University Center for Inherited Disease Research (CIDR), which is fully funded through a federal contract from the National Institutes of Health to the Johns Hopkins University (contract number HHSN268200782096C). Assistance with data cleaning was provided by the GENEVA Coordinating Center (U01 HG 004446; PI Bruce S Weir). Study recruitment and collection of datasets were supported by the Office of Research and Development, Medical Research Service, and the Baltimore Geriatrics Research, Education, and Clinical Center of the Department of Veterans Affairs and by grants from the National Institute of Neurological Disorders and Stroke (NINDS) and the NIH Office of Research on Women's Health (R01 NS45012, U01 NS069208-01).

GRAZ: Genetic studies of the Austrian Stroke Prevention Study are supported by the Austrian Science Fund (P20545).

ISGS/SWISS: This work was supported in part by the Intramural Research Program of the National Institute on Aging, National Institutes of Health, Department of Health and Human Services; project number Z01 AG000954-06. This work used samples and clinical data from the NINDS Human Genetics Resource Center DNA and Cell Line Repository (<http://ccr.coriell.org/ninds>), human subjects protocol numbers 2003-081 and 2004-147. The inclusion of Baltimore Longitudinal Study of Aging samples was supported in part by the Intramural Research Program of the National Institute on Aging, National Institutes of Health, Department of Health and Human Services; project number Z01 AG000015-50, human subjects protocol number 2003-078. The ISGS study was funded by a grant from the National Institute for Neurological Disorders and Stroke (US), Grant no. R01 NS42733 (J. Meschia, PI). The SWISS study was funded by a grant from the National Institute for Neurological Disorders and Stroke (US), Grant no. R01 NS39987 (J. Meschia, PI). This study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the National Institutes of Health, Bethesda, Md. (<http://biowulf.nih.gov>).

RACE: The RACE study is supported by grants from the National Institute of Neurological Disorders And Stroke (Award Number 21NS064908) and educational awards to investigators at the Center for Non-Communicable Diseases, Pakistan.

VISP: The GWAS component of the VISP study was supported by the National Human Genome Research Institute (NHGRI), Grant U01 HG005160 (PI Michele Sale & Bradford Worrall), as part of the Genomics and Randomized Trials Network (GARNET). Genotyping services were provided by the Johns Hopkins University Center for Inherited Disease Research (CIDR), which is fully funded through a federal contract from the National Institutes of Health to the Johns Hopkins University. Assistance with data cleaning was provided by the GARNET

Coordinating Center (U01 HG005157; PI Bruce S Weir). Study recruitment and collection of datasets for the VISP clinical trial were supported by an investigator-initiated research grant (R01 NS34447; PI James Toole) from the United States Public Health Service, National Institute of Neurological Disorders and Stroke of the National Institutes of Health, Bethesda, Maryland.

Disclosures

None

References

1. Rosamond W, Flegal K, Furie K, Go A, Greenlund K, Haase N, et al. Heart disease and stroke statistics--2008 update: A report from the american heart association statistics committee and stroke statistics subcommittee. *Circulation*. 2008;117:e25-146
2. Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, et al. Heart disease and stroke statistics--2010 update: A report from the american heart association. *Circulation*. 2010;121:e46-215
3. Adams HP, Jr., Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. Toast. Trial of org 10172 in acute stroke treatment. *Stroke*. 1993;24:35-41
4. Jerrard-Dunne P, Cloud G, Hassan A, Markus HS. Evaluating the genetic component of ischemic stroke subtypes: A family history study. *Stroke*. 2003;34:1364-1369
5. Polychronopoulos P, Gioldasis G, Ellul J, Metallinos IC, Lekka NP, Paschalis C, et al. Family history of stroke in stroke types and subtypes. *J Neurol Sci*. 2002;195:117-122
6. MacClellan LR, Mitchell BD, Cole JW, Wozniak MA, Stern BJ, Giles WH, et al. Familial aggregation of ischemic stroke in young women: The stroke prevention in young women study. *Genetic Epidemiology*. 2006;30:602-608
7. Erdmann J, Grosshennig A, Braund PS, Konig IR, Hengstenberg C, Hall AS, et al. New susceptibility locus for coronary artery disease on chromosome 3q22.3. *Nat Genet*. 2009;41:280-282

8. Kathiresan S, Voight BF, Purcell S, Musunuru K, Ardissino D, Mannucci PM, et al. Genome-wide association of early-onset myocardial infarction with single nucleotide polymorphisms and copy number variants. *Nat Genet.* 2009;41:334-341
9. McPherson R, Pertsemlidis A, Kavaslar N, Stewart A, Roberts R, Cox DR, et al. A common allele on chromosome 9 associated with coronary heart disease. *Science.* 2007;316:1488-1491
10. Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature.* 2007;447:661-678
11. Helgadottir A, Thorleifsson G, Manolescu A, Gretarsdottir S, Blondal T, Jonasdottir A, et al. A common variant on chromosome 9p21 affects the risk of myocardial infarction. *Science.* 2007;316:1491-1493
12. Samani NJ, Erdmann J, Hall AS, Hengstenberg C, Mangino M, Mayer B, et al. Genomewide association analysis of coronary artery disease. *N Engl J Med.* 2007;357:443-453
13. Gudbjartsson DF, Bjornsdottir US, Halapi E, Helgadottir A, Sulem P, Jonsdottir GM, et al. Sequence variants affecting eosinophil numbers associate with asthma and myocardial infarction. *Nat Genet.* 2009;41:342-347
14. Gschwendtner A, Bevan S, Cole JW, Plourde A, Matarin M, Ross-Adams H, et al. Sequence variants on chromosome 9p21.3 confer risk for atherosclerotic stroke. *Ann.Neurol.* 2009;65:531-539
15. Fleiss JL. The statistical basis of meta-analysis. *Stat Methods Med Res.* 1993;2:121-145

16. Yoshikawa Y, Satoh T, Tamura T, Wei P, Bilasy SE, Edamatsu H, et al. The m-ras-ra-gef-2-rap1 pathway mediates tumor necrosis factor- α dependent regulation of integrin activation in splenocytes. *Mol. Biol. Cell.* 2007;18:2949-2959
17. Arndt S, Melle C, Mondal K, Klein G, von Eggeling F, Bosserhoff A-K. Interactions of tango and leukocyte integrin cd11c/cd18 regulate the migration of human monocytes. *Journal of Leukocyte Biology.* 2007;82:1466-1472
18. Wild PS, Zeller T, Schillert A, Szymczak S, Sinning CR, Deiseroth A, et al. A genome-wide association study identifies lipa as a susceptibility gene for coronary artery disease. *Circulation: Cardiovascular Genetics.* 2011;4:403-412
19. Erdmann J, Willenborg C, Nahrstaedt J, Preuss M, König IR, Baumert J, et al. Genome-wide association study identifies a new locus for coronary artery disease on chromosome 10p11.23. *European Heart Journal.* 2011;32:158-168
20. The Coronary Artery Disease (C4D) Genetics Consortium. A genome-wide association study in europeans and south asians identifies five new loci for coronary artery disease. *Nat Genet.* 2011;43:339-344
21. Schunkert H, König IR, Kathiresan S, Reilly MP, Assimes TL, Holm H, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet.* 2011;43:333-338

Table 1. Previous published SNPs that were associated with MI or CAD

| SNP | Non-risk/ risk allele* | Frequency of risk allele | Region | Position | Reported gene | Previous reported odds ratio on MI | Reference |
|------------|---------------------------|-----------------------------|--------|-------------|------------------------|--|-----------|
| rs11206510 | C/T | 0.81 | 1p32 | 55,268,627 | PCSK9 | 1.15 | 8 |
| rs646776 | C/T | 0.81 | 1p13 | 109,620,053 | CELSR2-PSRC1-SORT1 | 1.17 | 8, 10 |
| rs17465637 | A/C | 0.72 | 1q41 | 220,890,152 | MIA3 | 1.13 | 8, 12 |
| rs6725887 | T/C | 0.14 | 2q33 | 203,454,130 | WDR12 | 1.17 | 8 |
| rs9818870 | C/T | 0.16 | 3q22.2 | 139,604,812 | MRAS | 1.15 | 7 |
| rs12526453 | G/C | 0.65 | 6p24 | 13,035,530 | PHACTR1 | 1.12 | 8 |
| rs4977574 | A/G | 0.56 | 9p21 | 22,088,574 | ANRIL/CDKN2B-AS1 | 1.28 | 8, 9, 11 |
| rs1746048 | T/C | 0.84 | 10q11 | 44,095,830 | CXCL12 | 1.14 | 8, 12 |
| rs3184504 | C/T | 0.39 | 12q24 | 110,368,991 | SH2B3 | 1.13 | 13 |
| rs1122608 | T/G | 0.75 | 19p13 | 11,024,601 | SMARCA4-LDLR | 1.15 | 8 |
| rs9982601 | C/T | 0.13 | 21q22 | 34,520,998 | SLC5A3-MRPS6- KCNE2 | 1.20 | 8 |

* Risk allele is the allele associated with increased risk of MI or CAD

Table 2. Association results of the 11 MI/CAD-risk loci and risk scores with ischemic stroke

| SNP | MI/CAD-risk allele | Heterogeneity between studies | | Association with overall IS | | Minimal OR detectable in current study [†] |
|------------|--------------------|-------------------------------|--------------------|-----------------------------|------|---|
| | | Q (P-value) | I ² , % | OR (95% CI) [*] | P | |
| rs11206510 | T | 8.12 (0.42) | 1.53 | 0.92 (0.86, 0.99) | 0.03 | 1.13 |
| rs646776 | T | 9.02 (0.34) | 11.34 | 0.95 (0.89, 1.01) | 0.12 | 1.13 |
| rs17465637 | C | 3.09 (0.93) | 0 | 1.03 (0.97, 1.09) | 0.37 | 1.11 |
| rs6725887 | C | 13.38 (0.06) | 47.67 | 1.03 (0.90, 1.16) | 0.69 | 1.13 |
| rs9818870 | T | 5.45 (0.61) | 0 | 1.03 (0.95, 1.11) | 0.47 | 1.12 |
| rs12526453 | C | 8.81 (0.36) | 9.2 | 1.00 (0.95, 1.06) | 0.92 | 1.10 |
| rs4977574 | G | 4.17 (0.84) | 0 | 1.02 (0.97, 1.07) | 0.54 | 1.09 |
| rs1746048 | C | 5.37 (0.72) | 0 | 1.03 (0.96, 1.10) | 0.38 | 1.14 |
| rs3184504 | T | 20.87 (0.01) | 61.67 | 1.11 (1.01, 1.21) | 0.03 | 1.09 |
| rs1122608 | G | 6.31 (0.61) | 0 | 1.00 (0.95, 1.06) | 0.92 | 1.11 |
| rs9982601 | T | 6.20 (0.52) | 0 | 1.00 (0.93, 1.09) | 0.95 | 1.13 |
| score1 | n/a | 12.31 (0.09) | 43.15 | 1.02 (0.99, 1.05) | 0.23 | n/a |

| | | | | | | |
|--------|-----|--------------|-------|-------------------|------|-----|
| score2 | n/a | 14.44 (0.04) | 51.52 | 1.02 (0.99, 1.06) | 0.23 | n/a |
|--------|-----|--------------|-------|-------------------|------|-----|

OR: odds ratio; CI: confidence interval; n/a: not applicable; *P*: p-value

* OR: odds ratios reflects the effect of the MI-associated risk allele with the previously published MI non-risk allele as the reference allele defined in Table 1.

† The minimal odds ratios that current sample size (6,865 overall IS cases) can detect with at least 80% power ($\alpha=0.0045$).

Table 3. Association results of the 11 MI/CAD-risk loci and risk scores with ischemic stroke, by TOAST subtypes

| SNP | Cardioembolic (n=1,216) | | Large Artery (n=1,137) | | Small Vessel (n=1,043) | | Other Known Causes (n=324) | | Undetermined causes (n=2,087) | |
|------------|----------------------------|----------|---------------------------|----------|---------------------------|----------|----------------------------------|----------|----------------------------------|----------|
| | OR | <i>P</i> | OR | <i>P</i> | OR | <i>P</i> | OR | <i>P</i> | OR | <i>P</i> |
| | (95% CI)* | | (95% CI)* | | (95% CI)* | | (95% CI)* | | (95% CI)* | |
| rs11206510 | 1.02 | | 0.91 | | 1.05 | | 0.89 | | 0.86 | |
| | (0.89, 1.18) | 0.78 | (0.80, 1.04) | 0.17 | (0.91, 1.21) | 0.50 | (0.72, 1.11) | 0.32 | (0.76, 0.99) | 0.03 |
| rs646776 | 0.93 | | 0.91 | | 0.95 | | 1.01 | | 0.93 | |
| | (0.83, 1.05) | 0.25 | (0.81, 1.03) | 0.12 | (0.84, 1.07) | 0.38 | (0.78, 1.31) | 0.94 | (0.82, 1.05) | 0.24 |
| rs17465637 | 1.01 | | 0.98 | | 1.12 | | 0.96 | | 1.01 | |
| | (0.89, 1.16) | 0.86 | (0.87, 1.09) | 0.69 | (1.00, 1.26) | 0.05 | (0.79, 1.17) | 0.69 | (0.92, 1.10) | 0.90 |
| rs6725887 | 0.98 | | 1.01 | | 0.91 | | 1.45 | | 0.96 | |
| | (0.83, 1.16) | 0.84 | (0.85, 1.19) | 0.93 | (0.76, 1.09) | 0.30 | (0.93, 2.26) | 0.10 | (0.83, 1.11) | 0.58 |
| rs9818870 | 1.17 | | 0.97 | | 0.91 | | 1.32 | | 0.99 | |
| | (1.02, 1.35) | 0.03 | (0.83, 1.13) | 0.66 | (0.77, 1.07) | 0.24 | (0.92, 1.89) | 0.13 | (0.87, 1.12) | 0.90 |

| | | | | | | | | | | |
|------------|--------------|------|--------------|------|--------------|------|--------------|------|--------------|------|
| rs12526453 | 1.06 | | 0.93 | | 0.90 | | 1.03 | | 1.00 | |
| | (0.95, 1.18) | 0.31 | (0.83, 1.03) | 0.15 | (0.80, 1.02) | 0.09 | (0.85, 1.25) | 0.74 | (0.91, 1.09) | 0.99 |
| rs4977574 | 1.02 | | 1.09 | | 1.03 | | 0.91 | | 1.01 | |
| | (0.89, 1.17) | 0.82 | (0.99, 1.20) | 0.09 | (0.94, 1.14) | 0.52 | (0.69, 1.18) | 0.47 | (0.94, 1.09) | 0.72 |
| rs1746048 | 0.99 | | 1.05 | | 1.01 | | 0.97 | | 1.00 | |
| | (0.88, 1.12) | 0.93 | (0.92, 1.2) | 0.44 | (0.86, 1.18) | 0.94 | (0.70, 1.33) | 0.85 | (0.91, 1.1) | 0.96 |
| rs3184504 | 1.05 | | 1.16 | | 1.03 | | 1.33 | | 1.12 | |
| | (0.91, 1.22) | 0.51 | (0.96, 1.39) | 0.11 | (0.89, 1.20) | 0.68 | (1.02, 1.72) | 0.04 | (0.98, 1.27) | 0.11 |
| rs1122608 | 0.98 | | 1.06 | | 1.03 | | 1.03 | | 0.99 | |
| | (0.88, 1.09) | 0.67 | (0.95, 1.18) | 0.33 | (0.92, 1.15) | 0.66 | (0.85, 1.26) | 0.75 | (0.91, 1.07) | 0.74 |
| rs9982601 | 1.05 | | 1.04 | | 1.06 | | 1.15 | | 1.02 | |
| | (0.89, 1.25) | 0.55 | (0.89, 1.22) | 0.63 | (0.88, 1.27) | 0.55 | (0.83, 1.60) | 0.39 | (0.85, 1.22) | 0.85 |
| score1 | 1.02 | | 1.02 | | 1.01 | | 1.06 | | 1.00 | |
| | (0.98, 1.07) | 0.29 | (0.97, 1.06) | 0.47 | (0.95, 1.07) | 0.85 | (0.98, 1.14) | 0.16 | (0.96, 1.04) | 0.97 |
| score2 | 1.04 | | 1.01 | | 1.00 | | 1.08 | | 1.00 | |
| | (0.99, 1.08) | 0.11 | (0.96, 1.08) | 0.63 | (0.94, 1.07) | 0.91 | (1.01, 1.17) | 0.03 | (0.96, 1.05) | 0.92 |

n: number of cases; OR: odds ratio; P: p-value

* OR: odds ratio reflects the effect of the MI-associated risk allele with the previously published MI non-risk allele as the reference allele defined in Table 1.