

## REFERENCES

- Adachi, J. and Hasegawa, M.** (1996). MOLPHY, Programs for Molecular Phylogenetics Based on Maximum Likelihood, (ed. Tokyo: Institute of Statistical Mathematics.
- Aerne, B. L., Baader, C. D. and Schmid, V.** (1995). Life stage and tissue-specific expression of the homeobox gene cnox1-Pc of the hydrozoan *Podocoryne carnea*. *Dev Biol* **169**, 547-56.
- Alberga, A., Boulay, J. L., Kempe, E., Dennefeld, C. and Haenlin, M.** (1991). The snail gene required for mesoderm formation in *Drosophila* is expressed dynamically in derivatives of all three germ layers. *Development* **111**, 983-92.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J.** (1990). Basic local alignment search tool. *J Mol Biol* **215**, 403-10.
- Amores, A., Force, A., Yan, Y. L., Joly, L., Amemiya, C., Fritz, A., Ho, R. K., Langeland, J., Prince, V., Wang, Y. L. et al.** (1998). Zebrafish hox clusters and vertebrate genome evolution. *Science* **282**, 1711-4.
- Arendt, D. and Nubler-Jung, K.** (1999). Comparison of early nerve cord development in insects and vertebrates. *Development* **126**, 2309-25.
- Ausubel, F. M., Brent, R., Kingston, R. E., Moore, D. D., Seidman, J. G., Smith, J. A. and Struhl, K.** (1996). Current protocols in molecular biology. New York, U.S.A: John Wiley & Sons, Inc.
- Ball, E. E., Hayward, D. C., Catmull, J., Reece-Hoyes, J. S., Hislop, N. R., Harrison, P. L. and Miller, D. J.** (2002). Molecular control of development in the reef coral, *Acropora millepora*. *Proc. 9th International Coral Reef Symposium, Bali*.
- Bally-Cuif, L. and Boncinelli, E.** (1997). Transcription factors and head formation in vertebrates. *BioEssays* **19**, 127-35.
- Barnes, L. M. E.** (2002) Characterising a putative methyl-CpG binding protein found in the staghorn coral *Acropora millepora* Biochemistry and Molecular Biology, James Cook University, Townsville
- Barnes, R.** (1987). Invertebrate Zoology. London: Saunders College Publishing.
- Beachy, P. A., Krasnow, M. A., Gavis, E. R. and Hogness, D. S.** (1988). An Ultrabithorax protein binds sequences near its own and the Antennapedia P1 promoters. *Cell* **55**, 1069-81.
- Beddington, R. S. and Robertson, E. J.** (1998). Anterior patterning in mouse. *Trends Genet* **14**, 277-84.
- Bell, E., Ensini, M., Gulisano, M. and Lumsden, A.** (2001). Dynamic domains of gene expression in the early avian forebrain. *Dev Biol* **236**, 76-88.

- Benton, W. and Davis, R.** (1977). Screening lambda-gt recombinant clones by hybridisation to single plaques in situ. *Science* **196**, 180-182.
- Berthelsen, J., Kilstrup-Nielsen, C., Blasi, F., Mavilio, F. and Zappavigna, V.** (1999). The subcellular localization of PBX1 and EXD proteins depends on nuclear import and export signals and is modulated by association with PREP1 and HTH. *Genes & Development* **13**, 946-953.
- Blitz, I. L. and Cho, K. W.** (1995). Anterior neurectoderm is progressively induced during gastrulation: the role of the Xenopus homeobox gene orthodenticle. *Development* **121**, 993-1004.
- Boncinelli, E., Gulisano, M. and Broccoli, V.** (1993). *Emx* and *Otx* homeobox genes in the developing mouse brain. *Journal of Neurobiology* **24**, 1356-1366.
- Bouwmeester, T. and Leyns, L.** (1997). Vertebrate head induction by anterior primitive endoderm. *BioEssays* **19**, 855-63.
- Bradford, M.** (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Analytical Biochemistry* **72**, 248-254.
- Brauer, A.** (1891). Ueber die Entwicklung von *Hydra*. *Z. wiss. Zool* **52**, 551-579.
- Bridge, D. M., Cunningham, C. W., DeSalle, R. and Buss, L. W.** (1995). Class-level relationships in the phylum Cnidaria: molecular and morphological evidence. *Molecular Biology and Evolution* **12**, 679-689.
- Bridge, D. M., Cunningham, C. W., Schierwater, B., DeSalle, R. and Buss, L. W.** (1992). Class level relationships in the phylum Cnidaria: evidence from mitochondrial genome structure. *Proceedings of The National Academy of Sciences* **89**, 8750-8753.
- Bridge, D. M., Stover, N. A. and Steele, R. E.** (2000). Expression of a novel receptor tyrosine kinase gene and a *paired*-like homeobox gene provides evidence of difference in patterning at the oral and aboral ends of hydra. *Development* **220**, 253-262.
- Briscoe, J., Sussel, L., Serup, P., Hartigan-O'Connor, D., Jessell, T. M., Rubenstein, J. L. and Ericson, J.** (1999). Homeobox gene Nkx2.2 and specification of neuronal identity by graded Sonic hedgehog signalling. *Nature* **398**, 622-7.
- Broun, M., Sokol, S. and Bode, H. R.** (1999). *Cngsc*, a homologue of *goosecoid*, participates in the patterning of the head, and is expressed in the organizer region of *Hydra*. *Development* **126**, 5245-5254.
- Brusca, R. and Brusca, G.** (1990). Invertebrates. Sunderland, Massachusetts: Sinauer Associates, Inc.
- Burglin, T. R.** (1997). Analysis of TALE superclass homeobox genes (MEIS, PBS, KNOW, Iroquois, RGIF) reveals a novel domain conserved between plants and animals. *Nucleic Acids Res* **25**, 4173-4180.

- Bürglin, T. R.** (1994). A comprehensive classification of homeobox gene. In *Guidebook to the homeobox genes*, (ed. Duboule, D.), pp. 27-71. Oxford: Oxford University Press.
- Burke, D., Dawson, D. and Stearns, T.** (2000). Methods in yeast genetics: A cold spring harbour laboratory course manual. New York: Cold Spring Harbor Laboratory Press.
- Campbell, G., Weaver, T. and Tomlinson, A.** (1993). Axis specification in the developing Drosophila appendage: the role of wingless, decapentaplegic, and the homeobox gene aristaless. *Cell* **74**, 1113-23.
- Cartwright, P., Bowsher, J. and Buss, L. W.** (1999). Expression of a Hox gene, Cnox-2, and the division of labor in a colonial hydroid. *Proc Natl Acad Sci U S A* **96**, 2183-6.
- Carvalho, A. B. and Clark, A. G.** (1999). Intron size and natural selection. *Nature* **401**, 344.
- Carver, E. A., Jiang, R., Lan, Y., Oram, K. F. and Gridley, T.** (2001). The mouse snail gene encodes a key regulator of the epithelial-mesenchymal transition. *Mol Cell Biol* **21**, 8184-8.
- Castillo-Davis, C. I., Mekhedov, S. L., Hartl, D. L., Koonin, E. V. and Kondrashov, F. A.** (2002). Selection for short introns in highly expressed genes. *Nat Genet* **31**, 415-8.
- Catmull, J., Hayward, D. C., McIntyre, N. E., Reece-Hoyes, J. S., Mastro, R., Callaerts, P., Ball, E. E. and Miller, D. J.** (1998). Pax-6 origins - implications from the structure of two coral Pax genes. *Development Genes and Evolution* **208**, 352-356.
- Chan, S.-K., Jaffe, L., Capovilla, M., Botas, J. and Mann, R. S.** (1994). The DNA binding specificity of Ultrabithorax is modulated by cooperative interactions with extradenticle, another homeoprotein. *Cell* **78**, 603-615.
- Chan, Y. M. and Jan, Y. N.** (1999). Conservation of neurogenic genes and mechanisms. *Curr Opin Neurobiol* **9**, 582-8.
- Chang, C. P., Shen, W. F., Rozenfeld, S., Lawrence, H. J., Largman, C. and Cleary, M. L.** (1995). Pbx proteins display hexapeptide-dependent cooperative DNA binding with a subset of Hox proteins. *Genes Dev* **9**, 663-74.
- Chia, F.-S. and Koss, R.** (1979). Fine structural studies of the nervous system and the apical organ in the planula larva of the sea anemone *Anthopleura elegantissima*. *Journal of Morphology* **160**.
- Chu, H., Parras, C., White, K. and Jimenez, F.** (1998). Formation and specification of ventral neuroblasts is controlled by vnd in Drosophila neurogenesis. *Genes Dev* **12**, 3613-24.
- Cohen, S. M. and Jurgens, G.** (1990). Mediation of *Drosophila* head development by gap-like segmentation genes. *Nature* **346**, 482-485.

- Cohen, S. M. and Jurgens, G.** (1991). Drosophila headlines. *Trends in Genetics* **7**, 267-271.
- Conway Morris, S.** (1998). Eggs and embryos from the Cambrian. *BioEssays* **20**, 676-82.
- Conway Morris, S.** (2000a). The Cambrian "explosion": slow-fuse or megatonnage? *Proceedings of The National Academy of Sciences* **97**, 4426-4429.
- Conway Morris, S.** (2000b). Nipping the Cambrian "explosion" in the bud? *BioEssays* **22**, 1053-6.
- Cornell, R. A. and Von Ohlen, T.** (2000). *Vnd/nkx*, *ind/gsh*, and *msh/msx*: conserved regulators of dorsoventral neural patterning? *Current Opinion in Neurobiology* **10**, 63-71.
- Cowden, J. and Levine, M.** (2002). The Snail repressor positions Notch signaling in the Drosophila embryo. *Development* **129**, 1785-93.
- Dalton, D., Chadwick, R. and McGinnis, W.** (1989). Expression and embryonic function of *empty spiracles*: a *Drosophila* homeo box gene with two patterning functions on the anterior-posterior axis of the embryo. *Genes & Development* **3**, 1940-1956.
- Daniel, A., Dumstrei, K., Lengyel, J. and Hartenstein, V.** (1999). The control of cell fate in the embryonic visual system by *atonal*, *tailless* and EGFR signaling. *Development* **126**, 2945-2954.
- Darmer, D., Hauser, F., Nothacker, H.-P., Bosch, T. C. G., Williamson, M. and Grimmelikhuijen, C. J. P.** (1998). Three different prohormones yield a variety of Hydra RFamide (Arg-Phe-NH<sub>2</sub>) neuropeptides in *Hydra magnipapillata*. *Journal of Biochemistry* **332**, 403-412.
- Derobert, Y., Germot, A., Spengler, T. and Mazan, S.** (2000). [Structural and functional diversity of homeodomain genes of the orthodenticle and empty spiracles classes in Craniata]. *J Soc Biol* **194**, 81-6.
- Derobert, Y., Plouhinec, J. L., Sauka-Spengler, T., Le Mentec, C., Baratte, B., Jaillard, D. and Mazan, S.** (2002). Structure and Expression of Three Emx Genes in the Dogfish *Scyliorhinus canicula*: Functional and Evolutionary Implications. *Dev Biol* **247**, 390-404.
- Ekker, S. C., Young, K. E., von Kessler, D. P. and Beachy, P. A.** (1991). Optimal DNA sequence recognition by the Ultrabithorax homeodomain of Drosophila. *Embo J* **10**, 1179-86.
- Essex, L. J., Mayor, R. and Sargent, M. G.** (1993). Expression of Xenopus snail in mesoderm and prospective neural fold ectoderm. *Dev Dyn* **198**, 108-22.
- Fautin, D. G. and Mariscal, R. N.** (1991). Cnidaria: Anthozoa. In *Microscopic Anatomy of Invertebrates*, vol. 2 (ed. Harrison, F. W. and J. A. Westfall), pp. 267-358. New York: Wiley-Liss, Inc.

- Feinberg, A. P. and Vogelstein, B.** (1984). A technique for radiolabelling DNA restriction endonuclease fragments to high specific activity. *Analytical Biochemistry* **137**, 266-267.
- Ferrier, D. E. K. and Holland, P. W. H.** (2001). Ancient origin of the HOX gene cluster. *Nature Review Genetics* **2**, 33-38.
- Finkelstein, R. and Boncinelli, E.** (1994). From fly head to mammalian forebrain: the story of *otd* and *Otx*. *Trends in Genetics* **10**, 310-315.
- Finkelstein, R. and Perrimon, N.** (1990). The *orthodenticle* gene is regulated by *bicoid* and *torso* and specifies *Drosophila* head development. *Nature* **346**, 485-488.
- Finley, K. D., Edeen, P. T., Foss, M., Gross, E., Ghbeish, N., Palmer, R. H., Taylor, B. J. and McKeown, M.** (1998). Dissatisfaction encodes a tailless-like nuclear receptor expressed in a subset of CNS neurons controlling *Drosophila* sexual behaviour. *Neuron* **21**, 1363-1374.
- Finnerty, J. R. and Martindale, M. Q.** (1997). Homeoboxes in sea anemones (Cnidaria:Anthozoa): a PCR-based survey of *Nematostella vectensis* and *Metridium senile*. *Biol Bull* **193**, 62-76.
- Finnerty, J. R. and Martindale, M. Q.** (1998). The evolution of the Hox cluster: insights from outgroups. *Curr Opin Genet Dev* **8**, 681-7.
- Finnerty, J. R. and Martindale, M. Q.** (1999). Ancient origins of axial patterning genes: Hox genes and ParaHox genes in the Cnidaria. *Evolution & Development* **1**, 16-23.
- Fisher, D. A. and Bode, H. R.** (1989). Nucleotide sequence of an actin-encoding gene from *Hydra attenuata*: structural characteristics and evolutionary implications. *Gene* **84**, 55-64.
- Force, A., Lynch, M., Pickett, F. B., Amores, A., Yan, Y. L. and Postlethwait, J.** (1999). Preservation of duplicate genes by complementary, degenerative mutations. *Genetics* **151**, 1531-45.
- Frasch, M., Hoey, T., Rushlow, C., Doyle, H. and Levine, M.** (1987). Characterization and localization of the even-skipped protein in *Drosophila*. *Embo J* **6**, 749-759.
- Freeman, G.** (1981a). The cleavage initiation site establishes the posterior pole of the hydrozoan embryo. *Roux's Arch. Dev. Biol.* **190**, 123-125.
- Freeman, G.** (1981b). The role of polarity in the development of the hydrozoan planula larva. *Roux's Arch. Dev. Biol.* **190**, 168-184.
- Freeman, G.** (1990). The establishment and role of polarity during embryogenesis in Hydrozoans. In *The Ceullular and Molecular Biology of Pattern Formation*, (ed. Stocum, D. and T. Karr), pp. 3-30. Oxford: Oxford University Press.

- Furukawa, T., Kozak, C. A. and Cepko, C. L.** (1997). *rax*, a novel paired-type homeobox gene, shows expression in the anterior neural fold and developing retina. *Proceedings of The National Academy of Sciences* **94**, 3088-3093.
- Fuse, N., Hirose, S. and Hayashi, S.** (1996). Determination of wing cell fate by the escargot and snail genes in Drosophila. *Development* **122**, 1059-67.
- Galliot, B.** (2000). Conserved and divergent genes in apex and axis development of cnidarians. *Current Opinions in Genetics & Development* **10**, 629-637.
- Galliot, B., de Vargas, C. and Miller, D.** (1999). Evolution of homeobox genes: Q<sub>50</sub> paired-like genes founded the paired class. *Development Genes and Evolution* **209**, 186-197.
- Galliot, B. and Miller, D.** (2000). Origin of anterior patterning. How old is our head? *Trends Genet* **16**, 1-5.
- Garcia-Fernandez, J. and Holland, P. W.** (1994). Archetypal organization of the amphioxus Hox gene cluster. *Nature* **370**, 563-6.
- Gauchat, D., Kreger, S., Holstein, T. and Galliot, B.** (1998). *prdl-a*, a gene marker for hydra apical differentiation related to triploblastic paired-like head-specific genes. *Development* **125**, 1637-1645.
- Gauchat, D., Mazet, F., Berney, C., Schummer, M., Kreger, S., Pawlowski, J. and Galliot, B.** (2000). Evolution of Antp-class genes and differential expression of *Hydra* Hox/paraHox genes in anterior patterning. *Proceedings of The National Academy of Sciences* **97**, 4493-4498.
- Geoffroy St Hilaire, E.** (1822). Considérations générales sur la vertébre. *Mém Mus. Hist. Nat.* **9**, 89-119.
- Gietz, R. and Schiestl, R.** (1995). Transforming yeast with DNA. *Methods Mol. Cell. Biol.* **5**.
- Glass, C. K.** (1994). Differential recognition of target genes by nuclear receptor monomers, dimers, and heterodimers. *Endocr Rev* **15**, 391-407.
- Goldstein and Freeman.** (1997). Axis specification in animal development. *BioEssays* **19**, 105-16.
- Grasso, L. C.** (1999) Cloning and characterization of novel nuclear receptors from the coral, *Acropora millepora* Research School of Biological Sciences, Australian National University, Canberra, ACT
- Grasso, L. C., Hayward, D. C., Trueman, J. W., Hardie, K. M., Janssens, P. A. and Ball, E. E.** (2001). The evolution of nuclear receptors: evidence from the coral Acropora. *Mol Phyl Evol* **21**, 93-102.
- Grens, A., Gee, L., Fisher, D. A. and Bode, H. R.** (1996). CnNK-2, an NK-2 homeobox gene, has a role in patterning the basal end of the axis in hydra. *Dev Biol* **180**, 473-88.

- Grimmelikhuijzen, C. J. P., Leviev, I. and Carstensen, K.** (1996). Peptides in the nervous system of cnidarians: structure, function and biosynthesis. *International Review of Cytology* **167**, 37-88.
- Grimmelikhuijzen, C. J. P. and Westfall, J. A.** (1995). The nervous system of Cnidarians. In *The nervous system of invertebrates: an evolutionary and comparative approach*, (ed. Breidbach, O. and W. Kutsch), pp. 7-24. Basel, Switzerland: Birkhauser Verlag.
- Groger, H., Callaerts, P., Gehring, W. J. and Schmid, V.** (1999). Gene duplication and recruitment of a specific tropomyosin into striated muscle cells in the jellyfish *Podocornyne carnea*. *J Exp Zool* **285**, 378-386.
- Gröger, H. and Schmid, V.** (2001). Larval development in Cnidaria: a connection to Bilateria? *Genesis* **29**, 110-114.
- Grossniklaus, U., Cadigan, K. M. and J., G. W.** (1994). Three maternal coordinate systems cooperate in the patterning of the *Drosophila* head. *Development* **120**, 3155-3171.
- Gulisano, M., Broccoli, V., Pardini, C. and Boncinelli, E.** (1996). *Emx1* and *Emx2* show different patterns of expression during proliferation and differentiation of the developing cerebral cortex in the mouse. *European Journal of Neuroscience* **8**, 1037-1050.
- Hadzi, J.** (1909). Über das Nervensystem von *Hydra*. *Arb. Yool. Ist. Wien* **17**, 225-268.
- Hammer, G. D., Krylova, I., Zhang, Y., Darimont, B. D., Simpson, K., Weigel, N. L. and Ingraham, H. A.** (1999). Phosphorylation of the nuclear receptor SF-1 modulates cofactor recruitment: integration of hormone signaling in reproduction and stress. *Mol Cell* **3**, 521-6.
- Hammerschmidt, M. and Nusslein-Volhard, C.** (1993). The expression of a zebrafish gene homologous to *Drosophila* snail suggests a conserved function in invertebrate and vertebrate gastrulation. *Development* **119**, 1107-18.
- Hansen, G. N., Williamson, M. and Grimmelikhuijzen, C. J.** (2000). Two-color double-labeling *in situ* hybridization of whole-mount *Hydra* using RNA probes for five different *Hydra* neuropeptide preprohormones: evidence for colocalization. *Cell Tissue Res* **301**, 245-53.
- Hansen, G. N., Williamson, M. and Grimmelikhuijzen, C. J.** (2002). A new case of neuropeptide coexpression (RGamide and LWamides) in *Hydra*, found by whole-mount, two-color double-labeling *in situ* hybridization. *Cell Tissue Res* **308**, 157-65.
- Harding, K., Hoey, T., Warrior, R. and Levine, M.** (1989). Autoregulatory and gap gene response elements of the even-skipped promoter of *Drosophila*. *Embo J* **8**, 1205-12.
- Harrison, P., Babcock, R. C., Bull, G., Oliver, J., Wallace, C. and Willis, B.** (1984). Mass spawning in tropical reef corals. *Science* **223**, 1186-1189.

- Harrison, P. L. and Wallace, C. C.** (1990). Reproduction, dispersal and recruitment of scleractinian corals. In *Coral reef ecosystem*, (ed. Dubinsky, Z.), pp. 133-207. Amsterdam: Elsevier.
- Hartmann, B., Hirth, F., Walldorf, U. and Reichert, H.** (2000). Expression, regulation and function of the homeobox gene *empty spiracles* in brain and ventral nerve cord development of *Drosophila*. *Mechanisms of Development* **90**, 143-153.
- Hartmann, B. and Reichert, H.** (1998). The genetics of embryonic brain development in *Drosophila*. *Mol Cell Neurosci* **12**, 194-205.
- Hartmann, B., Reichert, H. and Walldorf, U.** (2001). Interaction of gap genes in the *Drosophila* head: tailless regulates expression of empty spiracles in early embryonic patterning and brain development. *Mech Dev* **109**, 161-72.
- Hayward, D. C., Catmull, J., Reece-Hoyes, J. S., Berghammer, H., Dodd, H., Hann, S. J., Miller, D. J. and Ball, E. E.** (2001). Gene structure and larval expression of *cnox-2Am* from the coral *Acropora millepora*. *Development Genes and Evolution* **211**, 10-19.
- Heinemeyer, T., Wingender, E., Reuter, I., Hermjakob, H., Kel, A. E., Kel, O. V., Ignatjeva, E. V., Ananko, E. A., Podkolodnaya, O. A., Kolpakov, F. A. et al.** (1998). Databases on Transcriptional Regulation: TRANSFAC, TRRD, and COMPEL. *Nucleic Acids Res* **26**, 364-370.
- Hemmati-Brivanlou, A., de la Torre, J. R., Holt, C. and Harland, R. M.** (1991). Cephalic expression and molecular characterization of Xenopus En-2. *Development* **111**, 715-24.
- Hermesz, E., Mackem, S. and Mahon, K. A.** (1996). Rpx: a novel anterior-restricted homeobox gene progressively activated in the prechordal plate, anterior neural plate and Rathke's pouch of the mouse embryo. *Development* **122**, 41-52.
- Higashijima, S., Shishido, E., Matsuzaki, M. and Saigo, K.** (1996). eagle, a member of the steroid receptor gene superfamily, is expressed in a subset of neuroblasts and regulates the fate of their putative progeny in the *Drosophila* CNS. *Development* **122**, 527-36.
- Hirth, F., Therrianos, S., Loop, T., Gehring, W. J., Reichert, H. and Furukubo-Tokunaga, K.** (1995). Developmental defects in brain segmentation caused by mutations of the homeobox genes *orthodenticle* and *empty spiracles* in *Drosophila*. *Neuron* **15**, 769-778.
- Hoch, M., Gerwin, N., Taubert, H. and Jäckle, H.** (1992). Competition for overlapping sites in the regulatory region of the *Drosophilai* gene *Krüppel*. *Science* **256**, 94-97.
- Hoffman, C. and Winston, R.** (1987). A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene* **57**, 267-272.

- Holland, L. Z.** (2000). Body-plan evolution in the Bilateria: early antero-posterior patterning and the deuterostome-protostome dichotomy. *Current Opinions in Genetics & Development* **10**, 434-442.
- Holland, P., Garcia-Fernandez, J., Williams, N. A. and Sidow, A.** (1994). Gene duplications and the origins of vertebrate development. *Development Supplement*, 125-133.
- Holland, P. W.** (1999). Gene duplication: past, present and future. *Semin Cell Dev Biol* **10**, 541-7.
- Holland, P. W. H.** (1998). Major transitions in animal evolution: a developmental genetic perspective. *American Zoologist* **38**, 829-842.
- Hollemann, T., Bellefroid, E. and Pieler, T.** (1998). The *Xenopus* homologue of the *Drosophila* gene *tailless* has a function in early eye development. *Development* **125**, 2425-2432.
- Holley, S. A., Jackson, P. D., Sasai, Y., Lu, B., De Robertis, E. M., Hoffmann, F. M. and Ferguson, E. L.** (1995). A conserved system for dorsal-ventral patterning in insects and vertebrates involving *sog* and *chordin*. *Nature* **376**, 249-53.
- Hughes, A. L.** (1998). Phylogenetic tests of the hypothesis of block duplication of homologous genes on human chromosomes 6, 9, and 1. *Mol Biol Evol* **15**, 854-70.
- Hughes, A. L.** (1999). Phylogenies of developmentally important proteins do not support the hypothesis of two rounds of genome duplication early in vertebrate history. *J Mol Evol* **48**, 565-76.
- Hyman, L. H.** (1940). Metazoa of the tissue grade of construction - the radiate phyla - Phylum Cnidaria. In *The Invertebrates: Protozoa through Ctenophora*, pp. 365-641. New York: McGraw-Hill Book Company, Inc.
- Ip, Y. T., Park, R. E., Kosman, D., Bier, E. and Levine, M.** (1992a). The dorsal gradient morphogen regulates stripes of rhomboid expression in the presumptive neuroectoderm of the *Drosophila* embryo. *Genes Dev* **6**, 1728-39.
- Ip, Y. T., Park, R. E., Kosman, D., Yazdanbakhsh, K. and Levine, M.** (1992b). dorsal-twist interactions establish snail expression in the presumptive mesoderm of the *Drosophila* embryo. *Genes Dev* **6**, 1518-30.
- Jackson, A., Panayiotidis, P. and Foroni, L.** (1998). The human homologue of the *Drosophila* *tailless* gene (TLX): characterization and mapping to a region of common deletion in human lymphoid leukemia on chromosome 6q21. *Genomics* **50**, 34-43.
- Janknecht, R., de Martynoff, G., Lou, J., Hipskind, R., Nordheim, A. and Stunnenberg, H.** (1991). Rapid and efficient purification of native histidine-tagged protein expressed by recombinant vaccinia virus. *Proc Natl Acad Sci U S A* **88**, 8972-8976.
- Johnson, F. B., Parker, E. and Krasnow, M. A.** (1995). extradenticle protein is a selective cofactor for the *Drosophila* homeotics: Role of the homeodomain and YPWM

amino acid motif in the interaction. *Proceedings of The National Academy of Sciences* **92**, 739-743.

**Jones, B. and McGinnis, W.** (1993). The regulation of *empty spiracles* by *Abdominal-B* mediates an abdominal segment identity function. *Genes & Development* **7**, 229-240.

**Jürgens, G., Wieschaus, E., Nusslein-Volard, C. and Kluding, H.** (1984). Mutations affecting the pattern of the larval cuticle in *Drosophila melanogaster*. *Roux's Arch. Dev. Biol.* **193**, 359-377.

**Kasai, Y., Nambu, J. R., Lieberman, P. M. and Crews, S. T.** (1992). Dorsal-ventral patterning in *Drosophila*: DNA binding of snail protein to the single-minded gene. *Proc Natl Acad Sci U S A* **89**, 3414-8.

**Katsuyama, Wada, Yasugi and Saiga.** (1995). Expression of the labial group Hox gene HrHox-1 and its alteration induced by retinoic acid in development of the ascidian *Halocynthia roretzi*. *Development* **121**, 3197-205.

**Kazanskaya, O. V., Severtzova, E. A., Barth, K. A., Ermakova, G. V., Lukyanov, S. A., Benyumov, A. O., Pannese, M., Boncinelli, E., Wilson, S. W. and Zaraisky, A. G.** (1997). Anf: a novel class of vertebrate homeobox genes expressed at the anterior end of the main embryonic axis. *Gene* **200**, 25-34.

**Kirschner, M. and Gerhart, J.** (1998). Evolvability. *Proc Natl Acad Sci U S A* **95**, 8420-8427.

**Klein, W. H. and Li, X.** (1999). Function and evolution of Otx proteins. *Biochemical and Biophysical Research Communications* **258**, 229-233.

**Knoepfler, P. S. and Kamps, M. P.** (1995). The pentapeptide motif of Hox proteins is required for cooperative DNA binding with Pbx1, physically contacts Pbx1, and enhances DNA binding by Pbx1. *Mol Cell Biol* **15**, 5811-9.

**Kuhn, K., Streit, B. and Schierwater, B.** (1996). Homeobox genes in the cnidarian *Eleutheria dichotoma*: evolutionary implications for the origin of *Atennapedia*-class (HOM/Hox) genes. *Molecular Phylogenetics and Evolution* **6**, 30-38.

**Kuhn, K., Streit, B. and Schierwater, B.** (1999). Isolation of Hox genes from the scyphozoan *Cassiopeia xamachana*: implications for the early evolution of Hox genes. *J Exp Zool* **285**, 63-75.

**Kurant, E., Pai, C.-Y., Sharf, R., Halachmi, N., Sun, Y. H. and Salzberg, A.** (1998). Dorsotonals/homothorax, the *Drosophila* homologue of meis 1, interacts with extradenticle in patterning of the embryonic PNS. *Development* **125**, 1037-1048.

**Laemmli, U.** (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.

**Lai, Z. C., Fortini, M. E. and Rubin, G. M.** (1991). The embryonic expression patterns of zfh-1 and zfh-2, two *Drosophila* genes encoding novel zinc-finger homeodomain proteins. *Mech Dev* **34**, 123-34.

- Langeland, J. A., Tomsa, J. M., Jackman, W. R., Jr. and Kimmel, C. B.** (1998). An amphioxus snail gene: expression in paraxial mesoderm and neural plate suggests a conserved role in patterning the chordate embryo. *Dev Genes Evol* **208**, 569-77.
- Lewin, B.** (1997). Genes VI. Oxford: Oxford University Press.
- Liaw, G.-J. and lengyel, J.** (1992). Control of *tailless* expression by *bicoid*, *dorsal* and synergistically interacting terminal system regulatory elements. *Mechanisms of Development* **40**, 47-61.
- Liu, I. S., Chen, J. D., Ploder, L., Vidgen, D., van der Kooy, D., Kalnins, V. I. and McInnes, R. R.** (1994). Developmental expression of a novel murine homeobox gene (Chx10): evidence for roles in determination of the neuroretina and inner nuclear layer. *Neuron* **13**, 377-93.
- Liu, J., Wilson, T. E., Milbrandt, J. and Johnston, M.** (1993). Identifying DNA-binding sites and analyzing DNA-binding domains using a yeast selection system. *Methods: A Companion to Methods in Enzymology* **5**, 125-137.
- Locascio, Aniello, Amoroso, Manzanares, Krumlauf and Branno.** (1999). Patterning the ascidian nervous system: structure, expression and transgenic analysis of the CiHox3 gene. *Development* **126**, 4737-48.
- Lockett, T. J.** (1990). A bacteriophage lambda DNA purification procedure suitable for the analysis of DNA from either large or multiple small lysates. *Analytical Biochemistry* **185**, 230-234.
- Luisi, B. F., Xu, W. X., Otwinowski, Z., Freedman, L. P., Yamamoto, K. R. and Sigler, P. B.** (1991). Crystallographic analysis of the interaction of the glucocorticoid receptor with DNA. *Nature* **352**, 497-505.
- Lundin, L. G.** (1999). Gene duplications in early metazoan evolution. *Semin Cell Dev Biol* **10**, 523-30.
- Lynch, M. and Conery, J. S.** (2000). The evolutionary fate and consequences of duplicate genes. *Science* **290**, 1151-5.
- Lynch, M. and Force, A.** (2000). The probability of duplicate gene preservation by subfunctionalization. *Genetics* **154**, 459-73.
- Mackie, G. O.** (1990). The elementary nervous system revisited. *American Zoologist* **30**, 907-920.
- Mailhos, C., Andre, S., Mollereau, B., Goriely, A., Hemmati-Brivanlou, A. and Desplan, C.** (1998). Drosophila Goosecoid requires a conserved heptapeptide for repression of paired-class homeoprotein activators. *Development* **125**, 937-47.
- Mangelsdorf, D. J. and Evans, R. M.** (1995). The RXR heterodimers and orphan receptors. *Cell* **83**, 841-50.

- Mangelsdorf, D. J., Thummel, C., Beato, M., Herrlich, P., Schutz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., Chambon, P. et al.** (1995). The nuclear receptor superfamily: the second decade. *Cell* **83**, 835-9.
- Martin, A.** (2001). Is tetralogy true? Lack of support for the "one-to-four rule". *Mol Biol Evol* **18**, 89-93.
- Martin, V. J.** (2000). Reorganization of the nervous system during metamorphosis of a hydrozoan planula. *Invertebrate Biology* **119**, 243-253.
- Martin, V. J., Littlefield, C. L., Archer, W. E. and Bode, H. R.** (1997). Embryogenesis in hydra. *Biological Bulletins* **192**, 345-363.
- Martindale, M., Finnerty, J. and Henry, J.** (2002). The Radiata and the evolutionary origins of the bilaterian body plan. *Mol Phylogenet Evol* **24**, 358.
- Martindale, M. Q. and Henry, J. Q.** (1998). The development of radial and biradial symmetry: the evolution of bilaterality. *American Zoologist* **38**, 672-684.
- Martinez, D. E., Dirksen, M. L., Bode, P. M., Jamrich, M., Steele, R. E. and Bode, H. R.** (1997). Budhead, a fork head/HNF-3 homologue, is expressed during axis formation and head specification in hydra. *Dev Biol* **192**, 523-36.
- Masuda-Nakagawa, L. M., Gröger, H., Aerne, B. L. and Schmid, V.** (2000). The HOX-like gene *Cnox-2* is expressed at the anterior region in all life cycle stages of the jellyfish *Podycornye carnea*. *Development Genes and Evolution* **210**, 151-156.
- Mathers, P. H., Grinberg, A., Mahon, K. A. and Jamrich, M.** (1997). The Rx homeobox gene is essential for vertebrate eye development. *Nature* **387**, 603-7.
- McDonald, J. A., Holbrook, S., Isshiki, T., Weiss, J., Doe, C. Q. and Mellerick, D. M.** (1998). Dorsoventral patterning in the *Drosophila* central nervous system: the vnd homeobox gene specifies ventral column identity. *Genes Dev* **12**, 3603-12.
- McGinnis, W. and Krumlauf, R.** (1992). Homeobox genes and axial patterning. *Cell* **68**, 283-302.
- McMillan, J., Yellowlees, D., Heyward, A. J., Harrison, P. and Miller, D.** (1988). Preparation of high molecular weight DNA from hermatypic corals and its use for DNA hybridization and cloning. *Marine Biology* **98**, 271-276.
- Medina, M., Collins, A. G., Silberman, J. D. and Sogin, M. L.** (2001). Evaluating hypotheses of basal animal phylogeny using complete sequences of large and small subunit rRNA. *Proceedings of The National Academy of Sciences* **98**, 9707-9712.
- Mellerick, D. M. and Nirenberg, M.** (1995). Dorsal-ventral patterning genes restrict NK-2 homeobox gene expression to the ventral half of the central nervous system of *Drosophila* embryos. *Dev Biol* **171**, 306-16.
- Miles, A. and Miller, D. J.** (1992). Genomes of diploblastic organisms contain homeoboxes: sequence of eveC, an even-skipped homologue from the cnidarian *Acropora formosa*. *Proc R Soc Lond B Biol Sci* **248**, 159-61.

- Miller, D. J. and Ball, E. E.** (2000). The coral *Acropora*: what it can contribute to our knowledge of metazoan evolution and the evolution of developmental processes. *BioEssays* **22**, 291-6.
- Miller, D. J. and Harrison, P.** (1990). Molecular and developmental biology of the Cnidaria - basic aspects and phylogenetic implications. *Aust. J. Biotech* **4**, 241-250.
- Miller, D. J., Hayward, D. C., Reece-Hoyes, J. S., Scholten, I., Catmull, J., Gehring, W. J., Callaerts, P., Larsen, J. E. and Ball, E. E.** (2000). Pax gene diversity in the basal cnidarian *Acropora millepora* (Cnidaria, Anthozoa): implications for the evolution of the *Pax* gene family. *Proceedings of The National Academy of Sciences* **97**, 4475-4480.
- Minguillon, C., Ferrier, D. E., Cebrian, C. and Garcia-Fernandez, J.** (2002). Gene duplications in the prototypical cephalochordate amphioxus. *Gene* **287**, 121-8.
- Mitgutsch, C., Hauser, F. and Grimmelikhuijen, C. J. P.** (1999). Expression and developmentl regulation of the hydra-RFamide and Hydra-LWamide preprohormone genes in *Hydra*: evidence for transient phases of head formation. *Developmental Biology* **207**, 189-203.
- Miura, M., Yanazawa, M., Kato, K. and Kitamura, K.** (1997). Expression of a novel *aristaless* related homeobox gene 'Arx' in the vertebrate telencephalon, diencephalon and floor plate. *Mechanisms of Development* **65**, 99-109.
- Miyamoto, N., Yoshida, M., Kuratani, S., Matsuo, I. and Aizawa, S.** (1997). Defects of urogenital development in mice lacking *Emx2*. *Development* **124**, 1653-1664.
- Mochizuki, K., Sano, H., Kobayashi, S., Nishimiya-Fujisawa, C. and Fujisawa, T.** (2000). Expression and evolutionary conservation of *nanos*-related genes in *Hydra*. *Development Genes and Evolution* **210**, 591-602.
- Mokady, O., Dick, M. H., Lackschewitz, D., Schierwater, B. and Buss, L. W.** (1998). Over one-half billion years of head conservation? Expression of an *ems* class gene in *Hydractinia symbiolongicarpus* (Cnidaria: Hydrozoa). *Proceedings of The National Academy of Sciences* **95**, 3673-3678.
- Monaghan, A. P., Bock, D., Gass, P., Schwager, A., Wolfer, D. P., Lipp, H. P. and Schütz, G.** (1997). Defective limbic system in mice lacking the tailless gene. *Nature* **390**, 515-517.
- Monaghan, A. P., Grau, E., Bock, D. and Schütz, G.** (1995). The mouse homolog of the orphan nuclear receptor *tailless* is expressed in the developing forebrain. *Development* **121**, 839-853.
- Mori, H., Miyazaki, Y., Morita, T., Nitta, H. and Mishina, M.** (1994). Different spatio-temporal expressions of three otx homeoprotein transcripts during zebrafish embryogenesis. *Brain Res Mol Brain Res* **27**, 221-31.
- Morita, T., Nitta, H., Kiyama, Y., Mori, H. and Mishina, M.** (1995). Differential expression of two zebrafish emx homeoprotein mRNAs in the developing brain. *Neuroscience Letters* **198**, 131-134.

- Morse, A. N. C., Iwao, K., Baba, M., Shimioke, K., Hayashibara, K. and Omori, M.** (1996). An ancient chemosensory mechanism brings new life to coral reefs. *Biol Bull* **191**, 149-154.
- Much, J. W., Slade, D. J., Klampert, K., Garriga, G. and Wightman, B.** (2000). The *fax-1* nuclear hormone receptor regulates axon pathfinding and neurotransmitter expression. *Development* **127**, 703-712.
- Muller, P., Yanze, N., Schmid, V. and Spring, J.** (1999). The homeobox gene *Otx* of the Jellyfish *Podocoryne carnea*: role of a head gene in striated muscle and evolution. *Developmental Biology* **216**, 582-594.
- Murtha, M. T., Leckman, J. F. and Ruddle, F. H.** (1991). Detection of homeobox genes in development and evolution. *Proc Natl Acad Sci U S A* **88**, 10711-5.
- Myojin, M., Ueki, T., Sugahara, F., Murakami, Y., Shigentani, Y., Aizawa, S., Hirano, S. and Kuratani, S.** (2001). Isolation of *Dlx* and *Emx* gene cognates in an agnathan species, *Lampetra japonica*, and their expression patterns during embryonic and larval development: conserved and diversified regulatory patterns of homeobox genes in vertebrate head evolution. *Journal of Experimental Zoology (Mol. Dev. Evol)* **291**, 68-84.
- Naito, M., Ishiguro, H., Fujisawa, T. and Kurosawa, Y.** (1993). Presence of eight distinct homeobox-containing genes in cnidarians. *FEBS Lett* **333**, 271-4.
- Naruse, K., Fukamachi, S., Mitani, H., Kondo, M., Matsuoka, T., Kondo, S., Hanamura, N., Morita, Y., Hasegawa, K., Nishigaki, R. et al.** (2000). A detailed linkage map of medaka, *Oryzias latipes*: comparative genomics and genome evolution. *Genetics* **154**, 1773-84.
- Neuteboom, S. T., Peltenburg, L. T., van Dijk, M. A. and Murre, C.** (1995). The hexapeptide LFPWMR in *Hoxb-8* is required for cooperative DNA binding with *Pbx1* and *Pbx2* proteins. *Proc Natl Acad Sci U S A* **92**, 9166-70.
- Nielsen, C.** (1995). Animal evolution: interrelationships of the living phyla. Oxford: Oxford University Press.
- Nieto, M. A., Bennett, M. F., Sargent, M. G. and Wilkinson, D. G.** (1992). Cloning and developmental expression of *Sna*, a murine homologue of the *Drosophila* snail gene. *Development* **116**, 227-37.
- Oda, I. and Saiga, H.** (2001). *Hremx*, the ascidian homologue of *ems/emx*, is expressed in the anterior and posterior-lateral epidermis but not in the central nervous system during embryogenesis. *Development Genes and Evolution* **211**, 291-298.
- Odorico, D. and Miller, D.** (1997). Internal and external relationships of the Cnidaria: implications of primary and predicted secondary structure of the 5'-end of the 23S-like rDNA. *Proc R Soc Lond B Biol Sci* **264**, 77-82.
- Ohno, S.** (1970). Evolution by Gene Duplication. Berlin: Springer-Verlag.

- Ohta, T.** (1988). Time for acquiring a new gene by duplication. *Proc Natl Acad Sci U S A* **85**, 3509-12.
- Pankratz, M. J., Busch, M., Hoch, M., Seifert, E. and Jackle, H.** (1992). Spatial control of the gap gene knirps in the Drosophila embryo by posterior morphogen system. *Science* **255**, 986-9.
- Pannese, M., Lupo, G., Kablar, B., Boncinelli, E., Barsacchi, G. and Vignali, R.** (1998). The *Xenopus Emx* genes identify presumptive dorsal telencephalon and are induced by head organizer signals. *Mechanisms of Development* **73**, 73-83.
- Papaioannou, V. E.** (2001). T-box genes in development: from hydra to humans. *Int Rev Cytol* **207**, 1-70.
- Patarnello, T., Bargelloni, L., Boncinelli, E., Spada, F., Pannese, M. and Broccoli, V.** (1997). Evolution of Emx genes and brain development in vertebrates. *Proc R Soc Lond B Biol Sci* **264**, 1763-1766.
- Paznekas, W. A., Okajima, K., Schertzer, M., Wood, S. and Jabs, E. W.** (1999). Genomic organization, expression, and chromosome location of the human SNAIL gene (SNAI1) and a related processed pseudogene (SNAI1P). *Genomics* **62**, 42-9.
- Pellegrini, M., Mansouri, A., Simeone, A., Boncinelli, E. and Gruss, P.** (1996). Dentate gyrus formation requires *Emx2*. *Development* **122**, 3893-3898.
- Pellizzari, L., Tell, G., Fabbro, D., Pucillo, C. and Damante, G.** (1997). Functional interference between contacting amino acids of homeodomains. *FEBS Lett* **407**, 320-4.
- Peltenburg, L. T. and Murre, C.** (1996). Engrailed and Hox homeodomain proteins contain a related Pbx interaction motif that recognizes a common structure present in Pbx. *Embo J* **15**, 3385-93.
- Perlmann, T., Rangarajan, P., N., Umesono, K. and Evans, R. M.** (1993). Determinants for selective RAR and TR recognition of direct repeat HREs. *Genes & Development* **7**, 1411-1422.
- Phelan, M. L., Rambaldi, I. and Featherstone, M. S.** (1995). Cooperative interaction between HOX and PBX proteins mediated by a conserved peptide motif. *Molecular and Cellular Biology* **15**, 3989-3997.
- Pignoni, F., Baldarelli, R., Steingrimsson, E., Diaz, R. J., Patapoutian, A., Merriam, J. and Lengyel, J.** (1990). The Drosophila gene *tailless* is expressed at the embryonic termini and is a member of the steroid receptor superfamily. *Cell* **62**, 151-163.
- Pignoni, F., Steingrimsson, E. and Lengyel, J. A.** (1992). bicoid and the terminal system activate tailless expression in the early Drosophila embryo. *Development* **115**, 239-51.
- Pitman, J. L., Tsai, C. C., Edeen, P. T., Finley, K. D., Evans, R. M. and McKeown, M.** (2002). DSF nuclear receptor acts as a repressor in culture and in vivo. *Dev Biol* **245**, 315-28.

- Quandt, K., Frech, K., Karas, H., Wingender, E. and Werner, T.** (1995). MatInd and MatInspector - new fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids Res* **23**, 4878-4884.
- Qui, M., Anderson, S., Chen, S., Meneses, J. J., Hevner, R., Kuwana, E., Pedersen, R. A. and Rubenstein, J. L. R.** (1996). Mutation of the *Emx-1* homeobox gene disrupts the corpus callosum. *Developmental Biology* **178**, 174-178.
- Rastinejad, F., Perlmann, T., Evans, R. M. and Sigler, P. B.** (1995). Structural determinants of nuclear receptor assembly on DNA direct repeats. *Nature* **375**, 203-11.
- Reece-Hoyes, J. S.** (2001) *Pax* genes from the anthozoan hard coral, *Acropora millepora*: insights into *Pax* evolution Department of Biochemistry and Molecular Biology, James Cook University, Townsville
- Reichert, H. and Simeone, A.** (2001). Developmental genetic evidence for a monophyletic origin of the bilaterian brain. *Philos Trans R Soc Lond B Biol Sci* **356**, 1533-44.
- Rogozin, I. B., Lyons-Weiler, J. and Koonin, E. V.** (2000). Intron sliding in conserved gene families. *Trends in Genetics* **16**, 430-432.
- Royer, J. and Finkelstein, R.** (1995). Pattern formation in *Drosophila* head development: the role of the *orthodenticle* homeobox gene. *Development* **121**, 3561-3572.
- Ruddle, F. H., Amemiya, C. T., Carr, J. L., Kim, C. B., Ledje, C., Shashikant, C. S. and Wagner, G. P.** (1999). Evolution of chordate hox gene clusters. *Ann N Y Acad Sci* **870**, 238-48.
- Rudolph, K. M., Liaw, G.-J., Green, P., Courey, A. J., Hartenstein, V. and Lengyel, J.** (1997). Complex regulatory region mediating *tailless* expression in early embryonic patterning and brain development. *Development* **124**, 4297-4308.
- Ruppert, E. E. and Barnes, R. D.** (1991). Invertebrate Zoology. New York: Saunders College Publishing.
- Sambrook, J., Fritsch, E. F. and Maniatis, T.** (1989). Molecular cloning: a laboratory manual: Cold Spring Harbor Laboratory Press.
- Samuel, G., Miller, D. and Saint, R.** (2001). Conservation of a DPP/BMP signaling pathway in the nonbilateral cnidarian *Acropora millepora*. *Evol Dev* **3**, 241-50.
- Sanger, F., Nicklen, S. and Coulson, A. R.** (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings of The National Academy of Sciences* **74**, 5463-5467.
- Sasai, Y. and De Robertis, E. M.** (1997). Ectodermal patterning in vertebrate embryos. *Dev Biol* **182**, 5-20.

- Schierwater, B. and Desalle, R.** (2001). Current problems with the zootype and the early evolution of Hox genes. *Journal of Experimental Zoology (Mol. Dev. Evol)* **291**, 169-174.
- Schierwater, B., Murtha, M., Dick, M., Ruddle, F. H. and Buss, L. W.** (1991). Homeoboxes in cnidarians. *J Exp Zool* **260**, 413-6.
- Schneider, K.** (1890). Histologie von *Hydra fusca* mit besonderer Berücksichtigung des Nervensystems der Hydropolyphen. *Arch. mikrosk. Anat.* **35**, 321-379.
- Schneitz, K., Spielmann, P. and Noll, M.** (1993). Molecular genetics of Aristaless, a prd-type homeo box gene involved in the morphogenesis of proximal and distal pattern elements in a subset of appendages in Drosophila. *Genes Dev* **7**, 911.
- Schröder, R., Eckert, C., Wolff, C. and Tautz, D.** (2000). Conserved and divergent aspects of terminal patterning in the beetle *Tribolium castaneum*. *Proceedings of The National Academy of Sciences* **97**, 6591-6596.
- Schummer, M., Scheurlen, I., Schaller, C. and Galliot, B.** (1992). HOM/HOX homeobox genes are present in hydra (*Chlorohydra viridissima*) and are differentially expressed during regeneration. *Embo J* **11**, 1815-23.
- Scott, M. P.** (2000). Development: the natural history of genes. *Cell* **100**, 27-40.
- Sefton, M., Sanchez, S. and Nieto, M. A.** (1998). Conserved and divergent roles for members of the Snail family of transcription factors in the chick and mouse embryo. *Development* **125**, 3111-21.
- Simple, C. and Wolfe, K. H.** (1999). Gene duplication and gene conversion in the *Caenorhabditis elegans* genome. *J Mol Evol* **48**, 555-64.
- Shen, W. F., Chang, C. P., Rozenfeld, S., Sauvageau, G., Humphries, R. K., Lu, M., Lawrence, H. J., Cleary, M. L. and Largman, C.** (1996). Hox homeodomain proteins exhibit selective complex stabilities with Pbx and DNA. *Nucleic Acids Res* **24**, 898-906.
- Shenk, M. A., Bode, H. R. and Steele, R. E.** (1993a). Expression of Cnox-2, a HOM/HOX homeobox gene in hydra, is correlated with axial pattern formation. *Development* **117**, 657-67.
- Shenk, M. A., Gee, L., Steele, R. E. and Bode, H. R.** (1993b). Expression of Cnox-2, a HOM/HOX gene, is suppressed during head formation in hydra. *Dev Biol* **160**, 108-18.
- Simeone, A., Acampora, D., Gulisano, M., Stornaiuolo, A. and Boncinelli, E.** (1992a). Nested expression domains of four homeobox genes in developing rostral brain. *Nature* **358**, 687-690.
- Simeone, A., D'Apice, M. R., Nigro, V., Casanova, J., Graziani, F., Acampora, D. and Avantaggiato, V.** (1994). Orthopedia, a novel homeobox-containing gene expressed in the developing CNS of both mouse and Drosophila. *Neuron* **13**, 83-101.

- Simeone, A., Gulisano, M., Acampora, D., Stornaiuolo, A., Rambaldi, M. and Boncinelli, E.** (1992b). Two vertebrate homeobox genes related to the *Drosophila empty spiracles* gene are expressed in the embryonic cerebral cortex. *The EMBO Journal* **11**, 2541-2550.
- Skrabanek, L. and Wolfe, K. H.** (1998). Eukaryote genome duplication - where's the evidence? *Curr Opin Genet Dev* **8**, 694-700.
- Slack, J. M. W., Holland, P. W. H. and Graham, C. F.** (1993). The zootype and the phylogentic stage. *Nature* **361**, 490-492.
- Smith, D. B. and Johnson, K. S.** (1988). Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* **67**, 31-40.
- Smith, K. M., Gee, L., Blitz, I. L. and Bode, H. R.** (1999). *CnOtx*, a member of the Otx gene family has a role in cell movement in Hydra. *Developmental Biology* **212**, 392-404.
- Smith, K. M., Gee, L. and Bode, H. R.** (2000a). *HyAlx*, an *aristaless*-related gene, is involved in tentacle formation in hydra. *Development* **127**, 4743-4752.
- Smith, S., Metcalfe, J. A. and Elgar, G.** (2000b). Identification and analysis of two snail genes in the pufferfish (*Fugu rubripes*) and mapping of human SNA to 20q. *Gene* **247**, 119-28.
- Smith, S. T. and Jaynes, J. B.** (1996). A conserved region of engrailed, shared among all en-, gsc-, Nk1-, Nk2 and msh-class homeoproteins, mediates active transcriptional repression in vivo. *Development* **122**, 3141-3150.
- Sorauf, J. E. and Savarese, M.** (1995). A lower Cambrian coral from South Australia. *Palaeontology* **38**, 757-770.
- Sprigg, R. C.** (1949). Early cambrian 'jellyfishes' of Ediacara, South Australia, and Mt. John, Kimberley District, Western Australia. *Trans. Roy. Soc. S. Aust.* **73**, 72-99.
- Spring, J., Yanze, N., Josch, C., Middel, A. M., Winninger, B. and Schmid, V.** (2002). Conservation of Brachyury, Mef2, and Snail in the myogenic lineage of jellyfish: a connection to the mesoderm of bilateria. *Dev Biol* **244**, 372-84.
- Stoltzfus, A., Logsdon, J. M., Palmer, J. D. and Doolittle, W. F.** (1997). Intron "sliding" and the diversity of intron positions. *Proceedings of The National Academy of Sciences* **94**, 10739-10744.
- Strecker, T., Kongsuwan, K., Lengyel, J. and Merriam, J.** (1986). The zygotic mutant *tailless* affects the anterior and posterior ectodermal regions of the *Drosophila* embryo. *Developmental Biology* **113**, 64-76.
- Streit, Stern and Cd.** (1999). Neural induction. A bird's eye view. *Trends Genet* **15**, 20-4.

- Sugiyama, T. and Fujisawa, T.** (1978). Genetic analysis of developmental mechanisms in Hydra. II. Isolation and characterization of an interstitial cell-deficient strain. *J Cell Sci* **29**, 35-52.
- Svendsen, P. C. and McGhee, J. D.** (1995). The *C. elegans* neuronally expressed homeobox gene ceh-10 is closely related to genes expressed in the vertebrate eye. *Development* **121**, 1253-62.
- Takahashi, T., Koizumi, O., Ariura, Y., Romanovitch, A., Bosch, T. C., Kobayakawa, Y., Mohri, S., Bode, H. R., Yum, S., Hatta, M. et al.** (2000). A novel neuropeptide, Hym-355, positively regulates neuron differentiation in Hydra. *Development* **127**, 997-1005.
- Taylor, H. S.** (1998). A regulatory element of the empty spiracles homeobox gene is composed of three distinct conserved regions that bind regulatory proteins. *Molecular Reproduction and Development* **49**, 246-253.
- Technau, U. and Bode, H. R.** (1999). HyBra1, a Brachyury homologue, acts during head formation in Hydra. *Development* **126**, 999-1010.
- Teissier, G.** (1931). Etude expérimentale du développement de quelques hydriaires. *Ann. Sci. Nat. Sér. X Zool.* **14**, 5-60.
- Thompson, J. D., Higgins, D. G. and Gibson, T. J.** (1994). CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res* **22**, 4673-4680.
- Towbin, H., Staehelin, T. and Gordon, J.** (1979). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedures and some applications. *Proceedings of The National Academy of Sciences* **76**, 4350-4354.
- Treisman, J., Gonczy, P., Vashishtha, M., Harris, E. and Desplan, C.** (1989). A single amino acid can determine the DNA binding specificity of homeodomain proteins. *Cell* **59**, 553-62.
- Tynan, M. G.** (1983). Coral-like microfossils from the Lower Cambrian of California. *J. Paleontology* **57**, 1188-1211.
- Udolph, G., Rath, P. and Chia, W.** (2001). A requirement for Notch in the genesis of a subset of glial cells in the *Drosophila* embryonic central nervous system which arise through asymmetric divisions. *Development* **128**, 1457-66.
- Umesono, K. and Evans, R. M.** (1989). Determinants of target gene specificity for steroid/thyroid hormone receptors. *Cell* **57**, 1139-46.
- van Dijk, M. A. and Murre, C.** (1994). *extradenticle* raises the DNA binding specificity of homeotic selector gene products. *Cell* **78**, 617-624.
- van Dijk, M. A., Peltenburg, L. T. and Murre, C.** (1995). Hox gene products modulate the DNA binding activity of Pbx1 and Pbx2. *Mech Dev* **52**, 99-108.

- Veron, J. E. N.** (1995). Corals in Space and Time: The Biogeography and Evolution of the Scleractinia. Sydney: University of New South Wales Press.
- von Ohlen, T. and Doe, C. Q.** (2000). Convergence of dorsal, dpp, and egfr signaling pathways subdivides the drosophila neuroectoderm into three dorsal-ventral columns. *Dev Biol* **224**, 362-72.
- Wada, H., Saiga, H., Satoh, N. and Holland, P. W.** (1998). Tripartite organization of the ancestral chordate brain and the antiquity of placodes: insights from ascidian Pax-2/5/8, Hox and Otx genes. *Development* **125**, 1113-22.
- Wada, S., Katsuyama, Y., Sato, Y., Itoh, C. and Saiga, H.** (1996). Hroth an orthodenticle-related homeobox gene of the ascidian, *Halocynthia roretzi*: its expression and putative roles in the axis formation during embryogenesis. *Mech Dev* **60**, 59-71.
- Wada, S. and Saiga, H.** (1999). Cloning and embryonic expression of Hrsna, a snail family gene of the ascidian *Halocynthia roretzi*: implication in the origins of mechanisms for mesoderm specification and body axis formation in chordates. *Dev Growth Differ* **41**, 9-18.
- Walldorf, U. and Gehring, W. J.** (1992). *Empty spiracles*, a gap gene containing a homeobox involved in *Drosophila* head development. *The EMBO Journal* **11**, 2247-2259.
- Walldorf, U., Kiewe, A., Wickert, M., Ronshaugen, M. and McGinnis, W.** (2000). *Homeobrain*, a novel paired-like homeobox gene is expressed in the *Drosophila* brain. *Mechanisms of Development* **96**, 141-144.
- Weiss, J. B., Von Ohlen, T., Mellerick, D. M., Dressler, G., Doe, C. Q. and Scott, M. P.** (1998). Dorsoventral patterning in the *Drosophila* central nervous system: the *intermediate neuroblasts defective* homeobox gene specifies intermediate column identity. *Genes & Development* **12**, 3591-3602.
- Williams, N. A. and Holland, P. W.** (2000). An amphioxus Emx homeobox gene reveals duplication during vertebrate evolution. *Mol Biol Evol* **17**, 1520-8.
- Williams, S. P. and Sigler, P. B.** (1998). Atomic structure of progesterone complexed with its receptor. *Nature* **393**, 392-6.
- Wilmer, P.** (1990). Invertebrate Relationships: Patterns in Animal Evolution. Cambridge: Cambridge University Press.
- Wu, L. H. and Lengyel, J. A.** (1998). Role of caudal in hindgut specification and gastrulation suggests homology between *Drosophila* amnioproctodeal invagination and vertebrate blastopore. *Development* **125**, 2433-42.
- Yanze, N., Spring, J., Schmidli, C. and Schmid, V.** (2001). Conservation of *Hox/ParaHox*-related genes in the early development of a cnidarian. *Developmental Biology* **236**, 89-98.

- Yoshida, M., Suda, Y., Matsuo, I., Miyamoto, N., Takeda, N., Kuratani, S. and Aizawa, S.** (1997). *Emx1* and *Emx2* functions in development of dorsal telencephalon. *Development* **124**, 101-111.
- Yu, R. T., Chiang, M.-Y., Tanabe, T., Kobayashi, M., Yasuda, K., Evans, R. M. and Umesono, K.** (2000). The orphan nuclear receptor Tlx regulates Pax2 and is essential for vision. *Proceedings of The National Academy of Sciences* **97**, 2621-2625.
- Yu, R. T., McKeown, M., Evans, R. M. and Umesono, K.** (1994). Relationship between *Drosophila* gap gene *tailless* and a vertebrate nuclear receptor Tlx. *Nature* **370**, 375-379.
- Zaraisky, A. G., Lukyanov, S. A., Vasiliev, O. L., Smirnov, Y. V., Belyavsky, A. V. and Kazanskaya, O. V.** (1992). A novel homeobox gene expressed in the anterior neural plate of the Xenopus embryo. *Dev Biol* **152**, 373-82.