

References:

- Alvarado Bremer JR, Baker AJ, Mejuto J (1995) Mitochondrial DNA control region sequences indicate extensive mixing of swordfish (*Xiphias gladius*) populations in the Atlantic Ocean. *Can. J. Fish. Aquatic Sci.* 52: 1720-1732
- Alvez-Gomes JA (1999) Systematic biology of gymnotiform and mormyrid electric fishes: Phylogenetic relationships, molecular clocks and rates of evolution in the mitochondrial rRNA genes. *J. Exp. Biol.* 202: 1167-1183
- Avise JC (2000) Phylogeography: the history and formation of species. Harvard University Press, Cambridge, MA.
- Avise JC, Walker D, Johns GC (1998) Speciation durations and Pleistocene effects on vertebrate phylogeography. *Proc. R. Soc. Lond. B* 265: 1707-1712
- Bannikov AF, Tyler JC (1992) *Caprovesposus* from the Oligocene of Russia: The pelagic *Acronurus* presettlement stage of a surgeonfish (Teleostei: Acanthuridae). *Proc. Biol. Soc. Wash.* 105: 810-820
- Bannikov AF, Tyler JC (1995) Phylogenetic revision of the fish families Luvaridae and Kushlukiidae (Acanthuroidei), with a new genus and two new species of Eocene luvarids. *Smithson. Contr. Paleobiol.* 81: pp. 45
- Barber PH, Palumbi SR, Erdmann MV, Moosa MK (2002) Sharp genetic breaks among populations of *Haptosquilla pulchella* (Stomatopoda) indicate limits to larval transport: patterns, causes, and consequences. *Mol. Ecol.* 11: 659-674
- Barraclough TG, Nee S (2001) Phylogenetics and speciation. *Trends Ecol. Evol.* 16: 391-399
- Barraclough TG, Vogler AP (2000) Detecting the geographical pattern of speciation from species-level phylogenies. *Am. Nat.* 155: 419-434
- Bay LK, Choat JH, van Herwerden L, Robertson DR (online) High genetic diversities and complex genetic structure in an Indo-Pacific tropical reef fish (*Chlorurus sordidus*): Evidence of an unstable evolutionary past? *Mar. Biol.* doi: 10.1007/s00227-003-1224-3
- Bellwood DR (1996) The Eocene fishes of Monte Bolca - the earliest coral reef fish assemblage. *Coral Reefs* 15: 11-19
- Bellwood DR (1997) Reef fish biogeography: Habitat associations, fossils and phylogenies *Proc. 8th Int. Coral Reef Symp., Panama*, pp 379-384

Bellwood DR (1998) What are reef fishes? - Comment on the report by D. R. Robertson: Do coral-reef fish faunas have a distinctive taxonomic structure? (in *Coral Reefs* 17: 179-186). *Coral Reefs* 17: 187-189

Bellwood DR, Wainwright PC (2002) The history and biogeography of fishes on coral reefs. In: Sale PF (ed) *Coral reef fishes, dynamics and diversity in a complex ecosystem*. Academic Press, San Diego, pp 5-32

Benzie JAH (1999) Major genetic differences between crown-of-thorns starfish (*Acanthaster planci*) populations in the Indian and Pacific Oceans. *Evolution* 53: 1782-1795

Benzie JAH (2000) Population genetic structure in penaeid prawns. *Aquaculture Research* 31: 95-119

Benzie JAH, Ballment E, Forbes AT, Demetriades NT, Sugama K, Haryanti, Moria S (2002) Mitochondrial DNA variation in Indo-Pacific populations of the giant tiger prawn, *Penaeus monodon*. *Mol. Ecol.* 11: 2553-2569

Bermingham E, Lessios HA (1993) Rate variation of protein and mitochondrial DNA evolution as revealed by sea urchins separated by the Isthmus of Panama. *Proc. Natl. Acad. Sci.* 90: 2734-2738

Bermingham E, McCafferty SS, Martin AP (1997) Fish biogeography and molecular clocks: Perspectives from the Panamanian Isthmus. In: Kocher TD, Stepien CA (eds) *Molecular systematics of fishes*. Academic Press, California, pp 113-128

Bernardi G (2000) Barriers to gene flow in *Embiotoca jacksoni*, a marine fish lacking a pelagic larval stage. *Evolution* 54: 226-237

Bernardi G, Crane NL (1999) Molecular phylogeny of the humbug damselfishes inferred from mtDNA sequences. *J. Fish Biol.* 54: 1210-1217

Bernardi G, Holbrook SJ, Schmitt RJ (2001) Gene flow at three spatial scales in a coral reef fish, the three-spot dascyllus, *Dascyllus trimaculatus*. *Mar. Biol.* 138: 457-465

Bernardi G, Holbrook SJ, Schmitt RJ, Crane NL (2003) Genetic Evidence for two distinct clades in a French Polynesian population of the coral reef three-spotted damselfish *Dascyllus trimaculatus*. *Mar. Biol.* 143: 485-490

Bernardi G, Holbrook SJ, Schmitt RJ, Crane NL, DeMartini E (2002) Species boundaries, populations and colour morphs in the coral reef three-spot damselfish (*Dascyllus trimaculatus*) species complex. *Proc. R. Soc. Lond. B* 269: 599-605

Bernardi G, Robertson DR, Clifton KE, Azzurro E (2000) Molecular systematics, zoogeography, and evolutionary ecology of the Atlantic parrotfish genus *Sparisoma*. *Mol. Phylogenet. Evol.* 15: 292-300

Blot J, Tyler JC (1991) New genera and species of fossil surgeon fishes and their relatives (Acanthuroidei, Teleostei) from the Eocene of Monte Bolca, Italy, with application of the Blot formula to both fossil and recent forms. *Studi e Ricerche sui Giacimenti Terziari di Bolca* 6: 13-92

Bohonak AJ (2002) IBD (Isolation by Distance): a program for analyses of isolation by distance. *J. Heredity* 93: 153-154

Borden WC (1998) Phylogeny of the unicornfishes (*Naso*, Acanthuridae) based on soft anatomy. *Copeia*: 104-113

Borden WC (1999) Comparative myology of the unicornfishes, *Naso* (Acanthuridae, Percomorpha), with implications for phylogenetic analysis. *J. Morph.* 239: 191-224

Bowen BW, Avis JC (1990) Genetic structure of Atlantic and Gulf of Mexico populations of sea bass, menhaden, and sturgeon: Influence of zoogeographic factors and life-history patterns. *Mar. Biol.* 107: 371-381

Bowen BW, Bass AL, Rocha LA, Grant WS, Robertson DR (2001) Phylogeography of the trumpetfishes (*Aulostomus*): Ring species complex on a global scale. *Evolution* 55: 1029-1039

Bowen BW, Grant WS (1997) Phylogeography of the sardines (*Sardinops* spp) - assessing biogeographic models and population histories in temperate upwelling zones. *Evolution* 51: 1601-1610

Briolay J, Galtier N, Brito RM, Bouvet Y (1998) Molecular phylogeny of Cyprinidae inferred from cytochrome *b* DNA sequences. *Mol. Phylogenet. Evol.* 9: 100-108

Brooks DR, McLennan DA (2002) The nature of diversity. The University of Chicago Press, Chicago

Cantantore P, Roberti M, Pesole G, Ludovico A, Milella F, Gadaleta MN, Saccone C (1994) Evolutionary analysis of cytochrome *b* sequences in some Perciformes: Evidence for a slower rate of evolution than in mammals. *J. Mol. Evol.* 39: 589-597

Chase MR, Etter RJ, Rex MA, Quattro JM (1998) Extraction and amplification of mitochondrial DNA from formalin-fixed deep-sea mollusks. *Biotechniques* 24: 243-247

Chen L-S (1999) Ontogenetic development in post-settlement scarids (Pisces: Scaridae): transition in feeding modes. PhD Thesis. School of Marine Biology & Aquaculture, Townsville, Australia

Chenoweth SF, Hughes JM, Connolly RC (2002) Phylogeography of the pipefish, *Urocampus carinirostris*, suggests secondary intergradation of ancient lineages. Mar. Biol. 141: 541-547

Choat JH, Clements KD (1998) Vertebrate herbivory in marine and terrestrial environments: A nutritional ecology perspective. Annu. Rev. Ecol. Syst. 29: 375-403

Choat JH, Clements KD, Robbins WD (2002) The trophic status of herbivorous fishes on coral reefs - I: Dietary analyses. Mar. Biol. 140: 613-623

Choat JH, Robertson DR (2002) Age-based studies. In: Sale PF (ed) Coral reef fishes, dynamics and diversity in a complex ecosystem. Academic Press, San Diego, pp 57-80

Clements KD, Choat JH (1995) Fermentation in tropical marine herbivorous fishes. Physiol. Zool. 68: 355-378

Clements KD, Gray RD, Choat JH (2003) Rapid evolutionary divergence in reef fishes of the family Acanthuridae (Perciformes; Teleostei). Mol. Phylogenet. Evol. 26: 190-201

Colborn J, Crabtree RE, Shaklee JB, Pfeiler E, Bowen BW (2001) The evolutionary enigma of bonefishes (*Albula* spp.): Cryptic species and ancient separations in a globally distributed shorefish. Evolution 55: 807-820

Craig MT, Pondella DJ, Frank JPC, Hafner JC (2001) On the status of the serranid fish genus *Epinephelus*: Evidence for paraphyly based upon 16S rDNA sequence. Mol. Phylogenet. Evol. 19: 121-130

Dayton C, Santayana AC, Lacson JM (1994) Genetic evidence for reproductive isolation of the recently described unicornfish *Naso caesius* and its sibling *N. hexacanthus*. Mar. Biol. 118: 551-554

Dayton CL (2001) Genetic Evolution among selected members of the genus *Naso* (Nasinae), "unicornfishes" from Guam. Mar. Biol. 139: 771-776

Dillon N, Austin AD, Bartowsky E (1996) Comparison of preservation techniques for DNA extraction from hymenopterous insects. Insect Mol. Biol. 5: 21-24

Doherty PJ, Mather P, Planes S (1994) *Acanthochromis polyacanthus*, a fish lacking larval dispersal, has genetically differentiated populations at local and regional scales on the Great Barrier Reef. Mar. Biol. 121: 11-21

Donaldson KA, Wilson RR (1999) Amphi-Panamic geminates of snook (Percoidei: Centropomidae) provide a calibration of the divergence rate in the mitochondrial DNA control region of fishes. Mol. Phylogenet. Evol. 13: 208-213

- Duda TF, Palumbi SR (1999) Population structure of the black tiger prawn, *Penaeus monodon*, among western Indian Ocean and western Pacific populations. Mar. Biol. 134: 705-710
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. Genetics 131: 479-491
- Fauvelot C, Bernardi G, Planes S (2003) Reductions in the mitochondrial DNA diversity of coral reef fish provide evidence of population bottlenecks resulting from Holocene sea-level change. Evolution 57: 1571-1583
- Feral JP (2002) How useful are the genetic markers in attempts to understand and manage marine biodiversity? J. Exp. Mar. Biol. Ecol. 268: 121-145
- Finnerty JR, Block BA (1992) Direct sequencing of mitochondrial DNA detects highly divergent haplotypes in blue marlin (*Makaira nigricans*). Mol. Mar. Biol. Biotechnol. 1: 206-214
- Glazko GV, Nei M (2003) Estimation of divergence times for major lineages of primate species. Mol. Biol. Evol. 20: 424-434
- Grant WS, Bowen BW (1998) Shallow population histories in deep evolutionary lineages of marine fishes: Insights from sardines and anchovies and lessons for conservation. J. Heredity 89: 415-426
- Grant WS, Clark AM, Bowen BW (1998) Why restriction fragment length polymorphism analysis of mitochondrial DNA failed to resolve sardine (*Sardinops*) biogeography: insights from mitochondrial DNA cytochrome b sequences. Can. J. Fish. Aquat. Sci. 55: 2539-2547
- Graves JE (1998) Molecular insights into the population structures of cosmopolitan marine fishes. J. Hered. 89: 427-437
- Hallam A (1984) Pre-Quaternary sea-level changes. Annu. Rev. Earth Planet. Sci. 12: 205-243
- Haq BU, Hardenbol J, Vail PR (1987) Chronology of fluctuating sea levels since the Triassic. Science 235: 1156-1167
- Hartl DL (2000) A primer of population genetics. Sinauer Assoc. Inc., Sunderland MA. USA
- Hartl DL, Clark AG (1997) Principles of population genetics. Sinauer Associations Inc., Sunderland, MA., pp 1-542

- Hewitt G (2003) A climate for colonization. *Heredity* online/in press
- Hewitt GM (2001) Speciation, hybrid zones and phylogeography - or seeing genes in space and time. *Mol. Ecol.* 10: 537-549
- Hillis DM, Mable BK, Moritz C (1996) Applications of molecular systematics: The state of the field and a look to the future. In: Hillis DM, Moritz C, Mable BK (eds) *Molecular systematics*. Sinauer Assoc. Inc., Sunderland, pp 515-543
- Huelsenbeck JP, Ronquist F (2001) MR. BAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17: 754-755
- Johnson JW (2002) *Naso mcdadei*, a new species of unicornfish (Perciformes: Acanthuridae), with a review of the *Naso tuberosus* species complex. *Aust. J. Zool.* 50: 293-311
- Jones RS (1968) Ecological relationships in Hawaiian and Johnston Island Acanthuridae (Surgeonfishes). *Micronesica* 4: 309-361
- Keeney DB, Heupel M, Hueter RE, Heist EJ (2003) Genetic heterogeneity among blacktip sharks, *Carcharhinus limbatus*, continental nurseries along the U.S. Atlantic and Gulf of Mexico. *Mar. Biol.* in press/online
- Kiessling W (2001a) Paleoclimatic significance of Phanerozoic reefs. *Geology* 29: 751-754
- Kiessling W (2001b) Phanerozoic reef trends based on the Paleoreef database. In: Stanley GD (ed) *The history and sedimentology of ancient reef systems*. Kluwer Academic/Plenum Publ., New York, pp 41-87
- Knowlton N (1993) Sibling species in the sea. *Annu. Rev. Ecol. Syst.* 24: 189-216
- Knowlton N (2000) Molecular genetic analyses of species boundaries in the sea. *Hydrobiologia* 420: 73-90
- Knowlton N, Weigt LA (1998) New dates and new rates for divergence across the Isthmus of Panama. *Proc. R. Soc. Lond. B* 265: 2257-2263
- Kocher TD, Thomas WK, Meyer A, Edwards SV, Paabo S, Villablanca FX, Wilson C (1989) Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *Proc. Natl. Acad. Sci.* 86: 6196-6200
- Kuiter RH, Debelius H (2001) Surgeonfishes, rabbitfishes and their relatives. TMC Publ., Chorleywood, U.K.
- Kumazawa Y, Nishida M (2000) Molecular phylogeny of osteoglossids: A new model for Gondwanian origin and plate tectonic transportation of the Asian Arowana. *Mol. Biol. Evol.* 17: 1869-1878

- Langley CH, Fitch WM (1974) An examination of the constancy of the rate of molecular evolution. *J. Mol. Evol.* 3: 161-177
- Leis JM, Carson-Ewart BM (1997) *In situ* swimming speeds of the late pelagic larvae of some Indo-Pacific coral-reef fishes. *Mar. Ecol. Prog. Ser.* 159: 165-174
- Leis JM, Carson-Ewart BM (2000) The larvae of Indo-Pacific coastal fishes: An identification guide to marine fish larvae. In: de Winter AJ (ed). Brill, Leiden, pp 1-850
- Leis JM, McCormick MI (2002) The biology, behavior, and ecology of the pelagic, larval stage of coral reef fishes. In: Sale PF (ed) *Coral reef fishes, dynamics and diversity in a complex ecosystem*. Academic Press, San Diego, pp 171-199
- Leis JM, Richards WJ (1984) Acanthuroidei: Development and relationships. Am. Soc. Ichthyologists & Herpetologists, La Jolla, California
- Lessios HA, Kane J, Robertson DR (2003) Phylogeography of the pantropical sea urchin *Tripneustes*: Contrasting patterns of population structure between oceans. *Evolution* 57: 2026-2036
- Lessios HA, Kessing BD, Pearse JS (2001) Population structure and speciation in tropical seas: Global phylogeography of the sea urchin *Diadema*. *Evolution* 55: 955-975
- Li JJ, Liao X, Yang H (2000) Molecular characterization of a parasitic tapeworm (*Ligula*) based on DNA sequences from formalin-fixed specimens. *Biochemical Genetics* 38: 309-322
- Lo Galbo AM, Carpenter KE, Reed DL (2002) Evolution of tropic types in emperor fishes (*Lethrinus*, Lethrinidae, Percoidei) based on cytochrome b gene sequence variation. *J Mol. Evol.* 54: 754-762
- Losos JB, Glor RE (2003) Phylogenetic comparative methods and the geography of speciation. *Trends Ecol. Evol.* 18: 220-227
- Lyons LA, Laughlin TF, Copeland NG, Jenkins NA, Womack JE, O'Brien SJ (1997) Comparative anchor tagged sequences (CATS) for integrative mapping of mammalian genomes. *Nature genetics* 15: 47-56
- Machordom A, Doadrio I (2001) Evidence of a Cenozoic Betic-Kabilian connection based on freshwater fish phylogeography (*Luciobarbus*, Cyprinidae). *Mol. Phylogenet. Evol.* 18: 252-263
- Maddison DR, Maddison WP (2001) *MacClade 4: Analysis of phylogeny and character evolution*. Sinauer Associates, Sunderland, Massachusetts

- Markert JA, Arnegard ME, Danley PD, Kocher TD (1999) Biogeography and population genetics of the Lake Malawi cichlid *Melanochromis auratus*: habitat transience, philopatry and speciation. Mol. Ecol. 8: 1013-1026
- Martin AP (1995a) Metabolic rate and directional nucleotide substitution in animal mitochondrial DNA. Mol. Biol. Evol. 12: 1124-1131
- Martin AP (1995b) Mitochondrial DNA sequence evolution in sharks - rates, patterns and phylogenetic inferences. Mol. Biol. Evol. 12: 1114-1123
- Martin AP, Naylor GJP, Palumbi SR (1992) Rates of mitochondrial DNA evolution in sharks are slow compared with mammals. Nature 357: 153-155
- Martin AP, Palumbi SR (1993a) Body size, metabolic rate, generation time, and the molecular clock. Proc. Natl. Acad. Sci. 90: 4087-4091
- Martin AP, Palumbi SR (1993b) Protein evolution in different cellular environments: Cytochrome *b* in sharks and mammals. Mol. Biol. Evol. 10: 873-891
- Mayr E (1954) Geographic speciation in tropical echinoids. Evolution 8: 1-18
- McMillan WO, Palumbi SR (1995) Concordant evolutionary patterns among Indo-West Pacific Butterflyfishes. Proc. R. Soc. Lond. B 260: 229-236
- McMillan WO, Palumbi SR (1997) Rapid rate of control-region evolution in Pacific Butterflyfishes (Chaetodontidae). J Mol. Evol. 45: 473-484
- Muss A, Robertson DR, Stepien CA, Wirtz P, Bowen BW (2001) Phylogeography of *Ophioblennius*: The role of ocean currents and geography in reef fish evolution. Evolution 55: 561-572
- Nei M (1987) Molecular evolutionary genetics. Columbia University Press, New York
- Nei M, Kumar S (2000) Molecular evolution and phylogenetics. Oxford University Press, Inc. New York
- Nelson JS, Hoddell RJ, Chou LM, Chan WK, Phang VPE (2000) Phylogeographic structure of false clownfish, *Amphiprion ocellaris*, explained by sea level changes on the Sunda shelf. Mar. Biol. 137: 727-736
- Nicholas KB, Nicholas HBJ (1997) GeneDoc: a tool for editing and annotating multiple sequence alignments
- Norris RD (2000) Pelagic species diversity, biogeography, and evolution. Paleobiology 26: 236-258

- Orti G, Bell MA, Reimchen TE, Meyer A (1994) Global survey of mitochondrial DNA sequences in the threespine stickleback: Evidence for recent migrations. *Evolution* 48: 608-622
- Ovenden JR, Lloyd J, Newman SJ, Keenan CP, Slater LS (2002) Spatial genetic subdivision between northern Australian and southeast Asian populations of *Pristipomoides multidens*: A tropical marine reef fish species. *Fish. Res.* 59: 57-69
- Palumbi SR (1992) Marine speciation on a small planet. *Trends Ecol. Evol.* 7: 114-118
- Palumbi SR (1994) Genetic divergence, reproductive isolation, and marine speciation. *Annu. Rev. Ecol. Syst.* 25: 547-572
- Palumbi SR (1997) Molecular biogeography of the Pacific. *Coral Reefs* 16 Suppl.: S47-S52
- Palumbi SR (2003) Population genetics, demographic connectivity, and the design of marine reserves. *Ecol. Appl.* 13: S146-S158
- Pickering KT (2000) The Cenozoic world. In: Culver SJ, Rawson PF (eds) *Biotic response to global change*. Cambridge University Press, pp 20-34
- Planes S, Doherty PJ (1997) Genetic and color interactions at a contact zone of *Acanthochromis polyacanthus*: A marine fish lacking pelagic larvae. *Evolution* 51: 1232-1243
- Planes S, Doherty PJ, Bernardi G (2001) Strong genetic divergence among populations of a marine fish with limited dispersal, *Acanthochromis polyacanthus*, within the Great Barrier Reef and the Coral Sea. *Evolution* 55: 2263-2273
- Planes S, Fauvelot C (2002) Isolation by distance and vicariance drive genetic structure of a coral reef fish in the Pacific Ocean. *Evolution* 56: 378-399
- Planes S, Galzin R, Bonhomme F (1996) A genetic metapopulation model for reef fishes in oceanic islands: The case of the surgeonfish, *Acanthurus triostegus*. *J Evol. Biol.* 9: 103-117
- Posada D, Crandall KA (1998) Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817-818
- Prothero DR (1994) The late Eocene-Oligocene extinctions. *Annu. Rev. Earth Planet. Sci.* 22: 145-165
- Randall JE (1994) Unicornfishes of the subgenus *Axinurus* (Perciformes: Acanthuridae: *Naso*), with description of a new species. *Copeia*: 116-124
- Randall JE (2001) *Naso reticulatus*, a new unicornfish (Perciformes: Acanthuridae) from Taiwan and Indonesia, with a key to the species of *Naso*. *Zool. Studies* 40: 170-176

Randall JE (2002) Surgeonfishes of Hawai'i and the world. Mutual Publishing & Bishop Museum Press

Robertson DR (1982) Fish feces as fish food on a Pacific coral reef. Mar. Ecol. Prog. Ser. 7: 253-265

Rocha LA, Bass AL, Robertson DR, Bowen BW (2002) Adult habitat preferences, larval dispersal, and the comparative phylogeography of three Atlantic surgeonfishes (Teleostei: Acanthuridae). Mol. Ecol. 11: 243-252

Ruber L, Van Tassell JL, Zardoya R (2003) Rapid speciation and ecological divergence in the American seven-spined gobies (Gobiidae, Gobiosomatini) inferred from a molecular phylogeny. Evolution 57: 1584-1598

Sambrook J, Fritsh EF, Maniatis T (1989) Molecular cloning, a laboratory manual. Cold Spring Harbour Laboratory Press

Sanderson MJ (1997) A nonparametric approach to estimating divergence times in the absence of rate constancy. Mol. Biol. Evol. 14: 1218-1231

Sanderson MJ (2003) r8s: Inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. Bioinformatics 19: 301-302

Schluter D (2001) Ecology and the origin of species. Trends Ecol. Evol. 16: 372-380

Schluter D, Boughman JW, Rundle HD (2001) Parallel speciation with allopatry. Trends Ecol. Evol. 16: 283-284

Schneider S, Roessli D, Excoffier L (2000) A software for population genetics data analysis (ARLEQUIN)

Shedlock AM, Haygood MG, Pietsch TW, Bentzen P (1997) Enhanced DNA extraction and PCR amplification of mitochondrial genes from formalin-fixed museum specimens. Biotechniques 22: 394, 396, 398, 400

Sheehan PM (2001) History of marine biodiversity. Geological J 36: 231-249

Shiozawa DK, Kudo J, Evans RP, Woodward SR, Williams RN (1992) DNA extraction from preserved trout tissues. Great Basin Naturalist 52: 29-34

Shulman MJ, Bermingham E (1995) Early life histories, ocean currents, and the population genetics of Caribbean reef fishes. Evolution 49: 897-910

- Simon C, Frati F, Beckenbach A, Crespi B, Liu H, Flook P (1994) Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved Polymerase Chain Reaction primers. *Anna. Entomol. Soc. Am.* 87: 651-701
- Slatkin M (1993) Isolation by distance in equilibrium and non-equilibrium populations. *Evolution* 47: 264-279
- Smith JLB (1966) Fishes of the sub-family Nasinae with a synopsis of the Prionurinae. *Ichthyological Bulletin* 32: 635-682
- Smith MM, Heemstra PC (1986) Smiths' sea fishes. Springer-Verlag, Berlin, pp 1-1047
- Sorbini L, Tyler JC (1998a) A new genus and species of Eocene surgeon fish (Acanthuridae) from Monte Bolca, Italy, with similarities to the recent *Zebrasoma*. *Studi e Ricerche sui Giacimenti Terziari di Bolca* 7: 7-19
- Sorbini L, Tyler JC (1998b) A new species of the Eocene surgeon fish genus *Pesciarichthys* from Monte Bolca, Italy (Acanthuridae), with comments on caudal peduncle armature and supraneurals in acanthurids. *Studi e Ricerche sui Giacimenti Terziari di Bolca* 7: 21-34
- Stanley GDJ (2001) The history and sedimentology of ancient reef systems. In: Landman NH, Jones DS (eds). Kluwer Academic/Plenum Publisher, New York, pp 1-458
- Stepien CA, Faber JE (1998) Population genetic structure, phylogeography and spawning philopatry in walleye (*Stizostedion vitreum*) from mitochondrial DNA control region sequences. *Mol. Ecol.* 7: 1757-1769
- Stepien CA, Rosenblatt RH, Bargmeyer BA (2001) Phylogeography of the spotted sand bass, *Paralabrax maculatusfasciatus*: Divergence of Gulf of California and Pacific coast populations. *Evolution* 55: 1852-1862
- Stobutzki IC (1997) Energetic cost of sustained swimming in the late pelagic stages of reef fishes. *Mar. Ecol. Prog. Ser.* 152: 249-259
- Streelman JT, Alfaro M, Weastneat MW, Bellwood DR, Karl SA (2002) Evolutionary history of the parrotfishes: biogeography, ecomorphology, and comparative diversity. *Evolution* 56: 961-971
- Swofford DL (1998) PAUP* Phylogenetic Analysis Using Parsimony (* and other methods). Sinauer Associates, Massachusetts, Sunderland
- Takahashi H, Goto A (2001) Evolution of east Asian ninespine sticklebacks as shown by mitochondrial DNA control region sequences. *Mol. Phylogenet. Evol.* 21: 135-155

- Tang KL (2001) Phylogenetic relationships among damselfishes (Teleostei: Pomacentridae) as determined by mitochondrial DNA data. *Copeia*: 591-601
- Tang KL, Berendzen PB, Wiley EO, Morrissey JF, Winterbottom R, Johnson GD (1999) The phylogenetic relationships of the suborder Acanthuroidei (Teleostei: Perciformes) based on molecular and morphological evidence. *Mol. Phylogenet. Evol.* 11: 415-425
- Terry A, Buccarelli G, Bernardi G (2000) Restricted Gene Flow and Incipient Speciation in Disjunct Pacific Ocean and Sea of Cortez Populations of a Reef Fish Species, *Girella nigricans*. *Evolution* 54: 652-659
- Thacker CE (2003) Molecular phylogeny of the gobioid fishes (Teleostei : Perciformes : Gobioidei). *Mol. Phylogenet. Evol.* 26: 354-368
- Thompson JD, Gibson TJ, Plewniak F, F. J, Higgins DG (1997) The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 24: 4876-4882
- Turelli M, Barton NH, Coyne JA (2001) Theory and speciation. *Trends Ecol. Evol.* 16: 330-343
- Tyler JC (1970) Osteological aspects of interrelationships of surgeon fish genera (Acanthuridae). *Proc. Acad. Nat. Sci. Phila.* 122: 87-124
- Tyler JC (1997) The Miocene fish *Marosichthys*, a putative Tetraodontiform, actually a perciform surgeon fish (Acanthuridae) related to the recent *Naso*. *Beaufortia* 47: 1-10
- Tyler JC (1999a) A new genus and species of surgeon fish (Acanthuridae) with a unique dorsal-fin pterygiophore arrangement from the Eocene of Monte Bolca, Italy. *Studi e Ricerche sui Giacimenti Terziari di Bolca* 7: 245-256
- Tyler JC (1999b) A new genus and species of surgeon fish (Acanthuridae) with four dorsal-fin spines from the Eocene of Monte Bolca, Italy. *Studi e Ricerche sui Giacimenti Terziari di Bolca* 7: 257-268
- Tyler JC (2000) *Arambourgthurus*, a new genus of hypurostegic surgeonfish (Acanthuridae) from the Oligocene of Iran, with a phylogeny of Nasinae. *Geodiversitas* 22: 525-537
- Tyler JC, Bannikov AF (2000) A new species of the surgeon fish genus *Tauichthys* from the Eocene of Monte Bolca, Italy (Perciformes, Acanthuridae). *Bollettino del Museo Civico di Storia Naturale di Verona* 24: 29-36

Tyler JC, Johnson GD, Nakamura I, Collette BB (1989) Morphology of *Luvarus imperialis* (Luvaridae), with a phylogenetic analysis of the Acanthuroidei (Pisces). Smithson. Contr. Zool. 485: 1-78

Tyler JC, Sorbini L (1998) On the relationships of *Eonaso*, an Antillean fossil surgeon fish (Acanthuridae). Studi e Ricerche sui Giacimenti Terziari di Bolca 7: 35-42

van Herwerden L, Benzie J, Peplow L, Davies C (2000) Microsatellite markers for coral trout (*Plectropomus laevis*) and red throat emperor (*Lethrinus miniatus*) and their utility in other species of reef fish. Mol. Ecol. 9: 1929-1931

van Herwerden L, Davies CR, Choat JH (2002) Phylogenetic and evolutionary perspectives of the Indo-Pacific grouper *Plectropomus* species on the Great Barrier Reef, Australia. J. Fish Biol. 60: 1591-1596

Via S (2001) Sympatric speciation in animals: the ugly duckling grows up. Trends Ecol. Evol. 16: 381-390

Wang HY, Tsai MP, Dean J, Lee SC (2001) Molecular phylogeny of gobioid fishes (Perciformes : Gobioidei) based on mitochondrial 12S rRNA sequences. Mol. Phylogenet. Evol. 20: 390-408

Waters JM, Wallis GP (2001) Cladogenesis and loss of the marine life-history phase in freshwater galaxiid fishes (Osmeriformes: Galaxiidae). Evolution 55: 587-597

Weir BS (1996) Genetic Data Analysis II. Sinauer Assoc. Inc., Sunderland, MA. USA

Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. Evolution 38: 1358-1370

Williams ST, Benzie JAH (1998) Evidence of a biogeographic break between populations of a high dispersal starfish: Congruent regions within the Indo-West Pacific defined by color morphs, mtDNA, and allozyme data. Evolution 52: 87-99

Wilson DT, McCormick MI (1999) Microstructure of settlement-marks in the otoliths of tropical reef Fishes. Mar. Biol. 134: 29-41

Winterbottom R (1993) Myological evidence for the phylogeny of recent genera of surgeonfishes (Percomorpha, Acanthuridae), with comments on the Acanthuroidei. Copeia 1993: 21-39

Winterbottom R, McLennan DA (1993) Cladogram versatility: Evolution and biogeography of acanthuroid fishes. Evolution 47: 1557-1571

Wirgin I, Maceda L, Stabile J, Mesing C (1997) An evaluation of introgression of Atlantic coast striped bass mitochondrial DNA in a Gulf of Mexico population using formalin-preserved museum collections. Mol. Ecol. 6: 907-916

- Xia X (2000) Data Analysis in Molecular Biology and Evolution (DAMBE)
- Xia XH, Xie Z, Salemi M, Chen L, Wang Y (2003) An index of substitution saturation and its application. *Mol. Phylogenetic Evol.* 26: 1-7
- Yoder AD, Irwin JA, Payseur BA (2001) Failure of the ILD to determine data combinability for Slow Loris phylogeny. *Syst. Biol.* 50: 408-424
- Zachos J, Pagani M, Sloan L, Thomas E, Billups K (2001a) Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292: 686-692
- Zachos JC, Shackleton NJ, Revenaugh JS, Paelke H, Flower BP (2001b) Climate response to orbital forcing across the Oligocene-Miocene boundary. *Science* 292: 274-278
- Zardoya R, Doadrio I (1999) Molecular evidence on the evolutionary and biogeographical patterns of European cyprinids. *J. Mol. Evol.* 49: 227-237