

JCU ePrints

This file is part of the following reference:

Braby, Michael (1994) *Life history strategies of tropical satyrine butterflies in north-eastern Australia*. PhD thesis, James Cook University.

Access to this file is available from:

<http://eprints.jcu.edu.au/10857>

References

- Ackerman, J.D. (1983) Diversity and seasonality of male euglossine bees (Hymenoptera: Apidae) in central Panama. *Ecology* **64**: 274-283.
- Ackery, P.R. (1984) Systematic and faunistic studies on butterflies. In *The Biology of Butterflies* (Eds R.I. Vane-Wright and P.R. Ackery), pp. 9-12. *Symposium of the Royal Entomological Society of London. Number 11*. Academic Press, London.
- Adler, P.H. and Pearson, D.L. (1982) Why do male butterflies visit mud puddles? *Can. J. Zool.* **60**: 322-325.
- Aiello, A. (1992) Dry season strategies of two Panamanian butterfly species, *Anartia fatima* (Nymphalinae) and *Pierella luna luna* (Satyrinae) (Lepidoptera: Nymphalidae). In *Insects of Panama and Mesoamerica: Selected Studies* (Eds Quintero, D. and Aiello, A.), pp. 573-575. Oxford University Press, Oxford.
- Akkawi, M.M. and Scott, D.R. (1984) The effect of age of parents on the progeny of diapaused and non-diapaused *Heliothis zea*. *Entomol. Exp. Appl.* **35**: 235-239.
- Alan, F.E. (1989) *Chemical Analyses of Ecological Materials*. Blackwell Scientific Publishing, Oxford.
- Angel, F.M. (1951) Notes on the Lepidoptera of the Northern Territory of Australia, with descriptions of new species. *Trans. R. Soc. Aust.* **74**: 6-14.
- Arms, K., Feeny, P. and Lederhouse, R.C. (1974) Sodium: stimulus for puddling behaviour by tiger swallowtail butterflies. *Science* **185**: 372-374.
- Atkins, A.F. (1976) New records for butterflies in southern, central and northern Queensland. *Aust. ent. Mag.* **3**: 1-4.
- Bailey, N.T.J. (1951) On estimating the size of mobile populations from recapture data. *Biometrika* **38**: 293-306.
- Bailey, N.T.J. (1952) Improvements in the interpretation of recapture data. *J. Anim. Ecol.* **21**: 120-127.
- Baker, H.G. and Baker, I. (1973) Amino-acids in nectar and their evolutionary significance. *Nature* **241**: 543-545.

- Baker, H.G. and Baker, I. (1982) Chemical constituents of nectar in relation to pollination mechanisms and phylogeny. In *Biochemical Aspects of Evolutionary Biology* (Ed M. Nitecki), pp. 126-171. University of Chicago Press, Chicago.
- Ballmer, G.R. and Pratt, G.F. (1989) Instar number and larval development in *Lycaena phlaeas hypophlaeas* (Boisduval) (Lycaenidae). *J. Lepid. Soc.* **43**: 59-65.
- Banno, H. (1990) Plasticity of size and relative fecundity in the aphidophagous lycaenid butterfly, *Taraka hamada*. *Ecol. Ent.* **15**: 111-113.
- Barrett, C. and Burns, A.N. (1951) *Butterflies of Australia and New Guinea*. N.H. Seward, Melbourne.
- Begon, M. (1985) A general theory of life history variation. In *Behavioural Ecology. Ecological Consequences of Adaptive Behaviour* (Eds R.M. Sibly and R.H. Smith), pp. 91-97. Blackwell Scientific Publications, Oxford.
- Begon, M., Harper, J.L. and Townsend, C.R. (1990) *Ecology. Individuals, Populations and Communities*. Second Edition. Blackwell Scientific Publications, Oxford.
- Begon, M. and Parker, G.A. (1986) Should egg size and clutch size decrease with age? *Oikos* **47**: 293-302.
- Bell, G. and Koufopanou, V. (1986) The cost of reproduction. *Oxford Surv. Evol. Biol.* **3**: 83-131.
- Berger, A. (1989) Egg weight, batch size and fecundity of the spotted stalk borer, *Chilo partellus* in relation to weight of females and time of oviposition. *Entomol. exp. Appl.* **50**: 199-207.
- Blueweiss, L., Fox, H., Kudzma, V., Nakashima, D., Peters, R. and Sam, S. (1978) Relationships between body size and some life history parameters. *Oecologia* **37**: 257-272.
- Boggs, C.L. (1981a) Nutritional and life history determinants of resource allocation in holometabolous insects. *Am. Nat.* **117**: 692-709.
- Boggs, C.L. (1981b) Selection pressures affecting male nutrient investment at mating in Heliconiine butterflies. *Evolution* **35**: 931-940.

- Boggs, C.L. (1986) Reproductive strategies of female butterflies: variation in and constraints on fecundity. *Ecol. Entomol.* **11**: 7-15.
- Boggs, C.L. (1987) Ecology of nectar and pollen feeding in Lepidoptera. In *Nutritional Ecology of Insects, Mites and Spiders* (Eds F. Slansky, Jr. and J.G. Rodriguez), pp. 369-391. John Wiley and Sons, New York.
- Boggs, C.L. (1990) A general model of the role of male-donated nutrients in female insect's reproduction. *Am. Nat.* **136**: 598-617.
- Boggs, C.L. and Gilbert, L.E. (1979) Male contribution to egg production in butterflies: evidence for transfer of nutrients at mating. *Science* **206**: 83-84.
- Boggs, C.L. and Jackson, L.A. (1991) Mud puddling by butterflies is not a simple matter. *Ecol. Entomol.* **16**: 123-127.
- Boggs, C.L. and Ross, C.L. (1993) The effect of adult food limitation on life history traits in *Speyeria mormonia* (Lepidoptera: Nymphalidae). *Ecology* **74**: 433-441.
- Boggs, C.L., Smiley, J.T. and Gilbert, L.E. (1981) Patterns of pollen exploitation by *Heliconius* butterflies. *Oecologia* **48**: 284-289.
- Boggs, C.L. and Watt, W.B. (1981) Population structure of Pierid butterflies. IV. Genetic and physiological investment in offspring by male *Colias*. *Oecologia* **50**: 320-324.
- Boyce, R.M. (1984) Restitution of *r*- and *K*-selection as a model of density-dependent natural selection. *Ann. Rev. Ecol. Syst.* **15**: 427-447.
- Boyce, M.S. and Perrins, C.S. (1987) Optimising Great tit clutch size in a fluctuating environment. *Ecology* **68**: 142-153.
- Braby, M.F. (1991) Migration of *Euploea core corinna* (W.S. Macleay) (Nymphalidae: Danainae) in northern Queensland, Australia. *Aust. ent. Mag.* **18**: 143-146.
- Braby, M.F. (1992) Conservation needs of lowland, coastal paperbark woodlands and eucalypt open forests in northern Queensland - notes on some rare and threatened butterflies. *News Bull. Ent. Soc. Qld.* **20**: 76-88.
- Braby, M.F. (1993) Early stages, biology and taxonomic status of *Tisiphone helena* (Olliff) (Lepidoptera: Nymphalidae: Satyrinae). *J. Aust. ent. Soc.*

32: 273-282.

Braby, M.F. and New, T.R. (1988a) Population biology of adult *Geitoneura klugii* and *G. acantha* (Lepidoptera: Satyrinae) near Melbourne, Australia. *Aust. J. Zool.* **36**: 141-158.

Braby, M.F. and New, T.R. (1988b) Adult reproductive biology of *Geitoneura klugii* and *G. acantha* (Lepidoptera: Satyrinae) near Melbourne, Australia. *Aust. J. Zool.* **36**: 397-409.

Braby, M.F. and New, T.R. (1994) The Xenicas, *Geitoneura klugii* and *G. acantha* (Nymphalidae: Satyrinae). In *The Biology of Australian Butterflies* (Eds R.L. Kitching, R.E. Jones, N.E. Pierce). Monographs of Australian Lepidoptera, C.S.I.R.O. Publications, Melbourne. (in press)

Bradshaw, A.D. (1965) Evolutionary significance of phenotypic plasticity in plants. *Adv. Genetics* **13**: 115-155.

Brakefield, P.M. (1984) The ecological genetics of quantitative characters in *Maniola jurtina* and other butterflies. In *The Biology of Butterflies* (Eds R.I. Vane-Wright and P.R. Ackery), pp. 167-190. *Symposium of the Royal Entomological Society, Number 11*. Academic Press, London.

Brakefield, P.M. (1987) Tropical dry and wet season polyphenism in the butterfly *Melanitis leda* (Satyrinae): phenotypic plasticity and climatic correlations. *Biol. J. Linn. Soc.* **31**: 175-191.

Brakefield, P.M. and Larsen, T.B. (1984) The evolutionary significance of dry and wet season forms in some tropical butterflies. *Biol. J. Linn. Soc.* **22**: 1-12.

Brakefield, P.M. and Reitsma, N. (1991) Phenotypic plasticity, seasonal climate and the population biology of *Bicyclus* butterflies (Satyridae) in Malawi. *Ecol. Entomol.* **16**: 291-303.

Brown, K.S. and Benson, W.W. (1974) Adaptive polymorphism associated with multiple Mullerian mimicry in *Heliconius numata*. *Biotropica* **6**: 205-228.

Bucher, E.H. (1982) Chaco and Caatinga - South American arid savannas, woodlands and thickets. In *Ecology of Tropical Savannas* (Eds Huntley, B.J. and Walker, B.H.), pp. 48-79. Ecological Studies 42, Springer-Verlag, New York.

- Bulmer, M.G. (1983) Models for the evolution of protandry in insects. *Theor. Pop. Biol.* **23**: 319-327.
- Bureau of Meterology (1988a) *Climatic Averages of Australia*. Revised edition. Australian Government Publishing Service, Canberra.
- Bureau of Meterology (1988b) *Climatic Atlas of Australia*. Department of Administrative Services.
- Buskirk, R.E. and Buskirk, W.H. (1976) Changes in arthropod abundance in a highland Costa Rica forest. *Am. Midland Nat.* **95**: 288-298.
- Calow, P. and Sibly, R.M. (1983) Physiological trade-offs and the evolution of life cycles. *Sci. Prog.* **68**: 177-188.
- Campbell, A., Frazer, B.D., Gilbert, N., Gutierrez, A.P. and Mackauer, M. (1974) Temperature requirements of some aphids and their parasites. *J. Appl. Ecol.* **11**: 431-438.
- Campbell, I.M. (1962) Reproductive capacity on the genus *Choristoneura* Led. (Lepidoptera: Tortricidae). I. Quantitative inheritance and genes as controllers of rates. *Can. J. Genet. Cytol.* **4**: 272-288.
- Capinera, J.L., Barbosa, P. and Hagedorn, J.H. (1977) Yolk and yolk depletion of gypsy moth eggs: implications for population quality. *Ann. Entomol. Soc. Am.* **70**: 40-42.
- Carroll, A.L. and Quiring, D.T. (1993) Interactions between size and temperature influence fecundity and longevity of a tortricid moth, *Zeiraphera canadensis*. *Oecologia* **93**: 233-241.
- Caswell, H. (1989) Life-history strategies. In *Ecological Concepts* (Ed J.M. Cherrett), pp. 285-307. Blackwell Scientific Publications, Oxford.
- Charlesworth, B. (1971) Selection in density-regulated populations. *Ecology* **52**: 469-474.
- Charlesworth (1980) *Evolution in Age-structured Populations*. Cambridge University Press, Cambridge.
- Charlesworth, B. (1984) The evolutionary genetics of life-histories. In *Evolutionary Ecology* (Ed. B. Sharrocks), pp. 117-133. Blackwell Scientific Publications, Oxford.

- Cheng, H.H. (1972) Oviposition and longevity of the dark sided cutworm, *Euxoa messoria* (Lepidoptera, Noctuidae) in the laboratory. *Can. Ent.* **104**: 919-925.
- Chew, F.S. (1977) Coevolution of pierid butterflies and their cruciferous foodplants. II. The distribution of eggs on potential foodplants. *Evolution* **31**: 568-579.
- Chew, F.S. and Robbins, R.K. (1984) Egg-laying in butterflies. In *The Biology of Butterflies* (Eds R.I.Vane-Wright and P.R. Ackery), pp. 65-79. *Symposium of the Royal Entomological Society of London, Number 11*. Academic Press, London.
- Claret, J. and Carton, Y. (1980) Diapause in a tropical species, *Cothonaspis boulardi* (Parasitic Hymenoptera). *Oecologia* **45**: 32-34.
- Clench, H.K. (1977) A list of butterflies of Andros, Bahamas. *Ann. Carneg. Mus.* **46**: 173-193.
- Clutton-Brock, T.H. and Harvey, P.H. (1979) Comparison and adaptation. *Proc. R. Soc. Lond. B* **205**: 547-565.
- Cody, M.L. (1966) A general theory of clutch size. *Evolution* **20**: 174-184.
- Cole, L.C. (1954) The population consequences of life history phenomena. *Quart. Rev. Biol.* **29**: 103-137.
- Coley, P.D. (1980) Effects of leaf age and plant life history patterns on herbivory. *Nature* **284**: 541-546.
- Common, I.F.B. and Waterhouse, D.F. (1981) *Butterflies of Australia*. Angus and Robertson, Sydney.
- Corbet, A.S. and Pendlebury, H.M. (1978) *The Butterflies of the Malay Peninsula*. Third edition, revised by J.N. Elliot. Malay Nature Society, Kuala Lumpur.
- Courtney (1981) Coevolution of pierid butterflies and their cruciferous foodplants. III. *Anthocharis cardamines* survival, development and oviposition on different hostplants. *Oecologia* **51**: 91-96.
- Courtney (1982) Coevolution of pierid butterflies and their cruciferous foodplants. V. Habitat selection, community structure and speciation. *Oecologia* **54**: 101-107.

D'Abrera, B. (1971) *Butterflies of the Australian Region.* Lansdowne, Melbourne.

D'Abrera, B. (1985) *Butterflies of the Oriental region - Part II.* Hill House, Melbourne.

Danthanarayana, W. (1975) Factors determining variation in fecundity in the light brown apple moth, *Epiphyas postvittana* (Walker) (Tortricidae). *Aust. J. Zool.* **23**: 439-451.

David, W.A.L. and Gardiner, B.O.C. (1962) Oviposition and the hatching of the eggs of *Pieris brassicae* in a laboratory culture. *Bull. Ent. Res.* **53**: 91-109.

De Baar, M. (1981) Host records for *Hypocysta adiante adiante* (Hubn.) and *Xois arctous arctous* (Fab.) (Lepidoptera: Nymphalidae). *News Bull. Ent. Soc. Qld.* **9**: 9.

De Baar, M. (1983) New food plants, life history notes, and distribution records for some Australian Lepidoptera. *Aust. ent. Mag.* **9**: 97-98.

De Baar, M. (1988) Insects collected during a trip to Torres Strait, 27 March to 10 April 1987. *News Bull. Ent. Soc. Qld.* **15**: 107-117.

Denlinger, D. (1978) The developmental response of flesh flies (Diptera: Sarcophagidae) to tropical seasons. *Oecologia* **35**: 105-107.

Denlinger, D.L. (1980) Seasonal and annual variation of insect abundance in the Nairobi National Park, Kenya. *Biotropica* **12**: 100-106.

Denlinger, D. (1986) Dormancy in tropical insects. *Ann. Rev. Entomol.* **31**: 239-264.

Dennitson, C. (1978) An incorrect definition of fitness revisited. *Ann. Human Genet. Lond.* **42**: 77-85.

Dingle, H. (1978) *Evolution of Insect Migration and Diapause* (Ed). Springer-Verlag, Berlin.

Dingle, H. (1990) The evolution of life histories. In Population Biology (Eds K. Wohrmann and S.K. Jain), pp. 267-289. Springer-Verlag, Berlin.

Dixon, A.F.G., Kundu, R. and Kindlmann, P. (1993) Reproductive effort and maternal age in iteroparous insects using aphids as a model group. *Funct. Ecol.* **7**: 267-272.

- Dobzhansky, T.H. (1950) Evolution in the tropics. *Am. Sci.* 38: 209-221.
- Downes, J.A. (1973) Lepidoptera feeding at puddle-margins, dung and carrion. *J. Lep. Soc.* 27: 89-99.
- Duckworth, B.G. and Mclean, J. (1986) Notes on a collection of butterflies from the islands of the Great Barrier Reef, Queensland. *Aust. ent. Mag.* 13: 43-48.
- Dunlap-Piaka, H.L. (1979) Ovarian dynamics in *Heliconius* butterflies: correlations among daily oviposition rates, egg weights, and quantitative aspects of oogenesis. *J. Insect. Physiol.* 25: 741-749.
- Dunlap-Piaka, H.L., Boggs, C.L. and Gilbert, L.E. (1977) Ovarian dynamics in heliconiine butterflies: programmed senescence versus eternal youth. *Science* 197: 487-490.
- Dunn, K.L. and Dunn, L.E. (1991) *Review of Australian Butterflies: distribution, life history and taxonomy. Part I: Introduction, Papilionidae, Pieridae and regional adult temporal data*, pp 1-196. *Part IV: Family Nymphalidae*, pp 513-660. Privately published by the authors, Melbourne.
- Edwards, E.D. (1973) Delayed ovarian development and aestivation in adult females of *Heteronympha merope merope* (Lepidoptera: Satyrinae). *J. Aust. ent. Soc.* 12: 92-98.
- Edwards, E.D. (1980) The early stages of *Adaluma urumelia* Tindale and *Candalides geminus* Edwards and Kerr (Lepidoptera: Lycaenidae). *Aust. ent. Mag.* 7: 17-20.
- Ehrlich, A.H. and Ehrlich, P.R. (1978) Reproductive strategies in the butterflies. I. Mating frequency, plugging and egg number. *J. Kansas Entomol. Soc.* 51: 666-697.
- Ehrlich, P.R. and Gilbert, L.E. (1973) Population structure and dynamics of the tropical butterfly *Heliconius ethilla*. *Biotropica* 12: 69-82.
- Ehrlich, P.R. and Raven, P.H. (1965) Butterflies and plants: a study in coevolution. *Evolution* 18: 586-608.
- Elgar, M.A. (1990) Evolutionary compromise between a few large and many small eggs: comparative evidence in teleost fish. *Oikos* 59: 283-287.
- Emmel, T.C. (1970) The population biology of the neotropical satyrid butterfly,

- Euptychia hermes*. I. Interpopulation movement, general ecology, and population sizes in lowland Costa Rica (dry season 1966). *J. Res. Lepid.* 7: 153-165.
- Emmel, T.C. and Leck, C.F. (1970) Seasonal changes in organisation of tropical rain forest butterfly populations in Panama. *J. Res. Lepid.* 8: 133-152.
- Fagerström, T. and Wiklund, C. (1982) Why do males emerge before females? Protandry as a mating strategy in male and female butterflies. *Oecologia* 52: 164-166.
- Feeny, P.P. (1976) Plant apparency and chemical defence. *Rec. Adv. Phytochem.* 10: 1-40.
- Fenemore, P.G. (1977) Oviposition of the potato tuber moth, *Phthorimaea operculella* Zell. (Lepidoptera: Gelechiidae): fecundity in relation to mated state, age and pupal weight. *N. Z. J. Zool.* 4: 187-191.
- Fisher, R.A. (1930) *The Genetical Theory of Natural Selection*. Oxford University Press, London.
- Fisher, R.H. (1978) *Butterflies of South Australia*. Government Printer, South Australia.
- Fogden, M.P.L. (1972) The seasonality and population dynamics of equatorial forest birds in Sarawak. *Ibis* 114: 307-343.
- Forrest, T.G. (1987) Insect size tactics and developmental strategies. *Oecologia* 73: 178-184.
- Forsberg, J. and Wiklund, C. (1988) Protandry in the green-veined white butterfly, *Pieris napi* L. *Funct. Ecol.* 2: 81-88.
- Fox, K.J. (1973) Migrant Lepidoptera in New Zealand 1972-1973. *N. Z. Ent.* 5: 268-271.
- Fox, P.J. (1992a) *Melanitis leda bankia* (Fabricius) (Lepidoptera: Nymphalidae) in the Sydney district. *Victorian Ent.* 22: 10-11.
- Fox, P.J. (1992b) New distribution, food plant and life history notes for some southern Queensland butterflies. *Victorian Ent.* 22: 112-113.
- Fox, R.M., Lindsey, A.W., Clench, H.K. and Miller L.D. (1965) The butterflies of Liberia. *Mem. Amer. ent. Soc.* No. 19.

- Frith, C.B. and Frith, D.W. (1985) Seasonality of insect abundance in an Australian upland tropical rainforest. *Aust. J. Ecol.* **10**: 237-248.
- Fujiiie, A. (1980) Ecological studies of on the population of the pear leaf minor, *Bucculatrix pyrivorella* Kuroko (Lepidoptera: Lyonetiidae). III. Fecundity fluctuation from generation to generation within a year. *Appl. Ent. Zool.* **15**: 1-9.
- Gadgil, M. and Solbrig, O.T. (1972) The concept of *r*- and *K*-selection: evidence from wild flowers and some theoretical considerations. *Am. Nat.* **106**: 14-31.
- García-Barros, E. (1988) Delayed ovarian maturation in the butterfly *Hipparchia semele* as a possible response to summer drought. *Ecol. Entomol.* **13**: 391-398.
- Gilbert, L.E. (1972) Pollen feeding and reproductive biology of *Heliconius* butterflies. *Proc. Natl. Acad. Sci. USA* **69**: 1403-1407.
- Gilbert, L.E. and Singer, M.C. (1975) Butterfly ecology. *Ann. Rev. Ecol. Syst.* **6**: 365-397.
- Gilbert, N. (1984) Control of fecundity in *Pieris rapae* III. Synthesis. *J. Anim. Ecol.* **53**: 599-609.
- Gilbert, N.E. (1986) Control of fecundity in *Pieris rapae*. IV. Patterns of variation and their ecological consequences. *J. Anim. Ecol.* **55**: 317-329.
- Godfray, H.C.J., Partridge, L. and Harvey, P.H. (1991) Clutch size. *Ann. Rev. Ecol.* **22**: 409-429.
- Goodman, D. (1982) Optimal life histories, optimal notation and the value of reproductive value. *Am. Nat.* **119**: 803.
- Gould, S.J. and Lewontin, R.C. (1979) The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proc. R. Soc. Lond. B* **205**: 581-598.
- Greenfield, M.D. (1982) The question of paternal investment in Lepidoptera: male-contributed proteins in *Plodia interpunctella*. *Int. J. Invert. Reproduc.* **5**: 323-330.
- Greenslade, P.J.M. (1983) Adversity selection and the habitat templet. *Am. Nat.* **122**: 352-365.

- Grime, J.P. (1977) Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *Am. Nat.* **111**: 1169-1194.
- Grime, J.P. (1979) *Plant Strategies and Vegetation Processes*. Wiley, New York.
- Grime, J.P. (1988) The C-S-R model of primary plant strategies - origins, implications and tests. In *Plant Evolutionary Biology* (Eds Gottlieb, L.D. and S.K. Jain), pp. 371-393. Chapman and Hall, London.
- Gu, H. and Danthanarayana, W. (1990) The role of availability of food and water to the adult *Epiphyas postvittana*, the light brown apple moth, in its reproductive performance. *Entomol. exp. appl.* **54**: 101-108.
- Harvey, G.T. (1977) Mean weight and rearing performance of successive egg clusters of eastern spruce budworm (Lepidoptera: Tortricidae). *Can. Ent.* **109**: 487-496.
- Harvey, P.H. and Pagel, M.D. (1991) *The Comparative Method in Evolutionary Biology*. Oxford University Press, Oxford.
- Haukioja, E. and Neuvonen, S. (1985) The relationship between size and reproductive potential in male and female *Epirrita autumnata* (Lep., Geometridae). *Ecol. Entomol.* **10**: 267-270.
- Hawkeswood, T.J. (1990) Parasitism of *Melanitis leda bankia* (Fabricius) (Lepidoptera: Nymphalidae: Satyrinae) by a tachinid fly (Diptera: Tachinidae) in Queensland, Australia. *Bull. Annls. Soc. r. belge. Ent.* **126**: 55-62.
- Hayes, J.L. (1982) A study of the relationships of diapause phenomena and other life history characters in temperate butterflies. *Am. Nat.* **120**: 160-170.
- Henle, K. (1991) Life history patterns in lizards of the arid and semiarid zone of Australia. *Oecologia* **88**: 347-358.
- Hildrew, A.G. and Townsend, C.R. (1987) Organization in freshwater benthic communities. In *Organisations of Communities Past and Present* (Eds J.H.R. Gee and P.S. Giller), 347-372. Blackwell Scientific Publications, Oxford.
- Hill, C.J. (1988) The temporal component of butterfly species richness in a subtropical region of Australia. *Proc. Ecol. Soc. Aust.* **15**: 115-126.

- Hill, C.J. (1989) The effect of adult diet on the biology of butterflies 2. The common crow butterfly, *Euploea core corinna*. *Oecologia* **81**: 258-266.
- Hill, C.J. (1992) Temporal changes in abundance of two lycaenid butterflies (Lycaenidae) in relation to adult food resources. *J. Res. Lepid.* **46**: 173-181.
- Hill, C.J. (1994) Communities of butterflies. In *The Biology of Australian Butterflies* (Eds R.L. Kitching, R.E. Jones, N.E. Pierce). Monographs of Australian Lepidoptera, C.S.I.R.O. Publications, Melbourne. (in press)
- Hill, C.J., Gillison, A.N. and Jones, R.E. (1992) The spatial distribution of rain forest butterflies at three sites in north Queensland, Australia. *J. Trop. Ecol.* **8**: 37-46.
- Hill, C.J. and Pierce, N.E. (1989) The effect of adult diet on the biology of butterflies 1. The common imperial blue, *Jalmenus evagoras*. *Oecologia* **81**: 249-257.
- Holloway, B.A. (1962) *Melanitis leda* (Linn.) and other migrant butterflies in New Zealand during 1962. *Records of the Dominion Museum* **4**: 79-82.
- Honek, A. (1993) Intraspecific variation in body size and fecundity in insects: a general relationship. *Oikos* **66**: 483-492.
- Hough, J.A. and Pimental, D. (1978) Influence of host foliage on development, survival and fecundity of the gypsy moth. *Environ. Ent.* **7**: 97-102.
- Huntley, B.J. and Walker, B.H. (1982) *Ecology of Tropical Savannas* (Eds). Ecological Studies 42, Springer-Verlag, Berlin, Heidelberg, New York.
- Istock, C.A. (1983) The extent and consequences of heritable variation for fitness characters. In *Population Biology: Retrospect and Prospect* (Eds C.E. King and S. Dawson), pp. 61-96. Columbia University Press, New York.
- Ito, K. and Noor, N.D. (1993) Reproductive diapause and quiescence in a tropical rice bug, *Leptocoris oratorius*. *Entomol. exp. appl.* **68**: 71-77.
- Iwasa, Y., Odendaal, F.J., Murphy, D.D., Ehrlich, P.R. and Launer, A.E. (1983) Emergence patterns in male butterflies: a hypothesis and a test. *Theor. Pop. Biol.* **23**: 363-379.
- James, D.G. (1988) Aestivation and reproductive dormancy in adult *Heteronympha merope merope* (Fabricius) (Lepidoptera: Nymphalidae).

Aust. ent. Mag. **15**: 67-72.

- Janzen, D.H. (1973) Sweep samples of tropical foliage insects: effects of season, vegetation types, time of day and insularity. *Ecology* **54**: 687-708.
- Janzen, D.H. and Schoener, T.W. (1968) Differences in insect abundance and diversity between wetter and drier sites during a tropical dry season. *Ecology* **49**: 96-110.
- Jones, K.N., Odendaal, F.J. and Ehrlich, P.R. (1986) Evidence against the spermatophore as paternal investment in checkerspot butterflies (*Euphydryas*: Nymphalidae). *Am. Midl. Nat.* **116**: 1-6.
- Jones, R.E. (1987) Reproductive strategies for the seasonal tropics. *Insect Sci. Applic.* **8**: 515-521.
- Jones, R.E. (1992) Phenotypic variation in Australian *Eurema* species. *Aust. J. Zool.* **40**: 371-383.
- Jones, R.E., Hart, J.R. and Bull, G.D. (1982) Temperature, size and egg production in the cabbage white butterfly, *Pieris rapae* L. *Aust. J. Zool.* **30**: 223-232.
- Jones, R.E. and Rienks, J. (1987) Reproductive seasonality in the tropical genus *Eurema* (Lepidoptera: Pieridae). *Biotropica* **19**: 7-16.
- Jones, R.E., Rienks, J. and Wilson, L. (1985) Seasonality and environmentally induced polyphenism in *Eurema laeta lineata* (Lepidoptera: Pieridae). *J. Aust. ent. Soc.* **24**: 161-167.
- Jones, R.E., Rienks, J., Wilson, L., Lokkers, C. and Churchill, T. (1987) Temperature, development and survival in monophagous and polyphagous tropical pierid butterflies. *Aust. J. Zool.* **35**: 235-246.
- Karlsson, B. (1987) Variation in egg weight, oviposition rate and reproductive reserves with female age in a natural population of the speckled wood butterfly, *Pararge aegeria*. *Ecol. Entomol.* **12**: 473-476.
- Karlsson, B. and Wickman, P.-O. (1989) The cost of prolonged life: an experiment on a nymphalid butterfly. *Funct. Ecol.* **3**: 399-405.
- Karlsson, B. and Wickman, P.-O. (1990) Increase in reproductive effort as explained by body size and resource allocation in the speckled wood butterfly, *Pararge aegeria* (L.). *Funct. Ecol.* **4**: 609-617.

- Karlsson, B. and Wiklund, C. (1984) Egg weight variation and lack of correlation between egg weight and offspring fitness in the wall brown butterfly, *Lasiommata megera*. *Oikos* **43**: 376-385.
- Karlsson, B. and Wiklund, C. (1985) Egg weight variation in relation to egg mortality and starvation endurance of newly hatched larvae in some satyrid butterflies. *Ecol. Entomol.* **10**: 205-211.
- Kautsky, L. (1988) Life strategies of aquatic soft bottom macrophytes. *Oikos* **53**: 126-135.
- Kimura, K. and Tsubaki, Y. (1985) Egg weight variation associated with female age in *Pieris rapae crucivora* Boisduval (Lepidoptera: Pieridae). *Appl. Entomol. Zool.* **20**: 500-501.
- Kimura, M. (1956) A model of genetic system which leads to closer linkage under natural selection. *Evolution* **10**: 278-287.
- Kimura, T. and Masaki, S. (1977) Brachypteryism and seasonal adaptation in *Orgyia thyellina* Bultler (Lepidoptera: Lymantriidae). *Kontyû* **45**: 97-106.
- Kirk, W.D. (1982) Variation of forewing spot number in the *mineus*-group of *Mycalesis* (Lepidoptera, Satyridae) in S.E. Asia. *Malay Nat. J.* **35**: 229-236.
- Kitching, R.L. (1981) The geography of the Australian Papilionoidea. In *Ecological Biogeography of Australia* (Ed A. Keast), pp. 979-1004. Dr W. Junk Publishers, The Hague.
- Kitching, R.L. and Zalucki, M.P. (1983) A cautionary note on the use of oviposition records as larval food plant records. *Aust. ent. Mag.* **10**: 64-66.
- Krebs, J.R. and Davies, N.B. (1981) *An Introduction to Behavioural Ecology*. Blackwell Scientific Publications, London.
- Labine, P.A. (1968) The population biology of the butterfly *Euphydryas editha*. VIII. Oviposition and its relation to patterns of oviposition in other butterflies. *Evolution* **22**: 799-805.
- Lacey, C.J., Walker, J. and Noble, I.R. (1982) Fire in Australian savannas. In *Ecology of Tropical Savannas* (Eds Huntley, B.J. and Walker, B.H.), pp. 246-272. Ecological Studies 42, Springer-Verlag, Berlin, Heidelberg, New York.

- Lack, D. (1947) The significance of clutch size. *Ibis* **89**: 302-352.
- Lambkin, T.A. and Knight, A.I. (1983) Butterflies recorded from the Torres Strait and northern Peninsula regions in July 1983. *News Bull. Ent. Soc. Qld.* **11**: 98-103.
- Lambkin, T.A. and Knight, A.I. (1990) Butterflies recorded from Murray Island, Torres Strait. *Aust. ent. Mag.* **17**: 101-112.
- Lande, R. (1982) A quantitative genetic theory of life history evolution. *Ecology* **63**: 607-615.
- Lande, R. and Arnold, S.J. (1983) The measurement of selection of correlated characters. *Evolution* **37**: 1210-1226.
- Law, R. (1979) Ecological determinants in the evolution of life histories. In *Population Dynamics* (Eds R.M. Anderson, B.D. Turner and L.R. Taylor), pp. 81-103. Blackwell Scientific Publications, Oxford.
- Le Souef, J.C. (1971) To Iron Range in 1968. *Victorian Ent.* **1**(1): 8-10.
- Leather, S.R. (1984) The effect of adult feeding on the fecundity, weight loss, and survival of the pine beauty moth, *Panolis flammea* (D & S). *Oecologia* **65**: 70-74.
- Leather, S.R. (1988) Size, reproductive potential and fecundity in insects: things aren't as simple as they seem. *Oikos* **51**: 386-389.
- Leather, S.R. and Burnand, A.C. (1987) Factors affecting life-history parameters of the pine beauty moth, *Panolis flammea* (D&S): the hidden costs of reproduction. *Funct. Ecol.* **1**: 331-338.
- Lederhouse, R.C., Ayres, M.P. and Scriber, J.M. (1989) Evaluation of spermatophore counts in studying mating systems of Lepidoptera. *J. Lep. Soc.* **43**: 93-101.
- Lederhouse, R.C., Ayres, M.P. and Scriber, J.M. (1990) Adult nutrition affects male virility in *Papilio glaucus* L. *Funct. Ecol.* **4**: 743-751.
- Leigh, E.G. and Windsor, D.M. (1982) Forest production and regulation of primary consumers on Barro Colorado Island. In *The Ecology of a Tropical Forest* (Eds E.G. Leigh, A.S. Rand and D.M. Windsor), pp. 111-122. Smithsonian Institution Press, Washington.

- Leslie, P.H. (1948) Some further notes on the use of matrices in population mathematics. *Biometrika* **35**: 213-245.
- Levings, S.C. and Windsor, D.M. (1982) Seasonal and annual variation in litter arthropod populations. In *The Ecology of a Tropical Forest* (Eds E.G. Leigh, Jr., A.S. Rand, and D.M. Windsor), pp. 355-387. Smithsonian Institution Press, Washington, DC.
- Lewontin, R.C. (1965) Selection for colonising ability. In *The Genetics of Colonising Species* (Eds H.G. Baker and G.L. Stebbins), pp. 79-94. Academic Press, New York.
- Lieberman, D. and Lieberman, M. (1984) The causes and consequences of synchronous flushing in a dry tropical forest. *Biotropica* **16**: 193-201.
- Lloyd, D.G. (1987) Selection of offspring size at independence and other size-versus-number strategies. *Am. Nat.* **129**: 800-817.
- Lotka, A.J. (1913) A natural population norm. II. *J. Wash. Acad. Sci.* **3**: 289-293.
- Lowman, M.D. (1982) Seasonal variation in insect abundance among three Australian rainforests, with particular reference to phytophagous types. *Aust. J. Ecol.* **7**: 353-361.
- Lucas, P.W. and Pereira, B. (1990) Estimation of the fracture toughness of leaves. *Funct. Ecol.* **4**: 819-822.
- MacArthur, R.H. (1962) Some generalised theorems of natural selection. *Proc. Nat. Acad. Sci., U.S.* **48**: 1893-1897.
- MacArthur, R.H. (1972) *Geographical Ecology. Patterns in the Distribution of Species*. Harper and Row, New York.
- MacArthur, R.H. and Wilson, E.O. (1967) *Theory of Island Biogeography*. Princeton University Press, Princeton.
- Manski, M.J. (1960) Food plants of some Queensland Lepidoptera. *Qd. Nat.* **16**: 68-73.
- Manskic, R.C. and Manskie, N. (1972) Collecting in north-eastern Australia. *Victorian Ent.* **2(5)**: 17-23.
- Marshall, L.D. (1988) Intraspecific variation in reproductive effort by female *Parapediasia teterrella* (Lepidoptera: Pyralidae) and its relation to body

- size. *Can. J. Zool.* **68**: 44-48.
- Masaki, S. (1980) Summer diapause. *Ann. Rev. Entomol.* **25**: 1-25.
- Masetti, N.diM. and Scali, V. (1972) Ecological adjustments of the reproductive biology in *Maniola jurtina* from Tuscany. *Atti della Accademia Nazionale dei Lincei memorie classe di Scienze Fisiche Matematiche e Naturali, ser. 8a* **53**: 460-470.
- Masuzawa, T., Suwa, H. and Nakasuji, F. (1983) Differences of oviposition preferences and survival rate of two skipper butterflies *Parnara guttatta* and *Pelopidas mathias* (Lepidoptera: Hesperiidae) on rice plant and cogon grass. *N. Entomol.* **32**: 47-56.
- Mattson, W.J. (1980) Herbivory in relation to plant nitrogen content. *Ann. Rev. Ecol. Syst.* **11**: 119-161.
- McCubbin, C. (1971) *Australian Butterflies*. Nelson, Melbourne.
- McLachlan, A.J. (1986) Survival of the smallest: advantages and costs of small size in flying animals. *Ecol. Entomol.* **11**: 237-240.
- McLeod, L. (1968) Controlled environment experiments with *Precis octavia* Cram. (Nymphalidae). *J. Res. Lep.* **7**: 1-18.
- McLeod, L. (1984) Seasonal polyphenism in African *Precis* butterflies. In *The Biology of Butterflies* (Eds R.I. Vane-Wright and P.R. Ackery), pp. 313-315. *Symposium of the Royal Entomological Society, Number 11*. Academic Press, London.
- Medawar, P.B. (1946) Old age and natural death. *Modern Quarterly* **1**: 30-56.
- Medawar, P.B. (1952) *An Unsolved Problem of Biology*. Lewis, London.
- Menaut, J.C. and Cesar, J. (1982) The structure and dynamics of a west African savanna. In *Ecology of Tropical Savannas* (Eds Huntley, B.J. and Walker, B.H.), pp. 80-100. *Ecological Studies 42*, Springer-Verlag, Berlin, Heidelberg, New York.
- Miller, L.D. (1968) The higher classification, phylogeny and zoogeography of the Satyridae (Lepidoptera). *Mem. Am. Ent. Soc.* **24**: 1-174.
- Miller, T.A., Cooper, W.J. and Highfill, J.W. (1983) Egg production in female *Callosamia promethea* (Lepidoptera: Saturniidae) as a function of pupal

- size and adult longevity. *Ann. Ent. Soc. Am.* **76**: 668-670.
- Minson, D.J. (1971) Influence of lignin and silicon on a summative system for assessing the organic matter digestibility of *Panicum*. *Aust. J. Agric. Res.* **22**: 589-598.
- Monteith, G.B. (1972) A list of butterfly records from the Iron Range area of Cape York Peninsula. *News Bull. Ent. Soc. Qld.* **85**: 9-14.
- Monteith, G.B. (1982) Dry season aggregations of insects in Australian monsoon forests. *Mem. Qd. Mus.* **20**: 533-540.
- Moore, G.J. (1985) *The behaviour and ecology of two tropical satyrine butterflies*. Unpublished honours thesis, James Cook University of North Queensland, Townsville.
- Moore, G.J. (1986) Host plant discrimination in tropical satyrine butterflies. *Oecologia* **70**: 592-595.
- Moore, R.A. and Singer, M.C. (1987) Effects of maternal age and adult diet on egg weight in the butterfly *Euphydryas editha*. *Ecol. Entomol.* **12**: 401-408.
- Moulds, M.S. and d'Apice, J.W.C. (1982) Butterflies of the Upper Jardine River, Cape York Peninsula. *Aust. ent. Mag.* **9**: 21-26.
- Murphy, D.D., Launer, A.E. and Ehrlich, P.R. (1983) The role of adult feeding in egg production and population dynamics of the checkerspot butterfly *Euphydryas editha*. *Oecologia* **56**: 257-263.
- Murphy, G.I. (1968) Pattern in life history and the environment. *Am. Nat.* **102**: 390-404.
- Murray, B.G. (1990) Population dynamics, genetic change, and the measurement of fitness. *Oikos* **59**: 189-199.
- Nakasuji, F. (1978) Life history of the satyrid butterfly *Mycalesis gotama fluginia* and its survival rates in paddies. *Jap. J. appl. Ent. Zool.* **22**: 152-161.
- Nakasuji, F. (1982) Seasonal changes in native host plants of a migrant skipper, *Parnara guttata* Bremner et Grey (Lepidoptera: Hesperiidae). *Appl. Ent. Zool.* **17**: 146-148.
- Nakasuji, F. (1987) Egg size of skippers (Lepidoptera: Hesperiidae) in relation

to their host specificity and to leaf toughness of host plants. *Ecol. Res.* **2**: 175-183.

Nakasui, F. and Kimura, M. (1984) Seasonal polymorphism of egg size in a migrant skipper, *Parnara guttata guttata* (Lepidoptera, Hesperiidae). *Kontyû* **52**: 253-259.

New, T.R. (1984) *Insect Conservation - An Australian Perspective*. Dr. W. Junk Publishers, Dordrecht.

Njoku, E. (1963) Seasonal periodicity in the growth and development of some forest trees in Nigeria. *J. Ecol.* **51**: 617-624.

Norris, M.J. (1935) A feeding experiment on the adults of *Pieris rapae*. *Entomol.* **68**: 125-127.

Nylin, S. (1988) Host plant specialisation and seasonality in a polyphagous butterfly, *Polygonia c-album* (Nymphalidae). *Oikos* **53**: 381-386.

Oberhauser, K.S. (1989) Effects of spermatophores on male and female monarch butterfly reproductive success. *Behav. Ecol. Sociobiol.* **25**: 237-246.

Odendaal, F.J. (1990) The dry season influences reproductive parameters in female butterflies. *Biotropica* **22**: 100-102.

Oive, J. (1978) New distribution records for Queensland butterflies. *Aust. ent. Mag.* **5**: 8.

Owen, D.F. (1971) *Tropical Butterflies: the Ecology and Behaviour of Butterflies in the Tropics with Special Reference to African Species*. Clarendon Press, Oxford.

Owen, D.F. and Chanter, D.O. (1970) Species diversity and seasonal abundance in tropical Ichneumonidae. *Oikos* **21**: 142-144.

Owen, D.F., Owen, J. and Chanter, D.O. (1970) Seasonal changes in relative abundance and estimates of species diversity in a family of tropical butterflies. *Oikos* **21**: 200-205.

Pagel, M.D. and Harvey, P.H. (1988) Recent developments in the analysis of comparative data. *Quart. Rev. Biol.* **63**: 413-440.

Parker, G.A. and Begon, M. (1986) Optimal egg size and clutch size: effects of environment and maternal phenotype. *Am. Nat.* **128**: 573-592.

- Parker, G.A. and Courtney, S.P. (1983) Seasonal incidence: adaptive variation in the timing of life history stages. *J. Theor. Biol.* **105**: 147-155.
- Parry, G.D. (1981) The meanings of r- and K-selection. *Oecologia* **48**: 260-264.
- Partridge, L. and Harvey, P.H. (1985) Costs of reproduction. *Nature, London*. **316**: 20.
- Partridge, L. and Harvey, P.H. (1988) The ecological context of life history evolution. *Science* **24**: 1449-1455.
- Pearse, F.K. (1978) *Geographic variation and natural selection in the common brown butterfly Heteronympha merope*. Ph.D. thesis, La Trobe University, Melbourne.
- Phipps, J. (1968) The ecological distribution and life cycles of some tropical African grasshoppers (Acridoidea). *Bull. Entomol. Soc. Nigeria* **1**: 71-97.
- Pianka, E.R. (1970) On r- and K-selection. *Am. Nat.* **104**: 592-597.
- Pianka, E.R. (1972) r and K selection or b and d selection? *Am. Nat.* **106**: 581-588.
- Pivnick, K.A. and McNeil, J.N. (1985) Effects of nectar concentration on butterfly feeding: measured feeding rates for *Thymelicus lineola* (Lepidoptera: Hesperiidae) and a general feeding model for adult Lepidoptera. *Oecologia* **66**: 226-237.
- Pivnick, K.A. and McNeil, J.N. (1987) Puddling in butterflies: sodium affects reproductive success in *Thymelicus lineola*. *Phys. Ent.* **12**: 461-472.
- Pollard, E. (1977) A method for assessing changes in the abundance of butterflies. *Biol. Conserv.* **12**: 115-134.
- Pollard, E. (1982) Monitoring butterfly abundance in relation to the management of a nature reserve. *Biol. Conserv.* **24**: 317-328.
- Primack, R.B. and Kang, H. (1989) Measuring fitness and natural selection in wild plant populations. *Ann. Rev. Ecol. Syst.* **20**: 367-396.
- Proshold, F.I., Karpenko, C.P. and Graham, C.K. (1982) Egg production and oviposition in the tobacco budworm: effect of age and mating. *Ann. Ent. Soc. Am.* **75**: 51-55.
- Quick, W.N.B. (1974) Some abnormal insect records for the summers of 1972-3, 1973-4. *Victorian Ent.* **4**: 66-71.

- Rae, D.J. and De'Ath, G. (1991) Influence of constant temperature on development, survival and fecundity of sugarcane mealybug, *Saccharicoccus sacchari* (Cockerell) (Hemiptera: Pseudococcidae). *Aust. J. Zool.* **39**: 105-122.
- Reiss, M.J. (1989) *The Allometry of Growth and Reproduction*. Cambridge University Press, Cambridge.
- Reznick, D.N. (1985) Cost of reproduction: an evaluation of the empirical evidence. *Oikos* **44**: 257-267.
- Richards, L.J. and Myers, J.H. (1980) Maternal influences on size and emergence time of the cinnabar moth. *Can. J. Zool.* **58**: 1452-1457.
- Rienks, J.H. (1985) Phenotypic response to photoperiod and temperature in a tropical pierid butterfly. *Aust. J. Zool.* **33**: 837-847.
- Roff, D.A. (1981) On being the right size. *Am. Nat.* **118**: 405-422.
- Rossiter, M.C. (1991) Maternal effects generate variation in life history: consequences of egg weight plasticity in the gypsy moth. *Funct. Ecol.* **5**: 386-393.
- Roughgarden, J. (1971) Density-dependent natural selection. *Ecology* **52**: 453-468.
- Rutowski, R.L., Gilchrist, G.W. and Terkanian, B. (1987) Female butterflies mated with recently mated males show reduced reproductive output. *Behav. Ecol. Sociobiol.* **20**: 319-322.
- Samson, P.R. (1991) The life history of *Everes lacturnus australis* Couchman (Lepidoptera: Lycaenidae). *Aust. ent. Mag.* **18**: 71-74.
- Sands, D.P.A. and Brancatini, V. (1991) A portable penetrometer for measuring leaf toughness in insect herbivory studies. *Proc. Ent. Soc. Washington* **93**: 786-788.
- SAS Institute, Inc. (1988) *SAS User's Guide: Statistics*. SAS, Cary, North Carolina.
- Scali, V. (1971) Imaginal diapause and gonadal maturation of *Maniola jurtina* (Lepidoptera: Satyridae) from Tuscany. *J. Anim. Ecol.* **40**: 467-472.

Schaffer, W.M. (1974) Optimal reproductive effort in fluctuating environments. *Am. Nat.* **108**: 783-790.

Scheermeyer, E. (1987) *Seasonality or opportunism in reproduction of Australian danaine butterflies: Euploea core, E. tulliolus and Tirumala hamata (Lepidoptera)*. Ph.D. thesis, Griffith University, Brisbane.

Scheermeyer, E. (1993) Overwintering of three Australian danaines: *Tirumala hamata hamata*, *Euploea tulliolus tulliolus* and, *E. core corrina*. In *Biology and Conservation of the Monarch Butterfly* (Eds S.B. Malcolm and M.P. Zalucki), pp 345-353. No. 38 Science Series, Natural History Museum of Los Angeles County.

Scriber, J.M. and Slansky, F. (1981) The nutritional ecology of immature insects. *Ann. Rev. Entomol.* **26**: 183-211.

Seymour, J.E. (1991) *Aspects of the ecology of Microgaster demolitor, a larval parasitoid of Helicoverpa punctigera and Helicoverpa armigera in Australia*. Ph.D. thesis, James Cook University of North Queensland, Townsville.

Shapiro, A.M. (1976) Seasonal polyphenism. *Evol. Biol.* **9**: 259-333.

Shima, H. (1988) Phyletic relationships of the genus *Ypthima* Hübner (Lepidoptera: Satyridae). In recent advances in Rhopalocerology: commemorative publication in honour of Takashi Schirozu. *Special Bull. Lep. Soc. Japan* **6**: 69-81.

Sibly, R.M. and Calow, P. (1985) Classification of habitats by selection pressures: a synthesis of life-cycle and *r/K* theory. In *Behavioural Ecology. Ecological Consequences of Adaptive Behaviour* (Eds R.M. Sibly and R.H. Smith), pp. 75-90. Blackwell Scientific Publications, Oxford.

Sibly, R.M. and Calow, P. (1986) *Physiological Ecology of Animals. An Evolutionary Approach*. Blackwell Scientific Publications, Oxford.

Sibly, R., Calow, P. and Nichols, N. (1985) Are patterns of growth adaptive? *J. Theor. Biol.* **112**: 553-574.

Singer, M.C. (1982) Sexual selection for small size in male butterflies. *Am. Nat.* **119**: 440-443.

Singer, M.C. (1984) Butterfly-hostplant relationships: host quality, adult choice and larval success. In *The Biology of Butterflies* (Eds R.I. Vane-Wright

- and P.R. Ackery), pp. 81-88. *Symposium of the Royal Entomological Society of London. Number 11.* Academic Press, London.
- Smith, A.M. (1986) Fecundity and survival of the common army worm, *Mythimna convecta*: effects of temperature and larval nutrition. *Entomol. Exp. Appl.* **42**: 31-37.
- Smith, C.C. and Fretwell, S.D. (1974) The optimal balance between size and number of offspring. *Am. Nat.* **108**: 499-506.
- Smith, R.H., Sibly, R.M. and Moller, H. (1987) Control of size and fecundity in *Pieris rapae rapae*: towards a theory of butterfly life cycles. *J. Anim. Ecol.* **56**: 341-350.
- Smithers, C.N. (1975) *Melanitis leda bankia* (Fabricius) (Lepidoptera: Nymphalidae) in Sydney and on Norfolk Island. *Aust. ent. Mag.* **2**: 110-111.
- Smithers, C.N. (1983) Migration records in Australia. 3. Danainae and Acraeinae (Lepidoptera: Nymphalidae) in Australia. *Aust. ent. Mag.* **10**: 21-27.
- Southwood, T.R.E. (1977) Habitat, the templet for ecological strategies? *Anim. Ecol.* **46**: 337-365.
- Southwood, T.R.E. (1988) Tactics, strategies and templets. *Oikos* **52**: 3-18.
- Specht, R.L. (1981) Major vegetation formations in Australia. In *Ecological Biogeography of Australia* (Ed A. Keast), pp. 163-298. Dr W. Junk Publishers, The Hague.
- Spitzer, K. (1983) Seasonality of the butterfly fauna in southeastern Vietnam (Papilionoidea). *J. Res. Lepid.* **22**: 126-130.
- Stearns, S.C. (1976) Life history tactics: a review of the ideas. *Quart. Rev. Biol.* **51**: 3-43.
- Stearns, S.C. (1977) The evolution of life history traits. A critique of the theory and a review of the data. *Ann. Rev. Ecol. Syst.* **8**: 145-171.
- Stearns, S.C. (1984) The effects of size and phylogeny on patterns of covariation in the life history traits of lizards and snakes. *Am. Nat.* **123**: 56-72.
- Stearns, S.C. (1989) The evolutionary significance of phenotypic plasticity. *Bioscience* **39**: 436-445.

Stearns, S.C. (1992) *The Evolution of Life Histories*. Oxford University Press, Oxford.

Stearns, S.C. and Koella, J.C. (1986) The evolution of phenotypic plasticity in life-history traits: predictions of reaction norms for age and size at maturity. *Evolution* **40**: 893-913.

Stern, V.M. and Smith, R.F. (1960) Factors affecting egg production and oviposition in populations of *Colias philodice eurytheme* Boisduval. *Hilgardia* **29**: 411-454.

Stewart, D.J. (1981) Travelling in "wait-a-while" country. *Victorian Ent.* **11**: 41-46.

Suzuki, Y. (1978) Adult longevity and reproductive potential of the small cabbage white, *Pieris rapae crucivora* Boisduval (Lepidoptera: Pieridae). *Appl. Ent. Zool.* **13**: 312-313.

Svärd, L. and Wiklund, C. (1988) Fecundity, egg weight and longevity in relation to multiple matings in females of the monarch butterfly. *Behav. Ecol. Sociobiol.* **23**: 39-43.

Svärd, L. and Wiklund, C. (1989) Mass and production rate of ejaculates in relation to monandry/polyandry in butterflies. *Behav. Ecol. Sociobiol.* **24**: 395-402.

Svärd, L. and Wiklund, C. (1991) The effects of ejaculate mass on female reproductive output in the European swallowtail butterfly, *Papilio machaon* (L.) (Lepidoptera: Papilionidae). *J. Insect Behav.* **4**: 33-42.

Takahashi, M. (1980) A comparison of the egg diameters of *Mycalesis gotama* Moore and *M. madjicosa* Butler (Lepidoptera: Satyridae). *Trans. Lep. Soc. Japan* **30**: 237-238.

Tani, S. (1984) Summer diapause of *Neope niphonica* Butler (Lepidoptera: Satyridae) in the Boso Peninsula and its ecological significance. *Kontyû, Tokyo* **52**: 119-127.

Tauber, M.J., Tauber, C. and Masaki, S. (1986) *Seasonal Adaptations of Insects*. Oxford University Press, New York, Oxford.

Taylor, D.R., Aarssen, L.W. and Loehle, C. (1990) On the relationship between r/K selection and environmental carrying capacity: a new habitat template for plant life history strategies. *Oikos* **58**: 239-250.

- Taylor, J.A. and Tulloch, D. (1985) Rainfall in the wet-dry tropics: extreme events at Darwin and similarities between years during the period 1870-1983. *Aust. J. Ecol.* **10**: 281-295.
- Telfer, W.H. and Rutberg, L.D. (1960) The effects of blood protein depletion on the growth of the oocytes in cecropia moth. *Biol. Bull.* **118**: 352-366.
- Thomas, S. and Corden, M. (1977) *Metric Tables of Composition of Australian Foods.* Commonwealth Department of Health, Australian Government Publishing Service, Canberra.
- Thornhill, R. and Alcock, J. (1983) *The Evolution of Insect Mating Systems.* Harvard University Press, Cambridge.
- Tilman, D. (1987) On the meaning of competition and the mechanisms of competitive superiority. *Funct. Ecol.* **1**: 304-315.
- Tracey, J.G. (1982) *The Vegetation of the Humid Tropical Region of north Queensland.* CSIRO, Melbourne.
- Turley, G. (1974) Southern limits? *Victorian Ent.* **4**: 47-48.
- Turner, A.J. (1925) The Lepidoptera of Yeppoon. *Aust. Zool.* **4**: 17-23.
- UNESCO,UNED,FAO (1978) *Tropical Forest Ecosystems: A State of Knowledge Report.* United Nations Press, Paris.
- Valentine, P.S. (1988) *Australian Tropical Butterflies.* Tropic Australia Graphics, Paluma via Townsville.
- Valentine, P.S. and Johnson, S.J. (1989) Observations on the life history of *Graphium aristaeus parmatum* (Gray) (Lepidoptera: Papilionidae). *Aust. ent. Mag.* **16**: 17- 20.
- Via, S. (1993) Adaptive phenotypic plasticity: target or by-product of selection in a variable environment? *Am. Nat.* **142**: 352-365.
- Vickery, M.L. (1984) *Ecology of Tropical Plants.* John Wiley & Sons, New York.
- Wagner, T.L., Wu, H.-I., Sharpe, P.J.H., Schoolfield, R.M. and Coulson, R.N. (1984) Modeling insect development rates: a literature review and application of a biophysical model. *Ann. Ent. Soc. Am.* **77**: 208-221.

- Walker, J. and Gillison, A.N. (1982) Australian Savannas. In *Ecology of Tropical Savannas* (Eds B.J. Huntley and B.H. Walker), pp. 5-24. Ecological Studies 42, Springer-Verlag, Berlin, Heidelberg, New York.
- Watanabe, M. (1980) Multiple matings increase the fecundity of the yellow swallowtail butterfly, *Papilio xuthus* L., in summer generations. *J. Ins. Behav.* **1**: 17-30.
- Waterhouse, G.A. (1923) Notes on the life histories of Australian butterflies. *Proc. Linn. Soc. N.S.W.* **48**: i-xxiv.
- Waterhouse, G.A. (1928) A second monograph of the genus *Tisiphone*, Hubner. *Aust. Zool.* **5**: 217-240.
- Waterhouse, G.A. (1932) *What Butterfly is That?* Angus and Robertson, Sydney.
- Waterhouse, G.A. and Lyell, G. (1914) *The Butterflies of Australia*. Angus and Robertson, Sydney.
- Watt, W.B., Hoch, P.C. and Mills, S.G. (1974) Nectar resource use by *Colias* butterflies. Chemical and visual aspects. *Oecologia* **14**: 353-374.
- Welden, C.W. and Slauson, W.L. (1986) The intensity of competition verses its importance: an overlooked distinction and some implications. *Quart. Rev. Biol.* **61**: 23-44.
- White, J.W. (1975) Composition of honey. In *Honey, a Comprehensive Survey* (Ed E. Crane), pp. 157-206. Heinemann, London.
- Whittaker, R.H. (1975) The design and stability of plant communities. In *Unifying Concepts in Ecology* (Eds W.H. van Dobben and R.H. Low-Connell), pp. 169-181. Junk, The Hague.
- Whittaker, R.H. and Goodman, D. (1979) Classifying species according to their demographic strategy. I. Population fluctuations and environmental heterogeneity. *Am. Nat.* **113**: 185-200.
- Wickman, P.-O. and Karlsson, B. (1987) Changes in egg colour, egg weight and oviposition rate with the number of eggs laid by wild females of the small heath butterfly, *Coenonympha pamphilus*. *Ecol. Entomol.* **12**: 109-114.
- Wickman, P.-O., Wiklund, C. and Karlsson, B. (1990) Comparative phenology of four satyrine butterflies inhabiting dry grasslands in Sweden. *Holarct. Ecol.* **13**: 238-346.

- Wiklund, C. (1984) Egg laying patterns in butterflies in relation to their phenology and the visual abundance of their hostplants. *Oecologia* **63**: 23-29.
- Wiklund, C. and Fagerström, T (1977) Why do males emerge before females? A hypothesis to explain the incidence of protandry in butterflies. *Oecologia* **31**: 153-158.
- Wiklund, C. and Forsberg, J. (1991) Sexual size dimorphism in relation to female polygamy and protandry in butterflies: a comparative study of Swedish Pieridae and Satyridae. *Oikos* **60**: 373-381.
- Wiklund, C., Kaitala, A., Lindfors, V. and Abenius, J. (1993) Polyandry and its effect on female reproduction in the green-veined white butterfly (*Pieris napi* L.). *Behav. Ecol. Sociobiol.* **33**: 25-33.
- Wiklund, C. and Karlsson, B. (1984) Egg size variation in satyrid butterflies: adaptive vs historical, "Bauplan", and mechanistic explanations. *Oikos* **43**: 391-400.
- Wiklund, C. and Karlsson, B. (1988) Sexual size dimorphism in relation to fecundity in some Swedish satyrid butterflies. *Am. Nat.* **131**: 132-138.
- Wiklund, C. and Karlsson, B. (1990) Sexual size dimorphism and fecundity in satyrid butterflies: a reply to Harvey's comment. *Am. Nat.* **136**: 268-269.
- Wiklund, C., Karlsson, B. and Forsberg, J. (1987) Adaptive versus constraint explanations for egg-to-body size relationships in two butterfly families. *Am. Nat.* **130**: 828-838.
- Wiklund, C., Nylin, S. and Forsberg, J. (1991) Sex-related variation in growth rate as a result of selection for large size and protandry in a bivoltine butterfly, *Pieris napi*. *Oikos* **60**: 241-250.
- Wiklund, C. and Persson, A. (1983) Fecundity, and the relation of egg weight variation to offspring fitness in the speckled wood butterfly *Pararge aegeria*, or why don't butterfly females lay more eggs? *Oikos* **40**: 53-63.
- Wiklund, C. and Solbreck, C. (1982) Adaptive versus incidental explanations for the occurrence of protandry in a butterfly, *Leptidea sinapis* L. *Evolution* **36**: 56-62.
- Wiklund, C., Wickman, P.-O. and Nylin, S. (1992) A sex difference in the propensity to enter direct/diapause development: a result of selection for

- protandry. *Evolution* **46**: 519-528.
- Wilbur, H.M., Tinkle, D.W. and Collins, J.P. (1974) Environmental certainty, trophic level, and resource availability in life history evolution. *Am. Nat.* **108**: 805-817.
- Williams, G.C. (1966) *Adaptation and Natural Selection*. Princeton University Press, Princeton.
- Wilson, P. (1987) *Australia's Butterflies*. Kangaroo Press, Kenthurst.
- Windig, W.W. (1993) Seasonal polyphenism in *Bicyclus safitza*, a continuous reaction norm. *Netherlands J. Zool.* **42**: 583-594.
- Wolda, H. (1978a) Fluctuations in abundance of tropical insects. *Am. Nat.* **112**: 1017-1045.
- Wolda, H. (1978b) Seasonal fluctuations in rainfall, food and abundance of tropical insects. *J. Anim. Ecol.* **47**: 369-381.
- Wolda, H. (1980) Seasonality of tropical insects. I. Leafhoppers (Homoptera) in Las Cumbres, Panama. *J. Anim. Ecol.* **49**: 277-290.
- Wolda, H. (1988) Insect seasonality: why? *Ann. Rev. Ecol. Syst.* **19**: 1-18.
- Wolda, H. and Fisk, F.W. (1980) Seasonality of tropical insects. II. Blattaria in Panama. *J. Anim. Ecol.* **50**: 827-838.
- Wood, G.A. (1984) The life history of *Elymnias agondas australiana* Fruhstorfer (Lepidoptera: Nymphalidae). *Aust. ent. Mag.* **11**: 41-42.
- Wood, G.A. (1987) The butterflies of Murray Island, Torres Strait, Queensland. *Aust. ent. Mag.* **14**: 39-42.
- Wood, G.A. (1988) The life history of *Hypocysta angustata angustata* Waterhouse & Lyell and *Hypocysta irius* (Fabricius) (Lepidoptera: Nymphalidae: Satyrinae). *Aust. ent. Mag.* **14**: 83-86.
- Worner, S. P. (1992) Performance of phenological models under variable temperature regimes: consequences of the Kaufmann or rate summation effect. *Environ. Entomol.* **21**: 689-699.
- Young, A.M. (1972) Breeding success and survivorship in some tropical butterflies. *Oikos* **23**: 318-326.

Young, A.M. (1979) The evolution of eyespots in tropical butterflies in response to feeding on rotting fruit: an hypothesis. *New York ent. Soc.* **87**: 66-67.

Young, A.M. (1980a) The interaction of predators and "eyespots butterflies" feeding on rotting fruits and soupy fungi in tropical forests. *Entomologist's Rec.* **90**: 63-69.

Young, A.M. (1980b) Seasonal adult emergence of cicadas (Homoptera: Cicadidae) in northwestern Costa Rica. *Contr. Biol. Geol. Milwaukee Publ. Mus.* **40**: 1-29.

Young, A.M. (1981) Notes on the seasonal distribution of *Anaea* butterflies (Nymphalidae) in tropical dry forests. *Acta Oecol.* **2**: 17-30.

Young, A.M. (1982) *Population Biology of Tropical Insects*. Plenum Press, New York.

Appendix 1. Distribution of satyrine butterflies across sampling sites in northeastern Queensland (see Figure 2.3). Sites are arranged into four broad geographic zones according to distance from coast and elevation: moist lowland areas 0-20 km from coast below 200 m (Zone 1); near coastal areas 20-50 km from coast below 500 m (Zone 2); dry inland areas 50-300 km from coast of variable altitude (Zone 3); and moist upland areas above 500 m (Zone 4). Observations of taxa recorded at particular localities of interest are also listed (as pers. obs.), together with literature records where comprehensive lists are available.

near Mt. Lamond (pers. obs.)	+		+		+		+	
Claudie River (Waterhouse & Lyell 1914)		+	+	+		+	+	+
Iron Range (Le Souef 1971)		+	+	+	+	+	+	+
Iron Range (Daniels 1975)	+	+	+	+	+	+	+	+
71			+	+	+		+	+
72		+	+	+	+		+	+
73			+	+	+			
74			+	+	+			+
75			+	+	+			+
<hr/>								
HOPE VALE- TOWNSVILLE								
78	+	+	+	+	+		+	+
79								+
80		+	+	+			+	
81		+	+	+			+	+
28 km S Cape Tribulation (pers. obs.)		+		+				+
82	+	+	+	+				+
83	+	+	+	+	+		+	
84		+		+				
100		+	+	+			+	+

101	+	+	+	+	+	+	+
102		+					
114	+		+	+	+		
115	+	+	+	+	+		+
116	+	+		+			+
117	+	+	+	+			+
168	+	+	+	+	+	+	+
169	+	+	+	+	+		+
170	+	+	+	+	+		
152		+	+	+			+
153		+	+	+	+		
154	+		+	+	+		
156		+		+			+
Ollera Creek, 10 km NW Rollingstone (pers. obs.)	+	+	+	+	+	+	+

TOWNSVILLE-
ST. LAWRENCE

Mt. Elliot Nat. Park (pers. obs.)	+	+	+	+	+	+	+
Cape Cleveland (pers. obs.)	+	+	+	+		+	+
118	+	+	+				+
119	+	+			+		

120				+ + +			
121							
122		+	+	+		+	+
123			+	+	+		+
124				+		+	+
125				+			
126					+		
1			+	+	+		
2		+	+	+	+		+
3		+		+			+
4		+		+			+
5		+	+	+	+		+
6		+	+	+	+		
7		+	+	+	+		+
8		+	+	+	+		+
9		+	+	+	+		+
10		+	+	+	+	+	+
11				+			
12		+	+	+	+		+
Mackay (Waterhouse & Lyell 1914)	+	+		+	+	+	+
13		+	+	+	+		

14			+	+	+		+								
15			+	+	+	+		+							
16			+	+	+										
33			+	+	+	+					+				
34			+	+			+								
<hr/>															
ST.LAWRENCE- ROCKHAMPTON															
21			+			+	+		+	+					
22			+			+				+					
23			+			+	+		+	+					
Byfield (pers. obs.)			+	+	+			+	+						
Byfield (Stewart 1981)			+	+	+	+		+		+					
Yeppoon (Turner 1925)			+			+	+		+	+		+			
Totals for Zone 1	3	1	19	56	52	67	46	5	42	36	3	3	1	0	0
Total sites =	76														

Zone 2

CAPE YORK-HOPE VALE

Upper Jardine River

(Moulds & d'Apice 1982)

+

+

+

+

+

+

61

+

62

+

63

Mt. Black, 25 km W
Townsville (pers. obs.)

+ + + + + +

TOWNSVILLE-
ST.LAWRENCE

165

+ + + + + +

166

+ + + + + +

167

+ + + + + +

49

+ + + + + +

48

+ + + + + +

41

+ + + + + +

42

+ + + + + +

40

+ + + + + +

35

+ + + + + +

36

+ + + + + +

37

+ + + + + +

ST.LAWRENCE-
ROCKHAMPTON

20

+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +
+ + + + + +

Totals for Zone 2

0 0 0 14 24 26 18 0 25 10 4 2 0 0 0

Total sites = 37

Zone 3

HOPE VALE-
TOWNSVILLE

+

+ + + + + + + + + + + +

$$+ + + + + + + + + + +$$

+ + + + + + + + + + + + + + +

| | | |
|--|----|----|
| 44 | + | + |
| 45 | + | + |
| 38 | + | + |
| 39 | + | + |
| ST.LAWRENCE-ROCKHAMPTON | | |
| 18 | + | + |
| 19 | + | + |
| 17 | + | + |
| 32 | + | + |
| 31 | + | + |
| 24 | + | + |
| 25 | + | + |
| 26 | + | + |
| 27 | + | + |
| Totals for Zone 3 | 0 | 0 |
| | 2 | 13 |
| | 23 | 22 |
| | 0 | 27 |
| Total sites = 53 | 0 | 0 |
| | 3 | 0 |
| | 0 | 0 |
| | 0 | 0 |
| Zone 4 | | |
| ATHERTON
TABLELANDS-PALUMA
RANGE | | |
| 85 | + | + |

Appendix 1. Larval host plant records of satyrine butterflies in northern Queensland, 1985, 1989-93. * = Host species introduced to Australia. ϕ = G. Moore (pers. comm.). Egg-laying times are E.S.T.

| Satyrine species | Larval host plant | Locality | Date | Observations |
|---------------------------|--|----------------------|-------------|--------------------------|
| <i>Mycalkesis perseus</i> | <i>Themedea triandra</i> Forsskål | Townsville | 1985 | 1 egg laid ϕ |
| " | " | 30km SE Cardwell | 12.iii.1989 | 1 larva (instar V) |
| " | " | Townsville | 15.v.1990 | 1 larva (instar IV) |
| " | " | " | 1.vi.1990 | 1 larva (instar V) |
| " | " | " | 19.iii.1991 | 1 egg laid |
| " | " | " | 9.iv.1991 | 1 pupal exuvia |
| " | " | " | 1.vii.1991 | 1 egg laid at 1630 |
| " | " | " | 29.i.1992 | 1 egg |
| " | " | 15km SE Cardwell | 10.vi.1992 | 1 pupal exuvia |
| " | " | Townsville | 20.iv.1989 | 1 egg laid at 1530 |
| " | " | " | 9.vi.1989 | 2 larvae (instar V) |
| " | " | " | 1.v.1990 | 1 egg laid at 1530 |
| " | " | " | 12.v.1991 | 1 egg laid at 1530 |
| " | " | " | 8.i.1991 | 1 larva (instar V) |
| " | " | 20km NW Townsville | 10.iii.1991 | 2 eggs laid at 1654-1716 |
| " | <i>H. contortis</i> (L.) Roemer & Schultes | Townsville | 1985 | 1 egg laid ϕ |
| " | * <i>Panicum maximum</i> Jacq. | " | 18.ii.1990 | 1 egg laid at 1757 |
| " | " | 10km NW Rollingstone | 18.i.1992 | 1 larva (instar V) |
| " | " | Townsville | 1985 | 1 egg laid ϕ |
| " | " | " | 1985 | 1 egg laid ϕ |
| " | " | " | 1985 | 1 egg laid ϕ |
| " | " | Seaforth | 25.iii.1992 | 1 egg laid at 1611 |
| " | grass sp. 1 | Mt. Elliot Nat. Park | 10.ii.1991 | 2 eggs laid at 1546 |

| | | | | |
|----------------------------|--|------------------------------------|--------------|-----------------------------|
| <i>Myccalesis terminus</i> | grass sp.2 | " | 10.i.1991 | 1 egg laid at 1610 |
| | <i>Opismenus aenulus</i> (P.Bt.) Roemer & Schultes | 10km NW Rollingstone | 30.x.1993 | 1 egg laid at 1137 |
| | <i>Opismenus</i> sp. | Townsville | 3.iv.1990 | 1 egg laid at 1520 |
| | " | " | 8.iv.1990 | 2 eggs laid at 1530 |
| | " | " | 22.i.1992 | 2 eggs laid at 1430-1445 |
| | " | Bartle Frere | 16.viii.1992 | 1 egg laid at 1322 |
| | <i>Dichanthium sericeum</i> | Townsville | 23.xi.1990 | 1 egg laid at 1638 |
| | " | " | 28.iv.1991 | 1 egg |
| | " | " | 6.v.1991 | 2 eggs & 1 larva (instar I) |
| | " | " | 1985 | 3 eggs laid ϕ |
| | * <i>Axonopus compressus</i> (Sw.) P. Beauv. | " | 1985 | 4 eggs laid ϕ |
| | * <i>Panicum maximum</i> | 10km NW Rollingstone | 30.x.1993 | 1 egg laid at 1052 |
| | " | Cardwell | 9.i.1990 | 1 egg laid at 1610 |
| | <i>Themeda triandra</i> | Townsville | 22.i.1992 | 1 egg laid at 1521 |
| | grass sp.1 | Seaforth | 25.iii.1992 | 1 egg laid at 1530 |
| | grass sp.2 | Wenlock River, Cape York Peninsula | 2.x.1990 | 1 egg laid at 1633 |
| | <i>Flagellaria indica</i> L. (Flagellariaceae) | | | |
| | <i>Mycellesis sirius</i> | 8km SE Cardwell | 9.vii.1989 | 2 eggs laid at 1406-1408 |
| | " | 10km SE Cardwell | 3.ix.1989 | 6 eggs laid at 1355-1440 |
| | " | Cardwell | 8.ix.1989 | 1 egg laid at 1300 |
| | * <i>Panicum maximum</i> | 12km NW Rollingstone | 25.v.1992 | 1 egg laid at 1518 |
| | " | 12km NW Rollingstone | 13.vi.1993 | 7 eggs laid at 1410-1423 |
| | grass sp. | 10km SE Cardwell | 3.ix.1989 | 1 egg laid at 1410 |
| | " | Townsville | 2.iii.1991 | 5 larvae |
| | * <i>Panicum maximum</i> | Cape Cleveland | 3.iii.1991 | 1 larva (instar V) |
| | " | Majors Creek | 20.iii.1991 | eggs, larvae & pupae |

| | | | | |
|---|---|----------------------|-------------|-----------------------------------|
| " | " | 11km W Kennedy | 7.iv.1991 | 3 eggs, ca.100 larvae, 20 pupae |
| " | " | 10km NW Rollingstone | 18.i.1992 | 2 larvae & 5 pupae |
| " | <i>Imperata cylindrica</i> (L.) P. Beauv. | 12km W Kennedy | 27.ii.1991 | 20 larvae |
| " | " | Cape Cleveland | 3.iii.1991 | 3 larvae (1 instar I, 2 instar V) |
| " | " | 15km SE Cardwell | 2.xii.1991 | 1 egg laid at 1832 |
| " | <i>Ophiuros exaltatus</i> (L.) Kuntze | Mt. Elliot Nat. Park | 10.ii.1991 | 3 pupal exuviae |
| " | " | 15km SE Cardwell | 26.ii.1991 | 3 larvae |
| " | " | " | 2.xii.1991 | 5 eggs laid at 1825 |
| " | <i>Themeda triandra</i> | Mt. Elliot Nat. Park | 10.iii.1991 | 1 pupal exuvia |
| " | " | Townsville | 20.iii.1991 | 3 eggs |
| " | " | Byfield | 29.iii.1992 | 1 egg laid at 1737 |
| " | <i>Heteropogon triticeus</i> | Cardwell | 26.iv.1990 | 1 pupa |
| " | " | 15km SE Cardwell | 27.v.1992 | 2 pupal exuviae |
| " | * <i>Melinis minutiflora</i> P. Beauv. | 12km W Kennedy | 27.iii.1991 | 20 larvae |
| " | <i>Sorghum</i> sp. | Cape Cleveland | 3.iii.1991 | 3 eggs |
| " | <i>Chrysopogon</i> sp. | Innot Hot Springs | 2.vi.1991 | 1 larva (instar V) |
| " | <i>Paspalidium</i> sp. | Byfield | 29.iii.1992 | 2 eggs laid at 1738 |
| " | grass sp. | Townsville | 6.v.1991 | 1 larva (instar V) |
| " | <i>Hypoxysta adiante</i> | 8km SE Cardwell | 24.i.1991 | 1 egg laid at 0833 |
| " | <i>Themeda triandra</i> | Paluma | 17.iii.1992 | 3 eggs laid at 1105-1125 |
| " | <i>Eriachne pallens</i> R.Br. | 10km W Paluma | 21.xi.1993 | 2 eggs laid at 1439-1457 |
| " | grass sp. | " | " | " |

Appendix 2. Adult host plant records of satyrine butterflies in northern Queensland, 1989-93. * = Host species introduced to Australia. Feeding times are E.S.T.

| Satyrine species | Adult host plant | Locality | Date | Observations |
|--------------------------|---|----------------------|-------------|-------------------------|
| <i>Mycaleis perseus</i> | <i>Melaleuca leucadendra</i> (L.) L. (Myrtaceae)
(nectar) | Townsville | 24.vii.1989 | 1♀ at 1300 |
| " | * <i>Mangifera indica</i> L. (Anacardiaceae) (nectar) | 18km N Coen | 2.x.1990 | 1♂ at 1145 |
| " | " | Townsville | 27.xii.1990 | ♀ ♀ at 0930 |
| " | <i>Alphitonia excelsa</i> (Fenzl) Benth.
(Rhamnaceae) (nectar) | Townsville | 5.iii.1991 | 2♂♂ & 1♀ |
| " | <i>Pandanus</i> sp. (Pandanaceae) (rotting fruit) | 15 km NW Bluewater | 24.vii.1993 | 1♀ at 1650 |
| " | <i>Pandanus whitei</i> Martelli (Pandanaceae)
(rotting fruit) | 10km SE Cardwell | 25.ix.1990 | ♀ ♀ at 1500-1520 |
| <i>Mycaleis terminus</i> | * <i>Mangifera indica</i> (rotting fruit) | Townsville | 31.x.1990 | 2♂♂ at 1500-1530 |
| " | " | " | 30.xi.1990 | 1♀ at 1515 |
| " | <i>Nauclea orientalis</i> (L.) L. (Rubiaceae) (rotting
fruit) | 10km NW Rollingstone | 16.iii.1991 | ♂♂ & ♀ ♀ at 1330 |
| " | " | " | 18.i.1992 | ♂♂ & ♀ ♀ at 1000-1200 |
| " | " | " | 5.ii.1994 | ♂♂ at 1630 |
| " | <i>Ficus racemosa</i> L. (Moraceae) (rotting fruit) | 15km W Rupertswood | 27.iv.1991 | ♂♂ & ♀ ♀ at 1520-1600 |
| " | <i>Planchonia careya</i> (F. Muell.) Knuth
(Lecythidaceae) (sap) | Townsville | 2.iii.1991 | 1♂ at 0830 |
| <i>Mycaleis sirius</i> | <i>Helichrysum rupicola</i> DC (Asteraceae)
(nectar) | 20km SE Cardwell | 25.ix.1990 | 1♂ & 1♀ at 1504 |
| " | <i>Melastoma affine</i> (Melastomataceae) (fruit) | 9km SE Cardwell | 15.ix.1991 | 1♂ at 1209-1212 |
| <i>Ypthima arctous</i> | <i>Helichrysum rupicola</i> (nectar) | 8km SE Cardwell | 3.viii.1990 | 1♀ at 1600-1710 |
| " | * <i>Bidens pilosa</i> (Asteraceae) (nectar) | 10km NW Rollingstone | 17.iii.1992 | 2♂♂ & 2♀ ♀ at 1624-1634 |
| " | " | Townsville | 8.i.1991 | 1♀ at 1356-1358 |
| " | * <i>Emilia sonchifolia</i> (L.) DC (Asteraceae)
(nectar) | 15km S Eton | 5.iv.1992 | 1♂ at 1426 |

| | | | | |
|---|--|--------------------------------------|--------------|--|
| " | <i>Melastoma affine</i> D. Don (Melastomataceae)
(nectar) | Cardwell | 20.viii.1990 | 1♀ |
| " | " | Kennedy | 26.x.1990 | 1♀ at 1245 |
| " | <i>Callistemon viminalis</i> (Gaertner) G. Don
(Myrtaceae) (nectar) | 70km N Duaringa | 8.ix.1990 | 1♂ & 1♀ at 1310 |
| " | <i>Melaleuca linariifolia</i> Smith (Myrtaceae)
(nectar) | 6km SE Petford | 19.x.1990 | ♂♂ & ♀♀ at 1550 |
| " | <i>Lepospermum polygalifolium</i> sens. lat. Salisb.
(Myrtaceae) (nectar) | 9km SE Cardwell | 16.ix.1991 | ♂♂ & ♀♀ at 1020-1100,
1220-1300, 1515 |
| " | <i>Stylosanthes</i> sp. (Fabaceae) (nectar) | 20km SW Mareeba | 19.x.1990 | 1♀ at 1155 |
| " | " | Bluewater | 14.v.1992 | 2♂♂ at 1157-1200 |
| " | Fabaceae (nectar) | Blackdown Tbld. Nat.
Park | 7.ix.1990 | 2♀♀ at 1210 |
| " | <i>Bursaria tenifolia</i> Bailey (Pittosporaceae)
(nectar) | 15km SE Cardwell | 16.ix.1991 | 1♀ at 1215 |
| " | * <i>Stachytarpheta jamaicensis</i> (L.) Vahl
(Verbenaceae) (nectar) | 9km SE Cardwell | 7.iii.1992 | 1♀ at 1311-1321 |
| " | <i>Ceratanthus longicornis</i> (F. Muell.) G. Taylor
(Lamiaceae) (nectar) | | 7.iii.1992 | 1♂ at 1435 |
| " | Lamiaceae (nectar) | Cardwell | 24.i.1991 | 1♀ |
| " | <i>Hybanthus cuneaspermus</i> (L.) F. Muell.
(Violaceae) (nectar) | 15km SE Cardwell | 8.iii.1992 | 1♀ at 1026 |
| " | " | " | 28.v.1992 | 2♂♂ at 1157-1200 |
| " | <i>Waltheria indica</i> L. (Sterculiaceae) (nectar) | Cape Upstart Nat. Park | 23.iv.1992 | 1♂ at 1239 |
| " | <i>Spermacoce latifolia</i> Aublet (Rubiaceae)
(nectar) | Harvey Range, 42km
WSW Townsville | 19.v.1992 | ♀♀ at 1215-1220 |
| " | " | 15km SE Cardwell | 5.vi.1992 | 1♂ at 1506 |
| " | <i>Eriocaulon</i> sp. (Eriocaulaceae) (nectar) | 9km SE Cardwell | 17.x.1992 | 2♂♂ at 1505-1512 |
| " | <i>Opilia armamentea</i> Roxb. (Opliliaceae) (nectar) | Mt. Wite, Coen | 1.x.1990 | 2♀♀ at 1540 |
| " | <i>Leptospermum polygalifolium</i> sens. lat.
(nectar) | 35km NWW Torrens
Creek | 4.x.1991 | 1♂ at 0740 |
| | <i>Hypocysta adiante</i> | | | |

| | | | | |
|--|--|---------------------------|-------------|----------------------|
| " | <i>Melaleuca linariifolia</i> (nectar) | 6km SE Peltford | 19.x.1990 | ♀♀ at 1550-1600 |
| " | * <i>Stachyarpheia jamaicensis</i> (nectar) | 8km SE Cardwell | 10.i.1991 | 1♀ at 1515 |
| " | <i>Rostellularia adscendens</i> (R. Br.) R. Barker
(Acanthaceae) (nectar) | Cardwell | 12.i.1991 | 1♂ & 1♀ at 1600 |
| " | " | " | 24.i.1991 | 1♂ |
| " | " | 15km S Eungella | 8.iv.1992 | 2♂♂ at 1250 |
| " | " | 16km W Paluma | 5.vi.1993 | 1♂ & 1♀ at 1259-1301 |
| " | <i>Bursaria tenifolia</i> (nectar) | 15km SE Cardwell | 8.ix.1991 | 1♂ at 1114 |
| " | " | 17km SW Bluewater | 27.iii.1993 | 1♂ at 1008-1010 |
| " | " | Hidden Valley | 17.iv.1993 | 4♂♂ at 1436-1444 |
| " | <i>Spermacoce latifolia</i> (nectar) | Innot Hot Springs | 2.vi.1991 | ♀♀ at 1130-1135 |
| " | <i>Canthium</i> sp. (Rubiaceae) (nectar) | 15km SE Cardwell | 16.x.1992 | 1♂ at 1202-1205 |
| " | <i>Euphorbia</i> sp. (Euphorbiaceae) (nectar) | Marlborough | 28.iii.1992 | 1♂ at 1345 |
| " | * <i>Emilia sonchifolia</i> (Asteraceae) (nectar) | 15km S Eton | 5.iv.1992 | 1♂ at 1424 |
| " | <i>Vernonia cinerea</i> Less. (Asteraceae) (nectar) | 8km E Paluma | 14.v.1992 | 1♀ at 1523-1526 |
| " | Asteraceae (nectar) | 6km W Grantleigh | 6.ix.1990 | 4♀♀ at 0950 |
| " | Fabaceae, Asteraceae (nectar) | Blackdown Tbld. Nat. Park | 7.ix.1990 | ♀♀ at 1100-1200 |
| <i>Hypocysta metirius</i> | " | Atherton | 21.x.1990 | 1♀ at 1655 |
| " | * <i>Bidens pilosa</i> (nectar) | Kirrima, Cardwell Range | 4.vi.1991 | 1♀ at 1505 |
| " | * <i>Lantana camara</i> L. (Verbenaceae) (nectar) | Eungella | 24.vi.1991 | 1♀ at 1105 |
| " | * <i>Rubus rosifolius</i> Smith (Rosaceae) (nectar) | " | 8.iv.1991 | 2♂♂ at 0837, 0917 |
| " | <i>Acronychia laevis</i> Forster & G. Forster
(Rutaceae) (nectar) | Seaview Range | 10.x.1991 | ♂♂ & ♀♀ at 1510 |
| " | <i>Maesa dependens</i> F. Muell. (Myrsinaceae)
(nectar) | Atherton | 21.x.1990 | 1♀ at 1655 |
| <i>Hypocysta irius</i> | " | 17km W Kennedy | 4.vi.1991 | 5♀♀ at 1430-1445 |
| * <i>Bidens pilosa</i> (Asteraceae) (nectar) | | | | |
| <i>Helichrysum rupicola</i> (nectar) | | | | |

| | | | |
|--|---|--------------------|----------------------------|
| <i>*Ageratum houstonianum</i> Miller, * <i>Congya</i> sp.
(Asteraceae); * <i>Stachytarpheta jamaicensis</i> ;
<i>Spermicoce latifolia</i> (nectar) | | 4.vi.1991 | 1♀ at 1435-1445 |
| <i>Hypocysta pseudirius</i> | Asteraceae (nectar) | 6km W Grantleigh | 6.ix.1990 1♂ at 0950 |
| " | " | 20km ESE Duaringa | 6.ix.1990 3♀ ♀ at 1155 |
| <i>Tisiphone helena</i> | <i>Polyscias australiana</i> (F. Muell.) Philipson
(Araliaceae) (nectar) | 26km SW Bluerwater | 30.xi.1992 1♂ at 1005-1007 |

Appendix 4. Dominant plant species at sampling sites A-D, Cardwell.

SITE A

Overstorey

Canarium australianum F.Muell., *Xanthostemon chrysanthus* (F.Muell.) Benth., *Pongamia pinnata* (L.) Pierre, *Scolopia braunii* (Klotzsch) Sleumer, *Buckinghamia celsissima* F.Muell., *Planchonella* sp, *Paraserianthes toona* (Bailey) I.Nielsen, *Syzygium tierneyanum* (F.Muell.) T.Hartley & Perry, *Schefflera actinophylla* (Endl.) Harms, *Dillenia alata* (DC.) Martelli, *Deplanchea tetraphylla* (R.Br.) F.Muell., *Bursaria tenuifolia* Bailey, *Lophostemon suaveolens* (Gaetner) Peter G.Wilson & Waterhouse, *Eucalyptus tereticornis* Smith, *E. tessellaris* F.Muell., *E. drepanophylla* Benth., *E. clarksoniana* D.J. Carr & S.G.M. Carr, *E. platyphylla* F.Muell., *E. intermedia* R.Barker, *Acacia mangium* Willd., *A. aulacocarpa* Benth.

Understorey

Acacia flavesrens Benth., *Guioa acutifolia* (F.Muell.) Radlk., *Mallotus polyadenos* F.Muell., *Chionanthis ramifloris* Roxb., *Pandanus* sp, *Alyxia spicata* R.Br., *Scolopia braunii*, *Planchonia careya* (F.Muell.) Knuth, *Dillenia alata*, *Glochidion ferdinandi* (Muell.Arg) Bailey, *Alphitonia excelsa* (Fenzl) Benth., *Lantana camara* L, *Jagera pseudorhus* (A.Rich.) Radlk., *Polyscias elegans* (C.Moore & F.Muell.) Harms, *Timonius timon* (Sprengel) Merr. *Cycas media* R.Br., *Alyxia spicata*, *Eustrephus latifolius* Ker Gawler, *Scleria sphacelata* F.Muell., *Lomandra longifolia* Labill.

SITE B

Overstorey

Eucalyptus platyphylla, *E. drepanophylla*, *E. clarksoniana*, *Lophostemon suaveolens* with some *E. acmenoides* Schauer.

Understorey

Acacia flavesrens, *Melaleuca nervosa* (Lindley) Cheel, *Grevillea parallela* J.Knight and *Planchonia careya*.

SITE C

Overstorey

Melaleuca viridiflora Gaetner and *M. quinquenervia* (Cav.) S.T.Blake with some *Eucalyptus pellita* F.Muell., *Pandanus whitei* Martelli and *Deplanchea tetraphylla*.

Understorey

Grevillea pteridifolia J.Knight, *Pandanus whitei* with *Gahnia sieberiana* Kunth.

SITE D

Overstorey

Lophostemon suaveolens, *Eucalyptus platyphylla*, *E. intermedia* and *E. acmenoides*. *Melaleuca dealbata* S.T. Blake, *Eucalyptus leptophleba* F.Muell. and *E. tessellaris* form minor components.

Understorey

Planchonia careya, *Lophostemon suaveolens*, *Acacia flavesrens*, *Canarium australasicum* (Bailey) Leenh., *Grevillea parallela*, *Helicteres semiglabra* Bailey; with some *Acacia crassicarpa* Benth., *Alphitonia excelsa*, *Melaleuca quinquinervia* and *Acacia leptocarpa*.