

*Corrigendum to*

**“Modelling basin-wide variations in Amazon forest productivity – Part 1: Model calibration, evaluation and upscaling functions for canopy photosynthesis” published in Biogeosciences, 6, 1247–1272, 2009**

**L. M. Mercado<sup>1,2</sup>, J. Lloyd<sup>3</sup>, A. J. Dolman<sup>4</sup>, S. Sitch<sup>5</sup>, and S. Patiño<sup>2,3,6</sup>**

<sup>1</sup>Centre for Ecology and Hydrology, Wallingford, Oxon, OX10 8BB, UK

<sup>2</sup>Max Planck Institute for Biogeochemistry, 07745 Jena, Germany

<sup>3</sup>School of Geography, University of Leeds, LS2 9JT, UK

<sup>4</sup>Free University of Amsterdam, 1081 HV Amsterdam, The Netherlands

<sup>5</sup>Met Office Hadley Centre, JCHMR, Wallingford, Oxon, OX10 8BB, UK


<sup>6</sup>Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Diagonal 27 No. 15-09, Bogotá D.C, Colombia

In the above mentioned manuscript, sites names in Figs. 2, 3 and 4 were missing. The new figures can be found on the following pages.

The service charges for this open access publication have been covered by the Max Planck Society.



Correspondence to: L. M. Mercado  
([lmme@ceh.ac.uk](mailto:lmme@ceh.ac.uk))



**Fig. 2.** Light response of observed (black) and simulated (grey) values of  $G_p$  during the seasons tested.





Fig. 3. Diurnal cycle of observed (black) and simulated (grey)  $G_p$ .



**Fig. 4.** Mean diurnal cycle of observed (black) and simulated (grey) evaporative fraction.