

*The following supplements accompany the article*

# **Ecosystem equivalence and the ability to generalise: insights from global consistencies in mangrove fish assemblages**

**Marcus Sheaves**

Centre for Tropical Water and Aquatic Ecosystem Research, School of Marine and Tropical Biology, James Cook University, Townsville, Queensland 4811, Australia

\*Email: marcus.sheaves@jcu.edu.au

*Marine Ecology Progress Series 461: 137–149 (2012)*

## Supplement S1. Sampling considerations and assumptions

The diverse derivation of the data on which the present study was based presented a range of difficulties, some of which were addressed at least partially by the inclusion of explanatory variables and some that required specific assumptions:

- 1) The different studies had used different sampling gears and sampling approaches because they had been conducted under different local conditions and for different purposes. There was no sensible way to standardise for this, and although ‘gear type’ was included as an explanatory variable, to account for any systematic difference due to sampling gear, it is likely that gear-related differences and selectivities would still add substantial random variation to the data set, both in the species captured and the extent to which catch per unit effort (CPUE) related to actual abundances. For this reason, rather than concentrating on species and CPUE, the present study focussed on the relative number of species per family (i.e. the proportion of species per family for each study), a robust approach that also helped reduce the impact of a number of other issues (see below).
- 2) The studies dealt with a range of structurally different systems, ranging from mangroves on fringing reefs through coastal mangrove forests to riverine mangroves spanning a range of salinity environments. Thus, the data included a variety of mangrove setting and types and a vast range of physical environments. Although this was captured to some extent in 2 variables, one indicating system type and the other salinity regime, the exact type of system or its equivalence to systems reported in other studies was not always clear, again adding additional unexplainable variation to the data set. For studies that reported data from ecosystems other than mangrove forests or mangrove systems, the non-mangrove data were excluded, to the extent that they could be identified. However, the information needed to do this was not always available.
- 3) The scales of the different studies varied in terms of spatio-temporal extent, size of systems sampled, size of sampling units and the range of habitats sampled. While it was difficult to account for all these differences because many studies did not provide the information necessary to do so, a ‘system scale’ variable was included in the explanatory data set.
- 4) One particularly important scale issue related to the focus of the study. A number of studies explicitly targeted mangrove forests, either sampling inside the forest or using gears such as fyke nets to capture fish leaving the forest; however, the majority of studies either sampled

the whole system or did not define an explicit focus. It may have been preferable to concentrate only on studies with a purely mangrove forest focus, but there were relatively few of these, and they did not provide a broad global coverage. Consequently, data of both types were included, and these differences were accounted for with an explanatory variable.

- 5) Sampling intensity varied greatly among the studies in the number of units sampled, the spatio-temporal extent of sampling and the amount of effort per sampling site. This was extremely variable and difficult to account for but again underlines the need to deal with the data only at the most robust level. At a larger conceptual scale, the intensity of research differed greatly among geographic regions, with a large number of studies in some parts of the world but few studies in many others. To account for this disparity, any studies from a single geographic area that included data sets from multiple mangrove systems that had been (a) sampled using equivalent methods by the same sampling team and (b) reported to have consistent faunal composition (e.g. Sheaves & Johnston 2009) were treated as a single data set. In contrast, other studies reported different assemblages from different locations, and these were treated as independent data sets.

#### Supplement S2. The 76 studies from which data sets were extracted

- Adite A, Winemiller KO (1997) Trophic ecology and ecomorphology of fish assemblages in coastal lakes of Benin, West Africa. *Ecoscience* 4:6–23
- Albaret JJ, Simier M, Darboe FS, Ecoutin JM, Raffray J, de Morais LT (2004) Fish diversity and distribution in the Gambia Estuary, West Africa, in relation to environmental variables. *Aquat Living Resour* 17:35–46
- Amadi A (1990) A comparative ecology of estuaries in Nigeria. *Hydrobiologia* 208:27–38
- Austin H (1971) A survey of the ichthyofauna of the mangroves of western Puerto Rico during December, 1967 - August, 1968. *Caribb J Sci* 11:27–39
- Barletta M, Blaber S (2007) Comparison of fish assemblages and guilds in tropical habitats of the Embley (Indo-West Pacific) and Caete (Western Atlantic) estuaries. *Bull Mar Sci* 80:647–680
- Bell JD, Pollard DA, Burchmore JJ, Pease BC, Middleton MJ (1984) Structure of a fish community in a temperature tidal mangrove creek in Botany Bay, New South Wales. *Aust J Mar Freshwater Res* 35:33–46
- Blaber SJM (1980) Fish of the Trinity Inlet system of north Queensland with notes on the ecology of fish fauna of tropical Indo-Pacific estuaries. *Aust J Mar Freshwater Res* 31:137–146
- Blaber SJM, Blaber TG (1980) Factors affecting the distribution of juvenile estuarine and inshore fish. *J Fish Biol* 17:143–162
- Blaber SJM, Milton DA (1990) Species composition, community structure and zoogeography of fishes of mangrove estuaries in the Solomon Islands. *Mar Biol* 105:259–267
- Blaber SJM, Young JW, Dunning MC (1985) Community structure and zoogeographic affinities of the coastal fishes of the Dampier Region of north-western Australia. *Aust J Mar Freshwater Res* 36:247–266
- Blaber SJM, Brewer DT, Salini JP (1989) Species composition and biomasses of fishes in different habitats of a tropical northern Australian estuary: their occurrence in the adjoining sea and estuarine dependence. *Estuar Coast Shelf Sci* 29:509–531
- Chaves P, Bouchereau JL (1999) Biodiversity and dynamics of ichthyic communities in the mangrove of Guaratuba, Brazil. *Oceanol Acta* 22:353–364
- Chua TE (1973) An ecological study of the Ponggol Estuary in Singapore. *Hydrobiologia* 43:505–533
- Clynick G, Chapman M (2002) Assemblages of small fish in patchy mangrove forests in Sydney Harbour. *Mar Freshw Res* 53:669–677

- Collette B (1983) Mangrove fishes of New Guinea. In: Teas H (ed) Tasks for vegetation science, Vol 8. Dr W Junk, The Hague, p 91–102
- Halliday IA, Young WR (1996) Density, biomass and species composition of fish in a subtropical *Rhizophora stylosa* mangrove forest. *Mar Freshw Res* 47:609–615
- Harrison TD (2005) Ichthyofauna of South African estuaries in relation to the zoogeography of the region. *Smithiana Bull* 6:2–27
- Hindell JS, Jenkins GP (2004) Spatial and temporal variability in the assemblage structure of fishes associated with mangroves (*Avicennia marina*) and intertidal mudflats in temperate Australian embayments. *Mar Biol* 144:385–395
- Hoq M, Islam M (2007) Composition and seasonal dynamics of postlarval and juvenile fishes in the Sundarbans mangrove waters, Bangladesh. *Indian J Mar Sci* 36:206–215
- Huxham M, Kimani E, Augley J (2004) Mangrove fish: a comparison of community structure between forested and cleared habitats. *Estuar Coast Shelf Sci* 60:637–647
- Hyslop EJ (1999) Longitudinal variation in fish species composition in the Angabanga river, Papua New Guinea with observations on the trophic status of certain fish species. *Ecol Freshwat Fish* 8:102–107
- Ikejima K, Tongnunui P, Medej T, Taniuchi T (2003) Juvenile and small fishes in a mangrove estuary in Trang province, Thailand: seasonal and habitat differences. *Estuar Coast Shelf Sci* 56:447–457
- Inger R (1955) Ecological notes on the fish fauna of a coastal drainage of North Borneo. *Fieldiana Zool* 37:47–90
- Iose PK, McConnaughey J (1993) Fishery resources in Pala Lagoon. Dept Mar Water Resour Biol Rep Ser 37, American Samoa Government, Pago Pago
- Islam M, Ikejima K (2010) Gear type, species composition and economic value of fisheries in the mangroves of Pak Phanang, Thailand. *Wetlands Ecol Manage* 18:27–36
- Kimani E, Mwatha G, Wakwabi E, Ntiba J, Okoth B (1996) Fishes of a shallow tropical mangrove estuary, Gazi, Kenya. *Mar Freshw Res* 47:857–868
- Krumme U, Saint-Paul U, Rosenthal H (2004) Tidal and diel changes in the structure of a nekton assemblage in small intertidal mangrove creeks in northern Brazil. *Aquat Living Resour* 17:215–229
- Kuo SR, Lin HJ, Shao KT (1999) Fish assemblages in the mangrove creeks of Northern and Southern Taiwan. *Estuaries* 22:1004–1015
- Lal PN (1984) Coastal fisheries and the management of mangrove resources in Fiji. *S Pac Comm Newsl Noumea*:15–23
- Laroche J, Rasoanandrasana N (1997) Temporal patterns in a fish assemblage of a semiarid mangrove zone in Madagascar. *J Fish Biol* 51:3–20
- Ley JA (2005) Linking fish assemblages and attributes of mangrove estuaries in tropical Australia: criteria for regional marine reserves. *Mar Ecol Prog Ser* 305:41–57
- Ley JA, McIvor C, Montague C (1999) Fishes in mangrove prop-root habitats of northeastern Florida Bay: distinct assemblages across an estuarine gradient. *Estuar Coast Shelf Sci* 48:701–723
- Lin HJ, Shao KT (1999) Seasonal and diel changes in a subtropical mangrove fish assemblage. *Bull Mar Sci* 65:775–794
- Little MC, Reay PJ, Grove SJ (1988) The fish community of an East African mangrove creek. *J Fish Biol* 32:729–747
- Lorenz JJ (1999) The response of fishes to physicochemical changes in the mangroves of northeast Florida Bay. *Estuaries* 22:500–517
- Mbande S, Whitfield A, Cowley P (2005) The ichthyofaunal composition of the Mngazi and Mngazana estuaries: a comparative study. *Smithiana Bull* 4:1–20

- Morton R (1990) Community structure, density and standing crop of fishes in a subtropical Australian mangrove area. *Mar Biol* 105:385–394
- Morton R, Pollock B, Beumer J (1987) The occurrence and diet of fishes in a tidal inlet to a saltmarsh in southern Moreton Bay, Queensland. *Aust J Ecol* 12:217–237
- Mwandya AW, Gullstrom M, Ohman MC, Andersson MH, Mgaya YD (2009) Fish assemblages in Tanzanian mangrove creek systems influenced by solar salt farm constructions. *Estuar Coast Shelf Sci* 82:193–200
- Nagelkerken I, van der Velde G (2002) Do non-estuarine mangroves harbour higher densities of juvenile fish than adjacent shallow-water and coral reef habitats in Curacao (Netherlands Antilles)? *Mar Ecol Prog Ser* 245:191–204
- Nagelkerken I, Dorenbosch M, Verberk W, Chocheret de la Moriniere E, Van der Velde G (2000) Importance of shallow-water biotopes of a Caribbean bay for juvenile coral reef fishes: patterns in biotope association, community structure and spatial distribution. *Mar Ecol Prog Ser* 202:175–192
- Odum W, Heald E (1972) Trophic analyses of an estuarine mangrove community. *Bull Mar Sci* 22:671–738
- Pinto L (1987) Environmental factors influencing the occurrence of juvenile fish in the mangroves of Pagbilao, Philippines. *Hydrobiologia* 150:283–301
- Pinto L, Punchihewa N (1996) Utilisation of mangroves and seagasses by fishes in the Negombo Estuary, Sri Lanka. *Mar Biol* 126:333–345
- Quinn J (1980) Analysis of temporal changes in fish assemblages in Serpentine Creek, Queensland. *Environ Biol Fishes* 5:117–133
- Quinn NJ, Kojis BJ (1985) Does the presence of coral reefs in proximity to a tropical estuary affect the estuarine fish assemblage? *Proc 5th Int Coral Reef Congr* 5:445–450
- Quinn J, Kojis B (1986) Annual variation in the nocturnal nekton assemblage of a tropical estuary. *Estuar Coast Shelf Sci* 22:63–90
- Rehage J, Loftus W (2007) Seasonal fish community variation in headwater mangrove creeks in the southwestern Everglades and examination of their role as dry-down refuges. *Bull Mar Sci* 80:625–645
- Robertson AI, Duke NC (1990) Mangrove fish-communities in tropical Queensland, Australia: spatial and temporal patterns in densities, biomass and community structure. *Mar Biol* 104:369–379
- Ronnback P (1999) The ecological basis for economic value of seafood production supported by mangrove ecosystems. *Ecol Econ* 29:235–252
- Rooker JR, Dennis GD (1991) Diel, lunar and seasonal changes in a mangrove fish assemblage off southwestern Puerto Rico. *Bull Mar Sci* 49:684–698
- Sedberry G, Carter J (1993) The fish community of a shallow tropical lagoon in Belize, Central America. *Estuaries* 16:198–215
- Serafy JE, Lindeman KC, Hopkins TE, Ault JS (1997) Effects of freshwater canal discharge on fish assemblages in a subtropical bay: field and laboratory observations. *Mar Ecol Prog Ser* 160:161–172
- Serafy J, Faunce C, Lorenz JJ (2003) Mangrove shoreline fishes of Biscayne Bay, Florida. *Bull Mar Sci* 72:161–180
- Sheaves MJ (2006) Scale-dependent variation in composition of fish fauna among tropical estuarine sandy embayments. *Mar Ecol Prog Ser* 310:173–184
- Sheaves M, Johnston R (2009) Ecological drivers of spatial variability among fish fauna of 21 tropical Australian estuaries. *Mar Ecol Prog Ser* 385:245–260
- Sheaves M, Collins J, Houston W, Dale P, Revill A, Johnston R, Abrantes K (2006) Contribution of floodplain wetland pools to the ecological functioning of the Fitzroy River Estuary. Cooperative Research Centre for Coastal Zone, Estuarine and Waterway Management, Brisbane

- Sheaves M, Johnston R, Abrantes K (2007a) Fish fauna of dry sub-tropical estuarine floodplain wetlands. *Mar Freshw Res* 58:931–943
- Sheaves M, Johnston R, Molony B, Shepard G (2007b) The effect of impoundments on the structure and function of fish fauna in a highly regulated dry tropics estuary. *Estuaries* 30:507–517
- Sheaves M, Duc NH, Khoa NX (2008) Ecological attributes of a tropical river basin vulnerable to the impacts of clustered hydropower developments. *Mar Freshw Res* 59:971–986
- Sheaves M, Johnston R, Connolly R (2010) Temporal dynamics of fish assemblages of natural and artificial tropical estuaries. *Mar Ecol Prog Ser* 410:143–157
- Sheridan P (1992) Comparative habitat utilization by estuarine macrofauna within the mangrove ecosystem of Rookery Bay Florida. *Bull Mar Sci* 50:21–39
- Simier M, Blanc L, Aliaume C, Diouf PS, Albaret JJ (2004) Spatial and temporal structure of fish assemblages in an “inverse estuary”, the Sine Saloum system (Senegal). *Estuar Coast Shelf Sci* 59:69–86
- Simier M, Laurent C, Ecoutin JM, Albaret JJ (2006) The Gambia River estuary: a reference point for estuarine fish assemblages studies in West Africa. *Estuar Coast Shelf Sci* 69:615–628
- Stoner A (1986) Community structure of the demersal fish species of Laguna Joyuda, Puerto Rico. *Estuaries* 9:142–152
- Taylor D, Reyier E, McIvor C, Davis W (2007) An assessment of ichthyofaunal assemblages within the mangal of the Belize offshore cays. *Bull Mar Sci* 80:721–737
- Thayer GW, Colby DR, Hettler WF Jr (1987) Utilization of the red mangrove prop root habitat by fishes in South Florida. *Mar Ecol Prog Ser* 35:25–38
- Thollot P (1988) The importance of mangroves to the ichthyofauna of the New Caledonia lagoon. South Pacific Commission, Noumea
- Thollot P (1993) Western Samoa mangrove fish survey, Vol 23. Orstom, Noumea
- Unsworth RKF, Garrard SL, De Leon PS, Cullen LC, Smith DJ, Sloman KA, Bell JJ (2009) Structuring of Indo-Pacific fish assemblages along the mangrove-seagrass continuum. *Aquat Biol* 5:85–95
- Vaslet A, Bouchon-Navaro Y, Charrier G, Louis M, Bouchon C (2010) Spatial patterns in mangrove shoreline fish communities in relation with environmental variables in Caribbean lagoons. *Estuaries Coasts* 33:195–210
- Vidy G (2000) Estuarine and mangrove systems and the nursery concept: which is which? The case of the Sine Saloum system (Senegal). *Wetlands Ecol Manage* 8:37–51
- Vidy G, Darboe FS, Mbye EM (2004) Juvenile fish assemblages in the creeks of the Gambia Estuary. *Aquat Living Resour* 17:56–64
- Warburton K (1978) Community structure, abundance and diversity of fish in a Mexican coastal lagoon system. *Estuar Coast Mar Sci* 7:497–519
- Williamson I, King C, Marther P (1994) A comparison of fish communities in unmodified and modified inshore habitats of Raby Bay, Queensland. *Estuar Coast Shelf Sci* 39:401–411
- Wright JM (1986) The ecology of fish occurring in shallow water creeks of a Nigerian mangrove swamp. *J Fish Biol* 29:431–441

Supplement S3. 170 families of fish reported from mangroves in 4 faunal regions of the world defined by mCART analysis. P indicates presence in a region. EC Atlantic: Eastern central Atlantic, Central IP: Central Indo-Pacific

Family	EC Atlantic	Australasian	Caribbean	Central IP
Adrianichthyidae	-	P	-	-
Acanthuridae	P	P	P	P
Achiridae	P	-	P	-
Acropomatidae	-	-	-	P
Albulidae	-	-	P	P
Alestidae	P	-	-	-
Ambassidae	-	P	-	P
Amiidae	-	-	P	-
Anabantidae	-	P	-	-
Anablepidae	P	-	-	-
Anguilidae	-	P	P	P
Antennariidae	P	P	P	P
Aploactinidae	-	P	-	-
Aplocheilidae	-	-	-	P
Apogonidae	-	P	P	P
Arcanidae	-	-	-	P
Ariidae	P	P	P	P
Arripidae	-	-	-	P
Aspredinidae	P	-	-	-
Atherinidae	P	P	P	P
Atherinopsidae	-	-	P	-
Auchenipteridae	P	-	-	-
Aulostomidae	-	-	P	-
Bagridae	P	P	-	P
Balistidae	P	P	P	P
Batrachoididae	P	P	P	-
Belonidae	P	P	P	P
Blenniidae	-	P	P	P
Bothidae	P	P	P	P
Bythitidae	-	-	P	-
Callionymidae	-	P	P	P
Carangidae	P	P	P	P
Carcharhinidae	-	P	P	P
Catostomidae	-	-	P	-
Centrarchidae	-	-	P	-
Centriscidae	-	P	-	P
Centrogenyidae	-	P	-	-

Centropomidae	P	P	P	P
Chaetodontidae	P	P	P	P
Chanidae	P	P	-	P
Channidae	-	P	-	-
Characidae	P	-	-	-
Chirocentridae	-	P	-	P
Cichlidae	P	P	P	P
Citharidae	-	-	-	P
Clariidae	P	P	P	-
Claroteidae	P	-	-	-
Clupeidae	P	P	P	P
Congridae	-	-	P	-
Cynoglossidae	P	P	P	P
Cyprinidae	-	P	P	P
Cyprinodontidae	P	P	P	P
Dactylopteridae	P	P	P	P
Dasyatidae	P	P	P	P
Diodontidae	P	-	P	P
Drepaneidae	P	P	-	P
Echeneidae	-	P	P	P
Elassomatidae	-	-	P	-
Eleotridae	P	P	P	P
Elopidae	P	P	P	P
Engraulidae	P	P	P	P
Ephippidae	P	P	P	P
Esocidae	-	-	P	-
Exocoetidae	P	-	-	-
Fistulariidae	-	P	P	P
Fundulidae	-	-	P	-
Galaxiidae	-	-	-	P
Gerridae	P	P	P	P
Ginglymostomatidae	-	-	P	-
Gobiesocidae	-	-	P	-
Gobiidae	P	P	P	P
Gymnotidae	P	-	-	-
Gymnuridae	P	-	P	-
Haemulidae	P	P	P	P
Hemigaleidae	-	P	-	-
Hemiramphidae	P	P	P	P
Hepsetidae	P	-	-	-
Holocentridae	-	-	P	P
Ictaluridae	-	-	P	-
Kraemeridae	-	-	-	P
Kuhliidae	-	-	-	P
Kurtidae	-	-	-	P
Kyphosidae	-	-	P	P
Labridae	-	P	P	P
Labridsomidae	-	-	P	-

Lactariidae	-	P	-	P
Lateolabracidae	-	P	-	-
Latidae	-	P	P	P
Leiognathidae	-	P	-	P
Lepisosteidae	-	-	P	-
Leptobramidae	-	P	-	P
Lethrinidae	-	P	-	P
Lobotidae	P	-	P	P
Loricariidae	P	P	-	-
Lutjanidae	P	P	P	P
Megalopidae	P	P	P	P
Melanotaeniidae	-	-	-	P
Mochokidae	P	-	-	-
Monacanthidae	P	P	P	P
Monodactylidae	P	P	-	P
Moringuidae	-	P	-	P
Mormyridae	P	-	-	-
Moronidae	P	-	-	-
Mugilidae	P	P	P	P
Mullidae	-	P	P	P
Muraenesocidae	-	P	-	P
Muraenidae	P	P	P	P
Myliobatidae	P	P	-	-
Narkidae	-	-	-	P
Nemipteridae	-	-	-	P
Ogcocephalidae	P	-	P	-
Ophichthidae	P	P	P	P
Opistognathidae	-	P	-	-
Orectolobidae	-	P	P	-
Ostraciidae	-	-	P	P
Pallostethidae	-	P	-	P
Paralichthyidae	P	P	P	P
Pegasidae	-	P	-	P
Pempheridae	-	P	-	P
Percichthyidae	-	-	-	P
Percophidae	-	-	-	P
Pimelodidae	P	-	-	-
Platycephalidae	-	P	-	P
Plesiopidae	-	P	-	-
Pleuronectidae	-	-	-	P
Plotosidae	-	P	-	P
Poeciliidae	P	P	P	P
Polynemidae	P	P	P	P
Pomacanthidae	-	-	P	P
Pomacentridae	-	P	P	P
Pomatomidae	P	P	-	P
Priacanthidae	-	-	-	P
Pristidae	-	P	-	-



Pristigasteridae	P	P	-	P
Psettodidae	P	P	-	-
Pseudaphritidae	-	-	-	P
Pseudomugilidae	-	P	-	P
Ptereleotridae	-	-	-	P
Rachycentridae	-	P	-	P
Rhinobatidae	P	P	-	P
Rivulidae	-	-	P	-
Scaridae	-	-	P	P
Scatophagidae	-	P	-	P
Schilbeidae	P	-	-	-
Sciaenidae	P	P	P	P
Scombridae	P	P	-	P
Scorpaenidae	-	P	P	P
Serranidae	P	P	P	P
Siganidae	-	P	-	P
Sillaginidae	-	P	-	P
Siluridae	-	P	-	-
Soleidae	P	P	P	P
Sparidae	P	P	P	P
Sphyraenidae	P	P	P	P
Sphyrnidae	-	P	-	P
Sternopygidae	P	-	-	-
Stromateidae	-	P	-	-
Synanceiidae	-	P	-	P
Synbranchidae	-	-	-	P
Syngnathidae	P	P	P	P
Synodontidae	P	P	P	P
Terapontidae	-	P	-	P
Tetraodontidae	P	P	P	P
Tetrarogidae	-	P	-	P
Toxotidae	-	P	-	P
Triacanthidae	P	P	-	P
Trichiuridae	P	P	-	P
Triglidae	P	-	P	-
Uranoscopidae	-	P	-	P
Urolophidae	-	-	P	P

Supplement S4. The 64 families of mangrove-dwelling fish evaluated as using nekton as a major part of their diet

Habit group	Family
<i>Ambush</i>	Achiridae
	Bothidae
	Citharidae
	Paralichthyidae
	Pleuronectidae
	Psettodidae
	Uranoscopidae
	Platycephalidae
	Antennariidae
	Batrachoididae
	Channidae
	Congridae
	Eleotridae
	Muraenidae
	Ophichthidae
	Percophidae
	Synanceiidae
Tetrarogidae	
<i>Forager</i>	Ginglymostomatidae
	Muraenesocidae
	Orectolobidae
	Priacanthidae
	Triglidae
	Scorpaenidae
	Hemigaleidae
	Centrogyidae
	Plesiopidae
	Characidae
	Hepsetidae
	Lethrinidae
	Lobotidae
	Moronidae
	Centropomidae
	Latidae
	Lutjanidae
	Polynemidae
	Sciaenidae
	Serranidae
	Nemipteridae
	Percichthyidae
	Synodontidae
Lactariidae	
Centrarchidae	

	Kuhliidae
	Kurtidae
	Lateolabracidae
<hr/>	
<i>Cursorial</i>	Elopidae
	Scombridae
	Rachycentridae
	Carcharhinidae
	Arripidae
	Leptobramidae
	Trichiuridae
	Carangidae
	Chirocentridae
	Megalopidae
	Pomatomidae
	Sphyrnidae
<hr/>	
<i>Static habituator</i>	Esocidae
	Fistulariidae
	Lepisosteidae
	Aulostomidae
	Belonidae
	Sphyraenidae
<hr/>	