

Bio-geochemical cycles in acidic soils from a granitic watershed

M.C. PIERRET*, P. STILLE, F. CHABAUX, D. VIVILLE,
J. PRUNIER AND D. LEMARCHAND

CGS/EOST, 1 rue Blessig - 67000 Strasbourg-France

(*correspondence: pierret@illite.u-strasbg.fr)

(pstille@illite.u-strasbg.fr, fchabaux@illite.u-strasbg.fr,

prunier@illite.u-strasbg.fr, lemarcha@illite.u-strasbg.fr)

To assess transfers between vegetation, waters and minerals we studied soil litter, trees, acidic soils and soil solutions from a small granitic watershed (<http://ohge.u-strasbg.fr>). Leaching experiments have been performed on soils in order to recover the adsorbed elements or those fixed in acid soluble mineral phases such as Fe- hydroxides and phosphates. The isotope data of the leachates point to the presence of important quantities of litter derived elements. With the exception of the litter, the soil (upper 50cm) is depleted in Ca due to dissolution of apatite and plagioclase. Rhabdophane appears as secondary replacement mineral of apatite. It caught up some of the mobilized LREE. The leached Ca has been removed out of this soil by soil solutions and plant uptake. Ca and Mg cycles in the upper soil profile are dominated by the vegetation causing Ca and LREE depletion in the soil solutions. Some of the LREE depletion might also be due to precipitation of rhabdophane. At the watershed scale the biological Ca removal flux is of the same order of magnitude as the weathering flux. The Sr isotopic data further indicate that also ion exchange plays a role in the transfer and migration of Ca, Mg and K. The Na cycle, however, is dominated by mineral/ solution interactions with little influence of ionic exchanges. We therefore distinguish between 1) mainly "biogenic" elements like Ca, Mg, Sr and Ba (entering the vegetation cycle) and 2) elements like Na or Al which are mainly controlled by mineral dissolution. The knowledge of all these processes finally allows to understand the geochemical signatures of stream water (especially their seasonal variations) and to evaluate the weathering rates at a watershed.

Episodic, mafic crust formation in the Slave craton, Canada

A. PIETRANIK^{1,2}, C.J. HAWKESWORTH², C.D. STOREY²,
T.I.S. KEMP³, K.N. SIRCOMBE⁴, M.J. WHITEHOUSE⁵ AND
W. BLEEKER⁶

¹Institute of Geological Sciences, University of Wrocław,
50-205 Wrocław, Poland

(*correspondence: anna.pietranik@ing.uni.wroc.pl)

²Department of Earth Sciences, University of Bristol, Bristol
BS8 1RJ, UK

³School of Earth and Environmental Sciences, James Cook
University, Townsville QLD 4811, Australia

⁴Onshore Energy and Minerals Division, Geoscience
Australia, Canberra, Australia

⁵Swedish Museum of Natural History, Stockholm, Sweden

⁶Geological Survey of Canada, Ottawa, Ontario, Canada

Hf model ages of zircons provide information on the timing of new crust extraction from mantle. However, many model ages are hybrid due to mixing in the sedimentary environment. Zircons with mantle $\delta^{18}\text{O}$ values (5.0-5.6‰) [1] are more likely to preserve Hf model ages that reflect actual crust forming events.

Hf and O isotopes have been analysed in 3.9-2.8 Ga detrital zircons from the ~2.8 Ga Mesoarchean cover succession in the Slave craton, Canada [2]. The zircons with mantle-like δO^{18} form two linear arrays in zircon crystallization age versus initial ϵHf plots consistent with two episodes of crust formation at ~3.45 Ga and ~3.75 Ga. Slopes of the linear arrays correspond to $^{176}\text{Lu}/^{177}\text{Hf}$ ratios of ~0.022 suggesting that the sources of the magmas from which zircons crystallized was mafic in composition. The zircons with the lowest initial ϵHf through time all have elevated $\delta^{18}\text{O}$ values, but they also scatter around a linear array with the slope corresponding to $^{176}\text{Lu}/^{177}\text{Hf}$ ratio of ~0.022 consistent with mafic crust derivation from the mantle at ~4.4-4.5 Ga. Zircons that crystallized during crust formation and crustal recycling events both show a range of initial ϵHf and often elevated $\delta^{18}\text{O}$ consistent with derivation of magmas from heterogeneous crust composed of both unaltered, igneous and weathered, probably sedimentary rocks. Mafic crust generated in three episodes in the Slave Province was the source of magmas throughout the Archean, and zircons from Gondwana indicate that similar unaltered mafic crust could have had even longer residence times of over 1.5 Ga [3].

AP acknowledges grants from Polish Science Foundation and SYNTHESIS.

[1] Valley *et al.* (2005) *Contrib. Min.Pet.* **150**, 561-580.

[2] Sircombe *et al.* (2001) *EPSL* **189**, 207-220. [3] Kemp *et al.* (2006) *Nature* **439**, 580-583.