

RESEARCH PAPER

Nocturnal versus diurnal CO₂ uptake: how flexible is *Agave angustifolia*?

Klaus Winter^{1,*}, Milton Garcia¹ and Joseph A. M. Holtum^{1,2}

¹ Smithsonian Tropical Research Institute, PO Box 0843-03092, Balboa, Ancon, Republic of Panama

² Tropical Biology, James Cook University, Townsville, Queensland 4811, Australia

* To whom correspondence should be addressed. E-mail: winterk@si.edu

Received 8 October 2013; Revised 31 January 2014; Accepted 10 February 2014

Abstract

Agaves exhibit the water-conserving crassulacean acid metabolism (CAM) photosynthetic pathway. Some species are potential biofuel feedstocks because they are highly productive in seasonally dry landscapes. In plants with CAM, high growth rates are often believed to be associated with a significant contribution of C₃ photosynthesis to total carbon gain when conditions are favourable. There has even been a report of a shift from CAM to C₃ in response to overwatering a species of *Agave*. We investigated whether C₃ photosynthesis can contribute substantially to carbon uptake and growth in young and mature *Agave angustifolia* collected from its natural habitat in Panama. In well-watered plants, CO₂ uptake in the dark contributed about 75% of daily carbon gain. This day/night pattern of CO₂ exchange was highly conserved under a range of environmental conditions and was insensitive to intensive watering. Elevated CO₂ (800 ppm) stimulated CO₂ fixation predominantly in the light. Exposure to CO₂-free air at night markedly enhanced CO₂ uptake during the following light period, but CO₂ exchange rapidly reverted to its standard pattern when CO₂ was supplied during the subsequent 24 h. Although *A. angustifolia* consistently engages in CAM as its principal photosynthetic pathway, its relatively limited photosynthetic plasticity does not preclude it from occupying a range of habitats, from relatively mesic tropical environments in Panama to drier habitats in Mexico.

Key words: *Agave*, biofuel, climate change, crassulacean acid metabolism, C₃ photosynthesis, CO₂ response, drought stress, temperature response.

Introduction

Most, if not all, of the approximately 200 species in the New World genus *Agave* (family Asparagaceae, subfamily Agavoideae; Chase *et al.*, 2009; Govaerts *et al.*, 2013) exhibit the nocturnal uptake of CO₂ and accumulation of malic acid characteristic of crassulacean acid metabolism (CAM) (Nobel, 2003, 1996a). In addition to a water-use-efficient carbon metabolism, these archetypal dry-land CAM plants sport a xerophytic, water-conserving morphology that includes succulent, angled, persistent leaves with thick cuticles, sunken stomata, and a rosette leaf configuration around a central stem that channels rain water and condensate to the base of the roots (Gentry, 1982).

As biomass becomes an increasingly valuable commodity for energy generation, there is realization that the seasonally dry or semi-arid landscapes inhabited by *Agave*, whilst not optimal for growing traditional water-demanding food crops, may nonetheless be suitable for biomass generation (Borland *et al.*, 2009; Chambers and Holtum, 2010; Davis *et al.*, 2011; Holtum *et al.*, 2011; Yan *et al.*, 2011). In such landscapes, water-use-efficient *Agave* can accumulate biomass at annual rates that approach those produced by C₄ plants like sugar cane and *Miscanthus* in higher rainfall regions (Nobel, 1991, 1996a; Somerville *et al.*, 2010). For example, productivities

Abbreviations: CAM, crassulacean acid metabolism; PFD, photon flux density.

© The Author 2014. Published by Oxford University Press on behalf of the Society for Experimental Biology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

of 25, 35, and 47–50 Mg dry weight ha⁻¹ year⁻¹ have been reported for the CAM species *Agave tequilana*, *Ananas comosus* (pineapple) and *Opuntia ficus-indica*, respectively (Nobel, 1996a). *A. tequilana* is now being trialed in the seasonally dry Australian tropics as a biofuel feedstock (Chambers and Holtum, 2010; Holtum *et al.*, 2011).

The extent and capacity for day-time CO₂ fixation in well-watered agaves tends to be limited, but its expression differs among species (see Nobel, 2003, for a review). Interestingly, well-watered plants of mature *Agave deserti* did not exhibit afternoon CO₂ fixation, but when ‘overwatered’ (watered daily for 10 weeks) they switched to overwhelmingly C₃ photosynthesis (Hartsock and Nobel, 1976).

Using continuous whole-plant gas exchange, we explored here the potential for photosynthetic plasticity in *Agave angustifolia*, a putative wild ancestor of the agronomically significant species *Agave fourcroydes* and *A. tequilana* (Gentry, 1982; Colunga-García Marín *et al.*, 1999). *A. angustifolia* was chosen because, across its range from Mexico to Panama, it is found in habitats as diverse as coastal dunes at sea level to oak-pine forests at 2200 m (García-Mendoza and Chiang, 2003), and it grows in Panama (the site of this study) in relatively mesic environments where CO₂ fixation in the light is expected to be more favoured than in drier habitats.

In order to determine photosynthetic pathway plasticity of *A. angustifolia*, we examined 24 plants. Our goal was to explore conditions under which plants would markedly upregulate C₃ photosynthetic CO₂ uptake in the light. In two young plants, net CO₂ exchange was continuously monitored for 234 and 281 day/night cycles, during which the plants were exposed to a range of perturbations (light, temperature, CO₂, and watering regime) that have been reported to affect CO₂ uptake in the light in other CAM plants. Furthermore, net CO₂ exchange was monitored in two extremely well-watered mature plants in a naturally illuminated gas-exchange chamber for up to 16 d each, in order to document the effect of natural day-to-day variation in photon flux density (PFD) on light and dark CO₂ fixation. Thirdly, a total of 20 young, well-watered plants were grown under two CO₂ concentrations and two nutrient regimes to manipulate the relative contributions of day and night CO₂ fixation to growth. Although the longer than 200 d gas-exchange experiments were not replicated in the strictest sense because of their duration, all experiments taken together permitted a reasonable assessment of the degree of phenotypic photosynthetic plasticity in *A. angustifolia*. The results indicated that *A. angustifolia* exhibits a conserved carbon fixation strategy rather than using the light and dark options of CO₂ uptake in a highly flexible manner.

Materials and methods

Plant material

A. angustifolia Haw. was collected from Playa Majagual, Panamá (8°43'N, 79°45'E), and grown outdoors in forest topsoil at the Smithsonian Tropical Research Institute, Santa Cruz Experimental Research Facility, Gamboa, Republic of Panama (9°07'N, 79°42'W). Opinions differ as to whether *A. angustifolia* is a synonym of *Agave vivipara* L. or whether the two are distinct species (Wijnands, 1983;

García-Mendoza and Chiang, 2003; Govaerts *et al.*, 2013). Vouchers of the *Agave* examined in this study were deposited in the herbarium of the University of Panama (JARANDA 4484A and 4484B).

Measurement of CO₂ exchange in the laboratory

Bulbils of between 5 and 10 cm in height, comprising two to three leaves, were enclosed in a Perspex cuvette (internal dimensions 11 × 11 × 10 or 20 × 20 × 15 cm) by passing the base of a plantlet through a hole in the cuvette base and sealing the plantlet–cuvette interface with a non-porous synthetic rubber sealant (Terostat VII; Henkel-Teroson, Heidelberg, Germany). The root-containing base of the plantlet outside the cuvette was planted in a 1 litre pot containing potting mix (Cactus, Palm and Citrus Soil; Miracle-Gro Lawn Products, Marysville, OH, USA) and 2 g of Osmocote Plus fertilizer (Scotts-Sierra Horticultural Products, OH, USA).

The gas-exchange cuvette was located inside a controlled-environment chamber (Environmental Growth Chambers, OH, USA) operating under 12 h light (28 °C)/12 h dark cycles (17 or 22 °C as specified). PFD was measured at the top of the cuvette. Air containing 200, 400, or 800 ppm CO₂ was generated by a CO₂/CO₂-free-air mixing unit (Walz GmbH, Effeltrich, Germany). Net CO₂ exchange of plantlets was measured in flow-through gas-exchange systems consisting of Walz components and LI-6252 CO₂ analysers (Li-Cor, Lincoln, NE, USA) (Holtum and Winter 2003). Normal watering involved supplying water at least once every second day, and intensive watering involved supplying water twice per day. Drought treatments were imposed by withholding irrigation altogether.

Measurement of whole-plant CO₂ exchange under natural light

A mature plant, established in forest topsoil containing 50 g of Osmocote Plus fertilizer (Scotts-Sierra Horticultural Products), in a 190 l pot was placed inside a ventilated, naturally illuminated chamber constructed of glass panels and an aluminium framework (internal volume 8.8 m³). A blower (model 4C054; Grainger Industrial Supply, OH, USA) supplied external air to the chamber at 10.5 m³ min⁻¹. Within the chamber, air was circulated by four fans and temperature was regulated by a split air-conditioning system (model V1124C2H; Innovair, FL, USA). Whole-plant gas exchange was quantified at 30 min intervals from the rate at which the CO₂ concentration inside the chamber changed when air flow into the chamber was blocked for 5 min, thereby converting the chamber into a closed system. Changes in the CO₂ concentration inside the chamber were measured using a LI-7500 open-path CO₂ analyser (Li-Cor). Calculations of net CO₂ exchange were based on chamber volume that had been corrected for the volumes of the pot, plant, and equipment inside the chamber, and the rate at which the CO₂ concentration changed during the period when the chamber was isolated. CO₂ measurements were corrected for changes of temperature and humidity. Measurements of PFD were taken outside the chamber. PFD inside the chamber was approximately 15% below that outdoors. For further details of methods, see Winter *et al.* (2009). The experiment was repeated for a second mature plant (data not shown).

Growth of plants at 280 and 800 ppm CO₂

Twenty plants in 19 l pots were grown with daily watering in forest topsoil for 166 d inside two naturally illuminated glasshouses (internal volume 37.8 m³ each). Five of the 10 pots in each chamber were supplemented with 8 g of Osmocote Plus fertilizer (Scotts-Sierra Horticultural Products). One glasshouse was maintained at 280 ± 10 ppm (range) CO₂ by passing chamber air through soda lime to lower [CO₂]. An above-ambient CO₂ concentration of 800 ± 10 (range) ppm was achieved in the second glasshouse by releasing pulses of CO₂ gas into the chamber from a high-pressure cylinder in conjunction with a feedback control system. Within each glasshouse, air was circulated by five fans, and a split air-conditioning system maintained temperatures at close to ambient (Cernusak *et al.*, 2011).

Titrate acidity

Leaf discs punched from the centre of fully expanded leaves using a cork borer at the end of the light and dark periods were frozen in liquid nitrogen. Organic acids were extracted by sequentially boiling samples in 50% ethanol and water for 5 min. Extracts were cooled to room temperature and titrated with 10 mM KOH to pH 6.5 (Holtum *et al.*, 2004).

Stable isotope analysis

The $\delta^{13}\text{C}$ values of finely ground homogenous powder from the pooled dried leaves of whole plants were measured in an isotope ratio mass spectrometer (Delta V; Thermo Fisher Scientific) in the Stable Isotope Laboratory of the Smithsonian Tropical Research Institute. The abundance of ^{13}C in each sample was calculated relative to the abundance of ^{13}C in standard CO_2 that had been calibrated against Pee Dee belemnite (*Belemnitella americana*). Relative abundance was determined using the relationship:

$$\delta^{13}\text{C} (\text{‰}) = \left[\left(\frac{^{13}\text{C}/^{12}\text{C} \text{ of sample}}{^{13}\text{C}/^{12}\text{C} \text{ of standard}} \right) - 1 \right] \times 1000.$$

The $\delta^{13}\text{C}$ values of two C_4 plant species, *Saccharum spontaneum* and *Portulaca oleracea*, grown in each glasshouse and outside in the open air were used to correct for differences in the $\delta^{13}\text{C}$ value of the source CO_2 (Cernusak *et al.*, 2011). The CO_2 purchased for CO_2 enrichment was from a natural CO_2 spring and a correction of 2‰ was applied.

Results

CO_2 fixation in the dark (CAM phase I; Osmond, 1978) contributed 78% of the carbon gain in a young well-watered

A. angustifolia for which CO_2 uptake was monitored continuously during 281 consecutive 12h light/12h dark cycles (Fig. 1). The predominance of dark fixation was maintained under 12h PFDs of $400 \mu\text{mol m}^{-2} \text{s}^{-1}$ ($17.3 \text{ mol m}^{-2} \text{d}^{-1}$), $1500 \mu\text{mol m}^{-2} \text{s}^{-1}$ ($64.8 \text{ mol m}^{-2} \text{d}^{-1}$), and $2300 \mu\text{mol m}^{-2} \text{s}^{-1}$ ($99.4 \text{ mol m}^{-2} \text{d}^{-1}$).

Uptake of CO_2 in the dark remained the principal contributor to net carbon gain during 234 day/night cycles of gas exchange when a young *A. angustifolia* was subjected to 300 or 900 $\mu\text{mol m}^{-2} \text{s}^{-1}$ light; 200, 400 or 800 ppm CO_2 ; when well-watered, overwatered, or non-watered; and when night temperatures were 17 or 22 °C (Fig. 2). CO_2 uptake in the afternoon (CAM phase IV) was also always present, whereas the contribution of dawn CO_2 fixation (CAM phase II) to plant carbon gain was minimal. Net CO_2 loss was consistently observed during midday stomatal closure (CAM phase III) except for small CO_2 gains on those days when CO_2 was withheld during the dark while the plant was exposed to 800 ppm CO_2 . Over the course of the experiment (Fig. 2), the contributions to net carbon gain by phases I, II, III, and IV were 78.5, 1.7, -4.2, and 24%, respectively.

The contributions of light and dark CO_2 uptake to net carbon gain under 200 or 800 ppm CO_2 differed (Fig. 2). When the CO_2 concentration was reduced from 400 to 200 ppm, CO_2 uptake at a PFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$ was reduced by 62% during phase IV and by 16% in the dark. The reductions were 56 and 25%, respectively, at 900 $\mu\text{mol m}^{-2} \text{s}^{-1}$. In contrast, when the atmospheric CO_2 concentration was increased from 400 to 800 ppm, CO_2 uptake by the plant at $300 \mu\text{mol m}^{-2} \text{s}^{-1}$ increased by




Fig. 1. Net CO_2 balance of the shoot of a young individual of *A. angustifolia* during 281 consecutive 12h light/12h dark cycles. Carbon balances are shown for the light and dark periods. The following treatments were applied: (A) PFD was $400 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ during the initial 103 d, $2300 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ between d 104 and 122, and $1500 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ thereafter. (B) Irrigation was increased from once every 2 d to twice per day between d 85 and 122 ('overwatered').




Fig. 2. Net CO₂ balance of the shoot of a young individual of *A. angustifolia* during 234 consecutive 12h light/12h dark cycles. Carbon balances are shown for the four phases of the 24 h CAM cycle. As indicated by the four horizontal bars in the upper part of the figure, the following treatments were imposed in the course of the experiment: (A) PFD was 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ up to d 98, and increased to 900 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ thereafter. (B) Temperature during the dark period was lowered from 22 °C (standard night-time condition) to 17 °C between d 77 and 93. (C) Irrigation was stopped between d 153 and 209, and irrigation was increased from once to twice per day between d 100 and 121 ('overwatered'). (D) Ambient [CO₂] was 200, 400, or 800 ppm as indicated by yellow, red, and blue, respectively. The two green arrows indicate net CO₂ exchange in the light period following exposure of the plant to CO₂-free air during the previous night (d 114, 115, and 142). Data for the initial 9 d of the experiment are not shown. (This figure is available in colour at *JXB* online.)

64% during phase IV and remained unchanged during phase I. At a PFD of 900 $\mu\text{mol m}^{-2} \text{s}^{-1}$, the increases in CO₂ uptake during phases IV and I were 73 and 10%, respectively

Withholding water from the plant for 54 d also differentially influenced CO₂ fixation in the light and in the dark (Fig. 2). Overall net carbon gain by the plant was initially stimulated but was subsequently reduced as water in the pot became limiting. During the initial 34 d without watering, as CO₂ uptake increased by 46% in the dark and decreased by 17% during phase IV in the light, the contribution of nocturnal uptake to net carbon gain rose from 74 to 85%. Subsequently, CO₂ exchange fell during the light and the dark. The decrease was proportionally less in the dark such that, at the end of the drought treatment, dark fixation contributed 88% to net carbon gain. In contrast to drought, intensive watering of the plant for 30 d did not affect the relative contributions of light and dark CO₂ fixation to carbon gain, nor did decreasing the night temperature from 22 to 17 °C.

CO₂ assimilation in the light was upregulated when CO₂ was removed from the air supply during the preceding dark period (Fig. 2, d 141 and 142). Fig. 3 details how both the extent and pattern of CO₂ exchange in the light differed following exposure to CO₂-free air during the night. CO₂ was assimilated throughout the light principally via the contribution of

an extended phase IV (Fig. 3B). Phase III was transient, and phase II did not increase in duration although the rate of CO₂ uptake did increase. During the subsequent dark/light cycle, CO₂ exchange returned to the patterns observed prior to the CO₂-free treatment (Fig. 3C).

Nocturnal CO₂ uptake was the principal contributor to carbon gain in fully mature, well-watered *A. angustifolia* grown outdoors under natural sunlight (Fig. 4). Day-time CO₂ uptake occurred mainly during the late afternoon (phase IV). The contribution of phase II CO₂ uptake was variable but generally small, whereas net CO₂ loss was consistently observed during phase III midday stomatal closure.

Nocturnal CO₂ uptake decreased substantially following overcast days (Fig. 4) For example, the extremely overcast d 4 light period was followed by a night during which there was no net CO₂ uptake. Less extreme examples of this trend were evident on d 2, 8, and 12. On days following overcast days, CO₂ uptake during the afternoon tended to be more pronounced.

Fig. 5 quantifies the relationships between day-time and night-time CO₂ exchange and daily PFD for the mature *A. angustifolia* illustrated in Fig. 4. Nocturnal CO₂ uptake was correlated with the integrated PFD during the preceding light period and contributed predominately to net CO₂




Fig. 3. Gas exchange by a shoot of a well-watered young individual of *A. angustifolia* (d 141, 142 and 143 of the shoot shown in Fig. 2.) supplied with 800 ppm CO₂ during the dark and during the light on d 141 (A), supplied with 0 ppm CO₂ during the dark and 800 ppm CO₂ during the light on d 142 (B), and supplied with 800 ppm CO₂ during the dark and during the light on d 143 (C). Shading indicates the night. Day-time PFD was 900 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$.

uptake at all light intensities that supported positive daily carbon gain. Following sunny days, when the integrated PFD exceeded about $30 \text{ mol m}^{-2} \text{ d}^{-1}$, CO₂ uptake at night was saturated, providing 70–85% of the daily carbon gain. Below $30 \text{ mol m}^{-2} \text{ d}^{-1}$, total carbon gain fell and the proportional contribution of nocturnal CO₂ uptake to 24h carbon gain rose. Day-time CO₂ exchange became negative at around $21 \text{ mol m}^{-2} \text{ d}^{-1}$, whereas night-time CO₂ exchange became negative at about $10 \text{ mol m}^{-2} \text{ d}^{-1}$.

The biomass of *A. angustifolia* grown in unfertilized soil at 800 ppm CO₂ was double that of plants grown at 280 ppm CO₂, whereas leaf acidities were similar on both mass and leaf-area bases (Table 1). Fertilization increased the biomass 2.5-fold in plants grown at 280 ppm CO₂ and 1.7-fold in plants grown at 800 ppm CO₂. For both unfertilized and fertilized treatments, the $\delta^{13}\text{C}$ values for plants at 800 ppm CO₂ were 1.8‰ more negative than for plants grown at 280 ppm CO₂. In comparison to unfertilized plants, fertilized




Fig. 4. PFD (upper panels) and net CO₂ exchange (lower panels) for the shoot (aboveground parts that included 51 leaves) of a mature *A. angustifolia* prior to the emergence of the reproductive stalk. Measurements were conducted from 27 July 2011 (d 1) to 11 August 2011 (d 16) in a naturally illuminated gas-exchange chamber. Shading indicates night.

plants exhibited greater nocturnal accumulation of H⁺ per unit leaf area, but H⁺ accumulation per unit leaf mass was unchanged.

Discussion

Despite an ability to occupy contrasting habitats, photosynthetic flexibility in *A. angustifolia* does not appear to be exceptional in terms of the proportional contributions to carbon gain of CO₂ uptake in the dark and light. Nocturnal CO₂ uptake was the principal source of carbon in mature *A. angustifolia* and in young plants once they had established. The proportional contribution to daily carbon gain of nocturnal CO₂ uptake remained a remarkably consistent 70–85%, although the amount of CO₂ fixed per day by well-watered plants varied with light intensity and nutrient status. In contrast to *A. deserti* (Hartsock and Nobel, 1976), the day/night pattern of CO₂ exchange in *A. angustifolia* did not shift towards a C₃ pattern when the supply of water was effectively unlimited.

Drought affected plant carbon gain and increased the proportional contribution of nocturnal CO₂ uptake to it (Fig. 2). Two weeks after the cessation of irrigation, drought stress manifested itself as a continuous decline in light CO₂ fixation. Most importantly, the initial 25 d of the decline of CO₂ uptake in the light was accompanied by an increase in the

rate of dark CO₂ fixation. This drought-induced upregulation of CAM is a typical feature of facultative CAM. Facultative CAM or facultative components of CAM are not restricted to metabolically flexible annuals such as *Mesembryanthemum crystallinum* (Winter and von Willert, 1972) and *Calandrinia polyandra* (Winter and Holtum, 2011), and perennials such as some species of *Clusia* (Winter et al., 2009), but have also been observed in juveniles of constitutive CAM succulents such as *O. ficus-indica* and *Opuntia elatior* (Winter et al., 2008, 2011).

An attempt to force *A. angustifolia* into a C₃-like photosynthetic pattern by exposing it to 800 ppm CO₂ was partially successful in that daily carbon gain was enhanced and the proportional contribution of CO₂ uptake in the light rose, for example from 29 to 40% during an 11 d treatment (Fig. 2). The ability to maintain this pattern of CO₂ exchange was confirmed following a 166 d exposure to 800 ppm CO₂ after which tissue δ¹³C values were close to those predicted by Winter and Holtum (2002) for a 40% contribution to carbon gain of CO₂ fixation in the light.

In the short-term 800 ppm CO₂ fumigation treatments, CO₂ uptake in the dark was not or was only slightly enhanced (Fig. 2); in the longer-term experiment, there was no enhancement (Table 1), as nocturnal acidification remained unchanged. The contribution of nocturnal CO₂ fixation to carbon gain in CAM tissues is variably responsive to environmental stimuli,




Fig. 5. (A) Day-time (open symbols) and night-time (closed symbols) CO₂ balances in response to daily integrated PFD of the mature *A. angustifolia* plant studied in Fig. 4. (B) Percentage contribution of day-time (open symbols) and night-time (closed symbols) CO₂ uptake to 24 h carbon gain. The symbol with open and closed halves represents d 4, an overcast day during which CO₂ balances during the light and the dark were negative.

which include night temperature, day-length, light intensity, and atmospheric CO₂ concentration (Neales, 1973; Drennan and Nobel, 2000; Borland *et al.*, 2011). Increased CO₂ assimilation in the light but not the dark has been reported in *Ananas* grown at 700 ppm CO₂ under a 30/20 °C day/night regime that was optimal for nocturnal CO₂ uptake at ambient CO₂ (Zhu *et al.*, 1999). However, when *Ananas* was grown at higher night temperatures that were less than optimal for dark CO₂ uptake at ambient CO₂, the growth of plants at 700 ppm CO₂ increased CO₂ uptake in both the light and the dark. Increases in both light and dark CO₂ fixation following exposure to high concentrations of atmospheric CO₂ have been reported for *A. deserti* (Graham and Nobel, 1996) and

Table 1. Dry mass, nocturnal increase in leaf tissue acidity, and $\delta^{13}\text{C}$ value (whole shoot) of *A. angustifolia* grown at 280 and 800 ppm CO₂

Values are means \pm standard deviation (n=5), except for $\delta^{13}\text{C}$ value at 280 ppm+fertilizer, where n=3. Means on the same row with different superscript letters differ significantly (two-way analysis of variance followed by Fisher's least significant difference test, P<0.05).

Parameter	CO ₂ concentration			
	280 ppm		800 ppm	
	- fertilizer	+ fertilizer	- fertilizer	+ fertilizer
Total dry mass (g)	26 \pm 4 ^A	65 \pm 12 ^B	51 \pm 5 ^C	88 \pm 11 ^D
Nocturnal H ⁺ increase:				
$\mu\text{mol g}^{-1}$	226 \pm 40 ^A	210 \pm 14 ^{AB}	184 \pm 8 ^B	207 \pm 31 ^{AB}
fresh weight				
$\mu\text{mol cm}^{-2}$	79 \pm 12 ^A	96 \pm 4 ^B	77 \pm 6 ^A	93 \pm 11 ^B
$\delta^{13}\text{C}$ (‰)	-14.6 \pm 0.2 ^A	-14.0 \pm 0.2 ^B	-16.4 \pm 0.2 ^C	-15.8 \pm 0.2 ^D

Agave salmiana (Nobel, 1996b; Nobel *et al.*, 1996), and for the stem succulents, *O. ficus-indica* (Nobel and Israel, 1994) and *Stenocereus queretaroensis* (Nobel, 1996b).

A. angustifolia could be shifted towards a C₃-like light-only CO₂ uptake pattern by an extreme treatment that required withholding CO₂ during the dark and supplying 800 ppm CO₂ in the light (Fig. 3). In the light, the duration of phase IV increased at the expense of phase III, presumably because small amounts of acid formed at night would be rapidly consumed and the inhibition of stomatal opening by the resulting high internal CO₂ concentration would be transitory. In the 24 h cycle during which CO₂-free air was supplied at night, daily carbon gain fell because the increase in CO₂ gain in the light did not offset the lack of uptake of atmospheric CO₂ during the night.

The stimulation of light fixation following exposure to CO₂-free air at night lasted only during the light period following the treatment. Remarkably, no evidence of a metabolic memory of the CO₂-free treatment was evident in the subsequent night and the day that followed it.

It has been suggested that the feasibility of cultivating strong-CAM plants such as *Agave* or *Opuntia* for biofuel feedstock in seasonally dry environments could be improved by developing plants that would fix a greater proportion of CO₂ in the light during the moister parts of the year (Borland *et al.*, 2011). In effect, the proposal is to push the proportional contribution of CO₂ fixation in the light into the vicinity of 50–60%, a proportion that large surveys of CAM plants have revealed as being uncommon in the natural environment (Winter and Holtum, 2002; Crayn *et al.*, 2004; Silvera *et al.*, 2010).

As in *A. angustifolia*, 24 h CO₂ exchange by the most highly productive CAM species grown in commercial plantations in warm and temperate subtropical dry-land environments, *A. tequilana*, *O. ficus-indica*, *A. salmiana* and *Agave mapisaga*, is dominated year round by nocturnal CO₂ fixation (Nobel,

1996a; Nobel *et al.*, 1992; Pimienta-Barrios *et al.*, 2001, 2006). In *A. tequilana*, environmental factors that limited growth and productivity in the field were water during the cool dry winter and PAR during the warm wet summer (Nobel and Valenzuela, 1987), not CO₂ fixation in the light. Under these field limitations, a shift towards an increase in the capacity for CO₂ fixation in the light is unlikely to result in significantly increased productivity. Rather, productivity might be expected to be higher if plants were grown at sites that were less cloudy in the summer and more moist in the winter.

Proposals to grow *Agave* as biofuel feedstocks in seasonally dry regions have emphasized their suitability for so-called marginal lands (Borland *et al.*, 2009; Somerville *et al.*, 2010; Davis *et al.*, 2011). Land that is marginal for growing traditional crops may well support high growth rates of water-use efficient, high-temperature-tolerant CAM species, but it remains to be seen whether *Agave* can produce commercially relevant yields on truly marginal lands that are nutrient poor and severely water limited.

Acknowledgements

The authors acknowledge the contributions of J. Aranda who grew and maintained the plants and A. Virgo who drew the illustrations. The research was supported by funds from the Smithsonian Tropical Research Institute. J.A.M.H. was supported by the JCU School of Marine and Tropical Biology.

References

- Borland AM, Griffiths H, Hartwell J, Smith JAC. 2009. Exploiting the potential of plants with crassulacean acid metabolism for bioenergy production on marginal lands. *Journal of Experimental Botany* **60**, 2879–2896.
- Borland AM, Zambrano VAB, Ceusters J, Shorrock K. 2011. The photosynthetic plasticity of crassulacean acid metabolism: an evolutionary innovation for sustainable productivity in a changing world. *New Phytologist* **191**, 619–633.
- Cernusak LA, Winter K, Martínez C, Correa E, Aranda J, Garcia M, Jaramillo C, Turner BL. 2011. Responses of legume versus nonlegume tropical tree seedlings to elevated CO₂ concentration. *Plant Physiology* **157**, 372–385.
- Chambers D, Holtum JAM. 2010. *Feasibility of Agave as a feedstock for biofuel production in Australia*. Canberra: Rural Industries Research and Development Corporation.
- Chase MW, Reveal JL, Fay MF. 2009. A subfamilial classification for the expanded asparagalean families Amaryllidaceae, Asparagaceae and Xanthorrhoeaceae. *Botanical Journal of the Linnean Society* **161**, 132–136.
- Colunga-GarcíaMarín P, Coello-Coello J, Eguiarte LE, Piñero D. 1999. Isozymatic variation and phylogenetic relationships between henequén (*Agave fourcroydes*) and its wild ancestor *A. angustifolia* (Agavaceae). *American Journal of Botany* **86**, 115–123.
- Crayn DM, Winter K, Smith JAC. 2004. Multiple origins of crassulacean acid metabolism and the epiphytic habit in the Neotropical family Bromeliaceae. *Proceedings of the National Academy of Sciences, USA* **101**, 3703–3708.
- Davis SC, Dohleman FG, Long SP. 2011. The global potential for *Agave* as a biofuel feedstock. *GCB Bioenergy* **3**, 68–78.
- Drennan PM, Nobel PS. 2000. Responses of CAM species to increasing atmospheric CO₂ concentrations. *Plant, Cell & Environment* **23**, 767–781.
- García-Mendoza A, Chiang F. 2003. The confusion of *Agave vivipara* L. and *A. angustifolia* Haw., two distinct taxa. *Brittonia* **55**, 82–87.
- Gentry HS. 1982. *Agaves of continental North America*. Tucson: University of Arizona Press.
- Govaerts R, Zonneveld BJM, Zona SF. 2013. *World Checklist of Asparagaceae*. Facilitated by the Royal Botanic Gardens, Kew. <http://apps.kew.org/wcsp/>. Accessed 17 January 2013.
- Graham EA, Nobel PS. 1996. Long-term effects of a doubled atmospheric CO₂ concentration on the CAM species *Agave deserti*. *Journal of Experimental Botany* **47**, 61–69.
- Hartsock TL, Nobel PS. 1976. Watering converts a CAM plant to daytime CO₂ uptake. *Nature* **262**, 574–576.
- Holtum JAM, Aranda J, Virgo A, Gehrig HH, Winter K. 2004. δ¹³C values and crassulacean acid metabolism in *Clusia* species from Panama. *Trees* **18**, 658–668.
- Holtum JAM, Chambers D, Morgan T, Tan DKY. 2011. *Agave* as a biofuel feedstock in Australia. *GCB Bioenergy* **3**, 58–67.
- Holtum JAM, Winter K. 2003. Photosynthetic CO₂ uptake in seedlings of two tropical tree species exposed to oscillating elevated concentrations of CO₂. *Planta* **218**, 152–158.
- Neales TF. 1973. The effect of night temperature on CO₂ assimilation, transpiration, and water use efficiency in *Agave americana* L. *Australian Journal of Biological Sciences* **26**, 705–714.
- Nobel PS. 1991. Achievable productivities of certain CAM plants: basis for high values compared with C₃ and C₄ plants. *New Phytologist* **119**, 183–205.
- Nobel PS. 1996a. High productivity of certain agronomic CAM species. In: Winter K, Smith JAC, eds. *Crassulacean acid metabolism: biochemistry, ecophysiology and evolution*. Berlin: Springer-Verlag, 255–265.
- Nobel PS. 1996b. Responses of some North American CAM plants to freezing temperatures and doubled CO₂ concentrations: implications of global climate change for extending cultivation. *Journal of Arid Environments* **34**, 187–196.
- Nobel PS. 2003. *Environmental biology of agaves and cacti*. Cambridge: Cambridge University Press.
- Nobel PS, García-Moya E, Quero E. 1992. High annual productivity of certain agaves and cacti under cultivation. *Plant, Cell & Environment* **15**, 329–335.
- Nobel PS, Israel AA, Wang N. 1996. Growth, CO₂ uptake, and responses of the carboxylating enzymes to inorganic carbon in two highly productive CAM species at current and doubled CO₂ concentrations. *Plant, Cell & Environment* **19**, 585–592.
- Nobel PS, Israel AA. 1994. Cladode development, environmental responses of CO₂ uptake, and productivity for *Opuntia ficus-indica* under elevated CO₂. *Journal of Experimental Botany* **45**, 295–303.
- Nobel PS, Valenzuela AG. 1987. Environmental responses and productivity of the CAM plant, *Agave tequilana*. *Agricultural and Forest Meteorology* **39**, 319–334.
- Osmond CB. 1978. Crassulacean acid metabolism: a curiosity in context. *Annual Review of Plant Physiology* **29**, 379–414.
- Pimienta-Barrios E, Zañudo-Hernández J, García-Galindo J. 2006. Seasonal photosynthesis in young plants of *Agave tequilana*. *Agrociencia* **40**, 699–709.
- Pimienta-Barrios E, Robles-Murguía C, Nobel PS. 2001. Net CO₂ uptake for *Agave tequilana* in a warm and a temperate environment. *Biotropica* **33**, 312–318.
- Silvera K, Santiago LS, Cushman JC, Winter K. 2010. The incidence of crassulacean acid metabolism in Orchidaceae derived from carbon isotope ratios: a checklist of the flora of Panama and Costa Rica. *Botanical Journal of the Linnean Society* **163**, 194–222.
- Somerville C, Youngs H, Taylor C, Davis SC, Long SP. 2010. Feedstocks for lignocellulosic biofuels. *Science* **329**, 790–792.
- Wijnands DO. 1983. *The botany of the Commelins*. Rotterdam: Balkema.
- Winter K, Garcia M, Holtum JAM. 2008. On the nature of facultative and constitutive CAM: environmental and developmental control of CAM expression during early growth of *Clusia*, *Kalanchoë*, and *Opuntia*. *Journal of Experimental Botany* **59**, 1829–1840.
- Winter K, Garcia M, Holtum JAM. 2009. Canopy CO₂ exchange of two neotropical tree species exhibiting constitutive and facultative CAM photosynthesis, *Clusia rosea* and *Clusia cylindrica*. *Journal of Experimental Botany* **60**, 3167–3177.

Winter K, Garcia M, Holtum JAM. 2011. Drought-stress-induced up-regulation of CAM in seedlings of a tropical cactus, *Opuntia elatior*, operating predominantly in the C₃ mode. *Journal of Experimental Botany* **62**, 4037–4042.

Winter K, Holtum JAM. 2002. How closely do the $\delta^{13}\text{C}$ values of crassulacean acid metabolism plants reflect the proportion of CO₂ fixed during day and night? *Plant Physiology* **129**, 1843–1851.

Winter K, Holtum JAM. 2011. Induction and reversal of crassulacean acid metabolism in *Calandrinia polyandra*: effects of soil moisture and nutrients. *Functional Plant Biology* **38**, 576–582.

Winter K, von Willert DJ. 1972. NaCl-induzierter Crassulaceensäurestoffwechsel bei *Mesembryanthemum crystallinum*. *Zeitschrift für Pflanzenphysiologie* **67**, 166–170.

Yan X, Tan DKY, Inderwildi OR, Smith JAC, King DA. 2011. Life cycle energy and greenhouse gas analysis for Agave-derived bioethanol. *Energy and Environmental Science* **4**, 3110–3121.

Zhu J, Goldstein G, Bartholomew DP. 1999. Gas exchange and carbon isotope composition of *Ananas comosus* in response to elevated CO₂ concentration and temperature. *Plant, Cell & Environment* **22**, 999–1007.