

## Gene Expression Profiles of Aralkylamine N-Acetyltransferase, B-cell Translocation Gene-2 and Fatty Acid Synthase in Pasture-based Primiparous Holstein-Friesian Dairy Cows Supplemented with Crude Degummed Canola Oil

Malau-Aduli AEO<sup>1,5\*</sup>, Otto JR<sup>1</sup>, Suybeng B<sup>2</sup>, Kashani A<sup>1</sup>, Lane PA<sup>1</sup>, Malau-Aduli BS<sup>3</sup> and Nichols PD<sup>4</sup>

<sup>1</sup>Animal Science and Genetics, Tasmanian Institute of Agriculture, School of Land and Food, Faculty of Science, Engineering and Technology, University of Tasmania, Private Bag 54 Sandy Bay, Hobart, Tasmania 7001, Australia

<sup>2</sup>Institut National Supérieur des Sciences Agronomiques de l'alimentation et de l'environnement (AGROSUP), 26 Bd du Docteur Petitjean - BP 87 999, 21079 Dijon Cedex, France

<sup>3</sup>College of Medicine and Dentistry, Division of Tropical Health and Medicine, James Cook University, Townsville, QLD 4811, Australia

<sup>4</sup>CSIRO Food and Nutrition, Oceans and Atmosphere Flagships, Hobart, TAS 7001, Australia

<sup>5</sup>Veterinary and Biomedical Sciences, College of Public Health, Medical and Veterinary Sciences, Division of Tropical Health and Medicine, James Cook University, Townsville, Queensland 4811, Australia

\*Corresponding author: Aduli Enoch Othniel Malau-Aduli, Veterinary and Biomedical Sciences, College of Public Health, Medical and Veterinary Sciences, Division of Tropical Health and Medicine, James Cook University, Townsville, Queensland 4811, Australia, Tel: +61-3-6226-2617; Fax: +61-3-6226-7444; E-mail: Aduli.MalauAduli@utas.edu.au

Rec date: Apr 5, 2015, Acc date: Jul 6, 2015, Pub date: Jul 10, 2015

Copyright: © 2015 Malau-Aduli AEO, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Abstract

The effect of oil-rich supplements on the expression of genes involved in lipogenesis and reproduction in pasture-based dairy cows is currently unknown, or at best, scanty and limited to impacts on cow liveweight, body condition score, milk composition, fatty acid and plasma metabolite profiles only. This research investigated the gene expression patterns of Aralkylamine N-acetyltransferase (AANAT), B-cell translocation gene-2 (BTG2) and Fatty Acid Synthase (FASN) genes in response to incremental levels of dietary crude degummed canola oil (CDCO). We tested the hypothesis that the relative mRNA abundance and gene expression profiles of AANAT, BTG2 and FASN in primiparous Holstein-Friesian cows will be up-regulated in response to post-partum dietary supplementation with CDCO in a typical pasture-based dairy production system. Thus, the primary objective of this study was to investigate the expression of AANAT, BTG2 and FASN genes in response to incremental levels of CDCO. A random allocation of primiparous Holstein-Friesian dairy cows into four treatment groups comprising wheat-based pelleted with no supplemental CDCO (control), or with CDCO added at 25 ml kg<sup>-1</sup> DM (low), 35 ml kg<sup>-1</sup> DM (medium) and 50 ml kg<sup>-1</sup> DM (high) was utilized in a ten-week experimental feeding trial including two weeks of adjustment. Both level of supplementation and their interaction with duration were significant sources of variation ( $P < 0.05$ ) that influenced BTG2 expression, while the expressions of AANAT and FASN genes were unaffected ( $P > 0.05$ ). The high (0.67 fold), medium (0.87 fold) and low (0.56 fold) treatments had suppressed BTG2 expressions compared to the control (1.0 fold) group. The low expression of BTG2 might be important when the reproductive system of cows is recovering from the effect of gestation and new cell growth is required.

**Keywords** Gene expression; mRNA; Canola oil; Aralkylamine N-acetyltransferase; B-cell translocation gene-2; Fatty acid synthase; Pasture-based Holstein-Friesian cows

### Abbreviations

AANAT: Aralkylamine N-acetyltransferase; ADF: Acid Detergent Fibre; APRO: Anti-Proliferative; CDCO: Crude Degummed Canola Oil; AI: Artificial Insemination; BCS: Body Condition Score; BLAST: Basic Local Alignment Search Tool; BTG2: B-cell Translocation Gene 2; cDNA: Complementary Deoxyribonucleic Acid; CLA: Conjugated Linoleic Acid; CP: Crude Protein; Ct: Cycle threshold; DM: Dry Matter; FASN: Fatty Acid Synthase; GLC: Gas Liquid Chromatography; "LC  $\omega$ -3" to "LC  $\omega$ -3": Long chain Omega-3; ME: Metabolisable Energy; NADPH: Nicotinamide Adenine Dinucleotide Phosphate; NDF: Neutral Detergent; PUFA: Polyunsaturated Fatty Acids; qPCR: Quantitative Real Time Polymerase Chain Reaction; RNA: Ribonucleic Acid; SAS: Statistical Analysis System; SFA: Saturated Fatty Acids; TIA: Tasmanian Institute of Agriculture

### Introduction

Nutritional attempts to remedy infertility are of interest to the dairy industry [1] because the antagonistic relationship between high milk production and fertility in modern, high genetic-merit cows has concomitantly led to a gradual but progressive decline in reproductive performance in diverse dairy production systems around the world. Prolonged calving intervals along with embryonic losses and postpartum anovulatory intervals are some of the major causes of infertility in cows [2]. In a typical pasture-based dairy system, different sources of lipids fed to lactating cows have been trialled to primarily increase the energy density of the diet in order to enhance milk production when negative energy balance peaks [3]. Research findings suggest that dietary supplementation with fat sources containing adequate proportions of unsaturated fats could potentially improve fertility in high merit dairy cows [4]. Therefore, a new, effective and long-term nutritional strategy that can assist in a better understanding of nutrition-fertility interactions in pasture-based systems is a potential solution to the subfertility problem in dairy cows.

Lipids also epitomize an effective nutritional approach for modifying milk fat composition [5] to favour an elevated profile of beneficial polyunsaturated fatty acids [6]. Lipids also play a crucial role in regulating the expression of genes essential for fertility and de novo fat synthesis in dairy cows. It has been shown that dietary fats containing trans-10, cis-12 conjugated linoleic acid (CLA) cause milk fat depression by inhibiting the expression of fatty acid synthase (FASN) gene [3] which is known to play a central role in the biosynthesis of fat in the mammary gland of mammals [7]. Although FASN is an important gene involved in lipogenesis, there is only limited published information about its expression in fat-supplemented cows in a pasture-based dairy production system. Where such studies were conducted, results have been conflicting and inconsistent [8] and warrant further research.

Arylalkylamine N-acetyltransferase (AANAT) is an essential gene for melatonin biosynthesis [9]. Melatonin is directly associated with optimal functioning of the ovary, where it regulates the hypothalamic-pituitary-gonadal axis to instigate folliculogenesis and steroidogenesis [10,11]. However, previous studies on AANAT have focused mainly on humans, in spite of AANAT being an important gene controlling reproduction in other mammals [12]. Currently, there is limited information on the expression of AANAT gene in dairy cows, especially with regards to fat supplementation, thus creating an important knowledge gap that this study intends to fill.

B-cell translocation gene-2 (BTG2) is an anti-proliferative gene that regulates cell cycle growth and BTG2 research investigations have been limited to cancer studies [13] demonstrating that the anti-proliferative characteristic of BTG2 gene is crucial during ovulation in mammals [14]. Published information on the expression and function of the BTG2 gene in dairy cattle, especially when supplemented with dietary lipids in a pasture-based system, are to our knowledge, either non-existent or at best, scanty. The above mentioned genes are related with differences in total fatty acid content in animal tissues and the protection of long chain ( $\geq C20$ ) polyunsaturated fatty acids through the prevention of peroxidation [15].

Understanding the mechanism underpinning the impact of dietary fat intake on the reproductive sequences from oestrous to conception in cows could revolutionise how nutrition is managed in dairy farms to improve reproductive performance. It will also assist researchers in unravelling the mystery behind the current global and gradual, but progressive, decline in dairy cow fertility. In this regard, further studies are required to unpack the intricate biological mechanisms involved with feeding dietary fats to grazing cows and their effects on lactation and fertility traits. This will enable dairy farmers make informed choices and tailored decisions when feeding lactating cows with specific dietary fat supplements. We hypothesized in this study that supplementation with CDCO would affect the expression of genes involved in reproductive functions (AANAT, BTG2) and de novo fatty acid synthesis (FASN) in primiparous Holstein-Friesian cows grazing under similar environmental conditions. Therefore, the primary objective of this study was to determine the relative abundance and expression of genes encoding proteins required for optimal reproduction and de novo lipogenesis in pasture-based lactating cows subjected to zero, low, medium and high levels of dietary supplementation with CDCO.

## Materials and methods

### Animal ethics

The use of animals and procedures performed in this study were all approved by the University of Tasmania Animal Ethics Committee (Permit No AA0012583), and were conducted in accordance with the 1993 Tasmanian Animal Welfare Act and the 2004 Australian Code of Practice for the Care and Use of Animals for Scientific Purposes.

### Experimental site and location

The experiment was carried out at the University of Tasmania's Dairy Research Centre, Tasmanian Institute of Agriculture (TIA) Elliot Dairy Research Farm in Somerset, North-Western Tasmania, Australia, from September to November 2012. Tasmania is Australia's smallest state with a land size of 68,000 square kilometers and located within the cool, temperate, climatic zone at latitude 42° South and longitude 147° East. It is characterized by four distinct seasons - winter, autumn, spring and summer. The experiment was carried out in spring when the annual rainfall was 2500 mm and humidity was approximately 60%.

### Experimental design, treatment groups and supplementary feeding trial

The physical condition and energy status of the experimental cows was visually assessed based on body condition score (BCS) on a scale of 1-8 [16]. Twenty primiparous, spring-calving, purebred, Holstein-Friesian cows (average liveweight of 400  $\pm$  40 kg, BCS 4  $\pm$  1.0 and 40  $\pm$  8 days in milk (DIM), were randomly allocated into 1 of 4 treatment groups of supplementation with wheat-based CDCO pellets classified as low (25 ml/ kgDM), medium (35 ml/ kgDM), high (50 ml/ kgDM) and the control (no CDCO- 0 ml/ kgDM). This replicated herd of cows (n=5 per treatment group) receiving CDCO supplements was placed under the same management and rotated in electric fenced paddocks with the Control cows offered wheat-based pellets without CDCO. Together, the animals had access to 3000 kgDM of forages, a mixture of ryegrass (*Lolium perenne*), cocksfoot (*Dactylis glomerata*), and white clover (*Trifolium repens*) pasture grazed at the two-leaf stage. Water was offered ad libitum. The current level of CDCO was calculated based on 7% total fat allowed in the diet of grazing cows. Each cow received 6 kg of the pelleted supplements daily for eight weeks, after two weeks of adjustment. Supplements were offered to cows in two splits; morning (3 kg) and evening (3 kg) milking sessions at 0500 hr and 1500 hr. There was no feed residual left over from any of the groups. The exact pasture intake was difficult to estimate as the case is under grazing conditions.

### Feed chemical composition analysis

The physical condition Dry matter (DM) content of the basal and experimental diets was determined by drying samples to a constant temperature at 65°C in a fan forced oven, finely ground to pass through a 2 mm sieve using Laboratory Mill (Thomas Model 4 Wiley® Mill; Thomas Scientific), and further drying at 105°C for 24 h. The DM was computed as the difference between the initial and final weights of samples. Ash content was determined by combusting samples in a furnace at 600°C for 8 hours. Neutral detergent (NDF) and acid detergent fibre (ADF) contents were measured using an Ankom Fiber Analyzer (ANKOM220; ANKOM Technology, USA). Nitrogen content was determined using a Thermo Finnigan EA 1112

Series Flash Elemental Analyzer and the values multiplied by 6.25 to give the crude protein (CP) percentage. Ether extract (EE) was determined using an Ankom fat/oil extractor (ANKOMXT15;

ANKOM Technology, USA). Metabolisable energy (ME) was calculated in accordance with Van Es [17]. The chemical compositions of the treatment, control and basal feeds are presented in Table 1.

<sup>a-b</sup> Chemical composition (%DM)	<sup>c</sup> Feeds		
	Control (No canola oil)	Treatment (canola oil)	Basal diet (Pasture)
MC	9.1	8.2	5.5
DM	90.9	91.8	94.5
ADF	9.0	8.0	27.7
NDF	21.1	20.0	45.9
EE	2.1	6.2	3.0
Ash	8.9	9.7	9.3
NFC	59.0	52.8	23.9
OM	91.1	90.3	90.7
CP	10.4	12.7	21.0
ME (MJ/kg DM)	4.07	4.08	3.99

<sup>a</sup> All feeds were analysed based on a dry weight basis; <sup>b</sup> MC: Moisture Content ; DM: Dry Matter; OM: Organic Matter; NDF: Neutral Detergent Fibre; ADF: Acid Detergent Fibre; NFC: Non-Fibrous Carbohydrate; EE: Ether Extract; CP: Crude Protein; ME: Metabolisable Energy; Treatment: feed with added canola oil. <sup>c</sup> Control: Feed without Canola Oil; Basal Diet: Mixed Ryegrass Pasture.

**Table 1:** Chemical composition of the experimental and basal feeds.

### Fatty acid analysis of basal and supplementary feeds

The fatty acid profiles of both basal and supplementary feeds were analysed by gas liquid chromatography (GLC) and presented in

Table 2. The detailed procedure had been previously described and published [18].

<sup>a</sup> Fatty acid	<sup>b</sup> Feed components		
	Control (No canola oil)%	Treatment (canola oil)%	Basal (Pasture)%
12:0	0.00	0.00	0.05
14:0	0.10	0.09	0.10
15:0	0.20	0.13	0.20
16:1	0.00	0.00	1.00
16:0	32.10	26.10	10.00
17:0	0.20	0.18	0.10
18:3 $\omega$ 6	0.00	0.03	0.00
18:4 $\omega$ 3	0.00	0.00	0.90
18:2 $\omega$ 6 LA	17.70	6.86	9.10
18:3 $\omega$ 3 ALA	1.60	0.48	64.30
18:1 $\omega$ 9c	16.50	41.90	4.40
18:1 $\omega$ 7t	0.20	0.10	0.20
18:0	3.80	3.83	2.20
18:2CLA	0.10	1.48	0.00

19:0	0.90	3.47	0.10
20:4 $\omega$ 6	0.00	0.01	0.00
20:5 $\omega$ 3 EPA	11.80	0.20	0.10
20:3 $\omega$ 6	0.40	1.82	0.80
20:4 $\omega$ 3 ARA	0.40	0.22	0.10
20:2 $\omega$ 6	1.40	1.45	0.00
20:0	0.80	1.38	0.40
22:5 $\omega$ 6	0.30	0.04	0.10
22:6 $\omega$ 3 DHA	0.20	0.03	0.00
22:4 $\omega$ 6	0.20	0.00	0.00
22:5 $\omega$ 3 DPA	0.90	0.00	0.00
22:0	1.80	1.86	1.50
24:0	1.10	1.30	0.90
tSFA	41.20	38.64	16.45
tMUFA	23.30	48.74	8.00
tPUFA	35.00	12.62	75.40
$\omega$ -3 PUFA	14.90	0.93	65.40
$\omega$ -6 PUFA	20.10	10.24	10.10
$\omega$ -3 LC-PUFA	13.30	0.45	0.20

<sup>a</sup> $\Sigma$ tSFA is the sum of 12:0, 13:0, i14:0,14:0, i15:0, a15:0,15:0, i16:0, 16:0, i17:0, 17:0, i18:0, 18:0, 19:0, 20:0, 20:0, 22:0, 24:0;  $\Sigma$ tMUFA is the sum of 14:1 $\omega$  -5c, 15:1 $\omega$ -6c, 16:1 $\omega$  -9c, 16:1 $\omega$  -7c, 16:1 $\omega$  -7t, 16:1 $\omega$  -5c, 16:1,17:1 $\omega$  -8+a17:0, 17:1 $\omega$ -6c, 18:1 $\omega$  -9c, 18:1 $\omega$  -7c, 18:1 $\omega$  -7t, 18:1 $\omega$  -5c, 18:1a, 18:1b, 20:1 $\omega$  -11c, 20:1 $\omega$  -9c, 20:1 $\omega$  -7c, 20:1 $\omega$  -5c, 22:1 $\omega$  -11c, 22:1 $\omega$  -9c, 22:1 $\omega$  -7c, 24:1 $\omega$  -11c, 24:1 $\omega$  -9c, 24:1 $\omega$  -7c;  $\Sigma$ tPUFA is the sum of 18:3 $\omega$ -6, 18:4 $\omega$ -3, 18:2 $\omega$ -6, 18:3 $\omega$ -3, 18:2CLA, 20:4 $\omega$ -6, 20:5 $\omega$ -3, 20:3 $\omega$ -6, 20:4 $\omega$ -3, 20:2 $\omega$ -6, 22:5 $\omega$ -6, 22:6 $\omega$ -3, 22:4 $\omega$ -6, 22:5 $\omega$ -3;  $\Sigma$  $\omega$ -3 LC-PUFA is the sum of 20:5 $\omega$ -3, 20:4 $\omega$ -3, 22:6 $\omega$ -3, 22:5 $\omega$ -3;  $\Sigma$  $\omega$ -3 PUFA is the sum of 18:4 $\omega$ -3, 18:3n-3, 20:4 $\omega$ -3, 20:5 $\omega$ -3, 22:6 $\omega$ -3, 22:5 $\omega$ -3;  $\Sigma$  $\omega$ -6 is the sum of 15:1 $\omega$ -6, 17:1 $\omega$ -6, 18:2 $\omega$ -6, 18:3 $\omega$ -6, 20:4 $\omega$ -6, 20:3 $\omega$ -6, 20:2 $\omega$ -6, 22:5 $\omega$ -6, 22:4 $\omega$ -6. tSFA= total Saturated Fatty Acids, tMUFA: total Monounsaturated Fatty Acids, tPUFA: Total Polyunsaturated Fatty Acids,  $\omega$ -3 FA: Total Omega-3 Fatty Acids,  $\omega$ -6 FA: Total Omega-6 Fatty acids,  $\omega$ -3 LC-FA: Total Omega-3 Long Chain Fatty Acids; <sup>b</sup>Control: feed with no Added Canola Oil; Treatment: Feed with Canola Added; basal: Mixed Ryegrass Pasture.

**Table 2:** Fatty acid concentration as a percentage of total fatty acids of basal and supplementary feeds.

### Blood sample collection

Blood samples were collected from all experimental cows by coccygeal venipuncture into vacutainers containing heparin after the morning milking (0500 hr) on the day before the initiation of supplementation with CDCO and in week eight at the conclusion of the experiment. More frequent blood sample collection interval was restricted by the terms and conditions of the Animal Ethics Permit No AA0012583 granted by the University of Tasmania Animal Ethics Committee. The samples were immediately frozen in -20°C and transported to the laboratory for further storage in -80°C until RNA extraction.

### RNA extraction and cDNA synthesis

Frozen blood samples were thawed and utilised for the isolation of total RNA using TRIzol® Plus RNA Purification Kit (Life Technologies Pty Ltd, Victoria, Australia). A tissue lyser (Qiagen Ltd., Crawley, UK) was used to homogenise the sample in TRIzol® Reagent. Total RNA quantity and quality was measured using the NanoDrop 8000

spectrophotometer (NanoDrop, Wilmington, DE, USA). RNA that had an absorbance (A260/280) reading between 1.8 and 2 was deemed of good quality. The RNA samples were treated with PureLinkTMDNase (Life Technologies Pty Ltd, Victoria, Australia) and purified using the RNeasy1 Mini Kit (Qiagen Ltd, NSW, Australia). DNase-treated and purified total RNA was then reverse transcribed to cDNA with Mixed Oligo dT/Random Hexamer Primers using the Tetro cDNA Synthesis Kit (Bioline Pty Ltd, NSW, Australia) according to the manufacturer's instructions.

### Primer design and reference gene selection

All candidate and reference gene primers (Table 3) were designed using the Primer3 web-based software program (<http://frodo.wi.mit.edu/primer3/>) from GeneWorks Pty Ltd, SA, Australia). Basic Local Alignment Search Tool (BLAST) from the National Centre for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/BLAST/>) was used to check for the specificity of the primers. The validity of all primers was confirmed using a serial dilution of pooled

cDNA to create a standard curve. Subsequently, the amplified PCR products were sequenced to confirm their primer specific identity (Beckman Coulter CEQTM 8000 Series Genetic Analysis System). The mRNA abundance was determined using highly stable reference genes. The normalisation of expression data for the target genes Aralkylamine N-acetyltransferase (AANAT), B-cell translocation gene

2 (BTG2), and Fatty acid synthase (FASN) utilised two reference genes, Ubiquitin C (UBC) and Peptidyl-prolyl cis-trans isomeraseA (PPIA). A good selection criterion of reference genes was an expression ratio that was constant across all samples. The software program geNorm, version 3.5 [19], was used to calculate, confirm and validate the expression stability (M-value) of the reference genes.

<sup>a</sup> Gene symbol	qPCR Primers		<sup>b</sup> Ta	Amplicon Size (bp)
	Forward Primer	Reverse Primer		
AANAT	ACTGACCTTCACGGAGATGC	TTCACCTATTCTCCCGTTC	60	211
ADRB3	TCAGTAGGAAGCGGGTCCGGG	GGCTGGGGAAGGGCAGAGTT	60	291
BTG2	CTGGAGGAGAAGTGGCTGTC	AAAACAATGCCCAAGGTCTG	60	194
FASN	GTGTGGTACAGCCCCTCAAG	ACGCACCTGAATGACCACTT	60	110
Reference genes				
UBC	CGTCTTAGGGGTGGCTGTTA	AAATTGGGGTAAATGGCTAGA	60	90
PPIA	TCATTTGCACTGCCAAGACTG	TCATGCCCTCTTTCACITTTGC	60	72

<sup>a</sup>AANAT: Aralkylamine N-Acetyltransferase; ADRB3: β3-Adrenergic Receptor; BTG2: B-cell Translocation Gene 2; FASN: Fatty Acid Synthase; UBC: Ubiquitin C; PPIA: Peptidyl-Prolyl Cis-Trans IsomeraseA; <sup>b</sup>Ta: Empirical Annealing Temperature.

**Table 3:** Primer pairs designed for real-time PCR (qPCR).

### Quantitative real time PCR (qPCR)

Following reverse transcription, cDNA quantity was determined and standardised to the required concentration for qPCR. Triplicate 20 µL reactions were carried out in a 72-well Rotor-Gene (QIAGEN GmbH, Hilden, Germany), containing 4 µL cDNA (50 ng), 10 µL 2× SensiFAST SYBR No-ROX Mix (Bioline Pty Ltd., NSW, Australia), 4.4 µL DEPC H<sub>2</sub>O, and 0.8 µL forward and reverse primers (100 fmol). Assays were performed using the Rotor-Gene 3000 (QIAGEN Pty Ltd., VIC, Australia) with the following cycling parameters: 95°C for 2 min polymerase activation; 40 cycles of 95°C for 5 s denaturation, 60°C for 10 s annealing and 72°C for 5 s extension. Gene expression levels were recorded as Ct values (i.e., the number of PCR cycles at which the fluorescence signal was detected above the threshold value) and all samples were run in triplicates. Amplification efficiencies were determined for all candidate and reference genes using the formula  $E = 10^{(-1/\text{slope})}$ , with the slope of the linear curve of cycle threshold (Ct) values plotted against the log dilution as per Higuchi et al., [20]. Primer concentrations were optimised for each gene and disassociation curves were examined for the presence of a single PCR product. The efficiency of the reaction was calculated using a 5-fold serial dilution of cDNA and generation of a standard curve. All PCR efficiency coefficients were between 1.7 and 1.8 and therefore deemed acceptable. The Rotor-Gene 3000 (version 6.0.16) (QIAGEN Pty Ltd., VIC, Australia) was used for efficiency correction of the raw Ct values. This process involved an inter-plate calibration based on a calibrator sample included on all plates, averaging of replicates, normalisation to the reference gene and the calculation of quantities relative to the highest Ct and log<sub>2</sub> transformation of the expression values for all genes. A PCR efficiency coefficient between 1.7 and 1.8 was considered adequate. The mathematic model used to determine the expression

level of the target gene in comparison to the reference gene is given below as per Pfaffl [21].

$$\text{Ratio} = \frac{(E_{\text{target}})^{\Delta C_{\text{Ptarget}}(\text{control} - \text{sample})}}{(E_{\text{ref}})^{\Delta C_{\text{Pref}}(\text{control} - \text{sample})}}$$

### Statistical analysis

Initially, summary statistics by level and week (duration) of CDCO supplementation were computed to give means, standard deviations, standard error, variance, minimum and maximum values that were scrutinised for any data entry errors. Testing for linear, cubic and quadratic orthogonal contrasts by regressing the dependent on explanatory variables was carried out using PROC REG (SAS 2009) [22]. However, the linear, quadratic and cubic orthogonal contrasts were all found to be inconsequential. Therefore, repeated measures analysis of variance using PROC MIXED (SAS 2009) [22] was employed fitting fixed effects of treatment, week of supplementation and their second-order interactions on the expressions of AANAT, BTG2 and FASN genes. The 1st-order autoregressive covariance structure and level of supplementation were fitted as the repeated effects and cows as random effects. The degrees of freedom were estimated by the Satterthwaite method (SAS, 2009) [22]. Variable means are presented in Tables and Figures as LSM ± SEM. Tukey's pairwise comparison test was utilized in establishing differences between means using the P<0.05 threshold for significance unless otherwise stated.

### Results

The effects of supplementation with CDCO on milk yield, milk fat and number of services per conception are presented in Figures 1-3, respectively. Cows receiving 50 mL kg<sup>-1</sup> DM (High) of canola oil



produced more milk ( $168.7 \pm 3.4$  vs.  $157.1 \pm 3.7$  Litres) with a lower fat percentage ( $3.3 \pm 0.1$  vs.  $4.0 \pm 0.2\%$ ) than unsupplemented cows in the control treatment group (0 mL kg<sup>-1</sup> DM) as depicted in Figures 1-3.

Figures 4-6 and Table 4 represent the relative mRNA abundance and expressions of AANAT, BTG2 and FASN genes, and these are individually highlighted as follows.

### Aralkylamine N-acetyltransferase (AANAT)

Dietary supplementation of primiparous Holstein-Friesian cows with CDCO had no effect ( $P > 0.05$ ) on the expression of AANAT gene (Figure 4). As the week (duration) of supplementation progressed, the impact of CDCO supplements and interaction with duration of supplementation was insignificant on the expression of AANAT gene (Table 4).

<sup>a</sup> Effect	<sup>b</sup> Genes		
	AANAT	BTG2	FASN
<sup>c</sup> TRT	0.2019	0.0495	0.9289
<sup>d</sup> Week	0.2713	0.1818	0.3314
<sup>e</sup> Week*TRT	0.4956	0.0107	0.6647

<sup>a</sup>All p-values in bold were significant ( $P < 0.05$ ). <sup>b</sup>AANAT: Arylalkylamine-N-acetyltransferase; BTG2: B-cell Translocation Gene-2; FASN: Fatty Acid Synthase; TRT: Treatment, <sup>c</sup>Week: Week of lactation, <sup>e</sup> Week\*TRT: Interaction between Week of Supplementation and Treatment.

**Table 4:** Multi-trait analysis of variance (p-values) for fixed and interaction effects of treatment and week of supplementation on the relative mRNA expression of AANAT: Arylalkylamine-N-acetyltransferase; BTG2: B-cell Translocation Gene-2; FASN: Fatty Acid Synthase genes in grazing Holstein-Friesian dairy.

### Fatty acid synthase (FASN)

Differences in CDCO content in the supplemented primiparous Holstein-Friesian cows had no significant effect on the relative mRNA abundance of FASN gene. Week of supplementation and week by treatment interaction were inconsequential sources of variation ( $P > 0.05$ ).

### Discussion

A complex regulatory network of lipogenic genes, genetic and dietary differences impact the utilisation of oil-rich supplements in ruminants. To our best knowledge, the molecular mechanisms underlying these systems remain to be fully understood and characterized in pasture-based dairy production systems. Therefore, an attempt at trying to fathom the factors affecting dietary lipid metabolism in lactating cows is of utmost scientific relevance. This study utilised an experimental feeding trial with oil-rich dietary supplementation with CDCO in grazing primiparous Holstein-Friesian cows and determined the mRNA expression levels of BTG2, AANAT and FASN genes.

### BTG2

The mammalian BTG2 gene belongs to the anti-proliferative (APRO) family of genes that regulate cell cycle progression in a variety of cell types [23,24]. BTG2 is a prototypical member of the BTG/TOB family with anti-proliferative properties. The protein encoded by this gene controls cell cycle progression and proneural gene expression by acting as a transcription co-regulator that enhances or inhibits the

### B-cell translocation gene 2 (BTG2)

It was evident that both treatment and treatment by week of supplementation interactions were significant sources of variation that influenced the expression of BTG2 gene ( $P < 0.05$ ; Table 4). However, week of supplementation alone had no influence ( $P > 0.05$ ) on BTG2 gene expression. Cows receiving dietary supplementation with CDCO experienced consistent suppression of BTG2 gene expression compared to the control group (Figure 5). The cows in the control group recorded the greatest mRNA abundance of BTG2 (1.00 fold), followed by the medium group (0.87 fold), then the high group (0.67 fold) and finally the low group (0.56 fold).

activity of transcription factors [23,24]. Thus, BTG2 has many functions involving regulation of cell growth, death, differentiation and survival [25].

The present study found that supplementation of lactating Holstein-Friesian cows with CDCO repressed the expression of BTG2 significantly. The current result lends credence to the report of Jeckel et al. [26] who demonstrated that in rats, BTG2 gene was down-regulated when dietary fatty acid was fed. Therefore, lack of BTG2 up-regulation in the in the blood of cows in the present study would be due to the fatty acid content of the fed diet. Although some of the BTG2 gene expression studies were on tumorous diseases in humans, significant expressions have been observed in pig muscle, uterus and heart where the gene appears to play a role in cell development [27]. A previous study also found that gonadotropin hormones stimulate the expression of BTG2 genes in the ovary during ovulation [28]. Usually, luteinising hormone surges during the pre-ovulatory period, stops further growth of immature follicles, culminating in the ovulation of matured follicles for fertilisation [14,29]. The BTG2 gene is thus an essential gene for normal reproduction in mammals. The greater number of services per conception observed in the CDCO supplemented group suggests that these cows ovulated later than the control cows (Figure 3). This statement is supported by the low expression of BTG2 in supplemented cows (Figure 5). A previous study found that BTG2 gene acts as a co-activator for regulating hepatic gluconeogenesis [30]. In other words, the overexpression of BTG2 gene triggers hepatic gluconeogenesis, thus generating a spike in glucose synthesis which provides surplus energy for cell proliferation. It is well known that optimal recovery of the cow's uterus following a normal gestation period requires growth, development and

maturation of the granulosa/follicular cells for the cow to be ready for the next cycle of gestation. However, the anti-proliferative activity of the BTG2 gene might impact negatively on cell growth and tissue repair of the reproductive machinery. Our current findings seem to suggest that supplementation of cows with lipid-rich feeds could be utilised as a dietary manipulation tool to repress the expression of BTG2 gene and its anti-proliferative attributes.


## AANAT

From published literature, the AANAT gene has been reported to be associated with long-chain omega-3 (LC  $\omega$ -3) polyunsaturated fatty acid (PUFA) synthesis [15]. A major cause for disparity between this report and our observation in the current study could be due to differences in dietary lipid sources, dosages fed and the relatively smaller proportion of LC  $\omega$ -3 PUFA in our experimental diet compared to the control diet (0.45% vs. 13.30%; Table 2). The AANAT gene is also known to encode an acetyltransferase superfamily protein [31] that catalyses the rate-limiting step in the synthesis of melatonin from serotonin [9] primarily found in the pineal gland [32]. Melatonin (N-acetyl-5-methoxytryptamine) is a hormone that plays a role in reproductive functions in mammals [33], particularly in the growth and maturation of oocytes in the ovary and steroidogenesis in the granulosa cells via the mitogen-activated protein kinase pathway [10,34]. Melatonin is also essential for the function of the circadian clock that influences activity and sleep [31,35]. The mechanism by which melatonin regulates reproduction has been reported to be through the control of gonadotropin releasing hormone (GnRH) and gonadotropin inhibiting hormone (GnIH) receptors primarily found in the hypothalamic-pituitary axis to release gonadotropin hormones [11,33,36]. AANAT transcripts have been found to be differentially expressed in high vs. low omega-3 index (O3I) muscles, suggesting a role for melatonin in reducing oxidative damage, including that to PUFA [9]. The ability of melatonin to protect against lipid peroxidation has been repeatedly documented in many studies using animal and plant tissues [37]. Spanish scientists reported that melatonin consumption assists in the control of weight gain since it stimulates the appearance of brown fat (beige), a type of fat cell that burns calories instead of storing them [38]. Their research demonstrated that melatonin treatment not only induced browning of inguinal white adipose tissue in Zucker diabetic fatty rats, but also increased thermogenic activity [38]. Taken together, these findings highlight the anti-obesity effect of melatonin and explain its metabolic benefits of protecting against oxidative degradation of PUFA in the muscle tissue thereby producing higher O3I levels [38].


The afore-mentioned body of evidence in the published literature indicates that the expression of AANAT gene could play multifaceted functions in regulating fertility in dairy cows through biosynthesis of melatonin. This makes AANAT an ideal gene to explore in terms of possible nutritional manipulation of its expression to assist in controlling seasonal breeding in pasture-based dairy systems. From the current study, the supplementation of Holstein-Friesian cows with CDCO was inconsequential to AANAT gene expression since no differential expression of the AANAT gene was observed between the treatment groups. This suggests that supplementing grazing primiparous cows with CDCO may not be an influential determinant of AANAT gene expression. However, further research studies are warranted with different dietary fat sources to establish ideal dosage levels that can strongly up-regulate/and or down-regulate AANAT in dairy cows before its use can be adopted by the industry.

## FASN

FASN encodes a multifunctional enzyme that catalyses fatty acid synthesis [39]. FASN is considered as a fundamental enzyme in de novo lipogenesis in mammals and its main function is to catalyse the synthesis of palmitate from acetyl-CoA and malonyl-CoA, in the presence of NADPH, into long-chain saturated fatty acids (LC-SFA) [39,40]. It has been shown that the FASN gene contributes to the regulation of body weight in humans, which results in the development of obesity [39,41]. Thus, FASN is a complex homodimeric gene that plays a pivotal role in de novo lipid biosynthesis and regulation of milk fat content [7]. The lack of significant effect of the experimental lipid-rich diet on FASN expression is in agreement with the report of Bichi et al. [42]. It is pertinent to state that an observed trend in the current study towards the suppression of FASN expression (Figure 6) agrees with previous research in cattle [32,43,44] and ewes [45].




**Figure 1:** Effect of supplementing primiparous Holstein-Friesian dairy cows with CDCO on milk yield (Mean  $\pm$  SEM).




**Figure 2:** Influence of supplementing Holstein-Friesian dairy cows with CDCO on milk fat percentage (Mean  $\pm$  SEM).

The observed suppression trend of FASN expression is consistent with the role of fats containing conjugated linoleic acid (CLA), especially trans-10, cis-12 18:2 [34]. The fatty acid profile of our treatment diet had a greater proportion of CLA than the control diet


(1.48% versus 0.10%). This could help to explain the down-regulation of FASN expression in cows supplemented with CDCO and also the suppression of the milk fat content (Figure 2). However, the discrepancy between the present and previous studies could be due to differences in the dietary fat source, dosage, level of polyunsaturated fatty acid intake, rumen biohydrogenation intermediates, and basal diet offered. One clear message from our results is the potential for supplementation of grazing cows with CDCO to down-regulate FASN expression; that way, the energy spared from reduced milk fat synthesis could be partitioned towards metabolic functions in non-mammary tissues, specifically, in reproductive tissues postpartum or milk production (Figure 1). This could prove highly significant to pasture-based dairy farmers in terms of managing the energy needs for production and reproduction in their herds. The downside to this assertion in pasture-based dairy systems is that milk fat is an economically important constituent of total milk solids upon which Tasmanian dairy farmers are paid in Australia.




**Figure 3:** Number of services per conception in primiparous Holstein-Friesian dairy cows supplemented with CDCO (Mean  $\pm$  SEM).



**Figure 4:** Effect of dietary supplementation of grazing cows with CDCO on mRNA expression of AANAT gene (Mean  $\pm$  SEM).



**Figure 5:** Effect of supplementation with CDCO on mRNA expression of the BTG2 gene in grazing cows (Mean  $\pm$  SEM).



**Figure 6:** mRNA expression of FASN gene in grazing cows supplemented with CDCO (Mean  $\pm$  SEM).

Furthermore, extreme suppression of FASN expression may be adverse to the butter manufacturing industries due to lower milk fat content. However, the beneficial impact is that the milk is being rid of mostly saturated fats, thus giving consumers a healthier product for which a premium can be charged to compensate for lower milk solids.

## Conclusions

Dietary supplementation of grazing primiparous Holstein-Friesian dairy cows with CDCO had a significant influence in down-regulating the expression of BTG2. This might be important when the reproductive system of cows is recovering from the effect of gestation and new cell growth is required, but the downside is that hepatic gluconeogenesis could be hampered. mRNA abundance and



expression of AANAT and FASN genes were not significantly impacted by CDCO supplementation in spite of the observed trends towards up-regulation and down-regulation, respectively. The observed trend toward the suppression of FASN gene expression can be beneficial in sparing energy from milk fat synthesis and re-directing the surplus to non-mammary tissues *in vivo*. However, severe milk fat depression may be economically undesirable to Tasmanian dairy farmers because of its contribution to total milk solids upon which milk prices are based. These findings highlight the important role of nutrition in altering reproductive and lipogenic genes in the blood of lactating cows. Further studies with more experimental animals and CDCO supplementation levels would be required to confirm the current findings.

### Competing Interests

The authors declare that they have no competing interests.

### Authors Contributions

This work was carried out in collaboration between all authors. Author JRO ran the feeding trial, laboratory analyses, collected experimental samples, performed feed chemical analysis and wrote the first draft of the manuscript as part of his PhD project. Authors JRO, BS, AK designed the primers, selected reference genes, extracted RNA, synthesised cDNA, and carried out quantitative real time PCR. Authors BSMA, PDN and AEOMA contributed in the reading and making needed changes to the draft manuscript. Author AEOMA conceived the research idea, wrote the funding grant, experimental design, read and made final changes to the final manuscript as a research article. All authors read and approved the final manuscript.

### Acknowledgements

This study was supported with grants from the University of Tasmania Graduate Research Scholarship and a Postgraduate Top-up scholarship from the Tasmanian Institute of Agriculture. We also acknowledge the collaborative research support of CopRice Feeds Pty Limited, Cobden, Victoria, for manufacturing the experimental oil-enriched pellets to specification, field supervisory role of Mark Freeman, and Peter Nish of TasHerd Pty Limited, Hadspen, Tasmania for milk composition analysis.

### References

1. Chagas LM, Bass JJ, Blache D, Burke CR, Kay JK, et al. (2007) Invited review: New perspectives on the roles of nutrition and metabolic priorities in the subfertility of high-producing dairy cows. *J Dairy Sci* 90: 4022-4032.
2. Rocha A, Martins A, Carvalheira J (2010) Fertility time trends in dairy herds in northern Portugal. *Reprod Domest Anim* 45: 896-899.
3. Hutchinson I, de Veth MJ, Stanton C, Dewhurst RJ, Lonergan P, et al. (2011) Effects of lipid-encapsulated conjugated linoleic acid supplementation on milk production, bioenergetic status and indicators of reproductive performance in lactating dairy cows. *J Dairy Res* 78: 308-317.
4. Santos JE, Bilby TR, Thatcher WW, Staples CR, Silvestre FT (2008) Long chain fatty acids of diet as factors influencing reproduction in cattle. *Reprod Domest Anim* 43 Suppl 2: 23-30.
5. Staples CR, Burke JM, Thatcher WW (1998) Influence of supplemental fats on reproductive tissues and performance of lactating cows. *J Dairy Sci* 81: 856-871.

6. Otto JR, Malau-Aduli BS, Nichols PD, Malau-Aduli AEO, et al. (2014) Influence of supplementing pasture-based primiparous Holstein-Friesian dairy cows with crude degummed canola oil on milk fatty acid composition. *J Nutr Ther* 3: 55-66.
7. Roy R, Ordovas L, Zaragoza P, Romero A, Moreno C, et al. (2006) Association of polymorphisms in the bovine FASN gene with milk-fat content. *Anim Genet* 37: 215-218.
8. Vahmani P, Glover KE, Fredeen AH (2014) Effects of pasture versus confinement and marine oil supplementation on the expression of genes involved in lipid metabolism in mammary, liver, and adipose tissues of lactating dairy cows. *J Dairy Sci* 97: 4174-4183.
9. Öner Y, Orman A, Üstuner H, Yilmaz A (2014) Investigation of polymorphisms on ABCG2, AA-NAT and FABP3 genes in the Kivircik sheep reared in three different provinces of Turkey. *J Fac Vet Med* 20: 649-54.
10. Fiske VM, Parker KL, Ulmer RA, Ow CH, Aziz N (1984) Effect of melatonin alone or in combination with human chorionic gonadotropin or ovine luteinizing hormone on the *in vitro* secretion of estrogens or progesterone by granulosa cells of rats. *Endocrinol* 114: 407-410.
11. Chowdhury VS, Ubuka T, Tsutsui K (2013) Review: Melatonin stimulates the synthesis and release of gonadotropin-inhibitory hormone in birds. *Gen Comp Endocrinol* 181: 175-178.
12. Soria V, Martínez-Amorós È, Escaramís G, Valero J, Crespo JM, (2010) Resequencing and association analysis of arylalkylamine N-acetyltransferase (AANAT) gene and its contribution to major depression susceptibility. *J Pineal Res* 49: 35-44.
13. Choi YW, Park TJ, Kim HS, Lim IK (2013) Signals regulating necrosis of cardiomyoblast by BTG2/(TIS21/PC3) via activation of GSK3 $\beta$  and opening of mitochondrial permeability transition pore in response to H<sub>2</sub>O<sub>2</sub>. *Biochem Biophys Res Commun* 434: 559-565.
14. Park JI, Kim SG, Baek MW, Park TJ, Lim IK, et al. (2013) B-cell translocation gene 2: expression in the rat ovary and potential association with adenine nucleotide translocase 2 in mitochondria. *Mol Cell Endocrinol* 367: 31-40.
15. Perez R, Cañón J, Dunner S (2010) Genes associated with long-chain omega-3 fatty acids in bovine skeletal muscle. *J Appl Genet* 51: 479-487.
16. Stockdale CR (2001) Body condition at calving and the performance of dairy cows in early lactation under Australian conditions: A review. *Anim Prod Sci* 41: 823-839.
17. Van Es A (1975) Feed evaluation for dairy cows. *Livest Prod Sci* 2: 95-107.
18. Otto JR, Malau-Aduli BS, Nichols PD, Malau-Aduli AEO (2014) Influence of supplementing pasture-based primiparous Holstein-Friesian dairy cows with crude degummed canola oil on milk fatty acid composition. *J Nutr Ther* 3: 55-66.
19. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, et al. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034.
20. Higuchi R, Fockler C, Dollinger G, Watson R (1993) Kinetic PCR analysis: real-time monitoring of DNA amplification reactions. *Biotechnology (N Y)* 11: 1026-1030.
21. Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res* 29: e45.
22. SAS (2009) Statistical Analysis System Inc, SAS Institute, North Carolina, USA, Version 9.2.
23. Reiter RJ, Tan DX, Galano A (2014) Melatonin reduces lipid peroxidation and membrane viscosity: a review. *J Pineal Res* 56: 225-237.
24. Roy R, Taourit S, Zaragoza P, Eggen A, Rodellar C (2005) Genomic structure and alternative transcript of bovine fatty acid synthase gene (FASN): comparative analysis of the FASN gene between monogastric and ruminant species. *Cytogenet Genome Res* 111: 65-73.
25. Mo XY, Lan J, Jiao QZ, Xiong YZ, Zuo B, et al. (2011) Molecular characterization, expression pattern and association analysis of the porcine BTG2 gene. *Mol Biol Rep* 38: 4389-4396.

26. Jeckel KM, Bouma GJ, Hess AM, Petrilli EB, Frye MA (2014) Dietary fatty acids alter left ventricular myocardial gene expression in Wistar rats. *Nutr Res* 34: 694-706.
27. Feng Z, Tang ZL, Li K, Liu B, Yu M, et al. (2007) Molecular characterization of the BTG2 and BTG3 genes in fetal muscle development of pigs. *Gene* 403: 170-177.
28. Schmidt J, de Avila J, McLean D (2006) Regulation of protein tyrosine phosphatase 4a1, B-cell translocation gene 2, nuclear receptor subfamily 4a1 and diacylglycerol O-acyltransferase 1 by follicle stimulating hormone in the rat ovary. *Reprod Fertil Dev* 18: 757-765.
29. Li F, Liu J, Park ES, Jo M, Curry TE Jr (2009) The B cell translocation gene (BTG) family in the rat ovary: hormonal induction, regulation, and impact on cell cycle kinetics. *Endocrinology* 150: 3894-3902.
30. Hwang SL, Kwon O, Lee SJ, Roh SS, Kim YD, et al. (2012) B-cell translocation gene-2 increases hepatic gluconeogenesis via induction of CREB. *Biochem Biophys Res Commun* 427: 801-805.
31. Forrest RH, Hickford JG, Hogan A, Frampton C (2003) Polymorphism at the ovine beta3-adrenergic receptor locus: associations with birth weight, growth rate, carcass composition and cold survival. *Anim Genet* 34: 19-25.
32. Piesiewicz A, Kedzierska U, Adamska I, Usarek M, Zeman M, et al. (2012) Pineal arylalkylamine N-acetyltransferase (Aanat) gene expression as a target of inflammatory mediators in the chicken. *Gen Comp Endocrinol* 179: 143-151.
33. Malpaux B, Daveau A, Maurice-Mandon F, Duarte G, Chemineau P (1998). Evidence that melatonin acts in the premammillary hypothalamic area to control reproduction in the ewe: Presence of binding sites and stimulation of luteinizing hormone secretion by in situ micro implant delivery. *Endocrinol* 139: 1508-1516.
34. El-Raey M, Geshi M, Somfai T, Kaneda M, Hirako M, et al. (2011) Evidence of melatonin synthesis in the cumulus oocyte complexes and its role in enhancing oocyte maturation in vitro in cattle. *Mol Reprod Dev* 78: 250-262.
35. Guo B, Kongsuwan K, Greenwood PL, Zhou G, Zhang W, et al. (2014) A gene expression estimator of intramuscular fat percentage for use in both cattle and sheep. *J Anim Sci Biotechnol* 5: 35.
36. Soares JM Jr, Masana MI, ErÅyahin C, Dubocovich ML (2003) Functional melatonin receptors in rat ovaries at various stages of the estrous cycle. *J Pharmacol Exp Ther* 306: 694-702.
37. Holman BWB, Kashani A, Malau-Aduli AEO (2012) Growth and body conformation responses of genetically divergent Australian sheep to *Spirulina* (*Arthrospira platensis*) supplementation. *American J Exp Agric* 2: 160-173.
38. Holman BWB, Kashani A, Malau-Aduli AEO (2014) Effects of *Spirulina* (*Arthrospira platensis*) supplementation level and basal diet on liveweight, body conformation and growth traits in genetically divergent Australian dual-purpose lambs during simulated drought and typical pasture grazing. *Small Rum Res* 120: 6-1.
39. Strosberg AD (1997) Structure and function of the beta 3-adrenergic receptor. *Annu Rev Pharmacol Toxicol* 37: 421-450.
40. Wu JL, Liu WZ, Liu JH, Qiao LY, Yuan YN (2011) Distribution and quantification of  $\beta$ -3 adrenergic receptor in tissues of sheep. *Animal* 5: 88-93.
41. Wu J, Qiao L, Liu J, Yuan Y, Liu W (2012) SNP variation in ADRB3 gene reflects the breed difference of sheep populations. *Mol Biol Rep* 39: 8395-8403.
42. Bichi E, Frutos P, Toral PG, Keisler D, Hervás G, Loor JJ (2013) Dietary marine algae and its influence on tissue gene network expression during milk fat depression in dairy ewes. *Anim Feed Sci Tech* 186: 36-44.
43. Thering B, Graugnard D, Piantoni P, Loor J (2009) Adipose tissue lipogenic gene networks due to lipid feeding and milk fat depression in lactating cows. *J Dairy Sci* 92: 4290-4300.
44. Qi L, Yan S, Sheng R, Zhao Y, Guo X (2014) Effects of saturated long-chain fatty acid on mRNA expression of genes associated with milk fat and protein biosynthesis in bovine mammary epithelial cells. *Asian-Austral J Anim Sci* 27: 414-421.
45. Hussein M, Harvatine KH, Weerasinghe WMPB, Sinclair LA, et al. (2013) Conjugated linoleic acid-induced milk fat depression in lactating ewes is accompanied by reduced expression of mammary genes involved in lipid synthesis. *J Dairy Sci* 96: 3825-3834.