

Discostroma fericola* sp. nov. (Amphisphaeriaceae) and a key to species of *Discostroma

Barbara C. Paulus^{1,2}, Paul A. Gadek¹ and Kevin D. Hyde³

¹ School of Tropical Biology, James Cook University, Cairns, QLD 4870, Australia

² Current address: Landcare Research, Private Bag 92170, Auckland, New Zealand

³ Centre for Research in Fungal Diversity, Department of Ecology & Biodiversity,
The University of Hong Kong, Pokfulam Road, Hong Kong SAR, The People's
Republic of China

Paulus B.C., Gadek P.A. & Hyde K.D. (2006) *Discostroma fericola* sp. nov.
(Amphisphaeriaceae) and a key to species of *Discostroma*. *Sydowia* 58 (1): 76 - 90.

An investigation of microfungal diversity in a tropical rainforest of North Queensland, Australia, yielded a new species of *Discostroma* on decaying leaves of *Ficus pleurocarpa*. *Discostroma fericola* is described and illustrated and compared with similar species. A synopsis of the genus *Discostroma* and a key to the genus are provided.

Key words: Ascomycetes, *Discostromopsis*, leaf litter fungi.

A study to estimate microfungal diversity in decaying leaves of four rainforest tree species in a tropical rainforest in North Queensland, Australia (Paulus *et al.* 2003a, 2003b) resulted in the discovery of a new species of *Discostroma* Clem. on decaying leaves of *Ficus pleurocarpa* F. Muell. We provide descriptions and illustrations of *Discostroma fericola* sp. nov., comparisons with similar species and a synopsis of the genus in this paper.

Clements (1909) introduced *Discostroma* with a new combination, *D. rehmii* (Schnabl) Clem. Von Arx (1974) considered *D. rehmii* synonymous with *Metasphaeria massarina* Sacc., and introduced a new combination, *D. massarinum* (Sacc.) Arx (as *D. massarina*; Arx 1974). Currently 28 species are included in this genus (Müller & Loeffler 1957, Eriksson 1974, Brockmann 1976, Barr 1983, Sivanesan 1983, Eriksson 1992, Huhndorf 1992, Barr 1993 1994, Yuan & Barr 1994, Okane *et al.* 1996, Hatakeyama & Harada 2004). Of these, four species were described by Swart (1979) in *Discostromopsis* H.J. Swart, but his genus was considered synonymous with *Discostroma* by Sivanesan (1983) and the four species were redispersed to the later genus (Sivanesan & Shivas 2002b). Keys to species of *Discostroma* were provided by Brockmann (1976) and Barr (1994). Fungi included in *Discostroma* are mainly biotrophic (Brooks & El Alaily 1939; Kang *et al.* 1998, Samuels & Blackwell

2001), but fruiting bodies of *Discostroma* have also been reported from decaying plant material such as twigs (Brockmann 1976), leaves (Barr 1993, this study) and cones of *Pinus* (Müller & Loeffler 1957). The placement of *Discostroma* in the Amphisphaeriaceae *sensu stricto*, as suggested by its morphology and its mode of life (Samuels *et al.* 1987), was supported by molecular phylogenetic analyses (Kang *et al.* 1998, Jeewon *et al.* 2002, Smith & Hyde 2003). Using a molecular approach, Jeewon *et al.* (2002) also confirmed the anamorph-teleomorph connections between *Discostroma* and *Seimatosporium* Corda, which had been previously known from cultural studies alone (Müller & Shoemaker 1965, Brockmann 1976, Okane *et al.* 1996).

Materials and Methods


The study site comprises of upland rainforest near Topaz, Queensland, Australia (700 m a.s.l.). The forest has been classified as complex mesophyll rainforest (Type 1b rainforest; Tracey 1982) and has a high diversity of tree species. Decaying leaves were collected under six individual trees of *Ficus pleurocarpa*.

Slides were prepared from ascocarps removed from decaying leaves following incubation in humid chambers, which contained tissue paper moistened with sterile distilled water. Some slides were stained with Melzer's reagent to test amyloidity of the ascocarp apical apparatus. The range of measurements is derived from examining a minimum of 25 ascospores 20 ascocarps, 10 ascocarps per specimen mounted in sterile distilled water. The arithmetic mean and standard deviation of ascospore, ascocarp and ascocarpal dimensions were calculated in Microsoft Excel (Microsoft Corporation, Redmond). Spores, ascocarps and ascocarps were measured at their widest point. Slides were rendered semipermanent by the addition of 90% lactic acid. Photographs were taken on an Olympus BX50 microscope using an Olympus CA35 AD-4 camera. Photos were scanned and plates were assembled using Adobe Photoshop 5.0 (Adobe Systems Inc., San Jose). The type specimen is deposited at the Plant Pathology Herbarium (BRIP), Queensland, Department of Primary Industries.

Taxonomy

Discostroma ficicola Paulus, P. Gadek & Hyde K. D. sp. nov. Figs. 1–6.

Ascomata numerosa, epiphylla, nigra, globosa vel subglobosa, immersa vel semi-immersa, clypeo circulari et nigro, 112.5–175 × 112.5–150 µm. Ascii cylindrici vel angusti ellipsoidei, unitunicati, 8 spori uniseriati vel biseriati, annulo amyloideo, 2.5 × 1.5 µm, pulvillo dicto atramento scriptorio coeruleescente praediti, (80–) 90–110 × 10–12 µm. Ascospores ellipsoidae, uniseptatae, hyalinæ, guttulatae, 17–20 × 5–6.5 µm. Anamorphosis ignota.



Figs. 1 - 6. *Discostroma ficicola* (BRIP 29181). 1. Ascocarps on leaf surface. 2. Ascospores. 3. Paraphyses. 4. Ascus. 5. Ascospores. 6. Amyloid ascus apparatus (arrowed) and ascospores (note slight constriction at septa). Scale bars: 1 = 100 µm; 2 = 25 µm; 3, 5 - 6 = 5 µm; 4 = 10 µm.

Holotypus. AUSTRALIA, Queensland, Atherton Tablelands, Topaz, Old Boonjie Road, in decaying leaves of *Ficus pleurocarpa*, 8 Feb 2002, leg. B. Paulus and I. Steer, BRIP 29181.

Etymology. *ficicola* (Latin – *Ficus* = dweller), referring to the strangler fig *Ficus pleurocarpa* from which this fungus was identified.

Habitat. in decaying leaves of *Ficus pleurocarpa*.

Ascocarps numerous, mostly epiphyllous, black, globose or subglobose with circular black clypeus visible on leaf surface, submerged when young, semisubmerged with central papilla emerging when mature, macroscopically appearing as pustules on leaf surface, $112.5 - 175 \times 112.5 - 150$ µm (mean \pm SD = $138.4 \pm 17.6 \times 133.7 \pm 10.8$ µm, n = 10), peridium composed of several layers of dark brown, irregular cells. Paraphyses abundant, persistent, hyaline, smooth, septate, sometimes inflated at base with long, tapering apical

part, 3–9 µm wide at base, tapering to 2 µm wide at apex, which is obtuse or slightly inflated. Ascii unitunicate, cylindrical or narrowly clavate (when ascospores have obliquely or overlapping uniseriate arrangement) or narrowly ellipsoid (when ascospores are biseriate), 8-spored, apex rounded with a J+ subapical annulus, approximately 2.5 µm wide and 1.5 µm high, pulvillus staining slightly blue with ink, pedicel short with inflated blunt base, (80–) 90–110 µm × 10–12 µm (mean ± SD = 97.9 ± 10.3 × 10.6 ± 0.8 µm, n = 20). Ascospores oblong to ellipsoid, sometimes inequilateral, with one median septum, slightly constricted or not constricted at the septum, hyaline, smooth-walled, usually guttulate, 17–20 × 5–6.5 µm (mean ± SD = 18.7 ± 1.2 × 5.7 ± 0.7 µm, n = 25). Anamorph not observed.

Notes. For the generic placement of our taxon, we considered genera developing under clypei, with hyaline ascospores, and asci with an amyloid apical ring. *Hyponectria* Sacc., *Physalospora* Niessl and *Rachidicola* Hyde K.D. & J. Fröhl. were excluded on the basis of ascospore septation, and *Charonectria* Sacc. and *Arwidssonia* B. Erikss. on ascomatal differences (Niessl 1876, Saccardo 1878, 1880, Hyde & Fröhlich 1995, Sivanesan & Shivas 2002a, Wang & Hyde 1999). Within the Amphisphaeriaceae *sensu stricto*, *Ellurema* Nag Raj & W.B. Kendr. also has hyaline ascospores and an amyloid annulus but differs from our specimens in ascospore septation, ascus shape and abundance of paraphyses (Nag Raj & Kendrick 1985, Kang *et al.* 1999.). With the exception of *Discostroma*, other genera within the Amphisphaeriaceae differ in ascospore pigmentation and/or septation or amyloidity of the ascus ring (Kang *et al.* 1998). Several species of *Discostroma* (e.g. *D. cupulum*, *D. empetri*, *D. hyperboreum*, *D. ledi*, *D. rhododendri*, *D. succineum* and *D. tostum*) resemble *D. ficicola* in having ellipsoid, hyaline ascospores with strictly one single median septum and an amyloid ascus ring (Table 1). Two species, *D. pachystimae* and *D. caninum*, also form hyaline ascospores with predominantly one septum, but they may form additional septa (Table 1). *Discostroma ficicola* differs clearly from all species with similar ascospores in their lengths (key, Table 1).

Table 1 provides a summary of characters of 28 species of *Discostroma*. A key to the genus is included below.

Key to *Discostroma*

- | | |
|---|------------------------------|
| 1. Ascospores always muriform | 2 |
| 1. Ascospores predominantly with transverse septa only | 4 |
| 2. Ascospores dark brown, mostly with 3 transverse and one longitudinal septum, 18–20 (–34) × 8–10 (–12.5) µm | <i>Discostroma muricatum</i> |
| 2. Ascospores hyaline | 3 |

- 3. Ascospores with 3 transverse septa and 3 – 4 longitudinal septa $19 - 22 (- 24) \times 10 - 13 \mu\text{m}$ *Discostroma propendulum*
- 3. Ascospores with 3 – 7 (mostly 5) transverse septa and up to 3 longitudinal septa, pinkish in mass, $18 - 25 \times 8.5 - 11 \mu\text{m}$ *Discostroma massarinum*
- 4. Ascospores always with one transverse septum 5
- 4. Ascospore septation variable 11
- 5. Ascospores $13 - 16 \times 5 - 6 \mu\text{m}$, asci $65 - 80 \times 7.5 - 11.5 \mu\text{m}$ *Discostroma succineum*
- 5. Ascii larger 6
- 6. Ascospore length $\geq 14 \mu\text{m}$ 7
- 6. Ascospore length predominantly $\leq 14 \mu\text{m}$ 8
- 7. Ascospores $17 - 20 \times 5 - 6.5 \mu\text{m}$, asci $(80)90 - 110 \times 10 - 12 \mu\text{m}$ *Discostroma ficicola*
- 7. Ascospores $14 - 17 \times 7 - 8 \mu\text{m}$, asci $90 - 100 \times 11 - 12 \mu\text{m}$ *Discostroma hyperboreum*
- 8. Ascospores becoming yellowish to dull brown at maturity, $10.5 - 13 (- 16.5) \times 5 - 6.5 \mu\text{m}$, finely verruculose as longitudinal striae *Discostroma cupulum*
- 8. Ascospores remain hyaline at maturity, smooth 9
- 9. Ascospores $12 - 13.5 (- 15) \times 5.5 - 7 \mu\text{m}$, asci $80 - 93 \times 9 - 10.5 \mu\text{m}$ *Discostroma empetri*
- 9. Ascospore width $\leq 5.5 \mu\text{m}$ 10
- 10. On dead stems of *Epilobium*, ascospores $8 - 13.5 \times 3 - 4.5 \mu\text{m}$, ellipsoid to slightly curved, asci $62 - 85 \times 5 - 7 \mu\text{m}$ *Discostroma tostum*
- 10. On leaves of *Ledum groenlandicum*, ascospores $10.5 - 12 \times 4 - 4.5 \mu\text{m}$, ellipsoid, asci $63 - 75 \times 6 - 7 \mu\text{m}$ *Discostroma ledi*
- 10. On leaves of *Rhododendron californicum*, ascospores $10 - 13 \times 4.5 - 5.5 \mu\text{m}$, oblong ellipsoid, asci $60 - 88 \times 6.5 - 9 \mu\text{m}$ *Discostroma rhododendri*
- 11. Ascospores predominantly 1-septate but additional septa may develop 12
- 11. Ascospores predominantly with more than one septum or with approximately equal proportions of 1, 2 and 3 septa 15
- 12. Ascospore length mostly $\geq 12 \mu\text{m}$ 13
- 12. Ascospore length mostly $\leq 12 \mu\text{m}$ 14
- 13. Ascospores mostly 1-septate, rarely 3-septate, ellipsoid to fusoid, hyaline, $12 - 14 \times 4.5 - 5 \mu\text{m}$, ascomata erumpent *Discostroma pachystimae*
- 13. Ascospores mostly 1-septate but also 2- or 3-septate, ellipsoid, hyaline to yellowish, $12 - 15 (- 17) \times 5.3 - 6.5 \mu\text{m}$, ascomata visible as round, raised dark brown shiny areas *Discostroma caninum*
- 14. Ascospores $10 - 12 (- 13) \times 4 - 5 (- 5.3) \mu\text{m}$, 1-septate, sometimes 2-3-septate, asci $75 - 100 \times 5.3 - 7.4 \mu\text{m}$ *Discostroma rosae*

14. Ascospores $7 - 12 \times 3 - 4 \mu\text{m}$, usually 1-septate, sometimes with further transverse septa, asci $55 - 70 \times 5 - 7 \mu\text{m}$ *Discostroma strobiligenum*
15. Ascospores mostly 2-septate, becoming 3-septate when mature, $16 - 20 \times 14 - 22 \mu\text{m}$, hyaline when young, brown when mature *Discostroma tricellulare*
15. Ascospores mostly 3-septate or approximately equal proportions of 1, 2 or 3 septa 16
16. Ascospores 1, 2, or 3 septa often in the same ascus, $11 - 14 \times 4.5 - 5 \mu\text{m}$ *Discostroma polymorphum*
16. Ascospores predominantly 3-septate 17
17. Ascal ring distinct and J+ 18
17. Ascal ring indistinct and J- 23
18. Ascospores hyaline to yellowish 19
18. Ascospores pigmented 22
19. Ascospores with predominantly 3 transverse septa and, rarely, one longitudinal septum, $16.5 - 20(-25) \times 7.5 - 9(-10) \mu\text{m}$, asci $105 - 130 \times 8.5 - 13.3 \mu\text{m}$ *Discostroma sanguineae*
19. Ascospores with only transverse septa 20
20. Ascospores $12 - 18 \times 4 - 6 \mu\text{m}$, asci $70 - 111 \times 6 - 10 \mu\text{m}$ *Discostroma botan*
20. Ascospore width $\geq 6 \mu\text{m}$ 21
21. Ascospores $18 - 26 \times 6 - 8 \mu\text{m}$, asci $90 - 130 \times 9 - 11 (-15)$, ascomata $200 - 250 (-350) \mu\text{m}$, on leaves of *Pentaphylloides fruticosa* *Discostroma fruticosum*
21. Ascospores $15.5 - 24 \times 6.5 - 9 \mu\text{m}$, asci $100 - 160 \times 9 - 13 \mu\text{m}$, ascomata $300 - 450 \times 250 - 495 \times 250 - 495 \mu\text{m}$, on leaves of *Rubus* spp. *Discostroma rubicola*
22. Ascospores olivaceous, $15 - 17 \times 9 \mu\text{m}$, asci $100 \times 9 - 10 \mu\text{m}$, ascomata $270 - 450 \times 200 - 300 \mu\text{m}$ *Discostroma fuscellum*
22. Ascospores light brown, $15 - 18 \times 6.5 - 9 \mu\text{m}$, asci $117 - 150 \times 8 - 10 \mu\text{m}$, ascomata $400 - 600 \times 210 - 400 \mu\text{m}$ *Discostroma saccardoanum*
23. Ascospores brown, echinulate, $11.5 - 13.8 \times 4 - 4.5 \mu\text{m}$ *Discostroma osyridis*
23. Ascospore hyaline 24
24. Ascospores $14 - 18 \times 4 - 5 \mu\text{m}$, anamorph *Seimatosporium kriegerianum* *Discostroma callistemonis*
24. Ascospore length $\geq 17 \mu\text{m}$ 25
25. On dead leaves of *Leptospermum juniperinum*, ascospores $17 - 23 \times 4.5 - 6 \mu\text{m}$, anamorph *Seimatosporium leptospermi* *Discostroma leptospermi*
25. Ascomata on dead leaves of *Melaleuca* 26
26. Anamorph *Seimatosporium elegans*, on leaves of *Melaleuca ericifolia*, ascospores $17 - 23 \times 4.5 - 6 \mu\text{m}$.. *Discostroma elegans*

26. Anamorph *Seimatosporium dilophosporum*, on dead leaves of *Melaleuca squarrosa*, ascospores 19–21 × 4–6 µm *Discostroma stoneae*

Discussion

Discostroma is defined by a relatively broad generic concept. It is characterized by the presence of stroma and/or clypeus, ostiolate peritheciium, unitunicate asci with an amyloid or inamyloid apical annulus, pulvillus or plug-like structure in the ascocal apex which stains with ink (not consistently noted by authors of new species descriptions) and ellipsoid, hyaline or pigmented ascospores, which may be transversely septate or muriform, and are usually arranged uniseriately within the ascus. For many species, a *Seimatosporium* anamorph (synonym: *Sporocadus* Corda *sensu* Brockmann 1976) with holoblastic conidial ontogeny and percurrent enteroblastic proliferation of the conidiogenous cell has been reported. Molecular data will show whether the genus is monophyletic as currently circumscribed.

Ascomatal morphology deserves further examination in the genus *Discostroma*, particularly as stromata and clypei development may vary depending on substratum texture and developmental stage (Swart 1979, Lu & Hyde 2000, this paper). Rudimentary information is currently available only for some *Discostroma* species (Table 1). They seem to have been described mainly on the basis of their natural substrata (e.g., *D. ledi*, *D. tostum* and *D. rhododendri*; Table 1, Key). Fungal plant pathogens and endophytes commonly display a degree of host-specificity (Zhou & Hyde 2001, Photita *et al.* 2004). Because some *Discostroma* species have been reported only from particular plant genera, the inclusion of substratum as an important taxonomic character may be justified. However, ascomata of *Discostroma* have also been reported from decaying plant material and do not seem to be host-specific (Müller & Loeffler 1957, Brockmann 1976, Swart 1979, Barr 1993, this study). In addition, we observed that *D. ficicola* was able to colonise sterilised leaves of *Ficus pleurocarpa* at different stages of decay (data not shown). Further studies are required to confirm whether some species of *Discostroma* are saprobic for the major part of their life cycle and to confirm or refuse substratum or host as a valid taxonomic character.

Acknowledgements

Funding for this project was provided by: the Centre for Research in Fungal Diversity, Department of Ecology & Biodiversity,

University of Hong Kong; the Cooperative Research Centre for Rainforest Ecology and Management and the School of Tropical Biology, James Cook University. B.P. thanks N. Tucker for providing access to one of the collection sites, S. McKenna for identification of plant species and C. Pearce and I. Steer for their company on collection trips. B. Bussaban and C. McCulloch are gratefully acknowledged for assistance with photography, and B. Bussaban and H. Leung for assistance in obtaining literature. E. McKenzie, S. Pennycook and two anonymous reviewers are thanked for critical comments on the manuscript.

References

- Arx von J.A. (1974) *The genera of fungi sporulating in pure culture*. J. Cramer, Vaduz, Switzerland.
- Barr M.E. (1983) Muriform ascospores in class ascomycetes. *Mycotaxon* **18**: 149 – 157.
- Barr M.E. (1993) Redisposition of some taxa described by J.B. Ellis. *Mycotaxon* **46**: 45 – 76.
- Barr M.E. (1994) Notes on the Amphisphaeriaceae and related families. *Mycotaxon* **51**: 191 – 224.
- Brockmann I. (1976) Untersuchungen über die Gattung *Discostroma* Clements (Ascomycetes). *Sydowia* **28**: 275 – 338.
- Brooks F.T., El Alaily Y.A.S. (1939) A canker and die-back of roses caused by *Griphosphaeria corticola*. *Annals Applied Biology* **26**: 231 – 226.
- Clements F.E. (1909) *Genera of fungi*. H. W. Wilson Company, Minneapolis, U.S.A.
- Eriksson B. (1974) On ascomycetes on Diapensiales and Ericales in Fennoscandia. 2. Pyrenomyces. *Svensk Botanisk Tidskrift* **68**: 192 – 234.
- Eriksson O.E. (1992) *The non-lichenized pyrenomyces of Sweden*. Lund, Sweden.
- Hatakeyama S., Harada Y. (2004) A new species of *Discostroma* and its anamorph *Seimatosporium* with two morphological types of conidia, isolated from the stems of *Paeonia suffruticosa*. *Mycoscience* **45**: 106 – 111.
- Huhndorf S. (1992) Systematics of *Leptosphaeria* species found on the Rosaceae. *Illinois Natural History Survey Bulletin* **34**: 479 – 534.
- Hyde K.D., Fröhlich J. (1995) Fungi from palms. XXIII. *Rachidicola* gen. et sp. nov. *Sydowia* **47**: 217 – 222.
- Jeewon R., Liew E.C.Y., Hyde K.D. (2002) Phylogenetic relationships of *Pestalotiopsis* and allied genera inferred from ribosomal DNA sequences and morphological characters. *Molecular Phylogenetics and Evolution* **25**: 378 – 392.
- Kang J.C., Hyde K.D., Kong R.Y.C. (1999) Studies on Amphisphaerales: The Amphisphaeriaceae (*sensu stricto*). *Mycological Research* **103**: 53 – 64.
- Kang J.C., Kong R.Y.C., Hyde K.D. (1998) Studies in the Amphisphaerales 1. Amphisphaeriaceae (*sensu stricto*) and its phylogenetic relationships inferred from 5.8S rDNA and ITS 2 sequences. *Fungal Diversity* **1**: 147 – 157.
- Lu B.-S., Hyde K.D. (2000) *A world monograph of Anthostomella*. Fungal Diversity Press, Hong Kong, SAR, China.
- Müller E., Loeffler W. (1957) Über die Gattung *Clathridium* (Sacc.) Berl. *Sydowia* **11**: 116 – 120.

- Müller E., Shoemaker R. A. (1965) The ascogenous state of *Seimatosporium* (= *Monoceras*) *kriegerianum* on *Epilobium* species. *Canadian Journal of Botany* **43**: 1343 – 1345.
- Nag Raj T.R., Kendrick W.B. (1985) *Ellurema* gen. nov., with notes on *Lepteutypus cisticola* and *Seiridium canariense*. *Sydowia* **38**: 178 – 193.
- Niessl von Mayendorf G. (1876) Notizen über neue und kritische Pyrenomyceten. *Verhandlungen des Naturforschenden Vereines in Brünn* **14**: 156 – 218.
- Okane I., Nakagiri A., Ito T. (1996) *Discostroma tricellulare* a new endophytic ascomycete with a *Seimatosporium* anamorph isolated from *Rhododendron*. *Canadian Journal of Botany* **74**: 1338 – 1344.
- Paulus B., Gadek P., Hyde K.D. (2003a) Estimation of microfungal diversity in tropical rain forest leaf litter using particle filtration: the effects of leaf storage and surface treatment. *Mycological Research* **107**: 748 – 756.
- Paulus B. (2003b) Two new species of *Dactylaria* (anamorphic fungi) and an update of species in *Dactylaria* *sensu lato*. *Fungal Diversity* **14**: 145 – 158.
- Photita W., Lumyong S., Lumyong P., McKenzie E.H.C., Hyde K.D. (2004) Are some endophytes of *Musa acuminata* latent pathogens? *Fungal Diversity* **16**: 131 – 140.
- Saccardo P.A. (1878) Fungi veneti novi vel critici. Series VIII. *Michelia* **1**: 239 – 275.
- Saccardo P.A. (1880) Fungi Gallici lecti a cl. viris P. Brunaud, Abb. Letendre, A. Malbranche, J. Therry, vel editi in *Mycotheca Gallica* C. Roumégouëti. Series II. *Michelia* **2**: 39 – 135.
- Samuels G.J., Blackwell M. (2001) Pyrenomycetes fungi with perithecia. In *The Mycota*. A comprehensive treatise on fungi as experimental systems for basic and applied research. (eds Esser K., Lemke P.A.) Springer-Verlag, Berlin, Germany: 221 – 255.
- Samuels G.J., Mueller E., Petrini O. (1987) Studies in the Amphisphaeriaceae (*sensu lato*) 3. New species of *Monographella* and *Pestalosphaeria*, and two new genera. *Mycotaxon* **28**: 473 – 499.
- Sivanesan A. (1983) Studies on Ascomycetes. *Transactions of the British Mycological Society* **81**: 313 – 332.
- Sivanesan A., Shivas R.G. (2002a) New species from each of the pyrenomycete genera *Hyponectria*, *Physalospora* and *Trichosphaeria* from Queensland, Australia. *Fungal Diversity* **9**: 169 – 174.
- Sivanesan A., Shivas R.G. (2002b) New species of foliicolous Loculoascomycetes on *Dysoxylum*, *Melaleuca* and *Syzygium* from Queensland, Australia. *Fungal Diversity* **11**: 151 – 158.
- Smith G.J.D., Hyde K.D. (2003) The Xylariales, a monophyletic order containing seven families. *Fungal Diversity* **13**: 175 – 208.
- Swart H.J. (1979) Australian leaf inhabiting fungi X. *Seimatosporium* species on *Callistemon*, *Melaleuca* and *Leptospermum*. *Transactions of the British Mycological Society* **73**: 213 – 221.
- Tracey J.G. (1982) *The vegetation of the humid tropical region of north Queensland*. CSIRO, Melbourne, Australia.
- Wang Y.-Z., Hyde K.D. (1999) *Hyponectria buxi* with notes on the *Hyponectriaceae*. *Fungal Diversity* **3**: 159 – 172.
- Yuan Z.-Q., Barr M.E. (1994) New ascomycetous fungi on bush cinquefoil from Xinjiang, China. *Sydowia* **46**: 329 – 337.
- Zhou D., Hyde K.D. (2001) Host-specificity, host-exclusivity and host-recurrence in saprobic fungi. *Mycological Research* **105**: 1449 – 1457.

(Manuscript accepted 20 Dec. 2005; Corresponding Editor: U. Peintner)

Table 1. Summary of morphological characters of species in *Discostroma* (in alphabetical order)

Species	Ascomata	Ascic ^a	Ascospores ^b	Habitat	Anamorph
<i>Discostroma botan</i> Sat. Hatakeyama & Y. Harada Mycoscience 45: 106–111. (2004)	200–250 × 300–360 µm, perithecia slightly papillate, immersed.	(62–)70–111 × 6–10 µm, J*, uniseriate.	12–18 × 4–6 µm, hyaline, 3 transverse septa.	Stems of <i>Paeonia</i> <i>suffruticosa</i>	<i>Sematosporium</i> <i>botan</i> Sat. Hatakeyama & Y. Harada
<i>Discostroma callistomoides</i> (H.J. Swart) Sivan. Trans. Br. mycol. Soc. 73: 217 (1979)	170–190 × 80–100 µm, perithecia under distinct dark clypeus.	65–75 × 9–10 µm, J*, partly biserrate.	14–18 × 4–5 (-4.5) µm, hyaline, 3 transverse septa.	Leaves of <i>Calispermum</i> <i>paludosus</i>	<i>Sematosporium</i> <i>kriegerianum</i> (Bres.) Morgan-Jones & B. Sutton
<i>Discostroma curinum</i> Brockmann Sydowia 28: 328 (1976)	230–360 × 150– 200 (300) µm, stroma of intraepidermal, thick- walled hyphae.	85–120 × 6–9 µm, J*.	12–15 (-17) × 5.3– 6.5 µm, hyaline to yellowish, mostly 1-septate, but also 2- or 3-septate.	Dead twigs of <i>Rosa</i> spp.	<i>Sporocadus curinus</i> (Brun.) Brockmann
<i>Discostroma cupulum</i> (Ellis) M. E. Barr Mycotaxon 46: 55 (1993)	200–280 × 100–180 µm, clypeus present.	55–75 × 6–10 µm, J*.	10.5–13 (-16.5) × 5–6.5 µm, hyaline becoming light yellowish brown to dull brown, 1-septate.	Dead overwintered, attached leaves of <i>Quercus coccinea</i>	Not reported
<i>Discostroma elegans</i> (H.J. Swart) Sivan. Trans. Br. mycol. Soc. 73: 217 (1979)	120–130 × 30–90 µm, perithecia frequently flattened by resistance of host tissue.	70–80 × 10 µm, J*, partly biserrate.	18–22 × 4–5.5 µm, hyaline, 3 transverse septa.	Dead leaves of <i>Metaleuca ericifolia</i>	<i>Sematosporium</i> <i>elegans</i> H.J. Swart
<i>Discostroma empetri</i> M. E. Barr Mycotaxon 51: 200 (1994)	200–290 µm in diameter, surrounded by circular, shiny clypeus.	80–93 × 9–10.5 µm, J*.	12–13.5 (-15) × 5.5–7 (-8) µm, hyaline, 1-septate.	Dead leaves of <i>Empetrum nigrum</i>	Not reported

Species	Ascomata	Ascii ^a	Ascospores ^b	Habitat	Anamorph
<i>Discostroma fericola</i> Paulus, Gadek & K.D. Hyde	112.5 – 175 × 112.5 – 150 µm, black, circular clypeus visible, at maturity eruptant and appearing as pustules.	(80 –) 90 – 110 × 10 – 12 µm, J*, uniseriate or biseriate	17 – 20 × 5 – 6.5 µm, hyaline, 1-septate.	Decaying leaves of <i>Ficus pfeurocorpa</i>	Not observed
<i>Discostroma fruticosum</i> Z. Q. Yuan & M. E. Barr	200 – 250(-350) µm, eruptant to superficial.	90 – 130 (- 150) µm, J*.	18 – 26 × 6 – 8 µm, hyaline, (1 –) 3(- 5) septate.	Twigs of <i>Pentaphylloides</i> <i>fruticosa</i>	Not reported
<i>Discostroma fuscellum</i> (Berk. & Broome) Huhndorf Pl. Nat. Hist. Surv. Bull. 34: 520 (1987)	300 – 500 µm diameter, 150 – 250 µm high, beneath a blackened clypeus.	(101) 115 – 132 (- 137) × 7 – 12 (- 14) µm, J*.	15 – 21 × 5 – 9.5 µm, hyaline, 3 (seldom 4 or more) transverse septa, sometimes with 1 longitudinal septum.	<i>Cornus</i> , <i>Prunus</i> , <i>Ribes</i> , <i>Ros</i> , <i>Rubus</i> , <i>Viburnum</i> and <i>Vaccinium</i> spp.	<i>Seimatosporium</i> <i>lichenicola</i> (Conda) Shoemaker & E. Müll.
<i>Discostroma hyperboricum</i> (P. Karst.) O.E. Erikss. Non-lichenized Pyronomyctetes of Sweden, p. 7 (1992)	250 µm in diameter, hypophyllous.	90 – 100 × 11 – 12 µm, J*.	14 – 17 × 7 – 8 µm, hyaline, 1-septate.	Leaves of <i>Andromeda</i> <i>tetragona</i>	Not reported
<i>Discostroma tedi</i> M. E. Barr	200 – 245 × 170 – 245 µm, beneath a shallow clypeus.	63 – 75 × 6 – 7 µm, J*.	10.5 – 12 × 4.5 – 5 µm, hyaline, 1-septate.	Leaves of <i>Ledum</i> <i>groenlandicum</i>	Probably <i>Sporocadus</i> sp.

Species	Ascocarpha	Asci ^a	Ascospores ^b	Habitat	Anamorph
<i>Discostroma leptospermii</i> (H. J. Swart) Sivan. Trans. Br. mycol. Soc. 73: 218 (1979)	140 – 156 × 120 µm, under a small clypeus, inversely pear-shaped.	60 – 80 × 8 – 10 µm, J, partly biserrate.	17 – 23 × 4.5 – 6 µm, hyaline, 3 transverse septa.	Dead leaves of <i>Leptospermum</i> <i>jumperinum</i>	<i>Séimatosporium</i> <i>leptospermi</i> (Corda) B. G. Bagn. & Sheridan
<i>Discostroma massarinum</i> (Sacc.) Arix Arx (1974) Genera of Fungi Sporulating in Pure Culture, 2 nd ed., p. 131	400 – 500 × 350 – 450 µm, stroma more or less developed, visible as pustules.	130 – 180 × 9 – 11 µm, J.	18 – 25 × 8.5 – 11 µm, hyaline, in mass pinkish, 3 – 7 transverse septa and up to 3 longitudinal septa.	Dead twigs of <i>Ribes</i> spp.	<i>Séimatosporium</i> <i>ribis-alpini</i> (Corda) Shoemaker & E. Müll.
<i>Discostroma muricatum</i> (Ellis & Everh.) M. E. Barr Mycotaxon 13: 151 (1983)	up to 550 × 365 µm, superficial with bases ingrown in periderm.	(80 –) 100 – 110 × 12 – 17 µm, J.	18 – 20 (- 34) × 8 – 10 (12.5) µm, dark brown, 3-(4 – 6)-septate, one longitudinal septum in one or more cells.	Periderm of unknown tree	Not reported
<i>Discostroma osyridis</i> Sivan. Trans. Br. mycol. Soc. 81: 325 (1983)	160 – 220 × 150 – 175 µm, sparsely developed clypeus and little or no stromatic tissue.	75 – 80 × 5 – 6 µm, J.	11.5 – 13.8 × 4 – 4.5 µm, brown, transversely 3-septate, not constricted at septa, echinulate under oil immersion.	Stem of <i>Osyris alba</i>	<i>Séimatosporium</i> sp. Conidiophores hyaline, annelidic, 10 – 22 × 1.5 – 2 µm, conidia ellipsoid, brown, 3-septate, 11 – 11.5 × 4 – 4.3 µm
<i>Discostroma pachystimae</i> M. E. Barr & Rogerson Mycotaxon 51: 201 (1994)	330 × 275 µm, erumpent.	80 – 100 × 7 – 7.5 µm, slowly J.	12 – 14 × 4.5 – 5 µm hyaline, 1-septate, rarely 3-septate.	Twigs of <i>Pachystima</i> <i>myrsinites</i>	Not reported

Species	Ascomata	Ascii ^a	Ascospores ^b	Habitat	Anamorph
<i>Discostroma polymorphum</i> Brockmann Sydowia 28: 306 (1976)	230 – 370 (400) × 150 – 240 µm, stroma above and below ascoma.	85 – 110 (117) × 7 – 10 µm, J*.	1.1 – 14 × 5.6 – 7 µm, greyish brown, 1-, 2- and 3-septate,	Dead twigs of <i>Rosa</i> spp.	Hyaline, annelidic conidiophore, conidia 2- to 3-septate, elongate or slightly clavate, brown, 11 – 15.5 × 4 – 5.5 µm
<i>Discostroma propinquum</i> (P. Karst.) Brockmann Sydowia 28: 334 (1976)	350 – 400 × 350 – 450 µm, with clypeus, erumpent through epidermis.	110 – 140 × 18 – 20 µm, J*.	19 – 22 (– 24) × 10 – 13 µm, hyaline, 3 transverse septa and Arctostaphylos longitudinal septum in each segment.	Bud scales and stems of <i>Arctostaphylos alpina</i> and stems of <i>A. nubra</i>	Not reported
<i>Discostroma rhododendri</i> M. E. Barr Mycotaxon 51: 202 (1994)	200 – 220 × 150 – 180 µm, with small closely adherent clypeus.	60 – 88 × 6.5 – 9 µm, J*.	10 – 13 × 4.5 – 5.5 µm, hyaline, 1-septate.	Leaves of <i>Rhododendron californicum</i>	Not reported
<i>Discostroma rosae</i> Brockmann Sydowia 28: 321 (1976)	260 – 370 (415) × 150 – 300 µm, intracarpidermal brown hyphae, visible as round to elongate, dark brown or black, shiny areas.	75 – 100 × 5.3 – 7.4 µm, J*.	10 – 12 (– 13) × 4 – 5 (– 5.3) µm, hyaline, mostly 1 septate, sometimes 2- or 3-septate.	Dead twigs of <i>Rosa</i> spp.	<i>Seminatosporium rosae</i> Corda
<i>Discostroma rubicola</i> (Ellis & Everh.) M. E. Barr Mycotaxon 46: 56 (1993)	300 – 350 × 250 – 495 µm, erumpent.	100 – 160 × 9 – 13 µm, J*.	15.5 – 24 × 6.5 – 9 µm, hyaline, (1 – 2) 3- (4 –) septate.	Canes of <i>Rubus</i> deliciosus and other <i>Rubus</i> spp.	Acervular conidio-mata, conidia 22 – 30 × 9 – 10 (– 12) µm, yellowish or clear brown, paler toward base, conidia 3-distoseptate

Species	Ascomata	Asci ^a	Ascospores ^b	Habitat	Anamorph
<i>Discostroma saccardoanum</i> (Jacz.) Brockmann Sydowia 28: 304 (1976)	400 – 600 × 210 – 400 µm, stroma not well developed, visible as pustules.	117 – 150 × 8 – 10 µm, J ^c .	15 – 18 × 2 – 6 µm, light brown, 3 transverse septa and rarely with 1 longitudinal septum.	Dead twigs of <i>Ribes</i> spp.	Hyaline, annelidic conidiophores, elongate conidia, 1.4 – 1.7 × 5 – 6.3 µm, light to medium brown, 3 septate, no appendages
<i>Discostroma sanguineae</i> Brockmann Sydowia 28: 309 (1976)	330 – 430 × 250 – 370 µm, stroma in epidermis more developed near ostiole..	105 – 130 × 8.5 – 13.3 µm, J ^c .	16.5 – 20 (~25) × 7.5 – 9 (~10) µm, hyaline to yellow, with 3 transverse septa, rarely 1 longitudinal septum.	Dead twigs of <i>Cornus sanguinea</i>	<i>Sporocadus fiedleri</i> Rabenh.
<i>Discostroma stoneae</i> (H. J. Swart) Sivan. Trans. Br. mycol. Soc. 73: 217 (1979)	110 – 120 (~150) × 80 – 100 µm, under a well-developed clypeus.	70 – 80 × 9 – 12 µm, J ^c ; partly biseriate.	19 – 21 × 4 – 6 µm, hyaline, 3 transverse septa.	Dead leaves of <i>Melaleuca squarrosa</i>	<i>Seimatosporium dilophosporum</i> (Cooke) B. Sutton
<i>Discostroma strobiligenum</i> (E. Mill. & Loeffler) Brockmann Sydowia 28: 334 (1976)	150 – 250 µm, stroma immersed in substrate, clypeus dark.	55 – 70 × 5 – 7 µm, J ^c .	7 – 12 × 3 – 4 µm, hyaline, usually 1-septate, sometimes with further transverse septa.	Fallen cones of <i>Pinus sylvestris</i>	Not reported
<i>Discostroma succinum</i> (Roberge ex Desm.) M. E. Barr Mycobion 51: 202 (1994)	Loosely dispersed or aggregated, submerged perithecia, in greyish spots.	65 – 80 × 7.5 – 11.5 µm, J ^c .	13 – 16 × 5 – 6 µm, hyaline, 1-septate.	Dry leaves of <i>Quercus</i> spp.	Not reported

Species	Ascomata	Asci ^a	Ascospores ^b	Habitat	Anamorph
<i>Discostroma tostum</i> (Berk. & Broome) Brockmann Sydowia 28: 319 (1976)	250 – 370 × 150 – 200 µm, well developed clypeus.	62 – 85 × 5 – 7 µm, J ^c .	8 – 13.5 × 3 – 4.5 µm, hyaline, 1 septate.	Dead stems of <i>Epilobium</i> spp.	<i>Seimatosporium</i> <i>passerini</i> (Sacc.) Brockmann
<i>Discostroma tricellulare</i> Okane, Nakagiri & Tad. Ito Can. J. Bot. 74: 1339 (1996)	116 – 174 × 120 – 180 µm (on CMA 140 – 220 × 160 – 275 µm).	52 – 70 × 14 – 22 µm (on CMA 56 – 90 × 16 – 26 µm), J ^c , biserrate.	16 – 20 × 6 – 9 µm (on CMA 17 – 22 × 7 – 10 µm), brown to olivaceous when mature, mostly 2-septate, a few 3-septate when mature.	Leaves of <i>Rhododendron</i> spp. <i>azaleae</i> Okane, Nakagiri & Tad. Ito	<i>Seimatosporium</i>

^a all ascii are 8-spored, spores uniseriate unless otherwise stated; J or J^c refers to amyloidity of the apical apparatus.

^b spores are smooth unless otherwise stated