



RESILIENCE TO DROUGHTS IN MAMMALS: A CONCEPTUAL FRAMEWORK FOR ESTIMATING VULNERABILITY OF A SINGLE SPECIES

TASMIN L. RYMER

College of Marine & Environmental Sciences and Centre for Tropical

Environmental & Sustainability Sciences, James Cook University

Cairns, Queensland 4870 Australia

School of Animal, Plant & Environmental Sciences, University of the Witwatersrand

Johannesburg 2050 South Africa

E-MAIL: TASMIN.RYMER@JCU.EDU.AU

NEVILLE PILLAY

School of Animal, Plant & Environmental Sciences, University of the Witwatersrand

Johannesburg 2050 South Africa

E-MAIL: NEVILLE.PILLAY@WITS.AC.ZA

CARSTEN SCHRADIN

School of Animal, Plant & Environmental Sciences, University of the Witwatersrand

Johannesburg 2050 South Africa

*Institut Pluridisciplinaire Hubert Curien—Département Ecologie,
Physiologie et Ethologie, Université de Strasbourg*

Strasbourg 67087 France

CNRS, UMR7178

Strasbourg 67087 France

E-MAIL: CARSTEN.SCHRADIN@IPHC.CNRS.FR

KEYWORDS

allostasis, homeostasis, metabolism, osmoregulation, preadaptation, thermoregulation

ABSTRACT

The frequency and severity of droughts in certain areas is increasing as a consequence of climate change. The associated environmental challenges, including high temperatures, low food, and water availability, have affected, and will affect, many populations. Our aims are to review the behavioral, physiological, and morphological adaptations of mammals to arid environments, and to aid researchers and nature conservationists about which traits they should study to assess whether or not their study species will be able to cope with droughts. We provide a suite of traits that should be considered when making predictions about species resilience to drought. We define and differentiate between general adaptations, specialized adaptations, and exaptations, and argue that specialized adaptations are of little interest in establishing how nondesert specialists will cope with droughts. Attention should be placed on general adaptations of semidesert species and assess whether these exist as exaptations in nondesert species. We conclude that phenotypic flexibility is the most important general adaptation that may promote species resilience. Thus, to assess whether a species will be able to cope with increasing aridity, it is important to establish the degree of flexibility of traits identified in semidesert species that confer a fitness advantage under drying conditions.

INTRODUCTION

ANTHROPOGENIC-induced environmental change, including habitat loss (Galbraith et al. 2002), fragmentation (Fahrig 2003), and climate change (Etterson and Shaw 2001; Alley et al. 2003), is accelerating at a rate unprecedented in Earth's history (Etterson and Shaw 2001; Alley et al. 2003). In fact, the rate of vertebrate species climatic niche evolution is considerably slower than the current predicted rates of anthropogenic environmental change (Quintero and Wiens 2013). The accelerated rates of human-induced environmental change could thus be driving a sixth mass extinction event, requiring conservation efforts to be focused on reducing the rate of extinctions (Pimm 2009; Black et al. 2011; Harley 2011). However, we are currently unable to predict which species will be threatened by extinction under increased environmental change and which species will survive (Moritz and Agudo 2013).

Many studies have assessed the probability and potential of species to adapt to rapid environmental change via evolutionary adaptation (e.g., evolutionary rescue; Gonzalez et al. 2013; Box 1). However, evolutionary adaptation is generally a slow process, occurring over generations (Hoffmann and Sgrò 2011), and may occur too slowly for species to respond to rapid environmental and climate change. Under such circumstances, species may become

extinct, change their distribution range, or show an active response to change through adaptive (Breed et al. 2011; Box 1) or exaptive (Ghalambor et al. 2007; Box 1) phenotypic plasticity. Interestingly, a number of local declines and extinctions due to environmental change have been associated with changes in species interactions (Cahill et al. 2013; Ockendon et al. 2014). However, an individual's ability to cope with its prevailing environment is also dependent on its behavior, physiology, and morphology. These responses to abiotic factors then likely also influence species interactions, for example, through changes in activity patterns, ultimately further impacting extinction risk.

One of the main consequences of global climate change is an increase in the frequency and intensity of droughts (IPCC 2007; Dai 2011). Regions where droughts have become, and are projected to become, longer and more frequent include the Mediterranean, central Europe, central North America, and southern Africa (IPCC 2007). Although the latest IPCC report (2013) mentions a lower confidence for the increased incidence of droughts since the 1950s, it is still predicted that droughts are likely to increase in some areas in the future. Droughts are a normal part of climate variation, which means that, independent of global climate change, natural populations have and will continue to face this problem repeatedly. For example, in the last 500 years,

BOX 1
Definition of terms

Term	Definition
<i>Adaptive phenotypic plasticity</i>	Alteration of some aspect of the phenotype in response to prevailing environmental conditions, which increases an individual's chance of survival and reproduction (maximizes fitness; Charmantier et al. 2008; Ozgul et al. 2009, 2010; Piersma and van Gils 2010) and that may facilitate population persistence and potentially evolutionary adaptation (Hoffmann and Sgrò 2011). Traits that specifically evolved in response to historical selection pressures and have current utility in the same context (phenotypic adaptation).
<i>Aridity index</i>	$I_m = \frac{(100S - 60D)}{PE_t}$; where I_m = index of moisture availability; S = total moisture surplus (mm) from all months with a water surplus; D = total of all monthly deficiencies (mm); and PE = annual potential evapotranspiration (mm; Meigs 1953). Both S and D are represented by the difference between monthly precipitation and monthly potential evapotranspiration (Nash and Engineering Group Working Party 2012). Deserts have an aridity index below 0.2, semideserts between 0.2 and 0.5.
<i>Behavioral evader</i>	Animals able to escape environmental extremes using primarily behavioral and physiological adaptations (Willmer et al. 2000).
<i>Developmental plasticity</i>	Nonreversible phenotypic plasticity occurring during ontogeny (Piersma and Lindström 1997).
<i>Drought</i>	Current precipitation falls significantly below the long-term mean of a defined area (Dai 2011). Is defined on a temporal scale and can occur in any environment.
<i>Drought-affected areas</i>	Areas that are neither deserts nor semideserts, but experience unpredicted droughts, which might occur more often and become more intense in the future due to climate change (Li et al. 2009). In the future, increasingly more areas might become drought prone due to climate change.
<i>Environmental stressor</i>	Any stimulus that could affect an individual negatively if it cannot respond to it via a specific stress response, which itself is costly (e.g., Charmandari et al. 2005). Typically uncontrollable for the individual, but can be predictable (e.g., seasonal changes in food) or unpredictable (e.g., a storm; Wingfield 2013).

continued

BOX 1
Continued

Term	Definition
<i>Evolutionary adaptation</i>	Change in the relative gene frequencies of alleles (and associated phenotypic changes) within a population over multiple generations, with a corresponding increase in species fitness over time (Pianka 2011; Rezende and Diniz-Filho 2012).
<i>Exaptation</i>	Formerly called preexisting adaptations or preadaptations. Traits that evolved in response to historical selection pressures, which have current utility but in a different context (i.e., different environment or selection pressures; Gould and Lewontin 1979).
<i>Exaptive phenotypic plasticity</i>	Alteration of traits in response to prevailing environmental conditions, which specifically evolved in response to historical selection pressures and have current utility in a different context (exaptation), increasing an individual's chance of survival and reproduction (increasing fitness) in novel environments (Ghalambor et al. 2007).
<i>Interannual rainfall variability</i>	Expressed as a percentage of the mean deviation of precipitation from the average over the average precipitation (Nash and Engineering Group Working Party 2012).
<i>Phenotypic flexibility</i>	Reversible phenotypic plasticity (Piersma and Lindström 1997).
<i>Physiological endurer</i>	Animals that tolerate environmental extremes using physiological and morphological adaptations (Willmer et al. 2000).
<i>Predictable environment</i>	Resources spatially and/or temporally heterogeneous; selection pressures imposed on populations over multiple generations are seasonal/cyclical (e.g., Negus et al. 1986; Ylönen and Ronkainen 1994); species show evolutionary adaptation and phenotypic plasticity in response to change (Ledón-Rettig et al. 2008; Rymer et al. 2013).
<i>Resilience</i>	The ability of a species (populations or individuals) to persist following a disturbance or disturbances (Hughes et al. 2007; Tanentzap et al. 2013; Hodgson et al. 2015).
<i>Unpredictable environment</i>	Resources spatially and/or temporally heterogeneous with random, rare, or sporadic environmental changes (Valladares et al. 2002); rapid environment pressure imposed on individuals; and species show phenotypic flexibility in response to change (Piersma and Drent 2003; Rymer et al. 2013).
<i>Vulnerability</i>	The predisposition or susceptibility of a species (population or individual) to be negatively affected by a disturbance (Smit et al. 2000; Williams et al. 2008; Oppenheimer et al. 2014).

megadroughts (extreme drought events persisting for several years) have occurred in North America, China, and West Africa (Dai 2011).

Droughts are characterized by periods of unusually low water availability due to reduced precipitation and/or increased evaporation due to increased temperatures (Dai 2011). Droughts create water shortages, leading to reduced plant primary productivity (McNaughton et al. 1983) and thus an associated food reduction for animals, ultimately leading to an increased risk of individual death and population extinction (Miller et al. 2005; Seager et al. 2007). Animals exposed to such environmental conditions may respond through changes in behavior, physiology, and/or morphology that facilitate survival and/or reproduction. If individual differences in survival/reproductive probabilities are due to genetic differences, genetic selection within the population will result in evolutionary adaptation (Rezende and Diniz-Filho 2012). Under drought conditions, reproduction is often terminated or forfeited, and dry seasons are typically the nonbreeding season (Pinter 1994; Schradin and Pillay 2005; Sperry and Weatherhead 2008). Droughts represent periods of low resource availability that individuals must survive in order to reach a season when reproduction again becomes possible. For this reason, our focus rests with survival and not on reproductive success.

Individuals can cope with droughts using a variety of morphological, physiological, and behavioral traits, and to predict the extent to which species are resilient to droughts requires some measure of these traits. Here, we review these traits to provide researchers and nature conservationists with a guide to which phenotypic traits should be studied in order to assess whether or not a species of interest may be able to cope with droughts. We describe three different ways species can cope with droughts, namely through evolutionary adaptation (future change), adaptive phenotypic plasticity (current change), and exaptations (including exaptive phenotypic plasticity, current change). We focus specifically on

adaptive phenotypic plasticity (as per Ghalambor et al. 2007) here, although we recognize that phenotypic plasticity is not necessarily always adaptive, as the environment can also induce changes that decrease fitness, for example, due to constraints or morphological changes indicating significant costs. Another example is evolutionary traps, whereby a fitness increasing ("good") choice changes into a fitness decreasing ("poor") one as previously reliable environmental cues become unreliable, leading to an evolutionary mismatch and thereby trapping the individual into continuing use of the poorer option (Robertson et al. 2013), which could be particularly common in predictable environments (Box 1). Within this context, we argue that whether a trait is specialized and fixed or flexible is important for determining the resilience (the ability to persist following a disturbance or disturbances; Hughes et al. 2007; Tanentzap et al. 2013; Hodgson et al. 2015) or vulnerability (predisposition or susceptibility to be negatively affected by a disturbance; Smit et al. 2000; Williams et al. 2008; Oppenheimer et al. 2014) of a species (or population) to droughts (Box 1). Species with fixed adaptations to droughts are generally well adapted to arid environments, although in species with flexible adaptations and exaptations, the degree of flexibility determines resilience and will likely facilitate adaptive evolution in novel environments (Ghalambor et al. 2007). Resilience to droughts requires individuals to be able to cope with three environmental stressors: water shortage, food shortage, and fluctuating (often very high) temperatures. We define the term "environmental stressor" as any stimulus that could affect an individual negatively if it cannot respond to it via a specific stress response, which itself is costly to some extent. A stress response can be an increase in energy mobilization (i.e., increased blood glucocorticoid levels to mobilize energy; Charmandari et al. 2005), but also comprise other physiological responses (for example, decrease of blood glucocorticoid levels, if the stressor is reduced energy availability and the individual has to reduce energy mobilization).

Stressors are typically uncontrollable for the individual. For example, the individual cannot control rainfall during a given period. Stressors can occur predictably (e.g., seasonal changes in food) or unpredictably (e.g., a storm; Wingfield 2013).

We then highlight some approaches to categorize behavioral, physiological, and morphological traits of mammals for existence in deserts and semideserts and introduce our own categorization system, the adaptive triquetra, which categorizes traits facilitating coping with aridity instead of categorizing the species themselves. The adaptive triquetra system considers the primary driving stressor, the body system mounting the adaptive response (behavior, physiology, and/or morphology), and the extent to which this trait is flexible. We suggest that the selection pressures operating in deserts (defined below) drive the evolution of specialized, nonplastic adaptations, whereas those in semideserts (defined below) promote phenotypic flexibility. We also stress that traits that could potentially facilitate future coping (which would be classified as exaptations) must not be ignored. We focus our explanation of the adaptive triquetra on mammals, which are one of the best studied taxa for desert adaptations. Although adaptations in other taxa, such as insects and birds, might be quite different, we believe that our general conclusions would apply to them as well. Finally, we conclude that the degree of flexibility of traits that exist as exaptations in nondesert species will determine how well they can cope with droughts, and comparing their degree of flexibility with those of species from semideserts will enable us to predict their resilience.

EVOLUTIONARY ADAPTATION, ADAPTIVE PHENOTYPIC PLASTICITY, AND EXAPTATIONS

Evolutionary adaptation involves genetic changes over generations (Rezende and Diniz-Filho 2012). Over time, species may show genetic changes, leading to changes in morphology (e.g., peppered moths *Biston betularia*; Saccheri et al. 2008), physiology

(e.g., red abalone *Haliotis rufescens*; de Wit and Palumbi 2013), and behavior (e.g., bird song; Podos et al. 2004). The degree of adaptive phenotypic plasticity also represents an evolutionary adaptation. These adaptations confer a fitness advantage and evolved in response to particular past selection pressures (Baum and Larson 1991; Coddington 1994). Evolutionary adaptation represents how species responded to previous environmental change, and raises the question of whether, and the degree to which, species can respond to prevailing environmental change via genetic change. Although it is becoming increasingly recognized that evolutionary adaptation can be quite rapid (Hoffmann and Sgrò 2011), the evolutionary response to a changing environment can be constrained by a lack of genetic variation or genetic correlations (i.e., genetic constraints; Chevin 2013), the rate at which individuals within a population reproduce (Lewontin 1970), and the rate and predictability at which the strength and direction of selection changes (due to environmental change) over time (Lande and Shannon 1996; Gomulkiewicz and Houle 2009; Chevin 2013; Michel et al. 2014).

Species might not be adapted to prevailing specific environmental pressures, but may still show some degree of resilience to change due to exaptations, also called pre-existing adaptations or preadaptations (e.g., Pekár et al. 2013). In contrast to evolutionary adaptations, exaptations are traits that evolved in response to specific past selection pressures, and provide a fitness advantage in the prevailing environment under different selection pressures, meaning that the past and current environments are different for at least some specific aspects (Gould and Lewontin 1979). In other words, these traits provide a function in their current role, but were not selected for this purpose (Gould and Vrba 1982). For example, Hoffman (2014) proposed that euryhalinity (i.e., salinity tolerance) in the green toad *Bufo viridis* evolved to facilitate migration across the variable habitats of Eurasia and the North African Mediterranean coastal regions. These toads now occupy arid and semiarid regions,

suggesting that euryhalinity is an exaptation important for drought resistance (Hoffman 2014). Euryhalinity in these toads did not evolve in response to direct selection pressure from an arid or semienvironment (i.e., evolved in response to past selection pressure), but it does have current utility (i.e., provides a use in the existing context). The usefulness of the term exaptation has frequently been criticized because exemplary traits are often not clearly distinguishable from adaptations (Larson et al. 2013). The distinction between adaptation and exaptation is particularly important to consider when assessing whether an individual could survive in a future environment not experienced previously, as is predicted under accelerated environmental change (Sih et al. 2011). Existing phenotypic plasticity is likely to be of importance in the prevailing environment, even if it has evolved under different selection pressures in a different environment (i.e., exaptive phenotypic plasticity; Box 1). Therefore, a population might be able to respond to increased severity of droughts, not because it previously experienced droughts and is adapted to droughts, but because it experienced different environmental conditions that imposed similar selection pressures (e.g., food shortage). This population could thus possess beneficial exaptations in response to droughts.

Adaptive phenotypic plasticity is an evolved trait, enabling a comparatively rapid response by individuals to cope with environmental change. Phenotypic traits are altered adaptively in response to prevailing environmental conditions, which increases individual fitness (Charmantier et al. 2008; Ozgul et al. 2009, 2010; Piersma and van Gils 2010). Adaptive phenotypic plasticity may determine an individual's survival and reproduction in the short term, which may facilitate population persistence and evolutionary adaptation in the long term (West-Eberhard 1989; Hoffmann and Sgrò 2011). This could be especially important for juveniles and subadults that might be more vulnerable to stressful conditions than adults (e.g., Rivers et al. 2012). Importantly, adaptive phenotypic plasticity itself is the product of evolution (Baldwin 1896). Phenotypic plasticity

can be divided into developmental plasticity and phenotypic flexibility (Piersma and van Gils 2010; Box 1). For developmental plasticity (Nettle and Bateson 2015), the environment during early ontogeny determines which one of two or more developmental pathways is adopted, leading to alternative adult phenotypes that cannot change to any of the other phenotypes (Ledón-Rettig et al. 2008). The resulting phenotype is often robust, being able to cope with environmental changes. For developmental plasticity to be adaptive, the population must have experienced particular environments in the past and, as a result, the developmental input to produce particular phenotypes under particular sets of conditions experiences positive selection (Nettle and Bateson 2015). In other words, individuals experience a particular environment during early development that is predictive of the environmental conditions they are likely to experience in later life. In the case of phenotypic flexibility, the trait can change back and forth (i.e., it is reversible), such as changes of the osmoregulatory hormone arginine vasopressin (AVP), whose levels increase when the organism has to save water, but later decrease again. Importantly, individuals may have both developmentally plastic traits and also phenotypically flexible traits.

One specific case of developmental plasticity is epigenetic variation (Piersma and van Gils 2010; Champagne 2013). Epigenetic mechanisms can also be influenced by hormones, which can link organizational effects of hormones to differences in behavior (Champagne and Rissman 2011). Importantly, epigenetic inheritance can occur via at least four different pathways, including cellular epigenetic inheritance (information transmitted via the gametes), developmental inheritance (information transmitted via maternal effects), behavioral inheritance (information transmitted via niche construction), and symbiotic inheritance (Jablonka and Lamb 2005, 2007; Piersma and van Gils 2010). Several recent reviews have focused on the links between epigenetics and phenotypic plasticity (e.g., Reik 2007; Champagne and Rissman 2011; Champagne 2013; Bale 2015; Dochtermann et al. 2015). However,

the majority of studies on phenotypic plasticity provide little or no information on the underlying genetic and epigenetic mechanisms. Therefore, in the context of this review, we consider plasticity in general, rather than in relation to the absolute underlying mechanisms, by determining how variable the trait can be under varying environmental conditions.

Phenotypic plasticity has been hypothesized to be costly because of the need for the development and continued maintenance of neural structures (Dukas 1999), and sensory and/or response pathways involved in facilitating the flexible response (DeWitt et al. 1998; Mery and Burns 2010). However, estimating the actual costs of plasticity is difficult, and these costs might be more easily detected under certain conditions. For example, costs of plasticity are said to be greater in stressful conditions, and if plasticity utilizes resources that become scarce in response to the stress, these costs would be easier to detect (Steiner and Van Buskirk 2008). Van Buskirk and Steiner (2009) argue that the costs of plasticity are actually relatively weak and there might be a compromise or tradeoff between the fitness benefits and costs of greater phenotypic plasticity. One of the main costs of plasticity might be the time needed to mount an appropriate plastic response. Because plasticity seems to have surprisingly low costs, it has been argued recently that relaxed and variable selection intensities are more important constraints for the evolution of plasticity than are its costs (Murren et al. 2015).

Phenotypic plasticity, enabling a coping response to droughts, can either be evolutionary adaptations (adaptive phenotypic plasticity) or exaptations (exaptive phenotypic plasticity). For example, all mammals have the hormone AVP, which is important in osmoregulation, but not all mammals have experienced droughts as a selection pressure modifying the AVP system. Therefore, an increase of AVP plasma levels to reduce water loss during droughts might be higher in desert species, representing an adaptation to previous droughts, but would also occur—although maybe at a lower level—in

species from nondesert environments when exposed to droughts, as an exaptation that evolved to save water under different environmental conditions, such as during winter or during periods when food has a lower water content due to seasonal changes. In sum, phenotypic plasticity can have very low flexibility (being fixed) or high flexibility, and represent adaptations or exaptations to droughts.

THREE PRIMARY STRESSORS OF ARID ENVIRONMENTS

Identifying the limits of phenotypic plasticity is foundational for assessing the degree to which individuals can respond to rapid environmental change. To evaluate which adaptations and exaptations could promote species persistence to increased periodicity and/or intensity of droughts, we will focus on species that currently inhabit areas characterized by periods of dryness (i.e., semideserts and deserts) with low annual precipitation and high aridity index (Box 1).

Deserts and semideserts are characterized by three principal environmental stressors, namely low water availability, low food availability, and extremes in temperature. The definition of “drought” is a relative one, taken to mean that current precipitation falls significantly below the long-term mean of a defined area (Dai 2011). In this context, drought is defined on a temporal scale and can occur in any environment. In contrast, naturally dry areas, such as deserts, experience permanent water shortage, and semideserts experience seasons with water shortage. An aridity index, used in conjunction with total rainfall, enables us to categorize habitats based on water availability (e.g., deserts versus semideserts; Nash and Engineering Group Working Party 2012). Hyperarid areas, such as the northeastern parts of the Negev Desert in Israel (Boroda et al. 2014), experience more than 12 months without rainfall, have less than 25mm rainfall/annum, and an average aridity index below 0.05, whereas arid areas, such as the Namib Desert in southwestern Africa (Viles and Goudie 2013), receive less than 100mm of rainfall per year, which is often very unreliable, and have an aridity index

below 0.2 (Nash and Engineering Group Working Party 2012). In our review, we group both arid categories into “deserts,” as the selection pressures in both habitat types will drive the establishment of specialized adaptations (see below). Semideserts, such as the Hardenveld of the Succulent Karoo in South Africa (Acocks 1988) and the Victoria River district in Australia (Webb et al. 2014), generally receive between 100 and 200mm of rainfall per year on average, have a moderate aridity index of between 0.2 and 0.5, and a 25–50% interannual rainfall variability (for definition see Box 1; Nash and Engineering Group Working Party 2012). We define “drought-affected areas” as those that are neither deserts nor semideserts, but experience unpredictable droughts, which might occur more often and become more intense in the future due to climate change (Box 1).

Deserts and semideserts are characterized by low food availability. Under these conditions, vegetation is typically sparse and patchily distributed across the landscape (Zhang et al. 2005). Consequently, food availability and quality is spatially and temporally variable and the availability of food resources might be unpredictable. For example, some areas might experience seasonal periods of dryness, which are relatively predictable (predictable environment; Box 1), while other areas might experience unpredictable, extreme climatic events leading to droughts (unpredictable environment; Box 1). Animals inhabiting these environments have to cope both with periods of low water availability and simultaneously low food availability (Willmer et al. 2000). One important consideration is that plants in deserts are also adapted to low water availability (Chaves et al. 2003), whereas plants in areas experiencing increased frequencies of droughts due to climate change may not be adapted to cope with dry conditions over long periods (Vicente-Serrano et al. 2013). As a result, food and water availability in drought-affected areas may have an even greater influence on animal populations than those in naturally dry areas.

Ambient temperature (T_a) is the third factor influencing species in naturally dry and in drought-affected areas. These habitats

are often characterized by extreme daily fluctuations in T_a that can range from very cold at night to very hot during the day, a consequence of reduced absorption of heat by vegetation (Sarma et al. 2001), leading to greater reflection of solar radiation by the ground during the day (i.e., increased albedo; Charney et al. 1977), and rapid loss of heat at night (Sarma et al. 2001). Animals living in naturally dry environments have evolved under selective pressures of fluctuating temperature and tend to show specific behavioral, physiological, and/or morphological adaptations to these environmental extremes. For example, several large desert artiodactyls (Table 1), such as camels *Camelus dromedarius* (Elkhawad 1992; Ouajd and Kamel 2009) and Arabian oryx *Oryx leucoryx* (Hatem et al. 2012), utilize the carotid rete, a network of interlacing arterioles in the venous sinus, for selective brain cooling (Hatem et al. 2012). Venous blood cooled through the nasal countercurrent exchange mechanism (Ouajd and Kamel 2009) drains into the venous sinus, cooling the arterial blood moving to the brain (Mitchell et al. 1987).

To identify flexible traits that could confer a fitness advantage under conditions of increasing aridity, it is useful to study species that inhabit deserts or semideserts. Categorizing species living in these dry environments can help us to: identify key adaptations that have arisen in response to aridity; establish a comparative database of traits that can be used to compare with other species for which we lack information; and recognize traits in nondesert-adapted species that could promote responses to increasing aridity and droughts.

CATEGORIZING SPECIES FROM DESERTS AND SEMIDESERTS AS ENDURERS OR EVADERS

Willmer et al. (2000) classified desert animals into two main groups—physiological endurers, such as the Arabian oryx, which are able to tolerate environmental extremes using physiological and morphological adaptations (Table 1) and behavioral evaders, such as Merriam’s kangaroo rat

TABLE 1
Characteristic desert adaptations of physiological endurers and behavioral evaders

Behavior	Physiology	Morphology	Principal examples and references
Mechanisms for coping with water restriction			
<ul style="list-style-type: none"> Can go for extended periods without drinking [1l,o], [2b,e,f], shows water independence [4e], or may survive on water gained through food [2d,c,f,g,h], [3f,k], [4j] Reduced food intake [1o] 	<ul style="list-style-type: none"> Constant [1a,b, but see 1o] or decreased [1f,s] plasma volume Increased plasma Na⁺ [1b,d,f,i,o,s], protein [1f,s], [2e], sugar [1s], cholesterol [1s], and urea [2g, but see 1g,k] concentration Increased plasma arginine vasopressin [1o], [3b,k], aldosterone [1i], and renin [1i,o] secretion Decreased insulin secretion [3k] Maintenance [2g] or increased plasma osmolality [2e], [3k] Increased urine concentration [1o,s], [2g], [3a,c,j,k], [4a,c,j] Reduced urine production [1f,o,s] and volume [1a,b,i], [2g], [3k] Low glomerular filtration rate [1f,s] Increased urine Na excretion [1o] High tolerance to salt [1c], [3k] Excretion of relatively dry feces [1a,b,f,i], [2g], [4b,j] Reduction in metabolic rate [1c], [2d,h], [3a,c,d,e,f,g,j,m], [4j] Reduction in evaporative water loss [1a,b,i], [2d,g], [4c,j] Production of metabolic water [2d], [4c] Low water turnover [1a,b,i] Increased urine creatinine concentration [2g] and decreased creatinine clearance [1s] Decreased respiration rate [1c,k,m,s], [2g] Reduction of digestive secretions [1s] 	<ul style="list-style-type: none"> Nasal countercurrent exchange system [1j,s], [4d,j] Well-developed nasal glands [1s] Nostrils can close completely [1s] Kidney with long loops of Henle and well-developed medulla [1s] Increased tubular reabsorption of the kidney [1s], [2g] Thin erythrocyte with increased surface area: volume ratio [1s] 	Physiological endurers <ul style="list-style-type: none"> ¹ Camel <i>Camelus dromedarius</i> ^a Schmidt-Nielsen et al. 1956 ^b Macfarlane et al. 1963 ^c Schmidt-Nielsen et al. 1967 ^d Siebert and Macfarlane 1971 ^e Maloiy 1972 ^f Siebert and Macfarlane 1975 ^g Emmanuel et al. 1976 ^h Young 1976 ⁱ Finberg et al. 1978 ^j Schmidt-Nielsen et al. 1981a ^k Yagil 1985 ^l Jihad et al. 1989 ^m Wilson 1989 ⁿ Elkhawad 1992 ^o Ben Goumi et al. 1993 ^p Faye 1997 ^q Selim et al. 1999 ^r Dereje and Udén 2005 ^s Ouajd and Kamel 2009 ² Arabian oryx <i>Oryx leucoryx</i> ^a Hetem et al. 2010 ^b Stewart 1963 ^c Spalton 1999 ^d Williams et al. 2001 ^e Ostrowski et al. 2002 ^f Ostrowski et al. 2003 ^g Ostrowski et al. 2006 ^h Gilad et al. 2008
Mechanisms for coping with food restriction			
<ul style="list-style-type: none"> Specialist feeder [1r,s], [4i] or favors specific food type [2c,d], [3i] Feeds on water-rich invertebrates [3i] or green matter [1s], [2g], or digs up water-rich underground plant storage organs [2c] Stores/caches food [2b] Exploits newly emergent food material [2g] Migratory behavior [1f,l,s], [2b] or shifting home range to areas with ephemeral high food abundance [2h], [4h] 	<ul style="list-style-type: none"> Maintenance [3t,m] or minimal loss of body mass [1a,f,s], [2g] for physiological endurers, but tolerance to loss of body mass [3j,k] for behavioral evaders Mobilization of hepatic lipids [1s] and fatty acids [2g] to fuel metabolism Decreased leptin production [2g], [3m] Good aptitude for recycling nitrogen [1g] Enters torpor [3l,m] 	<ul style="list-style-type: none"> Reduction in organ size/volume [2g] Strong digestive efficiency [1q,s], [2g] Storage of fat/lipids [1r], [3m] 	Behavioral evaders <ul style="list-style-type: none"> ³ Golden spiny mouse <i>Acomys russatus</i> ^a Shkolnik and Borut 1969 ^b Castel and Abraham 1972 ^c Haim and Borut 1981 ^d Rubal et al. 1992 ^e Haim and Izhaki 1993 ^f Kam and Degen 1993 ^g Merkt and Taylor 1994 ^h Haim and Rozenfeld 1995 ⁱ Mendelssohn and Yom-Tov 1999 ^j Ron and Haim 2001 ^k Shanas and Haim 2004 ^l Ehrhardt et al. 2005 ^m Gutman et al. 2006

continued

TABLE 1
Continued

Behavior	Physiology	Morphology	Principal examples and references
Mechanisms for coping with extreme temperatures			
<ul style="list-style-type: none"> Utilizes burrows/nests [4f,j], cover [3h], or retreats to shade when available [1s], [2d,g] Utilizes substrate surface for conductance of body heat [2b,d] Primary activity: crepuscular or nocturnal [2b,d,g] Decreased activity [2g] Bipedal locomotion [2b] Minimizes exposure to sun by altering body posture [1p,s] Does not pant [1c,s] 	<ul style="list-style-type: none"> Adaptive heterothermy [1o], [2a,f] Adaptive hyperthermia [1c] Storage of body heat during the day and dissipation at night [1c] Higher stable hematocrit [1p], [2c] Increased erythrocyte circulation efficiency [1s], erythrocyte volume flexibility [1s], and enhanced erythrocyte life span [1s] Low cardiac rate and low blood pressure [1s] Selective brain cooling [1n,s] Enters torpor [3l,m] 	<ul style="list-style-type: none"> Subdivision of nasal sinus [1j], [4d,g] Limited subcutaneous fat deposition [1h,s] Short coarse pelage [2b] Light-colored pelage for reflecting solar radiation and reducing heat uptake [2b] Maintenance of nonshivering thermogenesis [3e] Dark pigmented skin [3h] 	⁴ Merriam's kangaroo rat <i>Dipodomys merriami</i> ^a Schmidt-Nielsen et al. 1948 ^b Schmidt-Nielsen and Schmidt-Nielsen 1951 ^c Schmidt-Nielsen and Schmidt-Nielsen 1952 ^d Jackson and Schmidt-Nielsen 1964 ^e Carpenter 1966 ^f Soholt 1974 ^g Schmidt-Nielsen et al. 1970 ^h Brown and Munger 1985 ⁱ Mares 1993 ^j Nagy 1994

Superscript numbers designate species (endurers: ¹*Camelus dromedarius*, ²*Oryx leucoryx*; evaders: ³*Acomys russatus*, ⁴*Dipodomys merriami*) and letters designate associated references, which appear in Appendix 1.

Dipodomys merriami, which are able to escape environmental extremes behaviorally (Table 1). Later, Ward (2009) recognized that endurers are typically large-bodied animals while evaders are typically small-bodied animals.

The endurer-evader concept is useful for categorizing desert specialists primarily in relation to one characteristic of arid environments, namely extremes in environmental temperature (T_a). However, this concept is a simplistic approach, based on the assumption that it is easier for small animals to find refuge in the shade than it is for large animals, while in fact we know that large animals will use any little shade they can find in their environment to avoid high T_a (Williams et al. 2001; Ostrowski et al. 2006; Ouajd and Kamel 2009). Although this concept, to a degree, considers the ability of animals to be behaviorally flexible (e.g., seeking shade), it generally assumes that animals have limited flexibility in their behavioral and morphological responses and neglects the important aspect


of exaptive phenotypic plasticity, which we consider to be a key mechanism to cope with change, including droughts. As such, the endurer-evader categorization provides an initial framework from which to further identify important traits to be studied in nondesert-adapted species for coping with heat, food, and water restriction as a consequence of droughts.

A NOVEL CLASSIFICATION: THE ADAPTIVE TRIQUETRA

Given the simplicity of the endurer-evader concept, we developed a three-tier system, the adaptive triquetra, that categorizes adaptations to aridity instead of categorizing species (Box 2). Having a species categorization is not helpful for generalizing adaptations across species, and importantly overlooks the underlying general adaptive responses to drought. We stress that providing this framework is only the first step. The framework we develop here must be validated in the future by data from a variety

BOX 2*Using the adaptive triquetra system*

The adaptive triquetra considers a three-step approach that can be used for either: directly assessing resilience in species for which good biological information already exists; and/or for identifying areas where more research may be required. For example, if someone is interested in a particular species, they might have good knowledge of its life history, but little knowledge about its environment, and could therefore focus on establishing which environmental stressor is more important before understanding how the species could respond. Similarly, someone might have good knowledge about the physiology of a species, but little about its behavior. Focus could be placed on how that species responds behaviorally to particular environmental stressors.

**Step 1: What is the Primary Environmental Stressor? (Water, Food, Temperature)**

Consider the environment in which the species of interest occurs, and what the primary stressor would be, taking microhabitat use into account (Tables 2a, b, and c). This is important because not all habitats will experience the same intensity of change and some environments will buffer some stressors better than others. For example, temperature extremes might be more critical in subtropical areas experiencing droughts than in temperate or polar climate zones. For a diurnal burrowing rodent, such as Brants's whistling rat *Parotomys brantsii* (Jackson et al. 2004), extreme temperatures may not represent a problem due to the buffering effects of burrows.

Step 2: What Traits Correspond to Each Body System? (Behavioral, Physiological, Morphological)

This step considers the presence of, and number of traits, corresponding to each system (Tables 2a, b, and c). The greater the number of traits allowing for coping with the stressors, the more likely it is that the species might be able to cope with rapid environmental changes. However, this will also heavily depend on the flexibility of the traits (Step 3).

Step 3: What is the Relative Flexibility of Each Trait? (Fixed, Developmentally Plastic, Flexible)

The ability of a species to persist under increasing likelihood of droughts is dependent on how individuals are able to respond with the traits they possess. An individual has a greater likelihood of responding to unpredictable events if it has greater flexibility (Tables 2a, b, and c). The interaction between two traits can also change the potential for resilience. For example, the interaction between a fixed morphological trait and a flexible behavioral trait is important for African striped mice *Rhabdomys pumilio* in the Succulent Karoo of South Africa. Striped mice have dark skin (Schradin and Pillay 2005), which facilitates thermoregulation through basking (Schradin et al. 2007), allowing mice to conserve energy in the early morning when temperatures are lower.

of different species and different research programs to determine its accuracy, and then modified as appropriate. Our comprehensive review of adaptations to droughts of specialist and nonspecialist species provides the foundational information needed for such future research programs on the effects of droughts on individual survival and population and species persistence.

The term “triquetra” is derived from the Latin *tri-* “three” and *quertrus* “cornered,” emphasizing the three-tiered nature of this classification system, as well as the interconnectedness of its parts. Our adaptive triquetra considers: which stressor is addressed (water restriction, food restriction, or extremes in T_a); which body system mounts an appropriate adaptive response (behavior, physiology, or morphology); and the nature of the trait (fixed versus flexible). Although these stressors are often interlinked (e.g., plants are dependent on water and, therefore, food is likely to be restricted if water is restricted), we argue that each stressor should first be examined independently, because some species might have adaptations or exaptations that facilitate better coping with one stressor compared to another. This is especially important for designing effective management strategies to support populations at risk. For example, during a severe drought period (1981–1984) in southern Africa, mortality of herbivores was quite high, but in some areas, such as the northeastern Tuli region of Botswana, mixed grazers-browsers (e.g., impala *Aepyceros melampus*) were little affected compared to grazers (e.g., blue wildebeest *Connochaetes taurinus* and zebra *Equus burchelli*), whereas in other areas, such as the central Kruger National Park in South Africa, the same grazers (blue wildebeest and zebra) were unaffected (Walker et al. 1987). Walker et al. (1987) suggest that mortality in Klaserie Reserve in South Africa (adjacent to the Kruger National Park) was heightened by close spacing of permanent water sources. Heavy grazing occurred around these areas, eliminating reserve stands. Instead, dropping fences and widely spacing water sources would have allowed animals to move between areas, accessing forage and promoting

survival. Therefore, supplemental feeding in Klaserie could have been highly beneficial to promote species survival.

WHICH STRESSOR IS ADDRESSED?

The first step of the adaptive triquetra model is to identify which of the three primary stressors has driven the evolution of the particular adaptation in question (Box 2). First, low water availability leads to the need for improved osmoregulation, resulting in adaptations that reduce water loss or increase water intake (Table 2a). Second, since food availability is a direct consequence of water restriction, there is a need for improved energy storage mechanisms, resulting in adaptations that minimize energy loss or maximize energy gain (Table 2b). These include both intrinsic adaptations, such as storage of body fat (e.g., Mongolian gerbils *Meriones unguiculatus*; Zhang and Wang 2007), and extrinsic adaptations, such as food caching (e.g., kangaroo rats *Dipodomys* spp.; Randall 1993). Third, high T_a leads to the need for improved thermoregulatory ability, resulting in adaptations that decrease the risk of overheating or increase the ability to tolerate heat (Table 2c).

These three stressors are not fully independent of each other. For example, in mammals, the need for improved thermoregulation is directly related to the need to save water because the ability to reduce water loss in mammals is constrained by the need for evaporative cooling to reduce overheating (Taylor 1970). Furthermore, water can be obtained from food directly (e.g., consumption of succulents by viscacha rats *Octomys mimax*; Bozinovic and Contreras 1990) and/or indirectly (e.g., use of metabolic water by the desert pocket mouse *Chaetodipus penicillatus*; Grubbs 1980). However, most animals cannot obtain sufficient water from their food to survive without access to freestanding water or condensed water. For species that are dependent on freestanding water, the primary stressor will be water availability, whereas for water-independent species, the primary stressor will be food availability.

TABLE 2A
The adaptive triquetra system for the environmental stressor water restriction

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
Behavior	Flexible	Consumes water-rich food resources or green matter	Semidesert <ul style="list-style-type: none">• African striped mouse <i>Rhabdomys pumilio</i>• Black-tailed tree rat <i>Thallomys nigricauda</i>• Springbok <i>Antidorcas marsupialis</i> Desert <ul style="list-style-type: none">• Arabian oryx <i>Oryx leucoryx</i>• Banner-tailed kangaroo rat <i>Dipodomys spectabilis</i>	Schradin and Pillay 2005b Frean et al. 1998 Nagy 1994		
		Reduces food intake	Semidesert <ul style="list-style-type: none">• African striped mouse <i>Rhabdomys pumilio</i>• Kirk's dik-dik <i>Madoqua kirkii</i>• Namaqua rock rat <i>Aethomys namaquensis</i> Desert <ul style="list-style-type: none">• Camel <i>Camelus dromedarius</i>• Jackrabbit <i>Lepus californicus</i>	Schradin et al. 2010 Maloiy 1973 Buffenstein 1985		
		Alters home range or changes distribution to increase access to water/take advantage of new water sources	Semidesert <ul style="list-style-type: none">• African buffalo <i>Synerus caffer</i>• Meheley's horseshoe bat <i>Rhinolophus mehelyi</i> Desert <ul style="list-style-type: none">• Desert mule deer <i>Odocoileus hemionus crooki</i>• Hairy-footed gerbil <i>Gerbillurus paeba</i>	Funston et al. 1994 Salsamendi et al. 2012		
		Drinks sea water or tolerates high salt content in food	Semidesert <ul style="list-style-type: none">• Mongolian gerbil <i>Meriones unguiculatus</i>• Tammar wallaby <i>Macropus eugenii</i> Desert <ul style="list-style-type: none">• Desert bighorn sheep <i>Ovis canadensis nelsoni</i>• Fawn hopping mouse <i>Notomys cervinus</i>	Winkelmann and Getz 1962 Purohit 1971		
		Can go for extended periods without drinking, but is not independent of water	Semidesert <ul style="list-style-type: none">• Eland <i>Tragelaphus oryx</i>• European free-tailed bat <i>Tadarida teniotis</i> Desert <ul style="list-style-type: none">• Camel <i>Camelus dromedarius</i>• Eastern Patagonian laucha <i>Elgmodontia typus</i>	King et al. 1975 Rainho 2007 Mares 1977a		
	Fixed	Increases arginine vasopressin secretion and/or replenishes AVP store	Semidesert <ul style="list-style-type: none">• African striped mouse <i>Rhabdomys pumilio</i>• Degu <i>Octodon degus</i>• Ethiopian Somali goat <i>Capra aegagrus hircus</i>	Schoepf and Schradin 2014 Bozinovic et al. 2003 Mengistu et al. 2007		
Physiology						

continued

TABLE 2A
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
		Desert				
		• Camel <i>Camelus dromedarius</i>	Ben Goumi et al. 1993			
		• Golden spiny mouse <i>Acomys russatus</i>	Castel and Abraham 1972			
		Semidesert				
		• Awassi sheep <i>Ovis aries</i>	Jaber et al. 2004			
		• Common spiny mouse <i>Acomys cahirinus</i>	Bukovetzky et al. 2012			
		Desert				
		• Black Bedouin goat <i>Capra aegagrus hircus</i>	Wittenberg et al. 1986			
		• Tarabul's gerbil <i>Gerbillus tarabuli</i>	Saadi and Lebaili 2012			
		Semidesert				
		• Awassi sheep <i>Ovis aries</i>	Jaber et al. 2011			
		• Cactus mouse <i>Peromyscus eremicus</i>	Hulbert et al. 1985			
		Desert				
		• Marwari sheep <i>Ovis aries</i>	Kataria and Kataria 2006			
		• Merriam's kangaroo rat <i>Dipodomys merriami</i>	Yousef and Johnson 1975			
		Semidesert				
		• African striped mouse <i>Rhabdomys pumilio</i>	Buffenstein 1984			
		• Bush karoo rat <i>Otomys unisulcatus</i>	Jackson et al. 2004			
		• Kirk's dik-dik <i>Madoqua kirkii</i>	Hoppe 1977			
		Desert				
		• Arabian oryx <i>Oryx leucoryx</i>	Ostrowski et al. 2006			
		• Merriam's kangaroo rat <i>Dipodomys merriami</i>	Nagy 1994			
		Semidesert				
		• Bushy-tailed hairy-footed gerbil <i>Gerbillus wallinus</i>	Downs and Perrin 1990			
		• Dorcas gazelle <i>Gazella dorcas</i>	Ghobrial 1970			
		Desert				
		• Arabian oryx <i>Oryx leucoryx</i>	Ostrowski et al. 2006			
		• Spinifex hopping mouse <i>Notomys alexis</i>	MacMillen and Lee 1969			
		Semidesert				
		• Kirk's dik-dik <i>Madoqua kirkii</i>	Maloiy 1973			
		• Springhare <i>Pedetes capensis</i>	Peinke and Brown 1999			
		Desert				
		• Agile kangaroo rat <i>Dipodomys agilis</i>	Schmidt-Nielsen and Schmidt-Nielsen 1952			
		• Desert bighorn sheep <i>Ovis canadensis nelsoni</i>	Turner 1970			
		Semidesert				
		• Eland <i>Tragelaphus oryx</i>	Taylor 1969			
		• Rock hyrax <i>Procavia capensis</i>	Rübsamen and Ketttembeil 1980			
		Desert				
		• Arabian oryx <i>Oryx leucoryx</i>	Ostrowski et al. 2006			
		• Spinifex hopping mouse <i>Notomys alexis</i>	Withers et al. 1979			

continued

TABLE 2A
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Utilization of metabolic water to reduce overall water loss	Semidesert	<ul style="list-style-type: none"> • Mongolian gerbil <i>Meriones unguiculatus</i> • Siorhi goat <i>Capra aegagrus hircus</i> 	Winkelmann and Getz 1962 Misra and Singh 2002		
		Desert	<ul style="list-style-type: none"> • Desert pocket mouse <i>Chaetodipus penicillatus</i> • Rüppell's fox <i>Vulpes rueppelli</i> 	Grubbs 1980 Williams et al. 2002		
	Increased tubular reabsorption of water from the kidney or the bladder	Semidesert	<ul style="list-style-type: none"> • Degu <i>Otocodon degus</i> • Kirk's dik-dik <i>Madoqua kirkii</i> 	Bozinovic et al. 2003 Maloiy et al. 1988		
		Desert	<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> • Agile kangaroo rat <i>Dipodomys agilis</i> 	Ostrowski et al. 2006 Howell and Gersh 1935		
	Maintenance or higher hematocrit	Semidesert	<ul style="list-style-type: none"> • Awassi sheep <i>Ovis aries</i> • Springhare <i>Pedetes capensis</i> 	Laden et al. 1987 Peinke and Brown 1999		
		Desert	<ul style="list-style-type: none"> • Pronghorn <i>Antilocapra americana</i> • Spinifex hopping mouse <i>Notomys alexis</i> 	McKean and Walker 1974 Heimeier and Donald 2006		
	Mobilization of hepatic lipids and fatty acids to fuel metabolism	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Least gerbil <i>Gerbillus pusillus</i> 	Fourie and Haim 1980 Buffenstein 1984		
	* And/or in response to food availability		<ul style="list-style-type: none"> • Awassi sheep <i>Ovis aries</i> 	Jaber et al. 2011		
		Desert	<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> • Hairy-footed gerbil <i>Gerbillurus paeba</i> 	Ostrowski et al. 2006 Buffenstein 1985		
	Enters torpor (or hibernation)	Semidesert	<ul style="list-style-type: none"> • Grey mouse lemur <i>Microcebus murinus</i> • Cactus mouse <i>Peromyscus eremicus</i> 	Schmid and Speakman 2009 MacMillen 1965		
	* And/or in response to temperature extremes and food restriction.	Desert	<ul style="list-style-type: none"> • Pinyon mouse <i>Peromyscus truei</i> • Stripe-faced dunnart <i>Sminthopsis macroura</i> 	Bradford 1974 Song and Geiser 1997		
	Information on this topic is limited and equivocal (Geiser 2010)					
	Information on large mammals is lacking in the literature					
Fixed	Shows water independence	Semidesert	<ul style="list-style-type: none"> • Drylands vesper mouse <i>Calomys musculinus</i> • Springbok <i>Antidorcas marsupialis</i> 	Mares 1977b Nagy and Knight 1994		
		Desert	<ul style="list-style-type: none"> • Desert leopard <i>Panthera pardus</i> • Peruvian desert mouse <i>Phyllotis gerbillus</i> 	Bothma and Riche 1984 Koford 1968		

continued

TABLE 2A
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Maintenance or tolerance of decreased plasma volume	Semidesert	<ul style="list-style-type: none"> • German mutton merino sheep <i>Ovis aries</i> • Springhare <i>Pedetes capensis</i> 	Degen 1977 Peinke and Brown 1999		
		Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • White-tailed antelope squirrel <i>Ammospermophilus leucurus</i> 	Siebert and Macfarlane 1975 Hartman and Morton 1973		
	Maintenance or increased plasma osmolality (sodium, protein, sugar, cholesterol, and/or urea)	Semidesert	<ul style="list-style-type: none"> • Degu <i>Octodon degus</i> • Tammar wallaby <i>Macropus eugenii</i> 	Bozinovic et al. 2003 Kinnear et al. 1968		
		Desert	<ul style="list-style-type: none"> • Barmer goat <i>Capra aegagrus hircus</i> • Desert wood rat <i>Neotoma lepida</i> 	Khan et al. 1978 MacMillen and Lee 1967		
	Tolerates increased urinary urea concentration	Semidesert	<ul style="list-style-type: none"> • Kirk's dik-dik <i>Madoqua kirki</i> • Namib brush-tailed gerbil <i>Gerbillurus setzeri</i> 	Maloiy 1973 Buffenstein et al. 1985		
		Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • Kowari <i>Dasyuroides byrnei</i> 	Emmanuel et al. 1976 Haines et al. 1974		
	Production of allantoin precipitate in place of urea	Semidesert	<ul style="list-style-type: none"> • Bushveld gerbil <i>Gerbilliscus leucogaster</i> • Cape gerbil <i>Gerbilliscus afra</i> 	Buffenstein et al. 1985 Buffenstein et al. 1985		
* Lack of evidence in larger mammals		Desert	<ul style="list-style-type: none"> • Dune hairy-footed gerbil <i>Gerbillurus tytonis</i> • Gerbil mouse <i>Malacothrix typica</i> 	Downs and Perrin 1991 Buffenstein et al. 1985		
	Low glomerular filtration rate	Semidesert	<ul style="list-style-type: none"> • Kirk's dik-dik <i>Madoqua kirki</i> • Least gerbil <i>Gerbillus pusillus</i> 	Rugangazi and Maloiy 1988 Buffenstein 1984		
		Desert	<ul style="list-style-type: none"> • Black Bedouin goat <i>Capra aegagrus hircus</i> • Merriam's kangaroo rat <i>Dipodomys merriami</i> 	Wittenberg et al. 1986 Schmidt-Nielsen and Schmidt-Nielsen 1952		
	Reduced metabolic rate	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Damara ground squirrel <i>Xerus princeps</i> • Kirk's dik-dik <i>Madoqua kirki</i> 	Haim and Fourie 1980a Haim et al. 1986 Hoppe 1977		
		Desert	<ul style="list-style-type: none"> • Euro <i>Macropus robustus erubescens</i> • Viscacha rat <i>Otomys mimax</i> 	Dawson et al. 1975 Bozinovic and Contreras 1990		

continued

TABLE 2A
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Reduction in evaporative water loss	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Kirk's dik-dik <i>Madoqua kirkii</i> • Panamint kangaroo rat <i>Dipodomys panamintinus</i> 	Christian 1978 Maloiy 1973 Hinds and MacMillen 1985		
		Desert	<ul style="list-style-type: none"> • Desert kangaroo rat <i>Dipodomys deserti</i> • Gemsbok <i>Oryx gazella</i> 	Hinds and MacMillen 1985 Taylor 1969		
	Low water turnover	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Shaw's jird <i>Meriones shawi</i> • Tammar wallaby <i>Macropus eugenii</i> 	Scantlebury et al. 2006 de Rouffignac and Morel 1965 Kinnear et al. 1968		
		Desert	<ul style="list-style-type: none"> • Namib desert golden mole <i>Eremitalpa granti namibensis</i> • Red kangaroo <i>Macropus rufus</i> 	Fielden et al. 1990 Dawson et al. 1975		
	Good aptitude for recycling (urea) nitrogen	Semidesert	<ul style="list-style-type: none"> • Rock hyrax <i>Procavia capensis</i> • Tammar wallaby <i>Macropus eugenii</i> 	Hume et al. 1980 Kennedy and Hume 1978		
		Desert	<ul style="list-style-type: none"> • Black Bedouin goat <i>Capra aegagrus hircus</i> • Fat sand rat <i>Psammomys obesus</i> 	Silanikove et al. 1980 Kaibling et al. 1975		
	Tolerates some level of dehydration	Semidesert	<ul style="list-style-type: none"> • Desert sheep <i>Ovis aries</i> • Springhare <i>Pedetes capensis</i> 	El-Hadi 1986 Peinke and Brown 1999		
		Desert	<ul style="list-style-type: none"> • Burro <i>Equus asinus</i> • Peruvian desert mouse <i>Phyllotis gerbillus</i> 	Yousef et al. 1970 Koford 1968		
Morphology	Flexible	Storage of water in the stomach/ rumen/ pharyngeal pouch	Semidesert	<ul style="list-style-type: none"> • African elephant <i>Loxodonta africana</i> 	Leggett 2004	
		* Lack of evidence in smaller mammals	Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> 	Elkhawad 1992	
	Fixed	Nasal countercurrent exchange system	Semidesert	<ul style="list-style-type: none"> • Kirk's dik-dik <i>Madoqua kirkii</i> 	Kamau et al. 1984	
		* Also provides a benefit against high temperatures	Desert	<ul style="list-style-type: none"> • Little red kaluta <i>Dasykaluta rosamondae</i> • Camel <i>Camelus dromedarius</i> • Greater bilby <i>Macrotis lagotis</i> 	Withers and Cooper 2009 Elkhawad 1992 Hulbert and Dawson 1974	

continued

TABLE 2A
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Kidney with well-developed medulla and long loops of Henle	Semidesert	<ul style="list-style-type: none"> • Salt flat mouse <i>Salinomys delicatus</i> Diaz and Ojeda 1999 • Tammar wallaby <i>Macropus eugenii</i> Kinnear et al. 1968 			
		Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> Ouajd and Kamel 2009 • Darwin's leaf-eared mouse <i>Phyllotis darwini darwini</i> Vermeulen and Nel 1988 			
	Elongated renal papilla	Semidesert	<ul style="list-style-type: none"> • Black-tailed tree rat <i>Thallomys nigricauda</i> Frean et al. 1998 			
* Large desert mammals do not show this characteristic (Mbassa 1988)			<ul style="list-style-type: none"> • Salt flat mouse <i>Salinomys delicatus</i> Diaz and Ojeda 1999 			
		Desert	<ul style="list-style-type: none"> • Egyptian gerbil <i>Gerbillus gerbillus</i> Khalil and Tawfic 1963 • Plains viscacha rat <i>Tympanoctomys barrerae</i> Diaz and Ojeda 1999 			

Systems providing an appropriate adaptive response (behavior, physiology, or morphology); flexibility of traits (flexible or fixed). Fixed traits also include traits that show developmental plasticity during ontogeny. Researchers should identify the relative plasticity of fixed traits in particular environments (i.e., the reaction norm). A list of traits that could confer a fitness advantage under increasing likelihood of droughts, in relation to the three parts of the triquetra. We have included examples of a large- and small-bodied semidesert and desert species for each trait. The final two columns can be used by researchers and nature conservationists to categorize their study species, to identify areas where information is lacking, and to reach conclusions about the relative status of their species. The selected references appear in Appendix 1.

TABLE 2B
The adaptive triquetra system for the environmental stressor food restriction.

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
Behavior	Flexible	Stores/caches food/ scatter hoards	Semidesert	<ul style="list-style-type: none"> • Black-backed jackal <i>Canis mesomelas</i> Nel 1984 • Giant kangaroo rat <i>Dipodomys ingens</i> Shaw 1934 		
		Desert	<ul style="list-style-type: none"> • Desert kit fox <i>Vulpes macrotis arsipus</i> Cypher 2003 • Wagner's gerbil <i>Gerbillus dasyurus dasyurus</i> Hatough-Bouran 1990 			
	Exploits newly emergent food material	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> Schradin and Pillay 2006 • Vicuña <i>Vicugna vicugna</i> Donadio et al. 2012 			
		Desert	<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> Ostrowski et al. 2006 • Piute ground squirrel <i>Spermophilus mollis</i> Rickart 1986 			

continued

TABLE 2B
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Migratory behavior or shifting home range (mobility) to areas with higher food abundance	Semidesert	<ul style="list-style-type: none"> • African buffalo <i>Synerus caffer</i> • African striped mouse <i>Rhabdomys pumilio</i> • Round-eared sengi <i>Macroscelides proboscideus</i> 	Funston et al. 1994		
		Desert	<ul style="list-style-type: none"> • Banner-tailed kangaroo rat <i>Dipodomys spectabilis</i> • Desert mule deer <i>Odocoileus hemionus crooki</i> 	Schradin and Pillay 2006		
	Coprophagy	Semidesert	<ul style="list-style-type: none"> • Feral horse <i>Equus caballus</i> • Southern mountain cavy <i>Microcavia australis</i> 	Krysl et al. 1984		
		Desert	<ul style="list-style-type: none"> • African elephant <i>Loxodonta africana</i> • Noki <i>Petromus typicus</i> 	Leggett 2004		
	Generalist feeder	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Ash-grey mouse <i>Pseudomys albocinereus</i> • Tammar wallaby <i>Macropus eugenii</i> 	Curtis and Perrin 1979		
		Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • Grey leaf-eared mouse <i>Graomys griseoflavus</i> 	Murray et al. 1999		
	Dietary flexibility	Semidesert	<ul style="list-style-type: none"> • Drylands vesper mouse <i>Calomys musculinus</i> • Springbok <i>Antidorcas marsupialis</i> 	Kinnear et al. 1968		
		Desert	<ul style="list-style-type: none"> • Brush-tailed mulgara <i>Dasyurus blythi</i> • Sand gazelle <i>Gazella marica</i> 	Dereje and Udén 2005		
Fixed	Specialist feeder	Semidesert	<ul style="list-style-type: none"> • Black-tailed tree rat <i>Thallomys nigricauda</i> • Springbok <i>Antidorcas marsupialis</i> 	Nagy and Knight 1994		
		Desert	<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> • Fat sand rat <i>Psammomys obesus</i> 	Spalton 1999		
Physiology	Flexible	Mobilization of hepatic lipids and fatty acids to fuel metabolism	Semidesert	Daly and Daly 1973		
		* And/or in response to water availability	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Least gerbil <i>Gerbillus pusillus</i> • Awassi sheep <i>Ovis aries</i> 	Fourie and Haim 1980		
			<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> • Hairy-footed gerbil <i>Gerbillurus paeba</i> 	Buffenstein 1984		
				Jaber et al. 2011		
				Ostrowski et al. 2006		
				Buffenstein 1985		

continued

TABLE 2B
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Enters torpor (or hibernation)	Semidesert	<ul style="list-style-type: none"> • Grey mouse lemur <i>Microcebus murinus</i> 	Schmid 2001; Schmid and Speakman 2009		
	* And/or in response to water restriction and temperature extremes.		<ul style="list-style-type: none"> • North African elephant shrew <i>Elephantulus rozeti</i> 	Lovegrove et al. 2001		
	Information on large mammals is lacking in the literature	Desert	<ul style="list-style-type: none"> • Round-tailed ground squirrel <i>Spermophilus tereticaudus</i> • Stripe-faced dunnart <i>Sminthopsis macroura</i> 	Hudson 1964 Song and Geiser 1997		
	Reduced leptin secretion in response to starvation, to reduce mobilization of fatty acid reserves	Semidesert	<ul style="list-style-type: none"> • Awassi sheep <i>Ovis aries</i> • Common spiny mouse <i>Acomys cahirinus</i> • Desert • Arabian oryx <i>Oryx leucoryx</i> • Golden spiny mouse <i>Acomys russatus</i> 	Jaber et al. 2011 Bukovetzky et al. 2012 Ostrowski et al. 2006 Gutman et al. 2008		
	Ecological leptin hypothesis	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Mongolian gerbil <i>Meriones unguiculatus</i> • Desert • Golden spiny mouse <i>Acomys russatus</i> 	Schradin et al. 2014 Zhang and Wang 2007 Gutman et al. 2006		
	* This dissociation between leptin and fattening is lacking in the literature					
Fixed	Maintenance, tolerance, or minimal loss of body mass	Semidesert	<ul style="list-style-type: none"> • Least gerbil <i>Gerbillus pusillus</i> • Springbok <i>Antidorcas marsupialis</i> 	Buffenstein 1984 Nagy 1994		
		Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • Sandy inland mouse <i>Pseudomys hermannsburgensis</i> 	Siebert and Macfarlane 1975 MacMillen and Lee 1967		
Morphology	Flexible	Flexibility in length/ mass of portions of the gut/intestine/and mass of other vital organs (e.g., liver, kidneys, cecum)	Semidesert	<ul style="list-style-type: none"> • Burro <i>Equus asinus</i> • Southern mountain cavy <i>Microcaviaustralis</i> 	Sneddon et al. 2006 Sassi et al. 2007	
		Desert	<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> • Darwin's leaf-eared mouse <i>Phyllotis darwini rufepstris</i> 	Ostrowski et al. 2006 Naya et al. 2005		
		Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Awassi sheep <i>Ovis aries</i> • Fat-tailed dunnart <i>Sminthopsis crassicaudata</i> 	Schradin and Pillay 2005a Khachadurian et al. 1966 Morton 1978		
		Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • Golden spiny mouse <i>Acomys russatus</i> 	Elkhawad 1992 Gutman et al. 2006		

continued

TABLE 2B
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
Fixed	Strong digestive efficiency	Semidesert	<ul style="list-style-type: none"> • Degu <i>Octodon degus</i> 	Veloso and Bozinovic 1993		
		Desert	<ul style="list-style-type: none"> • Kirk's dik-dik <i>Madoqua kirkii</i> • Arabian oryx <i>Oryx leucoryx</i> • Great Basin pocket mouse <i>Perognathus parvus</i> 	Hoppe 1977 Ostrowski et al. 2006 Schreiber 1979		

Systems providing an appropriate adaptive response (behavior, physiology, or morphology); flexibility of traits (flexible or fixed). Fixed traits also include traits that show developmental plasticity during ontogeny. Researchers should identify the relative plasticity of fixed traits in particular environments (i.e., the reaction norm). A list of traits that could confer a fitness advantage under increasing likelihood of droughts, in relation to the three parts of the triquetra. We have included examples of a large- and small-bodied semidesert and desert species for each trait. The final two columns can be used by researchers and nature conservationists to categorize their study species, to identify areas where information is lacking, and to reach conclusions about the relative status of their species. The selected references appear in Appendix 1.

WHICH BODY SYSTEM MOUNTS AN APPROPRIATE ADAPTIVE RESPONSE?

The second step of the adaptive triquetra model is to identify which body systems have changed in response to the environmental challenges associated with increasing droughts (Box 2). We expect that all species may use multiple adaptive strategies at the behavioral, physiological, and morphological levels to adapt to, and cope with, the prevailing environment. First, animals may use specific behaviors to avoid harsh environmental conditions. Animals can reduce water stress by choosing to consume water-rich food, by reducing their food intake (thereby minimizing water required for digestion), and/or by reducing respiration to minimize evaporative water loss (Table 2a). Animals can reduce food stress by consuming newly emergent food (e.g., plant/insect), by altering the diet to include a wider variety of food types (opportunistic feeders), or by storing or caching food (Table 2b). To avoid temperature extremes, animals can employ a range of behavioral strategies to minimize overheating, such as utilizing burrows/cover/shade that buffer temperature extremes and also minimize water loss, or altering body posture to minimize exposure to the sun (Tables 2a, b, and c). Altering activity levels by shifting to nocturnal activity is another strategy minimizing exposure to

lethal diurnal temperatures, as well as minimizing water loss (Tables 2a and c).

Second, physiological traits of osmoregulation, metabolism, and thermoregulation enable individuals to offset the limited resources of water and food, and to respond to thermal stress (Tables 2a, b, and c). We need to differentiate between physiological mechanisms of the three processes and the physiological markers indicating deviation from the optimal physiological state, which could suggest that the individual is struggling to maintain homeostasis. For example, decreased thyroxine levels could indicate a mechanism to save metabolic energy (Yousef and Johnson 1975), while reduced blood glucose levels might not be an adaptation, but rather a physiological marker of deviation (stress) from the optimal physiological state (McCue 2010).

Recently, the term allostasis has been used to describe the physiological mechanisms that maintain stability (homeostasis of blood glucose, pH, and O₂ levels) through change of the osmoregulatory, metabolic, and/or thermoregulatory system (McEwen and Wingfield 2003). Importantly, allostasis depends on the energy availability in the environment, which decreases during droughts, leading to allostasis overload type 1 (energy expenditure to cope with environmental stressors is greater than energy intake; McEwen and

TABLE 2C
The adaptive triquetra system for the environmental stressor extremes in environmental temperature (T_a)

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
Behavior	Flexible	Utilizes burrows/ nests, cover, or retreats to shade when available	Semidesert <ul style="list-style-type: none">• African striped mouse <i>Rhabdomys pumilio</i>• Giant kangaroo rat <i>Dipodomys ingens</i>• Southern hairy-nosed wombat <i>Lasiorhinus latifrons</i> Desert <ul style="list-style-type: none">• Euro <i>Macropus robustus erubescens</i> Dawson and Brown 1970• Sundevall's jird <i>Meriones crassus crassus</i> Lewis et al. 1965	Schradin 2005 Randall 1997 Walker et al. 2007		
		Utilizes substrate surface for conductance of body heat	Semidesert <ul style="list-style-type: none">• African striped mouse <i>Rhabdomys pumilio</i>• Cape ground squirrel <i>Xerus inauris</i>• Suricate <i>Suricata suricatta</i> Hinton and Dunn 1967 Desert <ul style="list-style-type: none">• Desert mule deer <i>Odocoileus hemionus crooki</i> Cain et al. 2006• White-tailed antelope squirrel <i>Ammospermophilus leucurus</i> Chappell and Bartholomew 1981a	Schradin 2006 Marsh et al. 1978		
		* Information is lacking for large semidesert species				
		Uses body parts to facilitate shading	Semidesert <ul style="list-style-type: none">• Cape ground squirrel <i>Xerus inauris</i>• Springbok <i>Antidorcas marsupialis</i> Fuller et al. 2005 Desert <ul style="list-style-type: none">• Camel <i>Camelus dromedarius</i> Ouajid and Kamel 2009• White-tailed antelope squirrel <i>Ammospermophilus leucurus</i> Chappell and Bartholomew 1981b	Bennett et al. 1984		
	Activity variable or polyphasic		Semidesert <ul style="list-style-type: none">• Mongolian gerbil <i>Meriones unguiculatus</i> Lewis et al. 1965• Black wildebeest <i>Connochaetes gnou</i> Maloney et al. 2005 Desert <ul style="list-style-type: none">• Arabian oryx <i>Oryx leucoryx</i> Stewart 1963• Lybian jird <i>Meriones libycus</i> Aulagnier et al. 2008			
	Primary activity—crepuscular		Semidesert <ul style="list-style-type: none">• African wild dog <i>Lycaon pictus</i> Woodroffe 2011• Bush karoo rat <i>Otomys unisulcatus</i> Vermeulen and Nel 1988 Desert <ul style="list-style-type: none">• King jird <i>Meriones rex</i> Alagaili et al. 2013• Red kangaroo <i>Macropus rufus</i> Dawson et al. 1975			

continued

TABLE 2C
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Decreased activity	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Degu <i>Octodon degus</i> • Feral pig <i>Sus scrofa</i> 	Schumann et al. 2005 Lagos et al. 1995 Dexter 2003		
		Desert	<ul style="list-style-type: none"> • Arabian oryx <i>Oryx leucoryx</i> • Greater bilby <i>Macrotis lagotis</i> 	Ostrowski et al. 2006 Hulbert and Dawson 1974		
	Social huddling at night to minimize heat loss	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Brants's whistling rat <i>Parotomys brantsii</i> • Vervet monkey <i>Chlorocebus pygerythrus</i> 	Scantlebury et al. 2006 du Plessis et al. 1992 Lubbe 2013		
		Desert	<ul style="list-style-type: none"> • Collared peccary <i>Pecari tajacu</i> • Spinifex hopping mouse <i>Notomys alexis</i> 	Zervanos and Hadley 1973 Withers et al. 1979		
	Bask to increase solar absorption	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Eland <i>Tragelaphus oryx</i> • Fat-tailed dunnart <i>Sminthopsis crassicaudata</i> 	Schradin et al. 2007 Hetem et al. 2011 Warneke et al. 2008		
		Desert	<ul style="list-style-type: none"> • Collared peccary <i>Pecari tajacu</i> • Fat-tailed antechinus <i>Pseudantechinus macdonnellensis</i> 	Zervanos and Hadley 1973 Geiser and Pavey 2007		
	Social flexibility or group flexibility	Semidesert	<ul style="list-style-type: none"> • African striped mouse <i>Rhabdomys pumilio</i> • Hamadryas baboon <i>Papio hamadryas hamadryas</i> • Round-eared sengi <i>Macroscelides proboscideus</i> 	Schradin et al. 2012 Schreier and Swedell 2012 Schubert et al. 2009		
		Desert	<ul style="list-style-type: none"> • Great gerbil <i>Rhomomys opimus</i> • Pronghorn <i>Antilocapra americana</i> 	Randall et al. 2005 Deblinger and Alldredge 1989		
Fixed	Primary activity—nocturnal	Semidesert	<ul style="list-style-type: none"> • Greater Egyptian jerboa <i>Jaculus orientalis</i> • Tammar wallaby <i>Macropus eugenii</i> 	El Ouezzani et al. 2001 Kinnear et al. 1968		
		Desert	<ul style="list-style-type: none"> • Desert bighorn sheep <i>Ovis canadensis nelsoni</i> • Yellow-rumped leaf-eared mouse <i>Phyllotis xanthopygus rupestris</i> 	Hansen 1982 Tirado et al. 2008		

continued

TABLE 2C
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
Physiology	Flexible	Primary activity—diurnal	Semidesert			
			• African striped mouse <i>Rhabdomys pumilio</i>	Haim and Fairall 1986		
			• Black wildebeest <i>Connochaetes gnou</i>	Mitchell et al. 2002		
			• Brants's whistling rat <i>Parotomys brantsii</i>	Jackson et al. 2004		
		Does not pant except in the final stages of heat stress	Desert			
		* Reflects a reflex to thermal extremes (i.e., may not have adaptive significance for all species; Woodroffe 2011)	• Fat sand rat <i>Psammomys obesus</i>	Ilan and Yom-Tov 1990		
			• Nubian ibex <i>Capra nubiana</i>	Hochman and Kotler 2006		
			Semidesert			
			• African buffalo <i>Synacerus caffer</i>	Taylor 1970a		
			• Lesser long-nosed bat <i>Leptonycteris yerbabuenae</i>	Carpenter and Graham 1967		
		Pants to reduce temperature	Desert			
			• Camel <i>Camelus dromedarius</i>	Ouajd and Kamel 2009		
			• Greater bilby <i>Macrotis lagotis</i>	Hulbert and Dawson 1974		
		Adaptive hypo- or hyperthermia	Semidesert			
			• Kirk's dik-dik <i>Madoqua kirkii</i>	Maloiy 1973		
			• Rock hyrax <i>Procavia capensis</i>	Rübsamen and Kettembeil 1980		
			Desert			
			• Arabian oryx <i>Oryx leucoryx</i>	Ostrowski et al. 2003		
			• Fennec fox <i>Vulpes zerda</i>	Noll-Banholzer 1979		
		Wide thermoneutral zone	Semidesert			
			• Grant's gazelle <i>Nanger granti</i>	Taylor 1970b		
			• Greater Egyptian jerboa <i>Jaculus orientalis</i>	El Hilali and Veillat 1975		
			Desert			
			• Camel <i>Camelus dromedarius</i>	Grigg et al. 2009		
			• Jackrabbit <i>Lepus californicus</i>	Shoemaker et al. 1976		
		Maintenance of nonshivering thermogenesis	Semidesert			
			• Littledale's whistling rat <i>Parotomys littledalei</i>	Jackson et al. 2004		
		* Information in large mammals is lacking	• Tammar wallaby <i>Macropus eugenii</i>	Kinnear et al. 1968		
			Desert			
			• Allenby's gerbil <i>Gerbillus allenbyi</i>	Haim 1984		
			• Camel <i>Camelus dromedarius</i>	Schmidt-Nielsen et al. 1967		
			Semidesert			
			• African lesser bushbaby <i>Galago moholi</i>	Nowack et al. 2013		
			• African striped mouse <i>Rhabdomys pumilio</i>	Haim and Fourie 1980b		
			• Damara ground squirrel <i>Xerus princeps</i>	Haim et al. 1986		
			Desert			
			• Golden spiny mouse <i>Acomys russatus</i>	Kronfeld-Schor et al. 2000		
			• Palestine mole rat <i>Spalax ehrenbergi</i> (chromosomal form 2n = 60)	Haim et al. 1984		

continued

TABLE 2C
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Enters torpor (or hibernation)	Semidesert	• Grey mouse lemur <i>Microcebus murinus</i>	Schmid 2001; Schmid and Speakman 2009		
	* And/or in response to food/water restriction.		• North African elephant shrew <i>Elephantulus rozeti</i>	Lovegrove et al. 2001		
	Information on large mammals is lacking in the literature	Desert	• Round-tailed ground squirrel <i>Spermophilus tereticaudus</i> • Stripe-faced dunnart <i>Sminthopsis macroura</i>	Hudson 1964; Song and Geiser 1997		
Fixed	Storage of body heat during the day and dissipation at night	Semidesert	• Grant's gazelle <i>Nanger granti</i>	Taylor 1970a		
	* The terms "adaptive heterothermy" and "heterothermy" are also used		• Little red kaluta <i>Dasykaluta rosamondae</i>	Withers and Cooper 2009		
		Desert	• Sand gazelle <i>Gazella marica</i>	Ostrowski and Williams 2006		
			• Harris's antelope squirrel <i>Ammospermophilus harrisii</i>	Osborn 1991		
	Increased erythrocyte circulation efficiency; erythrocyte volume flexibility and enhanced erythrocyte life span	Semidesert	• Guanaco <i>Lama guanicoe</i>	Cornelius and Kaneko 1962		
	* Information is lacking in small mammals and semidesert species	Desert	• Camel <i>Camelus dromedarius</i> • Desert bighorn sheep <i>Ovis canadensis nelsoni</i>	Oyewale et al. 2011; Turner 1979		
	Selective brain cooling	Semidesert	• Beisa oryx <i>Oryx beisa</i>	Fuller et al. 2004		
	* The incidence is lacking in the literature, and is debated in large mammals		• Springbok <i>Antidorcas marsupialis</i>	Mitchell et al. 1997		
		Desert	• Arabian oryx <i>Oryx leucoryx</i>	Hetem et al. 2012		
			• Camel <i>Camelus dromedarius</i>	Elkhawad 1992		
	Narrow but high thermoneutral zone	Semidesert	• African striped mouse <i>Rhabdomys pumilio</i>	Haim and Fourie 1980a		
	* Information in large mammals is lacking		• Cape short-eared gerbil <i>Desmodillus auricularis</i>	Downs and Perrin 1994		
			• Palmer's chipmunk <i>Tamias palmeri</i>	Lowrey and Longshore 2005		
		Desert	• Naked-footed gerbil <i>Gerbillus nanus</i>	Haim 1984		
			• Spiny mouse <i>Acomys spinosissimus</i>	Perrin and Downs 1994		

continued

TABLE 2C
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
Morphology	Flexible	High thermal conductance	Semidesert			
			• African striped mouse <i>Rhabdomys pumilio</i>	Haim and Fairall 1986		
			• Awassi sheep <i>Ovis aries</i>	Degen and Shkolnik 1978		
			• Western barred bandicoot <i>Perameles bougainville</i>	Larcombe and Withers 2006		
			Desert			
			• Collared peccary <i>Pecari tajacu</i>	Zervanos and Hadley 1973		
			• Desert kit fox <i>Vulpes macrotis arsipus</i>	Golightly and Ohmart 1983		
		Limited subcutaneous fat deposition or localized storage	Semidesert			
			• Boer goat <i>Capra aegagrus hircus</i>	Casey and van Niekerk 1988		
			• Fat-tailed dunnart <i>Sminthopsis crassicaudata</i>	Morton 1978		
	Fixed		Desert			
			• Camel <i>Camelus dromedarius</i>	Young 1976		
			• Soda Spring Valley kangaroo mouse <i>Microdipodops pallidus</i>	Bartholomew and MacMillen 1961		
		Bipedal locomotion (or tendency)	Semidesert			
			• Great jerboa <i>Allactaga major</i>	Schmidt-Nielsen and Schmidt-Nielsen 1952		
			• Tammar wallaby <i>Macropus eugenii</i>	Baudinette et al. 1992		
			Desert			
			• Dusky hopping mouse <i>Notomys fuscus</i>	Schmidt-Nielsen and Schmidt-Nielsen 1952		
			• Red kangaroo <i>Macropus rufus</i>	Dawson and Taylor 1973		
		Short coarse pelage	Semidesert			
			• African buffalo <i>Synerus caffer</i>	Marai and Haeb 2010		
			• Gulf coast kangaroo rat <i>Dipodomys compactus</i>	Baumgardner 1991		
			Desert			
			• Arabian oryx <i>Oryx leucoryx</i>	Stewart 1963		
			• Round-tailed ground squirrel <i>Spermophilus tereticaudus</i>	Walsberg 1988		
	Light colored pelage for reflecting solar radiation and reducing heat uptake		Semidesert			
			• Panamint kangaroo rat <i>Dipodomys panamintinus</i>	Intress and Best 1990		
			• Springbok <i>Antidorcas marsupialis</i>	Mitchell et al. 1997		
			Desert			
			• Desert bighorn sheep <i>Ovis canadensis nelsoni</i>	McCutchen 1981		
			• Eastern Patagonian laucha <i>Eligmodontia typus</i>	Giannoni et al. 2001		

continued

TABLE 2C
Continued

System	Nature of the trait	Trait	Example species	Selected references	Present in my study species?	Degree of plasticity or flexibility?
	Dark pigmented skin	Semidesert	<ul style="list-style-type: none"> • African buffalo <i>Synerus caffer</i> • African striped mouse <i>Rhabdomys pumilio</i> • Damara ground squirrel <i>Xerus princeps</i> 	Marai and Haeeb 2010 Timm and Kermott 1982 Waterman and Herron 2004		
		Desert	<ul style="list-style-type: none"> • Addax <i>Addax nasomaculatus</i> • Golden spiny mouse <i>Acomys russatus</i> 	Portas et al. 2003 Haim and Rozenfeld 1995		
	Presence of a carotid rete	Semidesert	<ul style="list-style-type: none"> • Kirk's dik-dik <i>Madoqua kirkii</i> • Springbok <i>Antidorcas marsupialis</i> 	Kamau et al. 1984 Mitchell et al. 2002		
	* Information is lacking for small mammals	Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • Pronghorn <i>Antilocapra americana</i> 			
	Nasal countercurrent exchange system	Semidesert	<ul style="list-style-type: none"> • Giraffe <i>Giraffa camelopardalis</i> • Little red kaluta <i>Dasykaluta rosamondae</i> 	Langman et al. 1979 Withers and Cooper 2009		
	* This concurrently provides a benefit against water loss	Desert	<ul style="list-style-type: none"> • Camel <i>Camelus dromedarius</i> • Merriam's kangaroo rat <i>Dipodomys merriami</i> 	Schmidt-Nielsen et al. 1981b Jackson and Schmidt-Nielsen 1964		

Systems providing an appropriate adaptive response (behavior, physiology, or morphology); flexibility of traits (flexible or fixed). Fixed traits also include traits that show developmental plasticity during ontogeny. Researchers should identify the relative plasticity of fixed traits in particular environments (i.e., the reaction norm). A list of traits that could confer a fitness advantage under increasing likelihood of droughts, in relation to the three parts of the triquetra. We have included examples of a large- and small-bodied semidesert and desert species for each trait. The final two columns can be used by researchers and nature conservationists to categorize their study species, to identify areas where information is lacking, and to reach conclusions about the relative status of their species. The selected references appear in Appendix 1.

Wingfield 2003). Whether this leads to permanent damage and pathology, and finally death, can be modeled using the reactive scope model, which takes into account the short- and long-term effects of physiological mediators of allostasis (Romero et al. 2009; Romero 2012).

Osmoregulation is regulated by the secretion of several hormones, especially AVP, aldosterone, and renin (Finberg et al. 1978; Ben Goumi et al. 1993), which regulate water reabsorption in the kidneys and water excretion via urine. Several physiolog-

ical markers can indicate a deviation from osmotic homeostasis, such as decreased plasma volume (Siebert and Macfarlane 1975; Ouajd and Kamel 2009) or increased concentration of proteins and sugars (Ouajd and Kamel 2009). Metabolism is regulated by corticosterone and thyroxine, mobilizing hepatic lipids and fatty acids in response to food restriction (Ostrowski et al. 2006; Ouajd and Kamel 2009). Thermoregulation is maintained by corticosterone, dexamethasone, and prednisolone, as well as sodium, and their influence on the sympathetic ner-

vous system (Maickel et al. 1967). Thermoregulatory enhancements are indicated by increased erythrocyte circulation efficiency and changes in erythrocyte volume flexibility (Ouajd and Kamel 2009), which would facilitate oxygen circulation in the body.

Third, morphological structures, both internal and external, can help an animal to cope with water and food shortage, as well as high T_a (Tables 2a, b, and c). Kidney morphology, in particular long loops of Henle, increases water reabsorption and minimizes water loss by increasing urine concentration (Ouajd and Kamel 2009). Organ size and volume can be altered in response to food restriction to minimize the energetic costs of maintaining those organs (Ostrowski et al. 2006). Body size itself can have a profound effect on resilience to climate, and can change both plastically or due to evolutionary adaptation (Brown and Brown 1998). In mammals, paleontologists have observed dwarfing (reduction in body size) during the Paleocene-Eocene boundary, a period of global warming (Gingerich 2006), and dwarfing might also occur in extant mammals in response to increasing T_a . Similarly, pelage color and pattern, and skin color can assist in thermoregulation, minimizing solar absorption and increasing solar radiation (Stewart 1963; Haim and Rozenfeld 1995).

HOW FLEXIBLE IS THE TRAIT?

The final stage of the adaptive triquetra model is to establish the flexibility of the trait under increasing drought conditions (Box 2). Specialized adaptations evolve in response to strong selection pressures imposed by particular habitats. These adaptations are likely to be species-specific because different species adopt different strategies, as well as a combination of strategies, for survival and reproduction under challenging environmental conditions. Furthermore, these adaptations are relatively fixed, with limited flexibility. Specialized adaptations, such as the carotid rete utilized in brain cooling, enables individuals of some species to cope with increasing heat

load (Mitchell et al. 2002). In sum, specialized adaptations are of little interest in establishing how a nondesert-adapted species would respond to the increasing frequency of droughts, because such species will not have specialized adaptations.

Traits that show adaptive developmental plasticity can vary between individuals due to environmental factors. However, although developmentally plastic traits are flexible during early life stages, they are generally fixed by adulthood (Piersma and van Gils 2010). Adaptive developmental plasticity is thus an important determinant of resilience to drought. Essentially, every fixed trait can be categorized according to the nature and degree of its developmental plasticity, also known as a reaction norm (Cassidy et al. 2014). This reaction norm can be narrow, with a slope close to zero (totally fixed trait), or it can be broad. Here, we include traits that show adaptive developmental plasticity as fixed traits, because they are not flexible in adults (Tables 2a, b, and c). As a consequence, to determine the resilience to droughts by a species, the level of adaptive developmental plasticity of every fixed trait should be estimated. Broad adaptive developmental plasticity indicates a greater likelihood that a generation exposed to droughts during early development (or to environmental stimuli predicting droughts) will develop traits enabling resilience to droughts.

General adaptations evolve in response to fluctuating and dynamic selection pressures in multiple habitats and are likely to be common among most (if not all) mammals. For example, all mammals secrete AVP from the pituitary to regulate water reabsorption in the kidneys (Acher 1993). Many general adaptations are likely to be phenotypically flexible. Exaptations must also be considered here because their utilization in the current context is flexible, even if the trait evolved as a specialized adaptation in another context. Recent evidence from the available vertebrate literature suggests that most phenotypic responses to climate change and increased droughts would be due to the flexibility of traits (Canale and Henry 2010). There-

fore, we suggest that identifying flexible traits representing general adaptations and exaptations of nondesert-adapted species is vital for determining whether a species has the potential to persist under increasing drought conditions (Tables 2a, b, and c).

USING THE ADAPTIVE TRIQUETRA TO ESTIMATE SPECIES RESILIENCE OR VULNERABILITY TO DROUGHT

We have attempted to summarize all known traits of the adaptive triquetra that mammals could use to cope with droughts (Tables 2a, b, and c). These can be used by ecologists and nature conservationists to assess the resilience or vulnerability of one or several species to drought. These tables reflect the variety of traits and species possessing these traits. Currently, in most cases, more research on a specific species would be required by ecologists/nature conservationists to accurately estimate the resilience to drought by a particular species. The tables enable a comparison to be made between different traits, as well as between different large- and/or small-bodied species. They provide a step-by-step approach that could help in making decisions on conservation action, especially of which species to focus on, and which of the three stressors is relevant. For example, for species A it might be most important to offer water at artificial water points during droughts, although for species B food shortage could be more important, so supplemental food should be the priority.

We have planned Tables 2a, b, and c in such a manner that they can be directly applied. In the second to last column, one can mark whether or not a specific trait is present in a species. In the last column, the flexibility of that trait for that species can be estimated. These two columns provide the first overview for a specific species and will give some indication of how resilient or vulnerable it could be. Nonetheless, we realize that, in nearly all cases, filling in these tables will generate more questions than answer the critical question of how resilient a species is to droughts. Thus, one conclusion is that to assess drought resilience of

a species, a detailed understanding of the biology and evolutionary history of the species is the requisite first step prior to large-scale decision-making processes.

One important aspect not covered by this process is the environment. By definition, drought occurs when the current precipitation falls significantly below the long-term mean of a defined area, leading to water shortage (Dai 2011). Thus, droughts can differ dramatically in intensity and duration, and a population that can cope well with, for example, a 30% reduction of precipitation over a few months, might not survive a 60% reduction in precipitation over a longer period. In addition, a population's habitat may provide suitable microclimatic refugia that could facilitate persistence, but not the habitat of another population of the same species. However, it is important to understand the complexity of the species first, and then focus on the environment and habitat later, as the biology of an animal will determine its vulnerability, while the modulating role of the environment may promote or hinder resilience.

CONCLUDING REMARKS: THE IMPORTANCE OF FLEXIBILITY

Phenotypic flexibility, rather than future evolutionary adaptation, will most likely facilitate adaptive responses to climate change and increased droughts in the short term (Canale and Henry 2010), which in the long term might lead to evolutionary adaptation (Lind et al. 2015; Scheiner et al. 2015). Identifying those traits that promote persistence in a period of change, as well as understanding the degree of their flexibility, is crucial for assessing whether or not a species will persist under increased frequency and/or periodicity of droughts. In addition to flexibility, resilience might also arise because of exaptation. The concept of exaptation has been criticized as not being useful to understand how species are successful in their current environment (Larson et al. 2013). Yet, the concept might be useful for understanding whether and how species will be able to cope with rapid global change because exaptations could provide important benefits for spe-

cies not currently accustomed to drought conditions.

Phenotypic flexibility operates at the individual rather than at the population level (Rymer et al. 2013), so the ability of species to cope with change will depend largely on the flexibility of its individuals. In addition, individuals that persist and reproduce transmit genes for trait flexibility to their offspring, and flexibility can be a strong driver of phenotypic evolution (Piersma and van Gils 2010; Standen et al. 2014). Thus, in the mid to long term, these traits, as well as the ability to be flexible, could become fixed through evolutionary adaptation, becoming general or specialized adaptations. However, because phenotypic flexibility can be costly, it is important to study to what degree those traits that are phenotypically flexible in order to establish the limits of this plasticity in promoting persistence.

To understand which traits could promote species persistence to increased periodicity of droughts and increased aridification, we should focus on those species currently inhabiting semideserts because they are likely to possess a suite of existing traits that could be advantageous during droughts and that

could be present in species from nondesert environments, in contrast to specialized traits of desert species. Comparative assessment of these species will facilitate identification of traits that could provide fitness advantages for other species encountering increasingly drier conditions. Furthermore, although a species may have a suite of existing phenotypic adaptations or exaptations to droughts, the interaction with other species in its environment, its habitat requirements, amount of available habitat, and human persecution/exploitation will all influence its persistence. Future empirical studies and conservation programs will test the usefulness and applicability of the adaptive triquetra and possibly modify it, enabling us to better predict which populations are vulnerable to drought-related extinction.

ACKNOWLEDGMENTS

The suggestions provided by the editor and two anonymous reviewers greatly improved the manuscript. We are grateful for support by the University of the Witwatersrand (Neville Pillay), James Cook University (Tasmin L. Rymer), the Université de Strasbourg's Institute for Advanced Study (Carsten Schradin), and the CNRS (Carsten Schradin).

REFERENCES

- Acher R. 1993. Neurohypophysial peptide systems: processing, machinery, hydroosmotic regulation, adaptation and evolution. *Regulatory Peptides* 45:1–13.
- Acocks J. P. H. 1988. Veld types of South Africa. *Memoirs of the Botanical Survey of South Africa* 57:1–146.
- Alley R. B., Marotzke J., Nordhaus W. D., Overpeck J. T., Peteet D. M., Pielke R. A. Jr., Pierrehumbert R. T., Rhines P. B., Stocker T. F., Talley L. D., Wallace J. M. 2003. Abrupt climate change. *Science* 299:2005–2010.
- Baldwin J. M. 1896. A new factor in evolution. *American Naturalist* 30:441–451.
- Bale T. L. 2015. Epigenetic and transgenerational reprogramming of brain development. *Nature Reviews Neuroscience* 16:332–344.
- Baum D. A., Larson A. 1991. Adaptation reviewed: a phylogenetic methodology for studying character macroevolution. *Systematic Zoology* 40:1–18.
- Ben Goumi M., Riad F., Giry J., de la Farge F., Safwate A., Davicco M.-J., Barlet J.-P. 1993. Hormonal control of water and sodium in plasma and urine of camels during dehydration and rehydration. *General and Comparative Endocrinology* 89:378–386.
- Black R., Bennett S. R. G., Thomas S. M., Beddington J. R. 2011. Climate change: migration as adaptation. *Nature* 478:447–449.
- Boorda R., Matmon A., Amit R., Haviv I., Arnold M., Aumaître G., Bourlès D. L., Keddadouche K., Eyal Y., Enzel Y. 2014. Evolution and degradation of flat-top mesas in the hyper-arid Negev, Israel revealed from ¹⁰Be cosmogenic nuclides. *Earth Surface Processes and Landforms* 39:1611–1621.
- Bozinovic F., Contreras L. C. 1990. Basal rate of metabolism and temperature regulation of two desert herbivorous octodontid rodents: *Octomys mimax* and *Tympanoctomys barrerae*. *Oecologia* 84:567–570.
- Breed M. F., Ottewell K. M., Gardner M. G., Lowe A. J. 2011. Clarifying climate change adaptation responses for scattered trees in modified landscapes. *Journal of Applied Ecology* 48:637–641.
- Brown C. R., Brown M. B. 1998. Intense natural selection on body size and wing and tail asymmetry in cliff swallows during severe weather. *Evolution* 52:1461–1475.
- Cahill A. E., Aiello-Lammens M. E., Fisher-Reid M. C., Hua X., Karanewsky C. J., Ryu H. Y., Sbeglia G. C.,

- Spagnolo F., Waldron J. B., Warsi O., Wiens J. J. 2013. How does climate change cause extinction? *Proceedings of the Royal Society B: Biological Sciences* 280:2012890.
- Canale C. I., Henry P.-Y. 2010. Adaptive phenotypic plasticity and resilience of vertebrates to increasing climatic unpredictability. *Climate Research* 43:135–147.
- Cassidy E. J., Bath E., Chenoweth S. F., Bonduriansky R. 2014. Sex-specific patterns of morphological diversification: evolution of reaction norms and static allometries in neriid flies. *Evolution* 68:368–383.
- Champagne F. A. 2013. Epigenetics and developmental plasticity across species. *Developmental Psychobiology* 55:33–41.
- Champagne F. A., Rissman E. F. 2011. Behavioral epigenetics: a new frontier in the study of hormones and behavior. *Hormones and Behavior* 59:277–278.
- Charmandari E., Tsigos C., Chrousos G. 2005. Endocrinology of the stress response. *Annual Review of Physiology* 67:259–284.
- Charmantier A., McCleery R. H., Cole L. R., Perrins C., Kruuk L. E. B., Sheldon B. C. 2008. Adaptive phenotypic plasticity in response to climate change in a wild bird population. *Science* 320:800–803.
- Charney J., Quirk W. J., Chow S.-H., Kornfield J. 1977. A comparative study of the effects of albedo change on drought in semi-arid regions. *Journal of the Atmospheric Sciences* 34:1366–1385.
- Chaves M. M., Maroco J. P., Pereira J. S. 2003. Understanding plant responses to drought—from genes to the whole plant. *Functional Plant Biology* 30:239–264.
- Chevin L.-M. 2013. Genetic constraints on adaptation to a changing environment. *Evolution* 67: 708–721.
- Coddington J. A. 1994. The roles of homology and convergence in studies of adaptation. Pages 53–78 in *Phylogenetics and Ecology*, edited by P. Eggleton and R. I. Vane-Wright. Linnean Society Symposium Series, Number 17. London (United Kingdom): Academic Press.
- Dai A. 2011. Drought under global warming: a review. *Wiley Interdisciplinary Reviews: Climate Change* 2:45–65.
- de Wit P., Palumbi S. R. 2013. Transcriptome-wide polymorphisms of red abalone (*Haliotis rufescens*) reveal patterns of gene flow and local adaptation. *Molecular Ecology* 22:2884–2897.
- DeWitt T. J., Sih A., Wilson D. S. 1998. Costs and limits of phenotypic plasticity. *Trends in Ecology and Evolution* 13:77–81.
- Dochtermann N. A., Schwab T., Sih A. 2015. The contribution of additive genetic variation to personality variation: heritability of personality. *Proceedings of the Royal Society B: Biological Sciences* 282:20142201.
- Dukas R. 1999. Costs of memory: ideas and predictions. *Journal of Theoretical Biology* 197:41–50.
- Elkhawad A. O. 1992. Selective brain cooling in desert animals: the camel (*Camelus dromedarius*). *Comparative Biochemistry and Physiology Part A: Physiology* 101:195–201.
- Etterson J. R., Shaw R. G. 2001. Constraint to adaptive evolution in response to global warming. *Science* 294:151–154.
- Fahrig L. 2003. Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology and Systematics* 34:487–515.
- Faye B. 1997. *Le guide de l'élevage du dromadaire*. Livourne (France): Sanofi Santé Nutrition Animale.
- Finberg J. P., Yagil R., Berlyne G. M. 1978. Response of the renin-aldosterone system in the camel to acute dehydration. *Journal of Applied Physiology* 44: 926–930.
- Galbraith H., Jones R., Park R., Clough J., Herrod-Julius S., Harrington B., Page G. 2002. Global climate change and sea level rise: potential losses of intertidal habitat for shorebirds. *Waterbirds* 25:173–183.
- Ghalambor C. K., McKay J. K., Carroll S. P., Reznick D. N. 2007. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology* 21:394–407.
- Gingerich P. D. 2006. Environment and evolution through the Paleocene-Eocene thermal maximum. *Trends in Ecology and Evolution* 21:246–253.
- Gomulkiewicz R., Houle D. 2009. Demographic and genetic constraints on evolution. *American Naturalist* 174:E218–E229.
- Gonzalez A., Ronce O., Ferriere R., Hochberg M. E. 2013. Evolutionary rescue: an emerging focus at the intersection between ecology and evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences* 368:20120404.
- Gould S. J., Lewontin R. C. 1979. The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proceedings of the Royal Society B: Biological Sciences* 205:581–598.
- Gould S. J., Vrba E. S. 1982. Exaptation—a missing term in the science of form. *Paleobiology* 8:4–15.
- Grubbs D. E. 1980. Tritiated water turnover in free-living desert rodents. *Comparative Biochemistry and Physiology Part A: Physiology* 66:89–98.
- Haim A., Rozenfeld F. M. 1995. Temporal segregation in co-existing *Acomys* species: the possible role of nest site. *Journal of Arid Environments* 29:505–509.
- Harley C. D. G. 2011. Climate change, keystone predation, and biodiversity loss. *Science* 334:1124–1127.
- Hetem R. S., Strauss W. M., Fick L. G., Maloney S. K., Meyer L. C. R., Fuller A., Shobrak M., Mitchell D. 2012. Selective brain cooling in Arabian oryx (*Oryx leucoryx*): a physiological mechanism for

- coping with aridity? *Journal of Experimental Biology* 215:3917–3924.
- Hodgson D., McDonald J. L., Hosken D. J. 2015. What do you mean, “resilient”? *Trends in Ecology and Evolution* 30:503–506.
- Hoffman J. 2014. Physiological response of *Bufo viridis* (Laurenti, 1768) populations across an aridity gradient. *Alytes—International Journal of Batrachology* 30:33–41.
- Hoffmann A. A., Sgrò C. M. 2011. Climate change and evolutionary adaptation. *Nature* 470:479–485.
- Hughes T. P., Rodrigues M. J., Bellwood D. R., Ceccarelli D., Hoegh-Guldberg O., McCook L., Moltschaniwskyj N., Pratchett M. S., Steneck R. S., Willis B. 2007. Phase shifts, herbivory, and the resilience of coral reefs to climate change. *Current Biology* 17:360–365.
- IPCC 2007. *Climate Change 2007: The Physical Science Basis: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. D. Solomon, Q. M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor, and H. L. Miller. Cambridge (United Kingdom): Cambridge University Press.
- IPCC 2013. *Climate Change 2013: The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker, D. Qin, G. K. Plattner, M. M. B. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley. Cambridge (United Kingdom): Cambridge University Press.
- Jablonska E., Lamb M. J. 2005. *Evolution in Four Dimensions: Genetic, Epigenetic, Behavioral, and Symbolic Variation in the History of Life*. Cambridge (Massachusetts): MIT Press.
- Jablonska E., Lamb M. J. 2007. Précis of *Evolution in Four Dimensions*. *Behavioral and Brain Sciences* 30:353–392.
- Jackson D. C., Schmidt-Nielsen K. 1964. Countercurrent heat exchange in the respiratory passages. *Proceedings of the National Academy of Sciences of the United States of America* 51:1192–1197.
- Jackson T. P., Bennett N. C., Spinks A. C. 2004. Is the distribution of the arid-occurring otomysine rodents of southern Africa related to physiological adaptation or refuge type? *Journal of Zoology* 264:1–10.
- Lande R., Shannon S. 1996. The role of genetic variation in adaptation and population persistence in a changing environment. *Evolution* 50:434–437.
- Larson G., Stephens P. A., Tehrani J. J., Layton R. H. 2013. Exaptation exaptation. *Trends in Ecology and Evolution* 28:497–498.
- Ledón-Rettig C. C., Pfennig D. W., Nascone-Yoder N. 2008. Ancestral variation and the potential for genetic accommodation in larval amphibians: implications for the evolution of novel feeding strategies. *Evolution and Development* 10:316–325.
- Lewontin R. C. 1970. The units of selection. *Annual Review of Ecology and Systematics* 1:1–18.
- Li Y., Ye W., Wang M., Yan X. 2009. Climate change and drought: a risk assessment of crop-yield impacts. *Climate Research* 39:31–46.
- Lind M. I., Yarlett K., Reger J., Carter M. J., Becker- man A. P. 2015. The alignment between phenotypic plasticity, the major axis of genetic variation and the response to selection. *Proceedings of the Royal Society B: Biological Sciences* 282:20151651.
- Maickel R. P., Stern D. N., Takabatake E., Brodie B. B. 1967. The sympathetic nervous system as a homeostatic mechanism. II. Effect of adrenocortical hormones on body temperature maintenance of cold-exposed adrenalectomized rats. *Journal of Pharmacology and Experimental Therapeutics* 157: 111–116.
- McCue M. D. 2010. Starvation physiology: reviewing the different strategies animals use to survive a common challenge. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 156:1–18.
- McEwen B. S., Wingfield J. C. 2003. The concept of allostasis in biology and biomedicine. *Hormones and Behavior* 43:2–15.
- McNaughton S. J., Wallace L. L., Coughenour M. B. 1983. Plant adaptation in an ecosystem context: effects of defoliation, nitrogen, and water on growth of an African C4 sedge. *Ecology* 64:307–318.
- Meigs P. 1953. World distribution of arid and semi-arid homoclimates. Pages 203–209 in *Reviews of Research on Arid Zone Hydrology*. Arid Zone Research Series, Volume 1. Paris (France): UNESCO.
- Mery F., Burns J. G. 2010. Behavioural plasticity: an interaction between evolution and experience. *Evolutionary Ecology* 24:571–583.
- Michel M. J., Chevin L.-M., Knouft J. H. 2014. Evolution of phenotype-environment associations by genetic responses to selection and phenotypic plasticity in a temporally autocorrelated environment. *Evolution* 68:1374–1384.
- Miller G. H., Fogel M. L., Magee J. W., Gagan M. K., Clarke S. J., Johnson B. J. 2005. Ecosystem collapse in Pleistocene Australia and a human role in megafaunal extinction. *Science* 309:287–290.
- Mitchell D., Laburn H. P., Nijland M. J. M., Zurovsky Y., Mitchell G. 1987. Selective brain cooling and survival. *South African Journal of Science* 83:598–604.
- Mitchell D., Maloney S. K., Jessen C., Laburn H. P., Kamerman P. R., Mitchell G., Fuller A. 2002. Adaptive heterothermy and selective brain cooling in arid-zone mammals. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 131:571–585.
- Moritz C., Agudo R. 2013. The future of species under climate change: resilience or decline? *Science* 341:504–508.

- Murren C. J., Auld J. R., Callahan H., Ghalambor C. K., Handelsman C. A., Heskel M. A., Kingsolver J. G., Maclean H. J., Masel J., Maughan H., Pfennig D. W., Reylea R. A., Seiter S., Snell-Rood E., Steiner U. K., Schlichting C. D. 2015. Constraints on the evolution of phenotypic plasticity: limits and costs of phenotype and plasticity. *Heredity* 115:293–301.
- Nash D. J., Engineering Group Working Party. 2012. Desert environments. *Geological Society, London, Engineering Geology Special Publications* 25:7–32.
- Negus N. C., Berger P. J., Brown B. W. 1986. Microtine population dynamics in a predictable environment. *Canadian Journal of Zoology* 64:785–792.
- Nettle D., Bateson M. 2015. Adaptive developmental plasticity: what is it, how can we recognize it and when can it evolve? *Proceedings of the Royal Society B: Biological Sciences* 282:20151005.
- Ockendon N., Baker D. J., Carr J. A., White E. C., Almond R. E. A., Amano T., Bertram E., Bradbury R. B., Bradley C., Butchart S. H. M., Doswald N., Foden W., Gill D. J. C., Green R. E., Sutherland W. J., Tanner E. V. J., Pearce-Higgins J. W. 2014. Mechanisms underpinning climatic impacts on natural populations: altered species interactions are more important than direct effects. *Global Change Biology* 20:2221–2229.
- Oppenheimer M., Campos M., Warren R., Birkmann J., Luber G., O'Neill B., Takahashi K. 2014. Emergent risks and key vulnerabilities. Pages 1039–1099 in *Climate Change 2014: Impacts, Adaptation, and Vulnerability: Part A: Global and Sectoral Aspects: Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge (United Kingdom): Cambridge University Press.
- Ostrowski S., Williams J. B., Mésochina P., Sauerwein H. 2006. Physiological acclimation of a desert antelope, Arabian oryx (*Oryx leucoryx*), to long-term food and water restriction. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 176:191–201.
- Ouajd S., Kamel B. 2009. Physiological particularities of dromedary (*Camelus dromedarius*) and experimental implications. *Scandinavian Journal of Laboratory Animal Science* 36:19–29.
- Ozgul A., Tuljapurkar S., Benton T. G., Pemberton J. M., Clutton-Brock T. H., Coulson T. 2009. The dynamics of phenotypic change and the shrinking sheep of St. Kilda. *Science* 325:464–467.
- Ozgul A., Childs D. Z., Oli M. K., Armitage K. B., Blumstein D. T., Olson L. E., Tuljapurkar S., Coulson T. 2010. Coupled dynamics of body mass and population growth in response to environmental change. *Nature* 466:482–485.
- Pekár S., Michalko R., Korenko S., Šedo O., Líznarová E., Sentenská L., Zdráhal Z. 2013. Phenotypic integration in a series of trophic traits: tracing the evolution of myrmecophagy in spiders (Araneae). *Zoology* 116:27–35.
- Pianka E. R. 2011. *Evolutionary Ecology*. Seventh Edition. Austin (Texas): E. R. Pianka.
- Piersma T., Drent J. 2003. Phenotypic flexibility and the evolution of organismal design. *Trends in Ecology and Evolution* 18:228–233.
- Piersma T., Lindström Å. 1997. Rapid reversible changes in organ size as a component of adaptive behaviour. *Trends in Ecology and Evolution* 12:134–138.
- Piersma T., van Gils J. A. 2010. *The Flexible Phenotype: A Body-Centred Integration of Ecology, Physiology, and Behaviour*. Oxford (United Kingdom): Oxford University Press.
- Pimm S. L. 2009. Climate disruption and biodiversity. *Current Biology* 19:R595–R601.
- Pinter A. J. 1994. Climatic factors, reproductive success and population dynamics in the montane vole, *Microtus montanus*. *University of Wyoming National Park Service Research Center Annual Report* 18:92–93.
- Podos J., Southall J. A., Rossi-Santos M. R. 2004. Vocal mechanics in Darwin's finches: correlation of beak gape and song frequency. *Journal of Experimental Biology* 207:607–619.
- Quintero I., Wiens J. J. 2013. Rates of projected climate change dramatically exceed past rates of climatic niche evolution among vertebrate species. *Ecology Letters* 16:1095–1103.
- Randall J. A. 1993. Behavioural adaptations of desert rodents (Heteromyidae). *Animal Behaviour* 45: 263–287.
- Reik W. 2007. Stability and flexibility of epigenetic gene regulation in mammalian development. *Nature* 447:425–432.
- Rezende E. L., Diniz-Filho J. A. F. 2012. Phylogenetic analyses: comparing species to infer adaptations and physiological mechanisms. *Comprehensive Physiology* 2:639–674.
- Rivers J. W., Liebl A. L., Owen J. C., Martin L. B., Betts M. G. 2012. Baseline corticosterone is positively related to juvenile survival in a migrant passerine bird. *Functional Ecology* 26:1127–1134.
- Robertson B. A., Rehage J. S., Sih A. 2013. Ecological novelty and the emergence of evolutionary traps. *Trends in Ecology and Evolution* 28:552–560.
- Romero L. M. 2012. Using the reactive scope model to understand why stress physiology predicts survival during starvation in Galápagos marine iguanas. *General and Comparative Endocrinology* 176:296–299.
- Romero L. M., Dickens M. J., Cyr N. E. 2009. The reactive scope model—a new model integrating homeostasis, allostasis, and stress. *Hormones and Behavior* 55:375–389.
- Rymer T. L., Pillay N., Schradin C. 2013. Extinction or survival? Behavioral flexibility in response to

- environmental change in the African striped mouse *Rhabdomys*. *Sustainability* 5:163–186.
- Saccheri I. J., Rousset F., Watts P. C., Brakefield P. M., Cook L. M. 2008. Selection and gene flow on a diminishing cline of melanic peppered moths. *Proceedings of the National Academy of Sciences of the United States of America* 105:16212–16217.
- Sarma V. V. L. N., Krishna G. M., Malini B. H., Rao K. N. 2001. Landuse/landcover change detection through remote sensing and its climatic implications in the Godavari delta region. *Journal of the Indian Society of Remote Sensing* 29:85–91.
- Scheiner S. M., Gomulkiewicz R., Holt R. D. 2015. The genetics of phenotypic plasticity. XIV. Coevolution. *American Naturalist* 185:594–609.
- Schradin C., Pillay N. 2005. Demography of the striped mouse (*Rhabdomys pumilio*) in the succulent karoo. *Mammalian Biology* 70:84–92.
- Schradin C., Krackow S., Schubert M., Keller C., Schradin B., Pillay N. 2007. Regulation of activity in desert-living striped mice: the importance of basking. *Ethology* 113:606–614.
- Seager R., Tang M., Held I., Kushnir Y., Lu J., Vecchi G., Huang H.-P., Harnik N., Leetmaa A., Lau N.-C., Li C., Velez J., Naik N. 2007. Model projections of an imminent transition to a more arid climate in southwestern North America. *Science* 316: 1181–1184.
- Siebert B. D., Macfarlane W. V. 1975. Dehydration in desert cattle and camels. *Physiological Zoology* 48:36–48.
- Sih A., Ferrari M. C. O., Harris D. J. 2011. Evolution and behavioural responses to human-induced rapid environmental change. *Evolutionary Applications* 4:367–387.
- Smit B., Burton I., Klein R. J. T., Wandel J. 2000. An anatomy of adaptation to climate change and variability. *Climate Change* 45:223–251.
- Sperry J. H., Weatherhead P. J. 2008. Prey-mediated effects of drought on condition and survival of a terrestrial snake. *Ecology* 89:2770–2776.
- Standen E. M., Du T. Y., Larsson H. C. E. 2014. Developmental plasticity and the origin of tetrapods. *Nature* 513:54–58.
- Steiner U. K., Van Buskirk J. 2008. Environmental stress and the costs of whole-organism phenotypic plasticity in tadpoles. *Journal of Evolutionary Biology* 21:97–103.
- Stewart D. R. M. 1963. The Arabian oryx (*Oryx leucoryx* Pallas). *African Journal of Ecology* 1:103–117.
- Tanentzap A. J., Lee W. G., Schulz K. A. C. 2013. Niches drive peaked and positive relationships between diversity and disturbance in natural ecosystems. *Ecosphere* 4:133.
- Taylor C. R. 1970. Dehydration and heat: effects on temperature regulation of East African ungulates. *American Journal of Physiology* 219:1136–1139.
- Valladares F., Balaguer L., Martinez-Ferri E., Perez-Corona E., Manrique E. 2002. Plasticity, instability and canalization: is the phenotypic variation in seedlings of sclerophyll oaks consistent with the environmental unpredictability of Mediterranean ecosystems? *New Phytologist* 156:457–467.
- Van Buskirk J., Steiner U. K. 2009. The fitness costs of developmental canalization and plasticity. *Journal of Evolutionary Biology* 22:852–860.
- Vicente-Serrano S. M., Gouveia C., Camarero J. J., Beguería S., Trigo R., López-Moreno J. I., Azorín-Molina C., Pasho E., Lorenzo-Lacruz J., Revuelto J., Morán-Tejeda E., Sanchez-Lorenzo A. 2013. Responses of vegetation to drought time-scales across global land biomes. *Proceedings of the National Academy of Sciences of the United States of America* 110: 52–57.
- Viles H. A., Goudie A. S. 2013. Weathering in the central Namib Desert, Namibia: controls, processes and implications. *Journal of Arid Environments* 93:20–29.
- Walker B. H., Emslie R. H., Owen-Smith R. N., Scholes R. J. 1987. To cull or not to cull: lessons from a southern African drought. *Journal of Applied Ecology* 24:381–401.
- Ward D. 2009. Morphological, physiological, and behavioural adaptations of desert animals to the abiotic environment. Pages 66–101 in *The Biology of Deserts*, by D. Ward. Oxford (United Kingdom): Oxford University Press.
- Webb J. K., Letnic M., Jessop T. S., Dempster T. 2014. Behavioural flexibility allows an invasive vertebrate to survive in a semi-arid environment. *Biology Letters* 10:20131014.
- West-Eberhard M. J. 1989. Phenotypic plasticity and the origins of diversity. *Annual Review of Ecology and Systematics* 20:249–278.
- Williams J. B., Ostrowski S., Bedin E., Ismail K. 2001. Seasonal variation in energy expenditure, water flux and food consumption of Arabian oryx *Oryx leucoryx*. *Journal of Experimental Biology* 204:2301–2311.
- Williams S. E., Shoo L. P., Isaac J. L., Hoffmann A. A., Langham G. 2008. Towards an integrated framework for assessing the vulnerability of species to climate change. *PLOS Biology* 6:e325.
- Willmer P., Stone G., Johnson I. 2000. *Environmental Physiology of Animals*. Malden (Massachusetts): Blackwell Science.
- Wingfield J. C. 2013. The comparative biology of environmental stress: behavioural endocrinology and variation in ability to cope with novel, changing environments. *Animal Behaviour* 85: 1127–1133.
- Ylönen H., Ronkainen H. 1994. Breeding suppression in the bank vole as antipredatory adaptation in a predictable environment. *Evolutionary Ecology* 8:658–666.

- Yousef M. K., Johnson H. D. 1975. Thyroid activity in desert rodents: a mechanism for lowered metabolic rate. *American Journal of Physiology* 229:427–431.
- Zhang Y. M., Chen Y. N., Pan B. R. 2005. Distribution and floristics of desert plant communities in the lower reaches of Tarim River, southern Xinjiang, People's Republic of China. *Journal of Arid Environments* 63:772–784.
- Zhang Z.-Q., Wang D.-H. 2007. Seasonal changes in thermogenesis and body mass in wild Mongolian gerbils (*Meriones unguiculatus*). *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 148:346–353.

Associate Editor: Andrew Sih
Handling Editor: John J. Wiens

APPENDIX 1

References associated with Tables 1, 2a, 2b, and 2c

- Alagaili A. N., Mohammed O. B., Bennett N. C., Oosthuizen M. K. 2013. A tale of two jirds: the locomotory activity patterns of the King jird (*Meriones rex*) and Lybian jird (*Meriones libycus*) from Saudi Arabia. *Journal of Arid Environments* 88:102–112.
- Albert S. K., Krausman P. R. 1993. Desert mule deer and forage resources in Southwest Arizona. *Southwestern Naturalist* 38:198–205.
- Aulagnier S., Granjon L., Shenbrot G., Bukhnikashvili A. 2008. *Meriones libycus*. IUCN 2014. IUCN Red List of Threatened Species, Version 2014.3. <http://www.iucnredlist.org>. (Accessed 3 December 2014).
- Bartholomew G. A., MacMillen R. E. 1961. Oxygen consumption, estivation, and hibernation in the kangaroo mouse, *Microdipodops pallidus*. *Physiological Zoology* 34:177–183.
- Baudinette R. V., Snyder G. K., Frappell P. B. 1992. Energetic cost of locomotion in the Tammar wallaby. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* 262:R771–R778.
- Baumgardner G. D. 1991. *Dipodomys compactus*. *Mammalian Species* 369:1–4.
- Ben Goumi M., Riad F., Giry J., de la Farge F., Safwate A., Davicco M.-J., Barlet J.-P. 1993. Hormonal control of water and sodium in plasma and urine of camels during dehydration and rehydration. *General and Comparative Endocrinology* 89:378–386.
- Bennett A. F., Huey R. B., John-Alder H., Nagy K. A. 1984. The parasol tail and thermoregulatory behavior of the Cape ground squirrel *Xerus inauris*. *Physiological Zoology* 57:57–62.
- Bothma J. du P., Riche E. A. N. le. 1984. Aspects of the ecology and the behaviour of the leopard *Panthera pardus* in the Kalahari desert. *Koedoe Supplement* (1984):259–279.
- Bozinovic F., Contreras L. C. 1990. Basal rate of metabolism and temperature regulation of two desert herbivorous octodontid rodents: *Octomys mimax* and *Typanoctomys barrerae*. *Oecologia* 84:567–570.
- Bozinovic F., Gallardo P. A., Visser G. H., Cortés A. 2003. Seasonal acclimatization in water flux rate, urine osmolality and kidney water channels in free-living degus: molecular mechanisms, physiological processes and ecological implications. *Journal of Experimental Biology* 206:2959–2966.
- Bradford D. F. 1974. Water stress of free-living *Peromyscus truei*. *Ecology* 55:1407–1414.
- Brown J. H., Munger J. C. 1985. Experimental manipulation of a desert rodent community: food addition and species removal. *Ecology* 66:1545–1563.
- Buffenstein R. 1984. Energy and water balance during torpor and hydropenia in the pygmy gerbil, *Gerbillus pusillus*. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 154:535–544.
- Buffenstein R. 1985. The effect of a high fibre diet on energy and water balance in two Namib desert rodents. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 155:211–218.
- Buffenstein R., Campbell W. E., Jarvis J. U. M. 1985. Identification of crystalline allantoin in the urine of African Cricetidae (Rodentia) and its role in their water economy. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 155:493–499.
- Bukovetzky E., Fares F., Schwimmer H., Haim A. 2012. Reproductive and metabolic responses of desert adapted common spiny male mice (*Acomys cahirinus*) to vasopressin treatment. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 162:349–356.
- Cain J. W. III, Krausman P. R., Rosenstock S. S., Turner J. C. 2006. Mechanisms of thermoregulation and water balance in desert ungulates. *Wildlife Society Bulletin* 34:570–581.
- Carpenter R. E. 1966. A comparison of thermoregulation and water metabolism in the kangaroo rats *Dipodomys agilis* and *Dipodomys merriami*. *University of California Publications in Zoology* 78:1–36.
- Carpenter R. E., Graham J. B. 1967. Physiological responses to temperature in the long-nosed bat, *Leptonycteris sanborni*. *Comparative Biochemistry and Physiology* 22:709–722.

- Casey N. H., van Niekerk W. A. 1988. The Boer goat. I. Origin, adaptability, performance testing, reproduction and milk production. *Small Ruminant Research* 1:291–302.
- Castel M., Abraham M. 1972. Effects of a dry diet on the antidiuretic hormone content of the neurohypophysis in spiny mice as compared to the albino rat and mouse. *General and Comparative Endocrinology* 19:48–55.
- Chappell M. A., Bartholomew G. A. 1981a. Activity and thermoregulation of the antelope ground squirrel *Ammospermophilus leucurus* in winter and summer. *Physiological Zoology* 54:215–223.
- Chappell M. A., Bartholomew G. A. 1981b. Standard operative temperatures and thermal energetics of the antelope ground squirrel *Ammospermophilus leucurus*. *Physiological Zoology* 54:81–93.
- Chen X., Dickman C. R., Thompson M. B. 1998. Diet of the mulgara, *Dasyurus cristicauda* (Marsupialia: Dasyuridae), in the Simpson Desert, central Australia. *Wildlife Research* 25:233–242.
- Christian D. P. 1978. Effects of humidity and body size on evaporative water loss in three desert rodents. *Comparative Biochemistry and Physiology Part A: Physiology* 60:425–430.
- Christian D. P. 1980. Vegetative cover, water resources, and microdistributional patterns in a desert rodent community. *Journal of Animal Ecology* 49:807–816.
- Cornelius C. E., Kaneko J. J. 1962. Erythrocyte life span in the guanaco. *Science* 137:673–674.
- Curtis B. A., Perrin M. R. 1979. Food preference of the vlei rat (*Otomys irroratus*) and the four-striped mouse (*Rhabdomys pumilio*). *South African Journal of Zoology* 14:224–229.
- Cypher B. L. 2003. Foxes (*Vulpes* species, *Urocyon* species, and *Alopex lagopus*). Pages 511–546 in *Wild Mammals of North America: Biology, Management, and Conservation*, edited by G. A. Feldhamer, B. C. Thompson, and J. A. Chapman. Second Edition. Baltimore (Maryland): Johns Hopkins University Press.
- Daly M., Daly S. 1973. On the feeding ecology of *Psammomys obesus* (Rodentia: Gerbillinae) in Wadi Saoura, Algeria. *Mammalia* 37:545–561.
- Dawson T. J., Brown G. D. 1970. A comparison of the insulative and reflective properties of the fur of desert kangaroos. *Comparative Biochemistry and Physiology* 37:23–38.
- Dawson T. J., Taylor C. R. 1973. Energetic cost of locomotion in kangaroos. *Nature* 246:313–314.
- Dawson T. J., Denny M. J. S., Russell E. M., Ellis B. 1975. Water usage and diet preferences of free ranging kangaroos, sheep and feral goats in the Australian arid zone during summer. *Journal of Zoology* 177:1–23.
- de Rouffignac C., Morel F. 1965. Etude comparée du renouvellement de l'eau chez quatre espèces de rongeurs, dont deux espèces d'habitat désertique. *Journal of Physiology* 58:309–322.
- Deblinger R. D., Alldredge A. W. 1989. Management implications of variations in pronghorn social behavior. *Wildlife Society Bulletin* 17:82–87.
- Degen A. A. 1977. Fat-tailed Awassi and German mutton merino sheep under semi-arid conditions. 1. Total body water, its distribution and water turnover. *Journal of Agricultural Science* 88:693–698.
- Degen A. A., Shkolnik A. 1978. Thermoregulation in fat-tailed Awassi, a desert sheep, and in German mutton merino, a mesic sheep. *Physiological Zoology* 51:333–339.
- Dereje M., Udén P. 2005. The browsing dromedary camel. I. Behaviour, plant preference and quality of forage selected. *Animal Feed Science and Technology* 121:297–308.
- Dexter N. 2003. The influence of pasture distribution, and temperature on adult body weight of feral pigs in a semi-arid environment. *Wildlife Research* 30:75–79.
- Diaz G. B., Ojeda R. A. 1999. Kidney structure and allometry of Argentine desert rodents. *Journal of Arid Environments* 41:453–461.
- Donadio E., Buskirk S. W., Novaro A. J. 2012. Juvenile and adult mortality patterns in a vicuña (*Vicugna vicugna*) population. *Journal of Mammalogy* 93:1536–1544.
- Downs C. T., Perrin M. R. 1990. The effect of diet on water and energy turnover rates of four *Gerbillurus* species in captivity. *Journal of Zoology* 222:215–233.
- Downs C. T., Perrin M. R. 1991. Urinary concentrating ability of four *Gerbillurus* species of southern Africa arid regions. *Journal of Arid Environments* 20:71–81.
- Downs C. T., Perrin M. R. 1994. Comparative aspects of the thermal biology of the short-tailed gerbil, *Desmodillus auricularis*, and the bushveld gerbil, *Taterillus leucogaster*. *Journal of Thermal Biology* 19:385–392.
- du Plessis A., Kerley G. I. H., deo Winter P. E. 1992. Refuge microclimates of rodents: a surface nesting *Otomys unisulcatus* and a burrowing *Parotomys brantsii*. *Acta Theriologica* 37:351–358.
- Ehrhardt N., Heldmaier G., Exner C. 2005. Adaptive mechanisms during food restriction in *Acomys russatus*: the use of torpor for desert survival. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 175:193–200.
- El-Hadi H. M. 1986. The effect of dehydration on Sudanese desert sheep and goats. *Journal of Agricultural Science* 106:17–20.
- El Hilali M., Veillat J.-P. 1975. *Jaculus orientalis*: a true hibernator. *Mammalia* 39:401–404.
- El Ouezzani S., Lafon P., Tramu G., Magoul R. 2001. Neuropeptide Y gene expression in the jerboa arcuate nucleus: modulation by food deprivation and relationship with hibernation. *Neuroscience Letters* 305:127–130.

- Elkhawad A. O. 1992. Selective brain cooling in desert animals: the camel (*Camelus dromedarius*). *Comparative Biochemistry and Physiology Part A: Physiology* 101:195–201.
- Emmanuel B., Howard B. R., Emady M. 1976. Urea degradation in the camel. *Canadian Journal of Animal Science* 56:595–601.
- Faye B. 1997. *Le guide de l'élevage du dromadaire*. Li-bourne (France): Sanofi Santé Nutrition Animale.
- Fielden L. J., Perrin M. R., Hickman G. C. 1990. Water metabolism in the Namib Desert golden mole, *Eremitalpa granti namibensis* (Chrysocloridae). *Comparative Biochemistry and Physiology Part A: Physiology* 96:227–234.
- Finberg J. P., Yagil R., Berlyne G. M. 1978. Response of the renin-aldosterone system in the camel to acute dehydration. *Journal of Applied Physiology* 44:926–930.
- Fourie F. le R., Haim A. 1980. Enzymatic activity of rodents acclimated to cold and long scotophase. *International Journal of Biometeorology* 24:237–241.
- Frean J. L., Downs C. T., Lovegrove B. G. 1998. Osmoregulatory capacity of an arboreal rodent *Thallomys nigricauda*. *Journal of Arid Environments* 40:97–107.
- Fuller A., Maloney S. K., Mitchell G., Mitchell D. 2004. The eland and the oryx revisited: body and brain temperatures of free-living animals. *International Congress Series* 1275:275–282.
- Fuller A., Kamerman P. R., Maloney S. K., Matthee A., Mitchell G., Mitchell D. 2005. A year in the thermal life of a free-ranging herd of springbok *Antidorcas marsupialis*. *Journal of Experimental Biology* 208:2855–2864.
- Funston P. J., Skinner J. D., Dott H. M. 1994. Seasonal variation in movement patterns, home range and habitat selection of buffaloes in a semi-arid habitat. *African Journal of Ecology* 32:100–114.
- Geiser F. 2010. Aestivation in mammals and birds. Pages 95–111 in *Aestivation: Molecular and Physiological Aspects*, edited by C. A. Navas and J. E. Carvalho. Berlin (Germany): Springer.
- Geiser F., Pavey C. R. 2007. Basking and torpor in a rock-dwelling desert marsupial: survival strategies in a resource-poor environment. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 177:885–892.
- Ghobrial L. I. 1970. The water relations of the desert antelope *Gazella dorcas dorcas*. *Physiological Zoology* 43:249–256.
- Giannoni S. M., Dacar M., Taraborelli P., Borghi C. E. 2001. Seed hoarding by rodents of the Monte Desert, Argentina. *Austral Ecology* 26:259–263.
- Giannoni S. M., Borghi C. E., Dacar M., Campos C. M. 2005. Main food categories in diets of sigmodontine rodents in the Monte (Argentina). *Mastozoología Neotropical* 12:181–187.
- Gihad El S. A., El Gallad T. T., Sooud A. E. O., Abou El Nasr H. M., Farid M. F. A. 1989. Feed and water intake, digestibility and nitrogen utilization by camels compared to sheep and goats fed low protein desert by-products. *Options Méditerranéennes—Série Séminaires* 2:75–81.
- Gilad O., Grant W. E., Saltz D. 2008. Simulated dynamics of Arabian oryx (*Oryx leucoryx*) in the Israeli Negev: effects of migration corridors and post-reintroduction changes in natality on population viability. *Ecological Modelling* 210:169–178.
- Golightly R. T. Jr., Ohmart R. D. 1983. Metabolism and body temperature of two desert canids: coyotes and kit foxes. *Journal of Mammalogy* 64:624–635.
- Grigg G., Beard L., Dörges B., Heucke J., Coventry J., Coppock A., Blomberg S. 2009. Strategic (adaptive) hypothermia in bull dromedary camels during rut; could it increase reproductive success? *Biology Letters* 5:853–856.
- Grubbs D. E. 1980. Tritiated water turnover in free-living desert rodents. *Comparative Biochemistry and Physiology Part A: Physiology* 66:89–98.
- Gutman R., Choshniak I., Kronfeld-Schor N. 2006. Defending body mass during food restriction in *Acomys russatus*: a desert rodent that does not store food. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* 290:R881–R891.
- Gutman R., Hacmon-Keren R., Choshniak I., Kronfeld-Schor N. 2008. Effect of food availability and leptin on the physiology and hypothalamic gene expression of the golden spiny mouse: a desert rodent that does not hoard food. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* 295:R2015–R2023.
- Haim A. 1984. Adaptive variations in heat production within gerbils (genus *Gerbillus*) from different habitats. *Oecologia* 61:49–52.
- Haim A., Borut A. 1981. Heat production and dissipation in golden spiny mice, *Acomys russatus* from two extreme habitats. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 142:445–450.
- Haim A., Fairall N. 1986. Geographical variations in heat production and dissipation within two populations of *Rhabdomys pumilio* (Muridae). *Comparative Biochemistry and Physiology Part A: Physiology* 84:111–112.
- Haim A., Fourie F. le R. 1980a. Heat production in nocturnal (*Praomys natalensis*) and diurnal (*Rhabdomys pumilio*) South African murids. *South African Journal of Zoology* 15:91–94.
- Haim A., Fourie F. le R. 1980b. Heat production in cold and long scotophase acclimated and winter acclimatized rodents. *International Journal of Biometeorology* 24:231–236.
- Haim A., Izhaki I. 1993. The ecological significance of resting metabolic rate and non-shivering thermogenesis for rodents. *Journal of Thermal Biology* 18:71–81.

- Haim A., Rozenfeld F. M. 1995. Temporal segregation in co-existing *Acomys* species: the possible role of nest site. *Journal of Arid Environments* 29:505–509.
- Haim A., Heth G., Avnon Z., Nevo E. 1984. Adaptive physiological variation in nonshivering thermogenesis and its significance in speciation. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 154:145–147.
- Haim A., Pelaot I., Sela A. 1986. Comparison of eco-physiological parameters between two *Apodemus* species coexisting in the same habitat. Pages 33–40 in *Environmental Quality and Ecosystem Stability*, Volume III A/B, edited by Z. Dubinsky and Y. Steinberger. Ramat-Gan (Israel): Bar-Ilan University Press.
- Haines H., Macfarlane W. V., Setchell C., Howard B. 1974. Water turnover and pulmocutaneous evaporation of Australian desert dasyurids and murids. *American Journal of Physiology* 227:958–963.
- Hansen M. C. 1982. Desert bighorn sheep: another view. *Wildlife Society Bulletin* 10:133–140.
- Hartman L. A., Morton M. L. 1973. Plasma volume changes in ground squirrels during water deprivation. *Comparative Biochemistry and Physiology Part A: Physiology* 46:79–94.
- Hatough-Bouran A. 1990. The burrowing habits of desertic rodents *Jaculus jaculus* and *Gerbillus dasyurus* in the Shaumari Reserve in Jordan. *Mammalia* 54:341–359.
- Heimeier R. A., Donald J. A. 2006. The response of the natriuretic peptide system to water deprivation in the desert rodent, *Notomys alexis*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 143:193–201.
- Hetem R. S., Strauss W. M., Fick L. G., Maloney S. K., Meyer L. C. R., Shobrak M., Fuller A., Mitchell D. 2010. Variation in the daily rhythm of body temperature of free-living Arabian oryx (*Oryx leucoryx*): does water limitation drive heterothermy? *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 180:1111–1119.
- Hetem R. S., Strauss W. M., Heusinkveld B. G., de Bie S., Prins H. H. T., van Wieren S. E. 2011. Energy advantages of orientation to solar radiation in three African ruminants. *Journal of Thermal Biology* 36:452–460.
- Hetem R. S., Strauss W. M., Fick L. G., Maloney S. K., Meyer L. C. R., Fuller A., Shobrak M., Mitchell D. 2012. Selective brain cooling in Arabian oryx (*Oryx leucoryx*): a physiological mechanism for coping with aridity? *Journal of Experimental Biology* 215:3917–3924.
- Hinds D. S., MacMillen R. E. 1985. Scaling of energy metabolism and evaporative water loss in heteromyid rodents. *Physiological Zoology* 58:282–298.
- Hinton H. E., Dunn A. M. S. 1967. *Mongooses: Their Natural History and Behaviour*. Edinburgh (United Kingdom): Oliver & Boyd.
- Hochman V., Kotler B. P. 2006. Effects of food quality, diet preference and water on patch use by Nubian ibex. *Oikos* 112:547–554.
- Hoppe P. P. 1977. Comparison of voluntary food and water consumption and digestion in Kirk's dikdik and suni. *African Journal of Ecology* 15:41–48.
- Howell A. B., Gersh I. 1935. Conservation of water by the rodent *Dipodomys*. *Journal of Mammalogy* 16:1–9.
- Hudson J. W. 1964. Temperature regulation in the round-tailed ground squirrel, *Citellus tereticaudus*. *Annales Academiae Scientiarum Fennicae, Series A IV* 71:217–233.
- Hulbert A. J., Dawson T. J. 1974. Thermoregulation in perameloid marsupials from different environments. *Comparative Biochemistry and Physiology Part A: Physiology* 47:591–616.
- Hulbert A. J., Hinds D. S., MacMillen R. E. 1985. Minimal metabolism, summit metabolism and plasma thyroxine in rodents from different environments. *Comparative Biochemistry and Physiology Part A: Physiology* 81:687–693.
- Hume J. D., Rübsamen K., Engelhardt W. V. 1980. Nitrogen metabolism and urea kinetics in the rock hyrax (*Procavia habessinica*). *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 138:307–314.
- Ilan M., Yom-Tov Y. 1990. Diel activity pattern of a diurnal desert rodent, *Psammomys obesus*. *Journal of Mammalogy* 71:66–69.
- Intriss C., Best T. L. 1990. *Dipodomys panamintinus*. *Mammalian Species* 354:1–7.
- Jaber L. S., Habre A., Rawda N., Abi Said M., Barbour E. K., Hamadeh S. 2004. The effect of water restriction on certain physiological parameters in Awassi sheep. *Small Ruminant Research* 54:115–120.
- Jaber L. S., Hanna N., Barbour E. K., Abi Said M., Rawda N., Chedid M., Hamadeh S. K. 2011. Fat mobilization in water restricted Awassi ewes supplemented with vitamin C. *Journal of Arid Environments* 75:625–628.
- Jackson D. C., Schmidt-Nielsen K. 1964. Countercurrent heat exchange in the respiratory passages. *Proceedings of the National Academy of Sciences of the United States of America* 51:1192–1197.
- Jackson T. P., Bennett N. C., Spinks A. C. 2004. Is the distribution of the arid-occurring otomyine rodents of southern Africa related to physiological adaptation or refuge type? *Journal of Zoology* 264:1–10.
- Kaißling B., de Rouffignac C., Barrett J. M., Kriz W. 1975. The structural organization of the kidney of the desert rodent *Psammomys obesus*. *Anatomy and Embryology* 148:121–143.
- Kam M., Degen A. A. 1993. Effect of dietary pre-formed water on energy and water budgets of two sympatric desert rodents, *Acomys russatus* and *Acomys cahirinus*. *Journal of Zoology* 231:51–59.

- Kamau J. M. Z., Maina J. N., Maloij G. M. O. 1984. The design and the role of the nasal passages in temperature regulation in the dik-dik antelope (*Rhynchotragus Kirkii*) with observations on the carotid rete. *Respiration Physiology* 56:183–194.
- Kataria N., Kataria A. K. 2006. Endocrine and metabolic responses of Marwari sheep in arid tract. *Slovenian Veterinary Research* 43:135–142.
- Kennedy P. M., Hume I. D. 1978. Recycling of urea nitrogen to the gut of the tammar wallaby (*Macropus eugenii*). *Comparative Biochemistry and Physiology Part A: Physiology* 61:117–121.
- Khachadurian A. K., Adrouni B., Yacoubian H. 1966. Metabolism of adipose tissue in the fat tail of the sheep in vivo. *Journal of Lipid Research* 7:427–436.
- Khalil F., Tawfic J. 1963. Some observations on the kidney of the desert *J. jaculus* and *G. gerbillus* and their possible bearing on the water economy of these animals. *Journal of Experimental Zoology Part A* 154:259–271.
- Khan M. S., Ghosh P. K., Sasidharan T. O. 1978. Effect of acute water restriction on plasma proteins and on blood and urinary electrolytes in Barmer goats of the Rajasthan desert. *Journal of Agricultural Science* 91:395–398.
- King J. M., Kingaby G. P., Colvin J. G., Heath B. R. 1975. Seasonal variation in water turnover by oryx and eland on the Galana Game Ranch Research Project. *African Journal of Ecology* 13:287–296.
- Kinnear J. E., Purohit K. G., Main A. R. 1968. The ability of the tammar wallaby (*Macropus eugenii*, Marsupialia) to drink sea water. *Comparative Biochemistry and Physiology* 25:761–782.
- Koford C. B. 1968. Peruvian desert mice: water independence, competition, and breeding cycle near the equator. *Science* 160:552–553.
- Kronfeld-Schor N., Haim A., Dayan T., Zisapel N., Klingenspor M., Heldmaier G. 2000. Seasonal thermogenic acclimation of diurnally and nocturnally active desert spiny mice. *Physiological and Biochemical Zoology* 73:37–44.
- Krysl L. J., Sowell B. F., Hubbert M. E., Plumb G. E., Jewett T. K., Smith M. A., Waggoner J. W. 1984. Horses and cattle grazing in the Wyoming red desert. II. Dietary quality. *Journal of Range Management* 37:252–255.
- Laden S., Nehmadi L., Yagil R. 1987. Dehydration tolerance in Awassi fat-tailed sheep. *Canadian Journal of Zoology* 65:363–367.
- Lagos V. O., Bozinovic F., Contreras L. C. 1995. Microhabitat use by a small diurnal rodent (*Octodon degus*) in a semiarid environment: thermoregulatory constraints or predation risk? *Journal of Mammalogy* 76:900–905.
- Langman V. A., Maloij G. M. O., Schmidt-Nielsen K., Schroter R. C. 1979. Nasal heat exchange in the giraffe and other large mammals. *Respiration Physiology* 37:325–333.
- Larcombe A. N., Withers P. C. 2006. Thermoregulatory, metabolic and ventilatory physiology of the western barred bandicoot (*Perameles bougainville bougainville*) in summer and winter. *Australian Journal of Zoology* 54:15–21.
- Leggett K. E. A. 2004. Coprophagy and unusual thermoregulatory behaviour in desert-dwelling elephants of north-western Namibia. *Pachyderm* 36:113–115.
- Lewis R. E., Lewis J. H., Harrison D. L. 1965. On a collection of mammals from northern Saudi Arabia. *Proceedings of the Zoological Society of London* 144:61–74.
- Lovegrove B. G., Raman J., Perrin M. R. 2001. Daily torpor in elephant shrews (Macroscelidea: *Elephantulus* spp.) in response to food deprivation. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 171:11–21.
- Lowrey C., Longshore K. 2005. *Long Term Conservation Strategy for Palmers Chipmunk (Tamias palmeri) within the Spring Mountains, Nevada*. Henderson (Nevada): United States Geological Survey.
- Lubbe A. 2013. Thermoregulatory plasticity in free-ranging velvet monkeys (*Chlorocebus pygerythrus*). [MSc. thesis]. Johannesburg (South Africa): University of the Witwatersrand.
- Macfarlane W. V., Morris R. J. H., Howard B. 1963. Turnover and distribution of water in desert camels, sheep, cattle and kangaroos. *Nature* 197:270–271.
- MacMillen R. E. 1965. Aestivation in the cactus mouse, *Peromyscus eremicus*. *Comparative Biochemistry and Physiology* 16:227–248.
- MacMillen R. E., Lee A. K. 1967. Australian desert mice: independence of exogenous water. *Science* 158:383–385.
- MacMillen R. E., Lee A. K. 1969. Water metabolism of Australian hopping mice. *Comparative Biochemistry and Physiology* 28:493–514.
- Maloij G. M. O. 1972. Comparative physiology of desert animals. *Symposia of the Zoological Society of London* 31:243–259.
- Maloij G. M. O. 1973. The water metabolism of a small East African antelope: the dik-dik. *Proceedings of the Royal Society B: Biological Sciences* 184:167–178.
- Maloij G. M. O., Rugangazi B. M., Clemens E. T. 1988. Physiology of the dik-dik antelope. *Comparative Biochemistry and Physiology Part A: Physiology* 91:1–8.
- Maloney S. K., Moss G., Cartmell T., Mitchell D. 2005. Alteration in diel activity patterns as a thermoregulatory strategy in black wildebeest (*Connochaetes gnou*). *Journal of Comparative Physiology A: Molecular and Integrative Physiology* 191:1055–1064.
- Marai I. F. M., Haeib A. A. M. 2010. Buffalo's biological functions as affected by heat stress—a review. *Livestock Science* 127:89–109.
- Mares M. A. 1977a. Water economy and salt balance in a South American desert rodent, *Eligmodontia typus*. *Comparative Biochemistry and Physiology Part A: Physiology* 56:325–332.

- Mares M. A. 1977b. Water independence in a South American non-desert rodent. *Journal of Mammalogy* 58:653–656.
- Mares M. A. 1993. Desert rodents, seed consumption, and convergence: evolutionary shuffling of adaptations. *BioScience* 43:372–379.
- Marsh A. C., Louw G., Berry H. H. 1978. Aspects of renal physiology, nutrition and thermoregulation in the ground squirrel *Xerus inauris*. *Madoqua* 2:129–135.
- Mbassa G. K. 1988. Mammalian renal modifications in dry environments. *Veterinary Research Communications* 12:1–18.
- McCutchen H. E. 1981. Desert bighorn zoogeography and adaptation in relation to historic land use. *Wildlife Society Bulletin* 9:171–179.
- McKean T., Walker B. 1974. Comparison of selected cardiopulmonary parameters between the pronghorn and the goat. *Respiration Physiology* 21:365–370.
- Mendelsohn H., Yom-Tov Y. 1999. *Mammalia of Israel*. Jerusalem (Israel): Israel Academy of Sciences and Humanities.
- Mengistu U., Dahlborn K., Olsson K. 2007. Effects of intermittent watering on water balance and feed intake in male Ethiopian Somali goats. *Small Ruminant Research* 67:45–54.
- Merkt J. R., Taylor C. R. 1994. “Metabolic switch” for desert survival. *Proceedings of the National Academy of Sciences of the United States of America* 91:12313–12316.
- Misra A. K., Singh K. 2002. Effect of water deprivation on dry matter intake, nutrient utilization and metabolic water production in goats under semi-arid zone of India. *Small Ruminant Research* 46:159–165.
- Mitchell D., Maloney S. K., Laburn H. P., Knight M. H., Kuhnen G., Jessen C. 1997. Activity, blood temperature and brain temperature of free-ranging springbok. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 167:335–343.
- Mitchell D., Maloney S. K., Jessen C., Laburn H. P., Kamerman P. R., Mitchell G., Fuller A. 2002. Adaptive heterothermy and selective brain cooling in arid-zone mammals. *Comparative Biochemistry and Physiology B: Biochemistry and Molecular Biology* 131:571–585.
- Mitchell J., Mitchell G., Lust A. 2009. Thermoregulatory anatomy of pronghorn (*Antilocapra americana*). *European Journal of Wildlife Research* 55:23–31.
- Morton S. R. 1978. An ecological study of *Sminthopsis carasicaudata* (Marsupialia: Dasyuridae). III. Reproduction and life history. *Australian Wildlife Research* 5:183–211.
- Murray B. R., Dickman C. R., Watts C. H. S., Morton S. R. 1999. The dietary ecology of Australian rodents. *Wildlife Research* 26:857–858.
- Nagy K. A. 1994. Seasonal water, energy and food use by free-living, arid-habitat mammals. *Australian Journal of Zoology* 42:55–63.
- Nagy K. A., Knight M. H. 1994. Energy, water, and food use by springbok antelope (*Antidorcas marsupialis*) in the Kalahari desert. *Journal of Mammalogy* 75:860–872.
- Naya D. E., Bacigalupe L. D., Bustamante D. M., Bozinovic F. 2005. Dynamic digestive responses to increased energy demands in the leaf-eared mouse (*Phyllotis darwini*). *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 175:31–36.
- Nel J. A. J. 1984. Behavioural ecology of canids in the south-western Kalahari. *Koedoe Supplement* (1984):229–235.
- Noll-Banholzer U. 1979. Body temperature, oxygen consumption, evaporative water loss and heart rate in the fennec. *Comparative Biochemistry and Physiology Part A: Physiology* 62:585–592.
- Nowack J., Dausmann K. H., Mzilikazi N. 2013. Non-shivering thermogenesis in the African lesser bushbaby, *Galago moholi*. *Journal of Experimental Biology* 216:3811–3817.
- Osborn S. D. 1991. Adaptive heterothermy in desert mammals. [PhD thesis]. Tucson (Arizona): University of Arizona.
- Ostrowski S., Williams J. B. 2006. Heterothermy of free-living Arabian sand gazelles (*Gazella subgutturosa marica*) in a desert environment. *Journal of Experimental Biology* 209:1421–1429.
- Ostrowski S., Williams J. B., Bedin E., Ismail K. 2002. Water influx and food consumption of free-living oryxes (*Oryx leucoryx*) in the Arabian desert in summer. *Journal of Mammalogy* 83:665–673.
- Ostrowski S., Williams J. B., Ismail K. 2003. Heterothermy and the water economy of free-living Arabian oryx (*Oryx leucoryx*). *Journal of Experimental Biology* 206:1471–1478.
- Ostrowski S., Williams J. B., Mésochina P., Sauerwein H. 2006. Physiological acclimation of a desert antelope, Arabian oryx (*Oryx leucoryx*), to long-term food and water restriction. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 176:191–201.
- Ouajd S., Kamel B. 2009. Physiological particularities of dromedary (*Camelus dromedarius*) and experimental implications. *Scandinavian Journal of Laboratory Animal Science* 36:19–29.
- Oywale J., Dzenda T., Yaquib L., Akanbi D., Ayo J., Owoyele O., Minka N., Dare T. 2011. Alterations in the osmotic fragility of camel and donkey erythrocytes caused by temperature, pH and blood storage. *Veterinarski Arhiv* 81:459–470.
- Peinke D. M., Brown C. R. 1999. Osmoregulation and water balance in the springhare (*Pedetes capensis*). *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 169:1–10.

- Perrin M. R., Downs C. T. 1994. Comparative aspects of the thermal biology of the Cape spiny mouse, *Acomys subspinosus*, and the common spiny mouse, *A. spinosissimus*. *Israel Journal of Zoology* 40:151–160.
- Portas T. J., Lynch M. J., Vogelnest L. 2003. Comparison of etorphine-detomidine and medetomidine-ketamine anesthesia in captive addax (*Addax nasomaculatus*). *Journal of Zoo and Wildlife Medicine* 34:269–273.
- Purohit K. G. 1971. Absolute duration of survival of tammar wallaby (*Macropus eugenii*, Marsupialia) on sea water and dry food. *Comparative Biochemistry and Physiology Part A: Physiology* 39:473–481.
- Rainho A. 2007. Summer foraging habitats of bats in a Mediterranean region of the Iberian Peninsula. *Acta Chiropterologica* 9:171–181.
- Randall J. A. 1997. Species-specific footdrumming in kangaroo rats: *Dipodomys ingens*, *D. deserti*, *D. spectabilis*. *Animal Behaviour* 54:1167–1175.
- Randall J. A., Rogovin K., Parker P. G., Eimes J. A. 2005. Flexible social structure of a desert rodent, *Rhombomys opimus*: philopatry, kinship, and ecological constraints. *Behavioral Ecology* 16:961–973.
- Rathbun G. B., Rathbun C. D. 2005. Noki or dassie-rat (*Petromus typicus*) feeding ecology and petrophily. *Belgian Journal of Zoology* 135(Supplement): 69–75.
- Rautenstrauch K. R., Krausman P. R. 1989. Influence of water availability and rainfall on movements of desert mule deer. *Journal of Mammalogy* 70:197–201.
- Reese J. B., Haines H. 1978. Effects of dehydration on metabolic rate and fluid distribution in the jack-rabbit, *Lepus californicus*. *Physiological Zoology* 51: 155–165.
- Rickart E. A. 1986. Serum thyroxine and seasonal fat-tenting of free-living piute ground squirrels, *Spermophilus mollis* (Rodentia: Sciuridae). *Comparative Biochemistry and Physiology Part A: Physiology* 85: 199–202.
- Ron U., Haim A. 2001. How dehydration affects the thermoregulatory and osmoregulatory abilities of the golden spiny mouse *Acomys russatus*. *Israel Journal of Zoology* 47:15–20.
- Rubal A., Choshniak I., Haim A. 1992. Daily rhythms of metabolic rate and body temperature of two murids from extremely different habitats. *Chronobiology International* 9:341–349.
- Rübsamen K., Kettembeil S. 1980. Effect of water restriction on oxygen uptake, evaporative water loss and body temperature of the rock hyrax. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 138:315–320.
- Rugangazi B. M., Maloij G. M. O. 1988. Renal excretion of urea in a small east African antelope, the dik-dik (*Rhynchotragus kirkii*). *Comparative Biochemistry and Physiology Part A: Physiology* 91:437–444.
- Saadi L., Lebaili N. 2012. Effect of a water-rich diet on adrenal zone glomerulosa in *Gerbillus tarabuli*. *Comptes Rendus Biologies* 335:96–102.
- Salsamendi E., Arostegui I., Aiharta J., Almenar D., Goiti U., Garin I. 2012. Foraging ecology in Mehely's horseshoe bats: influence of habitat structure and water availability. *Acta Chiropterologica* 14:121–132.
- Sassi P. L., Borghi C. E., Bozinovic F. 2007. Spatial and seasonal plasticity in digestive morphology of cavies (*Microcavia australis*) inhabiting habitats with different plant qualities. *Journal of Mammalogy* 88:165–172.
- Sassi P. L., Caviedes-Vidal E., Anton R., Bozinovic F. 2010. Plasticity in food assimilation, retention time and coprophagy allow herbivorous cavies (*Microcavia australis*) to cope with low food quality in the Monte desert. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 155:378–382.
- Scantlebury M., Bennett N. C., Speakman J. R., Pillay N., Schradin C. 2006. Huddling in groups leads to daily energy savings in free-living African four-striped grass mice, *Rhabdomys pumilio*. *Functional Ecology* 20:166–173.
- Schmid J. 2001. Daily torpor in free-ranging gray mouse lemurs (*Microcebus murinus*) in Madagascar. *International Journal of Primatology* 22:1021–1031.
- Schmid J., Speakman J. R. 2009. Torpor and energetic consequences in free-ranging grey mouse lemurs (*Microcebus murinus*): a comparison of dry and wet forests. *Naturwissenschaften* 96:609–620.
- Schmidt-Nielsen B., Schmidt-Nielsen K. 1951. A complete account of the water metabolism in kangaroo rats and an experimental verification. *Journal of Cellular and Comparative Physiology* 38:165–181.
- Schmidt-Nielsen B., Schmidt-Nielsen K., Houpt T. R., Jarnum S. A. 1956. Water balance of the camel. *American Journal of Physiology* 185:185–194.
- Schmidt-Nielsen K., Schmidt-Nielsen B. 1952. Water metabolism of desert mammals. *Physiological Reviews* 32:135–166.
- Schmidt-Nielsen K., Schmidt-Nielsen B., Brokaw A. 1948. Urea excretion in desert rodents exposed to high protein diets. *Journal of Cellular and Comparative Physiology* 32:361–379.
- Schmidt-Nielsen K., Crawford E. C. Jr., Newsome A. E., Rawson K. S., Hammel H. T. 1967. Metabolic rate of camels: effect of body temperature and dehydration. *American Journal of Physiology* 212:341–346.
- Schmidt-Nielsen K., Hainsworth F. R., Murrish D. E. 1970. Counter-current heat exchange in the respiratory passages: effect on water and heat balance. *Respiration Physiology* 9:263–276.
- Schmidt-Nielsen K., Schroter R. C., Shkolnik A. 1981a. Desaturation of exhaled air in camels. *Proceedings of the Royal Society B: Biological Sciences* 211:305–319.

- Schmidt-Nielsen K., Crawford E. C., Hammel H. T. 1981b. Respiratory water loss in camels. *Proceedings of the Royal Society B: Biological Sciences* 211:291–303.
- Schoepf I., Schradin C. 2014. Arginine vasopressin plasma levels change seasonally in African striped mice but do not differ between alternative reproductive tactics. *General and Comparative Endocrinology* 204:43–48.
- Schradin C. 2005. Nest-site competition in two diurnal rodents from the succulent karoo of South Africa. *Journal of Mammalogy* 86:757–762.
- Schradin C. 2006. Whole-day follows of striped mice (*Rhabdomys pumilio*), a diurnal murid rodent. *Journal of Ethology* 24:37–43.
- Schradin C., Pillay N. 2005a. Demography of the striped mouse (*Rhabdomys pumilio*) in the succulent karoo. *Mammalian Biology* 70:84–92.
- Schradin C., Pillay N. 2005b. Intraspecific variation in the spatial and social organization of the African striped mouse. *Journal of Mammalogy* 86:99–107.
- Schradin C., Pillay N. 2006. Female striped mice (*Rhabdomys pumilio*) change their home ranges in response to seasonal variation in food availability. *Behavioral Ecology* 17:452–458.
- Schradin C., Krackow S., Schubert M., Keller C., Schradin B., Pillay N. 2007. Regulation of activity in desert-living striped mice: the importance of basking. *Ethology* 113:606–614.
- Schradin C., Schmohl G., Rödel H. G., Schoepf I., Treffler S. M., Brenner J., Bleeker M., Schubert M., König B., Pillay N. 2010. Female home range size is regulated by resource distribution and intraspecific competition: a long-term field study. *Animal Behaviour* 79:195–203.
- Schradin C., Lindholm A. K., Johannessen J., Schoepf I., Yuen C.-H., König B., Pillay N. 2012. Social flexibility and social evolution in mammals: a case study of the African striped mouse (*Rhabdomys pumilio*). *Molecular Ecology* 21:541–553.
- Schradin C., Raynaud J., Arrivé M., Blanc S. 2014. Leptin levels in free ranging striped mice (*Rhabdomys pumilio*) increase when food decreases: the ecological leptin hypothesis. *General and Comparative Endocrinology* 206:139–145.
- Schreiber R. K. 1979. Coefficients of digestibility and caloric diet of rodents in the northern Great Basin desert. *Journal of Mammalogy* 60:416–420.
- Schreier A. L., Swedell L. 2012. Ecology and sociality in a multilevel society: ecological determinants of spatial cohesion in Hamadryas baboons. *American Journal of Physical Anthropology* 148:580–588.
- Schubert M., Pillay N., Ribble D. O., Schradin C. 2009. The round-eared sengi and the evolution of social monogamy: factors that constrain males to live with a single female. *Ethology* 115:972–985.
- Schulz E., Fraas S., Kaier T. M., Cunningham P. L., Ismail K., Wronski T. 2013. Food preferences and tooth wear in the sand gazelle (*Gazella marica*). *Mammalian Biology* 78:55–62.
- Schumann D. M., Cooper H. M., Hofmeyr M. D., Bennett N. C. 2005. Circadian rhythm of locomotor activity in the four-striped field mouse, *Rhabdomys pumilio*: a diurnal African rodent. *Physiology and Behavior* 85:231–239.
- Selim H. M., Imai S., El Sheik A. K., Attia H., Okamoto E., Miyagawa E., Maede Y. 1999. Rumen ciliate protozoal fauna of native sheep, friesian cattle and dromedary camel in Libya. *Journal of Veterinary Medical Science* 61:303–305.
- Shanas U., Haim A. 2004. Diet salinity and vasopressin as reproduction modulators in the desert-dwelling golden spiny mouse (*Acomys russatus*). *Physiology and Behavior* 81:645–650.
- Shaw W. T. 1934. The ability of the giant kangaroo rat as a harvester and storer of seeds. *Journal of Mammalogy* 15:275–286.
- Shkolnik A., Borut A. 1969. Temperature and water relations in two species of spiny mice (*Acomys*). *Journal of Mammalogy* 50:245–255.
- Shoemaker V. H., Nagy K. A., Costa W. R. 1976. Energy utilization and temperature regulation by jackrabbits (*Lepus californicus*) in the Mojave Desert. *Physiological Zoology* 49:364–375.
- Siebert B. D., Macfarlane W. V. 1971. Water turnover and renal function of dromedaries in the desert. *Physiological Zoology* 44:225–240.
- Siebert B. D., Macfarlane W. V. 1975. Dehydration in desert cattle and camels. *Physiological Zoology* 48:36–48.
- Silanikove N., Tagari H., Shkolnik A. 1980. Gross energy digestion and urea recycling in the desert black Bedouin goat. *Comparative Biochemistry and Physiology Part A: Physiology* 67:215–218.
- Sneddon J. C., Boomker E., Howard C. V. 2006. Mucosal surface area and fermentation activity in the hind gut of hydrated and chronically dehydrated working donkeys. *Journal of Animal Science* 84:119–124.
- Soholt L. F. 1974. Environmental conditions in an artificial burrow occupied by Merriam's kangaroo rat, *Dipodomys merriami*. *Journal of Mammalogy* 55:859–864.
- Song X., Geiser F. 1997. Daily torpor and energy expenditure in *Sminthopsis macroura*: interactions between food and water availability and temperature. *Physiological Zoology* 70:331–337.
- Spalton J. A. 1999. The food supply of the Arabian oryx (*Oryx leucoryx*) in the desert of Oman. *Journal of Zoology* 248:433–441.
- Stewart D. R. M. 1963. The Arabian oryx (*Oryx leucoryx* Pallas). *African Journal of Ecology* 1:103–117.
- Taylor C. R. 1969. Metabolism, respiratory changes, and water balance of an antelope, the eland. *American Journal of Physiology* 217:317–320.

- Taylor C. R. 1970a. Dehydration and heat: effects on temperature regulation of East African ungulates. *American Journal of Physiology* 219:1136–1139.
- Taylor C. R. 1970b. Strategies of temperature regulation: effect on evaporation in East African ungulates. *American Journal of Physiology* 219:1131–1135.
- Timm R. M., Kermott L. H. 1982. Subcutaneous and cutaneous melanins in *Rhabdomys*: complementary ultraviolet radiation shields. *Journal of Mammalogy* 63:16–22.
- Tirado C., Cortés A., Bozinovic F. 2008. Water balance in two South American *Phyllotis* desert rodents, *P. xanthopygus rupestrис* and *P. darwini darwini*. *Journal of Arid Environments* 72:664–670.
- Turner J. C. 1970. Water consumption of desert bighorn sheep. *Desert Bighorn Council Transactions* 14:189–197.
- Turner J. C. 1979. Osmotic fragility of desert bighorn sheep red blood cells. *Comparative Biochemistry and Physiology Part A: Physiology* 64:167–175.
- Veloso C., Bozinovic F. 1993. Dietary and digestive constraints on basal energy metabolism in a small herbivorous rodent. *Ecology* 74:2003–2010.
- Vermeulen H. C., Nel J. A. J. 1988. The bush Karoo rat *Otomys unisulcatus* on the Cape West coast. *South African Journal of Zoology* 23:103–111.
- Walker F. M., Taylor A. C., Sunnucks P. 2007. Does soil type drive social organization in southern hairy-nosed wombats? *Molecular Ecology* 16:199–208.
- Walsberg G. E. 1988. Consequences of skin color and fur properties for solar heat gain and ultraviolet irradiance in two mammals. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 158:213–221.
- Warneke L., Turner J. M., Geiser F. 2008. Torpor and basking in a small arid zone marsupial. *Naturwissenschaften* 95:73–78.
- Waterman J. M., Herron M. D. 2004. *Xerus princeps*. *Mammalian Species* 751:1–3.
- Williams J. B., Ostrowski S., Bedin E., Ismail K. 2001. Seasonal variation in energy expenditure, water flux and food consumption of Arabian oryx *Oryx leucoryx*. *Journal of Experimental Biology* 204:2301–2311.
- Williams J. B., Lenain D., Ostrowski S., Tielman B. I., Seddon P. J. 2002. Energy expenditure and water flux of Rüppell's foxes in Saudi Arabia. *Physiological and Biochemical Zoology* 75:479–488.
- Wilson R. T. 1989. *Ecophysiology of the Camelidae and Desert Ruminants*. Berlin (Germany): Springer-Verlag.
- Winkelmann J. R., Getz L. L. 1962. Water balance in the Mongolian gerbil. *Journal of Mammalogy* 43:150–154.
- Withers P. C., Cooper C. E. 2009. Thermal, metabolic, and hygric physiology of the little red kaluta, *Dasykaluta rosamondae* (Dasyuromorphia: Dasyuridae). *Journal of Mammalogy* 90:752–760.
- Withers P. C., Lee A. K., Martin R. W. 1979. Metabolism, respiration and evaporative water loss in the Australian hopping-mouse *Notomys alexis* (Rodentia: Muridae). *Australian Journal of Zoology* 27:195–204.
- Wittenberg C., Choshniak I., Shkolnik A., Thurau K., Rosenfeld J. 1986. Effect of dehydration and rapid rehydration on renal function and on plasma renin and aldosterone levels in the black Bedouin goat. *Pflügers Archiv* 406:405–408.
- Woodroffe R. 2011. Ranging behaviour of African wild dog packs in a human-dominated landscape. *Journal of Zoology* 283:88–97.
- Yagil R. 1985. The desert camel: comparative physiological adaptation. *Comparative Animal Nutrition* 5:1–164.
- Young R. A. 1976. Fat, energy and mammalian survival. *American Zoologist* 16:699–710.
- Yousef M. K., Johnson H. D. 1975. Thyroid activity in desert rodents: a mechanism for lowered metabolic rate. *American Journal of Physiology* 229:427–431.
- Yousef M. K., Dill D. B., Mayes M. G. 1970. Shifts in body fluids during dehydration in the burro, *Equus asinus*. *Journal of Applied Physiology* 29:345–349.
- Zervanos S. M., Hadley N. F. 1973. Adaptational biology and energy relationships of the collared peccary (*Tayassu tajacu*). *Ecology* 54:759–774.
- Zhang Z.-Q., Wang D.-H. 2007. Seasonal changes in thermogenesis and body mass in wild Mongolian gerbils (*Meriones unguiculatus*). *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 148:346–353.