

# ResearchOnline@JCU


This is the **Accepted Version** of a paper published in the  
journal: *Atherosclerosis*

Moxon, Joseph V., Jones, Rhondda E., Norman, Paul E., Clancy,  
Paula, Flicker, Leon, Almeida, Osvaldo P., Hankey, Graeme J.,  
Yeap, Bu B., and Golledge, Jonathan (2016) *Plasma ferritin  
concentrations are not associated with abdominal aortic aneurysm  
diagnosis, size or growth*. *Atherosclerosis*, 251. pp. 19-24

<http://dx.doi.org/10.1016/j.atherosclerosis.2016.05.022>

© 2015. This manuscript version is made available under  
the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>



*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

**Plasma ferritin concentrations are not associated with abdominal aortic aneurysm diagnosis, size or growth**

Joseph V. Moxon,<sup>1</sup> Rhondda E. Jones,<sup>2</sup> Paul E. Norman,<sup>3</sup> Paula Clancy<sup>1</sup>, Leon Flicker,<sup>4,6,8</sup>  
Osvaldo P. Almeida,<sup>7,8,9</sup> Graeme J. Hankey,<sup>4,5</sup> Bu B. Yeap,<sup>4,10</sup> Jonathan Golledge<sup>1,11\*</sup>

1. The Queensland Research Centre for Peripheral Vascular Disease, College of Medicine and Dentistry, James Cook University, Townsville, QLD 4811, Australia.
2. The Australian Institute of Tropical Health and Medicine, James Cook University, Townsville, QLD, Australia.
3. School of Surgery, University of Western Australia, Perth, WA, Australia.
4. School of Medicine and Pharmacology, University of Western Australia, Perth, Australia.
5. Department of Neurology, Sir Charles Gardiner Hospital, Nedlands, Perth, Australia
6. Department of Geriatric Medicine, Royal Perth Hospital, Perth, Australia.
7. School of Psychiatry and Clinical Neurosciences, University of Western Australia, Perth, Australia.
8. WA Centre for Health and Ageing, Centre for Medical Research, Perth, Australia.
9. Department of Psychiatry, Royal Perth Hospital, Perth, Australia.
10. Department of Diabetes and Endocrinology, Fiona Stanley Hospital, Perth, Australia.
11. Department of Vascular and Endovascular Surgery, The Townsville Hospital, Townsville, QLD, Australia.

\* Correspondence

Phone: 0061 7 4781 4130; Fax: 0061 7 4725 8250; Email: [jonathan.golledge@jcu.edu.au](mailto:jonathan.golledge@jcu.edu.au)

## **Abstract**

**Background and aims:** Experimental studies using a rodent model have suggested that iron overload may contribute to abdominal aortic aneurysm (AAA) pathogenesis.

**Methods:** We assessed the association of total body iron, as measured by plasma ferritin, with AAA diagnosis, size and growth in 4024 community-dwelling older men screened for AAA, using logistic regression and linear mixed effects models.

**Results:** Plasma ferritin concentrations were similar in men who did (n=293) and did not (n=3731) have an AAA (median [inter-quartile range] concentrations 115.4 [63.0-203.1] and 128.5 [66.1-229.1] ng/mL respectively,  $p=0.124$ ). There was no association between plasma ferritin concentration and AAA diagnosis in unadjusted logistic regression (odds ratio (OR) for a 1 standard deviation increase: 0.880 [95%CI: 0.764-1.015];  $p=0.078$ ), or when adjusting for AAA risk factors and factors known to influence circulating ferritin (OR for a 1 standard deviation increase: 0.898 [95% CI: 0.778-1.035];  $p=0.138$ ). Iron overload prevalence (plasma ferritin concentrations >200ng/mL) was lower in men with an AAA (25.3%) than those without (30.8%;  $p=0.048$ ), but was not associated with AAA diagnosis after adjusting as above (OR: 0.781 [95% CI:0.589-1.035];  $p=0.086$ ). The association of iron overload with AAA growth was investigated in 265 men with small AAAs who received at least 1 repeat ultrasound scan in the 3 years following screening. We saw no difference in AAA growth between men who did and did not have iron overload (n=65 and 185 respectively,  $p=0.164$ ).

**Conclusions:** Our data suggest that iron overload is unlikely to be important in AAA pathogenesis.

**Short title:** Ferritin, iron overload and AAA

**Key words:** Abdominal aortic aneurysm; Ferritin; Iron overload; Biomarker

**Abbreviations:** AAA: Abdominal aortic aneurysm; IQR: inter-quartile range; 95% CI: 95% confidence interval.

## **Introduction**

Abdominal aortic aneurysm (AAA) is an abnormal progressive dilatation of the infra-renal aorta diagnosed by aortic diameter  $\geq 30$ mm (1). AAA affects ~2% of men, and 1% of women aged over 65 years and is a leading cause of death in the elderly through aortic rupture and associated cardiovascular events (1-3). Recent data suggest that global mortality rates due to AAA increased by ~10% over the last 20 years (4). No currently available medications have been shown to slow AAA progression, and surgery is the only current treatment (1, 5). Several randomised control trials have demonstrated that surgery does not improve outcomes for patients with small (<55 mm) AAAs (6-9), and patients with small AAAs are managed conservatively via repeated imaging assessments to monitor AAA growth (1, 5). Approximately 60% of small AAAs ultimately expand to a size where surgical intervention is recommended (6, 7). Thus, there remains an unmet need to identify an effective non-surgical therapy to improve management of patients with small AAAs.

A large body of data suggest that aortic inflammation, oxidative stress, vascular smooth muscle cell death and extracellular matrix degradation are important in AAA pathogenesis (1, 5, 10). Several separate studies have identified iron deposits within infra-renal aortic biopsies recovered from patients with large AAAs (11-14). More recently, Sawada and colleagues reported that an iron-restricted diet reduced the development of AAA within a mouse model and posed the hypothesis that iron overload may be important in AAA pathogenesis (13). If this hypothesis was confirmed then iron restriction could be a treatment for AAA. This hypothesis has not yet been tested in large populations (13). Plasma ferritin concentration has been demonstrated to be a robust marker to assess body iron stores in older people (reviewed by Fairweather-Tait *et al.*(15)). In the current study we investigated the association of iron overload (as assessed by plasma ferritin concentration) with AAA diagnosis, size and growth in a cohort of 4024 community-dwelling men screened for AAA.

## **Materials and methods**

### Participants and definitions used:

This study included 4024 participants of the Health In Men Study (HIMS) for whom blood samples were available (16). The cohort characteristics and protocols of the HIMS have been previously described (16). HIMS developed from a randomized trial designed to examine if screening older men for AAA reduced mortality. Clinical information collected from each HIMS participant at recruitment included age, medical history and smoking status. Anthropomorphic measurements including height, weight and hip and waist circumference were also recorded. Body mass index (BMI) was calculated as the weight in kilograms divided by the height in metres<sup>2</sup>. Participants provided written informed consent, and this study was conducted under institutional ethics approval.

AAA was defined as a maximum infra-renal aortic diameter  $\geq 30$ mm on ultrasound. Coronary heart disease was defined as prior history of myocardial infarction, angina or coronary artery revascularisation. Diabetes was defined as prior diagnosis or treatment for hyperglycaemia. Hypertension was defined by treatment for, or previous diagnosis of high blood pressure. Men found to have AAA were offered a follow up ultrasound scan at 6-monthly intervals if aortic diameter at screening was  $\geq 40$  mm, or at 12 month intervals if aortic diameter was 30-39 mm, as described previously (17). Previous data have demonstrated that ultrasound measurement of AAA diameter for these patients has excellent reproducibility (95% confidence interval  $< 3$ mm, detailed in (17)). Circulating concentrations of C-reactive protein, low- and high-density lipoprotein cholesterol, creatinine and homocysteine were measured using reproducible assays as previously described (17-19).

### Measurement of plasma ferritin:

Plasma ferritin was measured using a commercially available enzyme-linked immunosorbent assay (ELISA; #2825-300, Monobind Inc. Lake Forest, CA, USA), as directed by the manufacturer. The inter assay coefficient of variation was 12.9%; intra assay coefficient of

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

variation was 4.5%. Plasma ferritin concentrations >200 µg/L (or >200 ng/mL) were considered indicative of iron overload, in line with World Health Organisation guidelines (20).

#### Statistical analyses:

Data were analysed using the SPSS v 22, and R statistical software packages. Continuous data are reported as median and inter-quartile range (IQR). Continuous variables were compared between 2 groups using the Mann-Whitney U test, and between 3 groups using the Kruskal-Wallis test. Correlation between continuous variables was assessed using the Spearman-rho test. Nominal data are shown as count and percent, and were compared using the chi-squared test.

The association of plasma ferritin concentration with AAA diagnosis was further assessed using univariate binary logistic regression and multivariate regression adjusted for i) AAA risk factors (age, hypertension, diabetes, coronary heart disease, smoking and BMI), and ii) AAA risk factors and dietary/non dietary factors known to influence circulating ferritin (aspirin use, meat consumption and frequency of vigorous and non-vigorous exercise) (21). For the purposes of this study meat consumption was categorised as i) not eating meat on a weekly basis, ii) eating meat up to 2 times a week and, iii) eating meat at least 3 times a week. Frequency of vigorous and non-vigorous exercise was categorised as i) no hours per week, ii) fewer than 4 hours per week, and iii) at least 4 hours per week. For nominal variables, men who had the risk factor were compared to those who did not. To assess the effect of exercise and meat consumption, men who did not perform the activity were used as the reference group. Odds ratios (OR) and 95% confidence intervals are reported for all covariates in each model.

Yearly AAA growth rate for men in the growth analysis was calculated as a percentage of the size of the AAA in the preceding year (e.g. AAA growth rate between screening and year 1 rescan =  $100 \times ((\text{Year 1 AAA diameter} / \text{AAA diameter at screening}) - 1)$ ). Using this approach,

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

AAA growth rate could only be completed for men with consecutive scans. Accordingly, men were removed from the analysis at the time of their last consecutive AAA measurement.

Yearly AAA growth was used as the response variable in a linear mixed effects model to investigate the association of iron overload with AAA growth using the R nlme and car packages. A random-intercept model was fitted using Iron overload, time (years, labelled as interval in Supplementary file 2) and initial AAA diameter as fixed effects. Variation between individual patients was treated as a random effect. More complex formulations of the random effects structure to allow random slopes did not improve the model as evaluated by the AIC or BIC criteria. Similarly, incorporation of interaction terms between the fixed effects did not improve the model. The distribution of residuals was examined via QQ-normal plots and scatter plots of fitted values vs standardised residuals and did not indicate problems with residual distributions.

For all analyses,  $p$ -values  $<0.05$  were considered to be significant.

## **Results**

### *Plasma ferritin concentrations were not associated with AAA diagnosis:*

Plasma ferritin concentrations were compared between men who did and did not have an AAA (n=293 and 3731, respectively). Demographic characteristics of all participants are shown in Table 1. Men with AAA were older, and had a higher prevalence of hypertension, ever smoking and coronary heart disease. Serum C-reactive protein (hsCRP) concentration was significantly higher in men that had AAAs than those who did not (median hsCRP [IQR] 2.5 [1.4-5.1] and 1.8 [1.0-3.7] mg/L respectively,  $p < 0.001$ ). Serum hsCRP concentrations were not correlated with plasma ferritin concentrations (Spearman  $r = 0.023$ ;  $p = 0.147$ ). Median ferritin concentrations were 115.4 [IQR: 63.0-203.1] ng/mL and 128.5 [IQR: 66.1-229.1] ng/mL in men who did and did not have an AAA, respectively ( $p = 0.124$ ). No significant association of plasma ferritin with AAA diagnosis was observed in univariate binary regression (OR for an increase in plasma ferritin of 175 ng/mL, i.e. ~1 std deviation, 0.880, 95% confidence intervals [95% CI] 0.764-1.015,  $p = 0.078$ ). Similarly, no significant relationship between plasma ferritin concentrations and AAA diagnosis was observed in regression models adjusted for AAA risk factors (OR: 0.871 (95% CI: 0.774-1.026);  $p = 0.110$ ), or AAA risk factors plus dietary and non-dietary factors known to influence circulating ferritin concentration (OR: 0.898 (95% CI: 0.778-1.035);  $p = 0.138$ ; Supplementary Table 1) (21). The prevalence of iron overload (circulating ferritin concentrations  $> 200$  ng/mL) was significantly lower in men that had an AAA (25.3%) than those who did not (30.8%,  $p = 0.048$ ), however this association was not significant after adjusting for AAA risk factors and factors known to influence circulating ferritin concentrations (OR: 0.781 (95% CI: 0.589-1.035);  $p = 0.086$ ; Table 2).

### *Plasma ferritin concentrations were not correlated with aortic diameter:*

No correlation between plasma ferritin concentration and aortic diameter was seen when assessing all men included in this analysis (n=4024; Spearman  $r = -0.018$ ;  $p = 0.246$ ), those without AAA only (n=3731; Spearman  $r = -0.009$ ;  $p = 0.603$ ), or those with AAA only



*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

(Spearman  $r = -0.006$ ;  $p=0.919$ , Supplementary Table 2). Men that had an AAA were grouped into tertiles according to their infra-renal aortic diameter (Supplementary Table 3). Median plasma ferritin concentrations were 113.4 [IQR: 60.8-219.8], 106.1 [IQR: 59.9-207.1] and 121.2 [IQR: 66.3-195.5] ng/mL in the first, second and third tertiles, respectively ( $p=0.996$ ). No difference in the prevalence of iron overload was observed between the tertiles ( $p=0.811$ ). Finally, no difference in circulating ferritin concentration was observed when comparing men who had large AAAs ( $>50$  mm,  $n=18$ ), with those who did not have an AAA ( $n=3731$ ; median serum ferritin concentration 141.0 [63.5-218.3] and 128.5 [66.1-229.1] ng/mL respectively;  $p=0.911$ ).

*Iron overload was not associated with AAA growth:*

The influence of iron overload on AAA growth was assessed in a sub-group of 250 men with small AAAs (median AAA diameter 33.2 [IQR 31.2-36.5] mm) who underwent at least one additional ultrasound scan to measure AAA diameter in the 12-36 months after initial screening. Sixty five men with iron overload, and 185 men without iron overload were included in this growth analysis (Table 3). Two hundred and 13 (85.2%) men received 3 additional scans after recruitment, 20 (8.0%) were rescanned twice after recruitment, and 17 (6.8%) received 1 scan after recruitment. AAA growth rate was calculated for each year and compared between groups by linear mixed effects models. No difference in % annual increase in AAA diameter was observed between men who did or did not have iron overload (Table 3, and Supplementary File 1).

## **Discussion**

Recent observations of i) iron deposits within the AAA wall; ii) co-localisation of aortic iron deposits with regions of oxidative stress and macrophage infiltration and iii), protection against AAA formation in experimental mice maintained on a low-iron diet (12, 13) have collectively stimulated the hypothesis that iron overload may be involved in AAA pathogenesis. To test this theory we investigated the association of iron overload with AAA diagnosis, size and growth, by measuring plasma ferritin concentrations in a large group of community-dwelling men screened for AAA. Plasma ferritin concentrations were not associated with AAA diagnosis or diameter, contrary to the assessed hypothesis. Moreover AAA growth was similar in men that did or did not have iron overload. These results collectively suggest that total body iron overload is not a major driver of AAA pathogenesis. This supposition is further supported by recent data from a small cohort of European patients with AAA which demonstrated no robust association of circulating concentrations of ferritin or hepcidin (the master regulator of iron retention) with AAA diagnosis (12).

Our results, however, do not rule out a pathological role for localised iron deposition in AAA. AAA is commonly accompanied by the formation of a large non-occlusive intraluminal thrombus (ILT) within the aneurysm sac, which has been suggested to be a major source of the iron deposited within the aortic wall (1, 12, 22, 23). The ILT remains in contact with the flowing blood and is continually remodelled allowing it to increase in volume in parallel with AAA expansion (24). Eventually the ILT forms a laminated neo-tissue comprising a blood-rich layer at the bloodstream interface and an acellular fibrinolysed region in contact with the AAA wall (25, 26). ILT remodelling results in the retention of circulating proteins, leukocytes and red blood cells (12, 22, 23, 27, 28). Relevant to the current study, lysis of red blood cells trapped within the ILT releases haem iron which is conveyed to the AAA wall (12, 22). This local aortic iron deposition could contribute to reactive oxidative species generation, which has been implicated in AAA (10, 12, 13).

The observation that iron reduced feed protected experimental mice against AAA, led Sawada and colleagues to suggest that reduction of total body iron through dietary intervention may have therapeutic potential for AAA by limiting the extent of aortic iron deposition (13). The current study suggests that total body iron overload is not present in patients with AAA. Furthermore, several studies have identified iron deficiency anaemia as a significant predictor for mortality in AAA patients (12, 29-32). It is therefore unlikely that iron restriction would be beneficial (and may even be harmful) for AAA patients. Our findings also highlight the need to carefully assess the relevance of rodent model findings to human AAA. It is likely that iron deposits within AAA are a result of remodelling of the large ILT not total body iron overload. It is plausible that anti-thrombotic therapy may provide a targeted means to reduce aortic iron accumulation in AAA patients. Moreover, as the ILT is also suggested to contribute to AAA pathogenesis by i) inducing hypoxia in ILT-covered regions of the aorta (33); and ii) providing a site for leukocyte and protease accumulation adjacent to the aneurysm sac (25, 26, 28, 34-36); anti-thrombotic therapy may provide additional benefits to AAA patients beyond reducing aortic iron deposition. This theory is supported by findings of studies demonstrating that experimental animals receiving platelet inhibitors show significant reductions in AAA diameter or rupture incidence coincident with lowered matrix metallo-proteinase activity and aortic inflammation, when compared to controls (37-39). It has also been reported in one surveillance study that patients with 40-49mm AAAs had slower AAA growth when prescribed aspirin (40). However, many AAA surveillance studies have reported no association between anti-platelet medication prescription and AAA growth although a current randomised trial is examining the benefit of ticagrelor in limiting AAA growth (41).

Findings of the current study should be considered in light of its limitations. First, total body iron was estimated by measuring plasma ferritin, which is known to rise under inflammatory conditions (15). AAA is an inflammatory disease (1), however, we observed no correlation between circulating hsCRP and ferritin concentrations, and inflammation appears an unlikely

confounder for this study. Moreover, circulating ferritin concentration is a well-accepted marker of iron status since it directly correlates with total body iron (15). Thus we believed that this protein was the most appropriate biomarker to include in the current study. Other markers of iron metabolism including circulating hepcidin, iron and haemoglobin concentration have been previously assessed in smaller groups of AAA patients, and have not been shown to be robustly associated with AAA (12). Accordingly, these other markers of iron metabolism were not measured in the current study. Second, the current study assessed community-dwelling men rather than a hospital cohort, and detected AAAs were mostly small (median AAA diameter 33.7 mm). In contrast, prior studies reporting an association of iron-related proteins with AAA were conducted in patients with large ( $\geq 55$ mm) AAAs (12, 13). It is possible that ferritin concentrations may differ between patients with large AAAs and non-aneurysmal controls, although we observed no correlation between AAA diameter and plasma ferritin, no difference in circulating ferritin concentrations between men with large AAAs  $>50$ mm and non-aneurysmal controls, and no association of iron overload with AAA growth. It is also important to note that our findings are based on the analysis of a single blood sample and it is possible that assessment of repeated blood samples may have shown different results. Finally, the samples assessed in the current study were collected from men aged over 65 years, and the possibility that iron overload earlier in life or in women may promote AAA formation cannot be discounted.

### **Acknowledgements**

We thank Australian Institute of Tropical Health and Medicine for providing research infrastructure.

**Declarations of interest:** None

### **Funding**

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

This work is supported by grants from James Cook University, the Australian National Health and Medical Research Council (NHMRC 1079193, 1079369, 1022752, 1021416, 1000967), The Townsville Hospital Private Practice Trust Fund and the Queensland Government. JG holds a Practitioner Fellowship from the NHMRC, Australia and a Senior Clinical Research Fellowship from the Queensland Government.

### **Author contributions**

Participant recruitment, screening and collection of biological samples and clinical data: PEN, LF, OPA, GJH, BBY and JG. Study design: JG, PEN, JVM. Laboratory assessment of plasma ferritin: PC. Statistical analysis and interpretation of data: JVM, REJ, JG. Manuscript preparation: JVM, JG. All authors provided critical evaluation of the final manuscript and agree with its contents and publication.

### **Clinical perspective**

There is no current medical treatment to slow abdominal aortic aneurysm (AAA) growth. Recent animal data have suggested that an iron-reduced diet protects against AAA formation, suggesting that i) total body iron overload may be involved in AAA pathogenesis, and ii) reduction of iron may be a treatment for AAA. Here, we assessed whether total body iron overload as assessed by circulating ferritin concentration was associated with AAA diagnosis, size and growth in a large cohort of community-screened men. Our data demonstrate no relationship between serum ferritin and AAA and do not support iron reduction as a therapy for AAA.

1 **References**

- 2 1. Moxon JV, Parr A, Emeto TI, Walker P, Norman PE, Golledge J. Diagnosis and  
3 monitoring of abdominal aortic aneurysm: Current status and future prospects. *Curr Probl*  
4 *Cardiol.* 2010,35:512-48.
- 5 2. Svensjo S, Mani K, Bjorck M, Lundkvist J, Wanhainen A. Screening for abdominal  
6 aortic aneurysm in 65-year-old men remains cost-effective with contemporary epidemiology  
7 and management. *Eur J Vasc Endovasc Surg.* 2014,47:357-65.
- 8 3. Khosla S, Morris DR, Moxon JV, Walker PJ, Gasser T, Golledge J. Meta-analysis of  
9 peak wall stress in ruptured, symptomatic and intact abdominal aortic aneurysms. *Br J Surg.*  
10 2014,101:1350-7.
- 11 4. Sampson UK, Norman PE, Fowkes FG, Aboyans V, Yanna S, Harrell FE, Jr.,  
12 Forouzanfar MH, Naghavi M, Denenberg JO, McDermott MM, Criqui MH, Mensah GA,  
13 Ezzati M, Murray C. Global and regional burden of aortic dissection and aneurysms:  
14 mortality trends in 21 world regions, 1990 to 2010. *Global Heart.* 2014,9:171-80 e10.
- 15 5. Golledge J, Muller J, Daugherty A, Norman P. Abdominal aortic aneurysm:  
16 pathogenesis and implications for management. *Arterioscler Thromb Vasc Biol.*  
17 2006,26:2605-13.
- 18 6. UK Small Aneurysm Trial Participants. Long-term outcomes of immediate repair  
19 compared with surveillance of small abdominal aortic aneurysms. *N Engl J Med*  
20 2002,346:1445-52.
- 21 7. Lederle FA, Wilson SE, Johnson GR, Reinke DB, Littooy FN, Acher CW, Ballard DJ,  
22 Messina LM, Gordon IL, Chute EP, Krupski WC, Busuttil SJ, Barone GW, Sparks S, Graham  
23 LM, Rapp JH, Makaroun MS, Moneta GL, Cambria RA, Makhoul RG, Eton D, Ansel HJ,  
24 Freischlag JA, Bandyk D, Aneurysm Detection and Management Veterans Affairs  
25 Cooperative Study Group. Immediate repair compared with surveillance of small abdominal  
26 aortic aneurysms. *N Engl J Med.* 2002,346:1437-44.

- 27 8. Cao P, CAESAR Trial Collaborators. Comparison of surveillance vs Aortic  
28 Endografting for Small Aneurysm Repair (CAESAR) trial: study design and progress. Eur J  
29 Vasc Endovasc Surg. 2005,30:245-51.
- 30 9. Ouriel K, Clair DG, Kent KC, Zarins CK. Endovascular repair compared with  
31 surveillance for patients with small abdominal aortic aneurysms. J Vasc Surg. 2010,51:1081-  
32 7.
- 33 10. Emeto T, Moxon JV, Au M, Golledge J. Oxidative stress and abdominal aortic  
34 aneurysm: Potential treatment targets. Clin Sci. 2015,130:301-15.
- 35 11. Hunter GC, Dubick MA, Keen CL, Eskelson CD. Effects of hypertension on aortic  
36 antioxidant status in human abdominal aneurysmal and occlusive disease. Proc Soc Exp  
37 Biol Med. 1991,196:273-9.
- 38 12. Martinez-Pinna R, Lindholt JS, Madrigal-Matute J, Blanco-Colio LM, Esteban-Salan  
39 M, Torres-Fonseca MM, Lefebvre T, Delbosc S, Laustsen J, Driss F, Vega de Ceniga M,  
40 Gouya L, Weiss G, Egido J, Meilhac O, Michel JB, Martin-Ventura J. From tissue iron  
41 retention to low systemic haemoglobin levels, new pathophysiological biomarkers of human  
42 abdominal aortic aneurysm. Thromb Haemost. 2014,112:87-95.
- 43 13. Sawada H, Hao H, Naito Y, Oboshi M, Hirotani S, Mitsuno M, Miyamoto Y, Hirota S,  
44 Masuyama T. Aortic iron overload with oxidative stress and inflammation in human and  
45 murine abdominal aortic aneurysm. Arterioscler Thromb Vasc Biol. 2015,35:1507-14.
- 46 14. Koksai C, Ercan M, Bozkurt AK, Cortelekoglu T, Konukoglu D. Abdominal aortic  
47 aneurysm or aortic occlusive disease: role of trace element imbalance. Angiology.  
48 2007,58:191-5.
- 49 15. Fairweather-Tait SJ, Wawer AA, Gillings R, Jennings A, Myint PK. Iron status in the  
50 elderly. Mech Ageing Dev. 2014,136-137:22-8.
- 51 16. Norman PE, Flicker L, Almeida OP, Hankey GJ, Hyde Z, Jamrozik K. Cohort profile:  
52 The Health In Men Study (HIMS). Int J Epidemiol. 2009,38:48-52.

- 53 17. Norman P, Spencer CA, Lawrence-Brown MM, Jamrozik K. C-reactive protein levels  
54 and the expansion of screen-detected abdominal aortic aneurysms in men. *Circulation*.  
55 2004,110:862-6.
- 56 18. Golledge J, van Bockxmeer F, Jamrozik K, McCann M, Norman PE. Association  
57 between serum lipoproteins and abdominal aortic aneurysm. *Am J Cardiol*. 2010,105:1480-  
58 4.
- 59 19. Wong YY, Golledge J, Flicker L, McCaul KA, Hankey GJ, van Bockxmeer FM, Yeap  
60 BB, Norman PE. Plasma total homocysteine is associated with abdominal aortic aneurysm  
61 and aortic diameter in older men. *J Vasc Surg*. 2013,58:364-70.
- 62 20. Organisation TWH. Serum ferritin concentrations for the assessment of iron status  
63 and iron deficiency in populations. WHO/NMH/NHD/MNM/112 2011,
- 64 21. Liu JM, Hankinson SE, Stampfer MJ, Rifai N, Willett WC, Ma J. Body iron stores and  
65 their determinants in healthy postmenopausal US women. *Am J Clin Nutr*. 2003,78:1160-7.
- 66 22. Nchimi A, Defawe O, Brisbois D, Broussaud TK, Defraigne JO, Magotteaux P,  
67 Massart B, Serfaty JM, Houard X, Michel JB, Sakalihan N. MR imaging of iron  
68 phagocytosis in intraluminal thrombi of abdominal aortic aneurysms in humans.  
69 *Radiology*.2010,254:973-81.
- 70 23. Moxon JV, Padula MP, Clancy P, Emeto TI, Herbert BR, Norman PE, Golledge J.  
71 Proteomic analysis of intra-arterial thrombus secretions reveals a negative association of  
72 clusterin and thrombospondin-1 with abdominal aortic aneurysm. *Atherosclerosis*.  
73 2011,219:432-9.
- 74 24. Parr A, McCann M, Bradshaw B, Shahzad A, Buttner P, Golledge J. Thrombus  
75 volume is associated with cardiovascular events and aneurysm growth in patients who have  
76 abdominal aortic aneurysms. *J Vasc Surg*. 2011,53:28-35.
- 77 25. Fontaine V, Jacob MP, Houard X, Rossignol P, Plissonnier D, Angles-Cano E, Michel  
78 JB. Involvement of the mural thrombus as a site of protease release and activation in human  
79 aortic aneurysms. *Am J Pathol*. 2002,161:1701-10.



- 80 26. Houard X, Rouzet F, Touat Z, Phillippe M, Dominiquez M, Fontaine V, Sarda-Mantel L,  
81 Meulemans A, Le Guludec D, Meilhac O, Michel JB. Topology of the fibrinolytic system  
82 within the mural thrombus of human abdominal aortic aneurysms. J Pathol. 2007,212:20-8.
- 83 27. Houard X, Leclercq A, Fontaine V, Coutard M, Martin-Ventura JL, Ho-Tin-Noe B,  
84 Touat Z, Meilhac O, Michel JB. Retention and activation of blood-borne proteases in the  
85 arterial wall: Implications for atherothrombosis. J Am Coll Cardiol. 2006,48:A3-9.
- 86 28. Martinez-Pinna R, Madrigal-Matute J, Tarin C, Burillo E, Esteban-Salan M, Pastor-  
87 Vargas C, Lindholt JS, Lopez JA, Calvo E, de Ceniga MV, Meilhac O, Egido J, Blanco-Colio  
88 LM, Michel JB, Martin-Ventura JL. Proteomic analysis of intraluminal thrombus highlights  
89 complement activation in human abdominal aortic aneurysms. Arterioscler Thromb Vasc  
90 Biol. 2013,33:2013-20.
- 91 29. Diehm N, Benenati JF, Becker GJ, Quesada R, Tsoukas AI, Katzen BT, Kovacs M.  
92 Anemia is associated with abdominal aortic aneurysm (AAA) size and decreased long-term  
93 survival after endovascular AAA repair. J Vasc Surg. 2007,46:676-81.
- 94 30. Saratzis A, Melas N, Hunter JP, Dixon H, Nightingale P, Kiskinis D, Saratzis N, Kitas  
95 GD. Anemia is associated with mortality following endovascular repair of abdominal aortic  
96 aneurysm. Vasc Endovascular Surg. 2012,46:223-8.
- 97 31. Ohrlander T, Dencker M, Dias NV, Gottsater A, Acosta S. Cardiovascular predictors  
98 for long-term mortality after EVAR for AAA. Vasc Med. 2011,16:422-7.
- 99 32. Koga Y, Mishima Y, Hara M, Hiraki T, Ushijima K. Prognostic factors associated with  
100 mortality in patients undergoing emergency surgery for abdominal aortic aneurysms. J  
101 Anesthesia. 2011,25:666-71.
- 102 33. Vorp DA, Lee PC, Wang DH, Makaroun MS, Nemoto EM, Ogawa S, Webster MW.  
103 Association of intraluminal thrombus in abdominal aortic aneurysm with local hypoxia and  
104 wall weakening. J Vasc Surg. 2001,34:291-9.
- 105 34. Coutard M, Touat Z, Houard X, Leclercq A, Michel JB. Thrombus versus wall  
106 biological activities in experimental aortic aneurysms. J Vasc Res. 2010,47:355-66.

- 107 35. Fontaine V, Touat Z, Mtairag el M, Vranckx R, Loudec L, Houard X, Andreassian B,  
108 Sebbaq U, Palombi T, Jacob MP, Meilhac O, Michel JB. Role of leukocyte elastase in  
109 preventing cellular re-colonization of the mural thrombus. *Am J Pathol.* 2004,164:2077-87.
- 110 36. Houard X, Ollivier V, Louedec L, Michel JB, Back M. Differential inflammatory activity  
111 across human abdominal aortic aneurysms reveals neutrophil-derived leukotriene B4 as a  
112 major chemotactic factor released from the intraluminal thrombus. *FASEB J* 2009,23:1376-  
113 83.
- 114 37. Dai J, Louedec L, Philippe M, Michel JB, Houard X. Effect of blocking platelet  
115 activation with AZD6140 on development of abdominal aortic aneurysm in a rat aneurysmal  
116 model. *J Vasc Surg.* 2009,49:719-27.
- 117 38. Owens AP, 3rd, Edwards TL, Antoniak S, Geddings JE, Jahangir E, Wei WQ, Denny  
118 JC, Boulaftali Y, Bergmeier W, Daugherty A, Sampson UK, Mackman N. Platelet Inhibitors  
119 Reduce Rupture in a Mouse Model of Established Abdominal Aortic Aneurysm. *Arterioscler*  
120 *Thromb Vasc Biol.* 2015,35:2032-41.
- 121 39. Liu O, Jia L, Liu X, Wang Y, Wang X, Qin Y, Du J, Zhang H. Clopidogrel, a platelet  
122 P2Y12 receptor inhibitor, reduces vascular inflammation and angiotensin II induced-  
123 abdominal aortic aneurysm progression. *PLoS One.* 2012,7:e51707.
- 124 40. Lindholt JS, Sorensen HT, Michel JB, Thomsen HF, Henneberg EW. Low-dose  
125 aspirin may prevent growth and later surgical repair of medium-sized abdominal aortic  
126 aneurysms. *Vasc Endovascular Surg.* 2008,42:329-34.
- 127 41. Golledge J, Norman PE. Current status of medical management for abdominal aortic  
128 aneurysm. *Atherosclerosis.*  
129 2011,217:57-63.
- 130
- 131
- 132

This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in *Atherosclerosis* 11-5-2016

133 **Table 1:** Characteristics of 4024 HIMS participants included in the current study.

134

<b>Characteristic</b>	<b>Whole cohort (n=4024)</b>	<b>No AAA (n=3731)</b>	<b>AAA (n=293)</b>	<b>p value<sup>a</sup></b>
Abdominal aortic diameter (mm)	21.6 (20.2-23.8)	21.4 (20.0-23.1)	33.7 (31.5-39.3)	<0.001
<b>AAA risk factors and co-morbidities</b>				
Age	70.0 (67.0-73.0)	70.0 (67.0-73.0)	71.0 (68.0-74.0)	<0.001
Hypertension	1715 (42.6%)	1564 (41.9%)	151 (51.5%)	0.001
Diabetes	332 (8.3%)	302 (8.1%)	30 (10.2%)	0.199
Ever smoking	2597 (64.5%)	2344 (62.8%)	253 (86.3%)	<0.001
CHD	1237 (30.7%)	1102 (29.5%)	135 (46.1%)	<0.001
BMI	26.5 (24.4-28.7) <sup>b</sup>	26.5 (24.4-28.7) <sup>b</sup>	27.3 (25.1-29.5)	<0.001
<b>Circulating concentrations of</b>				
LDL-C (mmol/L)	2.8 (2.3-3.4) <sup>c</sup>	2.9 (2.3-3.4) <sup>c</sup>	2.6 (2.1-3.3)	<0.001
HDL-C (mmol/L)	1.3 (1.1-1.6)	1.4 (1.1-1.6)	1.2 (1.0-1.5)	<0.001

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

Creatinine (µmol/L)	89.0 (78.0-102.0)	88.0 (78.0-101.0)	96.0 (83.0-115.5)	<0.001
hsCRP (mg/L)	1.9 (1.0-3.8) <sup>d</sup>	1.8 (1.0-3.7) <sup>d</sup>	2.5 (1.4-5.1)	<0.001
Homocysteine (mol/L)	12.4 (10.3-15.1) <sup>d</sup>	12.3 (10.2-14.9) <sup>d</sup>	13.9 (11.4-17.5)	<0.001
Ferritin ng/mL	128.0 (65.7-227.3)	128.5 (66.1-229.1)	115.4 (63.0-203.1)	0.124
Iron overload <sup>e</sup>	1222 (30.4%)	1148 (30.8%)	74 (25.3%)	0.048

135

136 CHD: Coronary heart disease; BMI: Body mass index; LDL-C: low density lipoprotein cholesterol; HDL-C: high density lipoprotein cholesterol;

137 hsCRP: C-reactive protein.

138

139 <sup>a</sup>Comparing men with and without AAA. Nominal variables are reported as count and %, and compared using the Chi-squared test. Continuous

140 variables are reported as median and IQR, and compared via Mann-Whitney U test.; <sup>b</sup> 3 missing values; <sup>c</sup> 2 missing values; <sup>d</sup> 1 missing value; <sup>e</sup>

141 Defined as circulating ferritin concentrations >200 ng/mL.

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

142 **Table 2:** Binary logistic regression testing the association of iron overload (plasma ferritin  
 143 >200ng/mL) with AAA adjusting for AAA risk factors and factors known to influence  
 144 circulating ferritin concentration in 4024 community screened men.

<b>Covariate</b>	<b>Odds Ratio</b>	<b>95% CI</b>	<b>p value</b>
<b>Model 1: Unadjusted analysis</b>			
Iron overload	0.760	0.579-0.998	0.049
<b>Model 2: Adjusted for AAA and AAA risk factors</b>			
Iron overload	0.774	0.586-1.022	0.071
Ever smoking	3.514	2.496-4.949	<0.001
History of hypertension	1.145	0.889-1.474	0.293
History of diabetes	1.066	0.708-1.603	0.760
History of CHD	1.861	1.449-2.390	<0.001
Older age <sup>a</sup>	1.375	1.189-1.591	<0.001
Body mass index [3] <sup>b</sup>	1.291	1.144-1.456	<0.001
<b>Model 3: Adjusted for AAA risk factors and factors affecting circulating ferritin<sup>c</sup></b>			
Iron overload	0.781	0.589-1.035	0.086
Ever smoking	3.516	2.493-4.958	<0.001
History of hypertension	1.302	1.008-1.681	0.043
History of diabetes	0.984	0.651-1.486	0.939
History of CHD	2.161	1.653-2.827	<0.001
Older age <sup>a</sup>	1.360	1.172-1.579	<0.001
Body mass index [3] <sup>b</sup>	1.242	1.099-1.404	0.001
Aspirin prescription [184]	1.157	0.883-1.516	0.292

*Hours non-vigorous exercise per week*

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

0		Reference	
1-4	0.701	0.520-0.946	0.020
>4	0.684	0.506-0.925	0.014
<i>Hours vigorous exercise per week</i>			
0		Reference	
1-4	0.853	0.619-1.177	0.334
>4	0.739	0.478-1.143	0.174
<i>Weekly meat consumption [184]</i>			
None		Reference	
1-2 servings	0.984	0.285-3.403	0.980
3 or more servings	1.261	0.373-4.260	0.709

---

145

146 **Abbreviations:** 95% CI: 95% confidence interval. CHD: Coronary heart disease.

147 Square brackets indicate number of missing data points for that variable. For nominal  
148 variables, men who had the risk factor were compared to those who did not. <sup>a</sup> Refers to a 5  
149 year increase in age. <sup>b</sup> Odds ratio for an increase of ~1 std deviation in BMI (3.4).

150 <sup>c</sup> Including dietary and non-dietary factors known to influence circulating ferritin  
151 concentrations (see (21)).

This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in *Atherosclerosis* 11-5-2016

152 **Table 3:** Characteristics of 250 HIMS participants with and without iron overload in whom AAA growth was assessed.

Characteristic	No iron overload (plasma ferritin <200 ng/mL)	Iron overload (plasma ferritin >200 ng/mL)	p value
Number	185	65	---
<b>AAA risk factors and co-morbidities:</b>			
Age	71 (68-75)	71 (68-74)	0.357
Hypertension	94 (50.8%)	31 (47.7%)	0.665
Diabetes	20 (10.8%)	7 (10.8%)	0.993
Ever smoking	155 (83.8%)	60 (92.3%)	0.088
CHD	76 (41.2%)	31 (47.7%)	0.354
BMI <sup>a</sup>	27.2 (25.1-29.3)	27.2 (25.2-29.4)	0.722
AAA diameter at screening	33.2 (31.2-36.5)	33.1 (31.2-35.8)	0.446
<b>Circulating concentrations of:</b>			
LDL-C (mmol/L)	2.6 (2.0-3.3)	2.7 (2.1-3.4)	0.940

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

HDL-C (mmol/L)	1.2 (1.0-1.4)	1.3 (1.0-1.5)	0.665
Creatinine (µmol/L)	95.0 (82.5-112.0)	90.0 (78.5-117.5)	0.411
hsCRP (mg/L)	2.3 (1.2-4.8)	2.7 (1.5-6.3)	0.338
Homocysteine (mol/L)	13.6 (11.4-17.9)	14.3 (11.3-17.2)	0.745
Ferritin (ng/mL)	85.4 (47.1-134.3)	312.5 (252.9-442.9)	<0.001
	<b>Percent growth per year:<sup>b</sup></b>		0.164 <sup>c</sup>
Screening to year 1	4.3 ± 0.4[183]	3.5 ± 0.6 [65]	
Years 1 to 2 (%)	3.4± 0.4[166]	2.7 ± 0.6 [62]	
Years 2 to 3 (%)	3.1± 0.4 [153]	2.9 ± 0.6 [62]	

153 CHD: Coronary heart disease; BMI: Body mass index; LDL-C: low density lipoprotein cholesterol; HDL-C: high density lipoprotein cholesterol;  
 154 hsCRP: C-reactive protein. <sup>a</sup> Denotes 3 missing data points. <sup>b</sup> Reported as mean and standard error (unadjusted for random effects), square  
 155 brackets denote number of patients included in each group for each time point. <sup>c</sup> *p* value comparing AAA growth rate between groups over the  
 156 follow up period assessed via linear mixed effects modelling. Nominal variables are reported as count and %, and compared using the Chi-  
 157 squared test. Continuous variables are reported as median and IQR and are compared via the Mann-Whitney U test.

158



*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

**Moxon *et al*: Plasma ferritin concentrations are not associated with abdominal aortic aneurysm diagnosis, size or growth. Supplementary material.**

**Supplementary Table 1:** Binary logistic regression testing the association of circulating ferritin concentration with AAA in 4024 community screened men.

**Supplementary Table 2:** The correlation of plasma ferritin concentration with infra-renal aortic diameter and demographic and biochemical characteristics of: A) All participants of the current study (n=4,024); B) Men who did not have AAA (n=3,731); and C) Men who had an AAA (n=293).

**Supplementary Table 3:** Characteristics of 293 participants in the current study who had an AAA, stratified into tertiles according to AAA diameter.

**Supplementary File 1:** Statistical output of linear mixed effect model assessing the effect of iron overload on AAA growth.

**Supplementary Table 1:** Binary logistic regression testing the association of circulating ferritin concentration with AAA in 4024 community screened men.

Covariate	Odds Ratio	95% CI	p value
<b>Model 1: Unadjusted analysis</b>			
Plasma ferritin <sup>a</sup>	0.880	0.764-1.015	0.078
<b>Model 2: Adjusted for AAA risk factors</b>			
Plasma ferritin <sup>a</sup>	0.891	0.774-1.026	0.110
Ever smoking	3.504	2.489-4.934	<0.001
History of hypertension	1.149	0.892-1.479	0.282
History of diabetes	1.065	0.708-1.603	0.761
History of CHD	1.866	1.453-2.396	<0.001
Older age <sup>b</sup>	1.374	1.188-1.590	<0.001
Body mass index [3] <sup>a</sup>	1.291	1.145-1.456	<0.001
<b>Model 3: Adjusted for AAA risk factors and potential confounders<sup>c</sup></b>			
Plasma ferritin <sup>a</sup>	0.898	0.778-1.035	0.138
Ever smoking	3.508	2.488-4.948	<0.001
History of hypertension	1.303	1.009-1.682	0.042
History of diabetes	0.986	0.653-1.489	0.948
History of CHD	2.159	1.651-2.823	<0.001
Older age <sup>b</sup>	1.359	1.171-1.577	<0.001
Body mass index [3] <sup>a</sup>	1.242	1.099-1.403	0.001
Aspirin prescription [184]	1.165	0.889-1.526	0.267
<i>Hours non-vigorous exercise per week</i>			
0		Reference	
1-4	0.700	0.519-0.944	0.020
>4	0.683	0.505-0.923	0.013
<i>Hours vigorous exercise per week</i>			
0		Reference	
1-4	0.856	0.620-1.180	0.342
>4	0.735	0.476-1.137	0.167
<i>Weekly meat consumption [184]</i>			
None		Reference	
1-2 servings	0.999	0.289-3.456	0.999
3 or more servings	1.285	0.380-4.343	0.687

**Abbreviations:** 95% CI: 95% confidence interval. CHD: Coronary heart disease.

Square brackets indicate number of missing data points for that variable. For nominal variables, men who had the risk factor were compared to those who did not.

<sup>a</sup>Odds ratios refers to an increase of ~1 std deviation in plasma ferritin (175 ng/mL) and BMI (3.4).

<sup>b</sup>Refers to a 5 year increase in age.

<sup>c</sup>Includes dietary and non-dietary factors known to influence circulating ferritin concentrations (detailed by Liu *et al.* (2003), *Am J Clin Nutr* 78:1160-7).

This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in *Atherosclerosis* 11-5-2016

**Supplementary Table 2:** The correlation of plasma ferritin concentration with infra-renal aortic diameter and demographic and biochemical characteristics of: A) All participants of the current study (n=4,024); B) Men who did not have AAA (n=3,731); and C) Men who had an AAA (n=293).

Characteristic	Plasma ferritin (ng/mL) in:					
	A) All participating men (n=4,024)		B) Men without AAA (n=3,731)		C) Men with AAA (n=293)	
	Spearman's rho	p value	Spearman's rho	p value	Spearman's rho	p value
Infra-renal aortic diameter (mm)	-0.18	0.246	-0.009	0.603	-0.006	0.919
Age (years)	-0.051	0.001	-0.044	0.007	-0.110	0.060
BMI	0.058 [3]	<0.0001	-0.063 [3]	<0.001	0.005	0.931
Circulating LDL-C (mMol/L)	0.042 [2]	0.008	0.047 [2]	0.004	-0.058	0.325
Circulating HDL-C (mMol/L)	-0.010	0.517	-0.016	0.322	0.026	0.657
Circulating homocysteine (Mol/L)	-0.077 [1]	<0.001	-0.079 [1]	<0.001	-0.026	0.663
Circulating creatinine (µMol/L)	-0.028	0.079	-0.024	0.135	-0.018	0.764
Circulating hsCRP (mg/L)	0.023 [1]	0.147	0.022 [1]	0.184	0.074	0.206

Abbreviations: LDL-C: Low density lipoprotein cholesterol; HDL-C: High density lipoprotein cholesterol.

Numbers in square brackets denote the number of missing data-points for that characteristic.

**Supplementary Table 3:** Characteristics of 293 participants in the current study who had an AAA, stratified into tertiles according to AAA diameter.

Characteristic	Tertile 1 (n=106)	Tertile 2 (n=95)	Tertile 3 (n=92)	p value <sup>a</sup>
Abdominal aortic diameter (mm)	31.0 (30.5-31.7)	34.0 (33.2-35.2)	43.5 (40.0-47.5)	<0.001
<b>AAA risk factors and co-morbidities</b>				
Age (years)	72.0 (68.0-75.0)	71.0 (68.0-73.0)	72.0 (68.0-75.0)	0.390
Hypertension	49 (46.2%)	50 (52.6%)	52 (56.5%)	0.340
Diabetes	9 (8.5%)	11 (11.6%)	10 (10.9%)	0.749
Ever smoking	92 (8.7%)	80 (84.2%)	81 (88.0%)	0.737
CHD	48 (45.3%)	40 (42.1%)	47 (51.1%)	0.459
BMI	27.1 (25.1-29.4)	27.2 (25.4-29.4)	27.7 (24.3-29.9)	0.897
<b>Circulating concentrations of</b>				
LDL-C (mmol/L)	2.7 (2.0-3.4)	2.7 (2.2-3.3)	2.4 (1.9-3.0)	0.034
HDL-C (mmol/L)	1.3 (1.0-1.5)	1.3 (1.1-1.4)	1.2 (1.0-1.4)	0.232
Creatinine (µmol/L)	100.0 (82.0-116.5)	90.0 (80.0-104.0)	101.5 (89.3-136.8)	0.001
hsCRP (mg/L)	2.6 (1.6-4.8)	2.4 (1.2-5.0)	2.4 (1.4-5.6)	0.757
Homocysteine (mol/L)	13.9 (11.6-17.5)	13.7 (11.2-17.9)	13.9 (10.7-17.4)	0.784
Ferritin ng/mL	113.4 (60.8-219.8)	106.1 (59.9-207.1)	121.2 (66.3-195.5)	0.996
Iron overload <sup>b</sup>	28 (26.4%)	25 (26.3%)	21 (22.8%)	0.811

<sup>a</sup>Calculated using the Kruskal-Wallis test (continuous variables) or Chi-squared test (nominal variables).

<sup>b</sup>Defined as plasma ferritin concentrations >200 ng/mL.

CHD: Coronary heart disease; BMI: Body mass index; LDL-C: low density lipoprotein cholesterol; HDL-C: high density lipoprotein cholesterol; hsCRP: C-reactive protein.

This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in *Atherosclerosis* 11-5-2016

**Supplementary File 1:** Statistical output of linear mixed effect model assessing the effect of iron overload on AAA growth.

This analysis was conducted using the freely available 'car', and 'nlme' R packages (1,2).

Linear mixed effects models were constructed to assess changes in AAA diameter over time between groups of patients with/without iron overload. Individual patients were treated as random effects. Fixed effects were Iron.overload, interval and startSize (assuming an interaction between Iron.overload and interval). Variables in this analysis were:

Rate: Growth rate of AAA for that period calculated as AAA diameter at start of period/AAA diameter at start of prior period

Sequence: Arbitrary participant ID

Iron Overload: Does the participant have iron overload evidenced by plasma ferritin concentration >200 ng/mL.(coded as 0=no, 1=yes)

Interval: Time period assessed (e.g. rate0.1 = time between screening and year 1 rescan)

startSize: AAA diameter at the start of the interval

### Creating models

```
AAA.growth.lme<-lme(rate~Iron.overload*interval + startSize,
                    random=~1|Sequence, data=AAA.growth,
                    na.action='na.omit')
```

```
anova(AAA.growth.lme)
```

##	numDF	denDF	F-value	p-value
## (Intercept)	1	436	308566	<.0001
## Iron.overload	1	246	2	0.1635
## interval	2	436	3	0.0569
## startSize	1	436	3	0.0710
## Iron.overload:interval	2	436	0	0.8275

### Calculating estimates and standard errors of AAA growth per interval in patients with and without iron overload (no adjustment for random effects).

These data will be generated by fitting and summarizing linear models to assess AAA growth rates in subsets of men without iron overload (first model), and with iron overload.

#### 1 - Model for men without iron overload

```
no.iron.overload.lm<-lm(rate~interval-1, data=AAA.growth,
                        subset=Iron.overload=='0', na.action=na.exclude)
```

```
summary(no.iron.overload.lm)
```

```
##
## Call:
## lm(formula = rate ~ interval - 1, data = AAA.growth, subset =
##      "0", na.action = na.exclude)
##
## Residuals:
```

##	Min	1Q	Median	3Q	Max
----	-----	----	--------	----	-----

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*

```
## -0.14605 -0.03412 -0.00694 0.02743 0.18405
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## intervalrate0.1 1.04275    0.00371    281 <2e-16 ***
## intervalrate1.2 1.03424    0.00390    265 <2e-16 ***
## intervalrate2.3 1.03091    0.00406    254 <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.0502 on 499 degrees of freedom
## (53 observations deleted due to missingness)
## Multiple R-squared:  0.998, Adjusted R-squared:  0.998
## F-statistic: 7.12e+04 on 3 and 499 DF, p-value: <2e-16

2 - Model for men with iron overload
iron.overload.lm<-lm(rate~interval-1, data=AAA.growth,
                    subset=Iron.overload=='1', na.action=na.exclude)
summary(iron.overload.lm)
##
## Call:
## lm(formula = rate ~ interval - 1, data = AAA.growth, subset =
##      "1", na.action = na.exclude)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -0.16445 -0.02707 -0.00471  0.02435  0.11740
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## intervalrate0.1 1.03496    0.00571    181 <2e-16 ***
## intervalrate1.2 1.02707    0.00576    178 <2e-16 ***
## intervalrate2.3 1.02931    0.00576    179 <2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.0454 on 184 degrees of freedom
## (8 observations deleted due to missingness)
## Multiple R-squared:  0.998, Adjusted R-squared:  0.998
## F-statistic: 3.22e+04 on 3 and 184 DF, p-value: <2e-16

#Note, to convert coefficients to % growth, all estimates were treated as
(1-estimate)*100. Std errors were multiplied by 100.
```

## References

1. Fox J and Weisberg S (2011). An R companion to applied regression, Second Edition. Thousand Oaks, CA: Sage.
2. Pinheiro J, Bates D, DebRoy S, Sarkar D and the R Development Core Team (2013). nlme: Linear and nonlinear mixed effects models. R package version 3.1-111.

*This is the text version of 'Plasma ferritin concentrations are not associated with AAA diagnosis, size or growth' – accepted for publication in Atherosclerosis 11-5-2016*