

What predicts medical lethality of suicide attempts in Asian youths?



Carol C. Choo^{a,*}, Keith M. Harris^{b,c}, Peter K.H. Chew^a, Roger C. Ho^d

^a College of Healthcare Sciences, James Cook University, Singapore

^b School of Medicine, University of Tasmania, Australia

^c University of Queensland, Australia

^d Yong Loo Lin School of Medicine, National University of Singapore, Singapore

ARTICLE INFO

Keywords:

Youth suicides
Medical lethality
Suicide risk
Protective factors

ABSTRACT

This study explores youth suicide attempts in Singapore using multiple databases of comprehensive archival records. Three years of medical records related to suicide attempters ($N = 666$) who were admitted to the emergency department of a large teaching hospital in Singapore were subjected to retrospective analysis. Compared to other age groups, a peak in suicide attempts ($n = 207$) was observed in youths aged between 15 and 24 years old (76.3% females; 23.7% males, mean age = 19.30, $SD = 2.89$). The model using recognized risk and protective factors was significant in predicting medical lethality, and correctly classified 62.8% of high lethality cases. Only resolution of precipitant made a unique statistical significant contribution. Analysis was separately conducted for males and females. Implications of the findings are discussed.

1. Introduction

Worldwide trends reveal that suicide rates increase with age from adolescence to young adulthood (Lynch et al., 2008; World Health Organization, 2014). Suicide under 14 years of age is rare, rates begin to rise through adolescence into young adulthood (Bertolote and Fleischmann, 2002; Malone et al., 2013; Pritchard, 1995), with corresponding heightened rates of suicidal ideation and attempts (Apter et al., 2009; World Health Organization, 2013). Trends might be associated with onset of schizophrenia and depression in late adolescence (Kerfoot and Butler, 1988; Limosin et al., 2007; Osborn et al., 2008; Palmer et al., 2005), and increased alcohol consumption in youths (Beautrais, 2000; Hawton, 1998; Klerman, 1988; Nock and Kessler, 2006).

Youth suicide rates have been rising faster compared to other age groups (Wasserman et al., 2005; Mittendorfer-Rutz and Wasserman, 2004), with a peak in those between 15 and 24 years old (Blum and Nelson-Mmari, 2004; Tiller et al., 1998). In most Asian countries, suicide rates increase with age in both genders, but with a smaller male preponderance compared to their Western counterparts (Liu and Yip, 2008; Maris et al., 2000; Peng and Choo, 1990). Young females have lower rates of completed suicides than males (Bridge et al., 2006; Li et al., 2008), but higher rates of suicidal ideation and attempts (Fergusson et al., 2000; Grunbaum et al., 2000; Lee and Tsang, 2004; Sung and Kyoung, 2012). This is often attributed to males having multiple risk factors, e.g., comorbid mood and alcohol abuse disorders,

using more lethal suicide methods (Hawton, 2000; Värnik et al., 2009), in lethal suicide attempts (Brent et al., 1998; Gould et al., 1996; Shaffer and Pfeffer, 2001).

Age plays an important role in determining suicide precipitants. Youths tend to be more impulsive, making them more prone to suicidal behaviour (Daniel and Goldston, 2009). Mental illness has been associated with youth suicide (Chia, 1981; Lo, 1992), and for those without mental illness, precipitants were mainly adverse interpersonal circumstances, and stressful life events (Brent et al., 1998; Dieserud et al., 2010). However, explanations of mechanisms leading to suicide remain inadequate. Many biopsychosocial factors interact and lead to suicide attempts (Cantor, 2000; Rohaimi, 2016), suicide epidemiology needs to include an expanded interdisciplinary perspective.

Youths have unique patterns of risk and protective factors. Risk factors include psychopathology e.g., mood and conduct disorders (Lee et al., 2010) and family history of psychopathology/suicide (Bridge et al., 2006; Cash and Bridge, 2009). Psychosocial variables include social isolation, help-rejection, lack of coping skills, hopelessness, impulsivity (Berman and Jobes, 1994), alcohol/drug abuse (Lee et al., 2010; Lester, 2000; Schilling et al., 2009; Sung and Kyoung, 2012), abuse, poor socio-economic circumstances (Gould et al., 2003), lack of support (Teft, 2000), family stresses (Pfeffer, 1986), academic problems (Lee et al., 2006), and chronic illness (Beautrais, 1998; Hoberman and Garkindel, 1988; Poteet, 1987). A combination of risk factors contributes to heightened risk, e.g., stressful life events in youths with conduct disorder, personality disorder, heightened hopelessness, im-

* Corresponding author.

E-mail address: carol.choo@jcu.edu.au (C.C. Choo).




Fig. 1. Graph of suicide attempts by age groups for females and males from 2004–2006.

pulsivity and anger contribute to elevated suicide risk (Males, 2000).

In Asia, prominent risk factors for youth suicides include mental illness, interpersonal conflicts, school problems and stress with military service (Cheok et al., 2000; Chia et al., 2008; Ho and Kua, 1998; Tsoi and Kok, 1982). Recent life and relationship stressors, family problems (Ang et al., 2006; Chia, 1999; Ung, 2003), and academic stress (Loh et al., 2012) predicted youth suicides in Singapore. Suicide attempters often reacted impulsively to a stressor (Kok, 1988). Suicide protective factors in youths is comparatively less researched, these include family cohesion, religious beliefs (Kyle, 2013), positive outlook, self-efficacy in coping, feelings for family (Borowsky et al., 2001; Chan, 1995; Eisenberg and Resnick, 2006; Young et al., 2011), and parental support (Ackard et al., 2006; Borowsky et al., 2001; Brausch and Gutierrez, 2010; Eisenberg et al., 2007; Logan et al., 2011; Taliaferro and Muehlenkamp, 2013).

Most suicide studies were conducted on suicide deaths, thus a study on suicide attempts would enhance understanding of protective factors to reduce the impact of risk factors. Analysis of the prediction of medical lethality could help in identifying high lethality attempts, with the ultimate aim to prevent lethal attempts resulting in death.

This study aims to explore age trends in suicide attempts in Singapore, and analyse the prediction of medical lethality of youth suicide attempts using available risk and protective factors.

The hypotheses are

- A peak in suicide attempts will be noted in youths between 15–24

years old

- Medical lethality in youth suicide attempts will be predicted by a combination of risk and protective factors. Based on evidence in both Western and Asian studies, analysis will be conducted on the following variables, available as part of the Suicide Risk Assessment Form utilized by medical officers at the local hospital where this study took place. The risk factors include: physical illness (e.g., Chong et al., 1992), mental illness (e.g. Judd et al., 2012), alcohol/drug use (e.g., LeardMann et al., 2013), interpersonal conflict (e.g., Chen et al., 2012), protective factors include: emotional support (e.g., Takahashi, 1998), willingness to seek help (e.g., Evans et al., 2005), resolution of precipitants (e.g., Schneidman, 2001), religion (e.g., Kok and Tseng, 1992), regret of the attempt (e.g., Bhugra, 2002), and positive future planning (e.g., Williams and Pollock, 2000).

2. Method

2.1. Procedure

Ethics approval was obtained from the Domains-Specific Review Board of a local hospital and the Human Research Ethics Committee at James Cook University. The inclusion criteria were suicide attempters admitted to the emergency department from January 2004 to December 2006. This is a retrospective study of archival data extracted from the Patient Psychiatric Assessment Form (PPAF), which includes the suicide

risk assessment form. This data set is the most comprehensive data set available from the hospital, as data containing variables of interest was not available outside the stipulated period. All cases of attempted suicide were assessed by medical officers in the emergency department under the supervision of a consultant psychiatrist, and the interview took approximately 20 min. At the time of the evaluation, the medical officer made a formal psychiatric/medical diagnosis. In this sample, 94% did not have a formal diagnosis during the time of evaluation. The method of suicide attempt was overdose for 78.5% of the suicide attempts. The rest of the attempts included cutting, jumping, stabbing, and hanging.

2.2. Measures

The Suicide Risk Assessment Form is a questionnaire used in a semi-structured interview by medical officers. The content includes: demographic information, details of the attempt, mental status examination and psychiatric diagnosis. Risk and protective factors are recorded on dichotomous scales and include: history of physical and mental illness, alcohol or drug abuse, interpersonal conflict; emotional support, willingness to seek help, resolution of precipitant, religion, regret, and positive future planning. It assesses medical severity of the attempt on a 4-point scale (high, moderate, low, no actual attempt). This questionnaire was devised for the collation of information deemed important for clinical usage in suicide risk assessment and recommendation of management plan, and psychometric properties were not available.

2.3. Cases of patients

666 patients were admitted for a suicide attempt from January 2004 to December 2006. Fig. 1 indicates a marked rise for suicide attempts in those aged between 15–24 years old in both females and males, with a sharper peak in females. Those aged between 15 and 24 were subjected to further analysis. A total of 264 suicide attempters were between the ages of 15–24. Patients with missing data on key variables were removed from the data set ($n = 62$), resulting in a sample of 207 participants (76.3% females; 53.1% Chinese, 20.3% Malays, 18.4% Indians, 8.2% Others) with mean age of 19.30 ($SD = 2.89$).

3. Results

The percentage of patients with risk factors, protective factors, and medical lethality are presented in Table 1. Medical severity of the attempt were recoded: 'high' and 'moderate' were recoded into 'high', while 'low' and 'no actual attempt' recoded into 'low'.

Direct logistic regression was performed to predict medical lethality. The full model containing 10 predictors was statistically significant, $\chi^2(10, N = 207) = 23.59, p < .05$, indicating that the model was able to distinguish high and low medical lethality. The model explained between 10.8% (Cox and Snell R^2) and 14.4% (Nagelkerke R^2) of the variance in medical lethality, and correctly classified 62.8% of the high lethality cases. As shown in Table 2, only resolution of precipitant made a unique statistically significant contribution to the model, with an odds ratio of .33. Those who had resolution of precipitant were .33 times less likely to attempt medically lethal suicide attempts than those who had no resolution of precipitant, controlling for all other predictors in the model.

Two direct logistic regressions were performed, one for each gender, to predict medical lethality. For males, the full model containing nine independent variables (physical illness was not included as a predictor because 0% of males had physical illness) was not statistically significant, $\chi^2(9, N = 49) = 14.97, p > .05$, indicating that the model was unable to distinguish attempts with high and low medical lethality.

For females, the full model containing 10 predictors was statistically significant, $\chi^2(10, N = 158) = 19.60, p < .05$, indicating that the model was able to distinguish attempts with high and low medical

Table 1
Percentage of patients with risk and protective factors (yes/no), and medical lethality (high/low).

Predictors and criterion	All ($n = 207$)		Males ($n = 49$)		Females ($n = 158$)	
	Yes	No	Yes	No	Yes	No
Risk factors						
1) Physical illness	1.9	98.1	0	100	2.5	97.5
2) Alcohol/drug abuse	11.6	88.4	22.4	77.6	8.2	91.8
3) Mental illness	17.9	82.1	20.4	79.6	17.1	82.9
4) Interpersonal conflict	45.9	54.1	51.0	49.0	44.3	55.7
Protective factors						
1) Emotional support	82.1	17.9	75.5	24.5	84.2	15.8
2) Willing to seek help	69.6	30.4	75.5	24.5	67.7	32.3
3) Resolution of precipitant	52.2	47.8	40.8	59.2	55.7	44.3
4) Religion	36.2	63.8	30.6	69.4	38.0	62.0
5) Expressed regret	84.5	15.5	81.6	18.4	85.4	14.6
6) Positive future planning	82.6	17.4	77.6	22.4	84.2	15.8
Criterion						
Medical lethality [high/low]	50.2	49.8	57.1	42.9	48.1	51.9

lethality. The model explained between 11.7% (Cox and Snell R^2) and 15.6% (Nagelkerke R^2) of the variance in medical lethality, and correctly classified 63.3% of the high lethality cases. As shown in Table 3, only two of the independent variables made a unique statistically significant contribution to the model (willing to seek help and resolution of precipitant). The strongest predictor of medical lethality was willingness to seek help, with an odds ratio of 2.27. Females who were willing to seek help were 2.27 times more likely to make medically lethal suicide attempts, compared to those who were not willing to seek help, controlling for all other predictors in the model. Another predictor was resolution of precipitant with an odds ratio of .39. Females who had resolution of precipitant were .39 times less likely to attempt medically severe suicide than those who had no resolution of precipitant, controlling for all other predictors in the model.

4. Discussion

The study examined age trends and prediction of medical lethality for youth suicide attempts in Singapore. Medical records of 666 suicide attempters were analysed. The hypotheses are firstly, a peak in suicide attempts will be noted in youths between 15–24 years old, and secondly, medical lethality in youth suicide attempts will be predicted by a combination of risk and protective factors. As hypothesized, there was a peak for youths between 15 and 24 years old. As hypothesized, the model using recognized risk and protective factors was significant in predicting medical lethality, and correctly classified 62.8% of the high lethality cases. Only resolution of precipitant made a unique statistical significant contribution, indicating that youths who had resolution of precipitants were less likely to make a medically lethal attempt.

The results suggest that current practice in assessing recognized risk and protective factors seems to suffice in suicide assessment, and indicates support for intervention strategies in helping vulnerable youths to identify and resolve precipitants to their suicide attempt (Ho et al., 1999; Kerr et al., 2006; Shilubane et al., 2012, 2014a,b; Zhang et al., 2011). Future research could explore the impact of

Table 2
Logistic regression predicting medical lethality.

Predictors	B	S.E.	Wald	df	p	Odds ratio	95.0% CI for odds ratio	
							Lower	Upper
Risk factors								
1) Physical illness	-.60	1.06	.32	1	.57	.55	.07	4.37
2) Alcohol/drug abuse	.21	.48	.19	1	.67	1.23	.48	3.18
3) Mental illness	.33	.42	.64	1	.42	1.40	.62	3.17
4) Interpersonal conflict	-.00	.31	.00	1	.99	1.00	.55	1.82
Protective factors								
1) Emotional support	-.45	.42	1.16	1	.28	.64	.28	1.44
2) Willing to seek help	.52	.34	2.40	1	.12	1.68	.87	3.25
<u>3) Resolution of precipitant</u>	-1.1	.34	10.72	1	.00	.33	.17	.64
4) Religion	.46	.32	2.09	1	.15	1.59	.85	2.96
5) Expressed regret	.74	.47	2.50	1	.11	2.09	.84	5.20
6) Positive future planning	-.55	.45	1.50	1	.22	.58	.24	1.39
Constant	.19	.60	.10	1	.75	1.21		

Note. Significant predictor(s) are underlined.

Table 3
Logistic regression predicting medical lethality (females).

Predictors	B	S.E.	Wald	df	p	Odds ratio	95.0% CI for odds ratio	
							Lower	Upper
Risk factors								
1) Physical illness	-.34	1.13	.09	1	.76	.71	.08	6.47
2) Alcohol/drug abuse	-.48	.67	.50	1	.48	.62	.17	2.32
3) Mental illness	.29	.50	.33	1	.57	1.33	.50	3.54
4) Interpersonal conflict	.02	.35	.00	1	.95	1.02	.51	2.05
Protective factors								
1) Emotional support	-.76	.51	2.26	1	.13	.47	.17	1.26
2) <u>Willing to seek help</u>	.82	.39	4.43	1	.04	2.27	1.06	4.88
<u>3) Resolution of precipitant</u>	-.94	.39	5.95	1	.02	.39	.18	.83
4) Religion	.21	.36	.33	1	.57	1.23	.61	2.50
5) Expressed regret	.57	.55	1.02	1	.31	1.74	.59	5.13
6) Positive future planning	-.76	.54	1.95	1	.16	.47	.16	1.36
Constant	.61	.77	.62	1	.43	1.84		

Note. Significant predictor(s) are underlined.

precipitants on vulnerable youths and factors that could mitigate their impact.

When results were analysed separately for both genders, the model was not significant for males, possibly due to the relatively small sample size. Future studies could use a larger sample size for males or use in-depth interviews to explore male suicidality.

The model was significant in predicting medical lethality for females, and correctly classified 63.3% of high lethality cases, suggesting that current practice in assessment seems to suffice for females. The strongest predictor of medical lethality for females was willingness to seek help. This result that females willing to seek help were more likely to make medically lethal suicide attempts may seem counterintuitive, but consideration of literature on youths' help-seeking behaviour offer further insight into female suicide attempts.

Youths tend to seek help from friends/family, rather than from professionals (Boldero and Fallon, 1995; Kalafat, 1997; Kalafat and Elias, 1995; Sawyer et al., 2000). However, help-seeking from peers may not always be beneficial (Offer et al., 1991) and youths may not receive the help that they require, as peers often might be poorly equipped to provide appropriate. What some youths view as peer support may reinforce antisocial behaviour and maladaptive emotional coping (Kerr et al., 2006). Suicidal youths often form poor quality friendship (Cole et al., 1992), which might increase suicide vulnerability or modelling of maladaptive suicidal behaviours (Deane et al., 2001; Prinstein et al., 2001).

Help-seeking behaviour requires self-awareness of the problem, willingness to seek assistance and social norms that encourage help-seeking (Rickwood et al., 2007). Youths willing to seek help might not have done so prior to the suicide attempt, due to stigma. Fear of stigma is high among youths and prevalent in Asian societies (Chong et al., 2007). Possible factors preventing youths from seeking help include lack of knowledge of available services and difficulty in taking the initiative to seek help (Curtis, 2010). Such factors could be explored in future research.

Although research is needed to further explore help seeking and suicide lethality, current findings offer interesting perspectives for suicide interventions in young females. Targeting help-seeking could be effective for school-based suicide prevention programs (Aseltine et al., 2007; Carlton and Deane, 2000), where information about informal and formal sources of help is disseminated (Strunk et al., 2014). Informal sources include parents and friends (Cauce and Srebnik, 2003) who could be gatekeepers and connect youths with professional help. Peers are often the first to know of a youth's suicidality, but are often unsure about what to do (Mazza, 2006). Programs could equip gatekeepers to support suicidal youths. Training in early identification of warning signs and referral options could reduce stigma on help-seeking (Budinger et al., 2014), enhance gatekeepers' preparedness to offer help (Baber and Bean, 2009; King and Smith, 2000; Wyman et al., 2008), improve gatekeepers' ability to assist distressed youths (Frederico and Davis, 1996), and ultimately prevent

suicide (Shilubane et al., 2014a,b). This is helpful when youths are disconnected from social support (Bourke and Evans, 2000).

Limitations of the study include the reliance on self-report, and the usage of discrete items on dichotomous scales, which place constraints on the depth and types of information obtained. Future research could employ qualitative interviews to uncover the processes underlying help seeking, suicide lethality, resolution of precipitants and other factors relevant for understanding youth suicide attempts.

It is desirable to have a multi-level modelling analysis to identify different layers of factors contributing to medical lethality in youth suicide, which take into account the social cultural environment, and other contextual factors, that might impact on suicide attempters, e.g. socioeconomic status (Gould et al., 2003), and whether the youths came from dysfunctional family backgrounds (e.g., Ackard et al., 2006; Borowsky et al., 2001; Brausch and Gutierrez, 2010; Eisenberg et al., 2007; Logan et al., 2011; Taliaferro and Muehlenkamp, 2013). As local youths aged between 15 and 24 years old may be either schooling or in employment or serving mandatory National Service, these factors may affect the kind of stress experienced and in turn contribute to suicide vulnerability (Cheok et al., 2000; Chia et al., 2008; Ho and Kua, 1998; Lee et al., 2006; Tsoi and Kok, 1982). However, there are constraints as such data was not collected. Future research could gather further information and employ multi-level modelling analysis to give further understanding into youth suicides.

In conclusion, the findings have implications for informing suicide assessment and primary prevention for youth suicides. A peak was seen in suicide attempts for youths aged between 15 and 24 years old, drawing attention to the need for primary suicide prevention for at-risk youths. The parsimonious model using recognized risk and protective factors was significant in predicting medical lethality in youths, and correctly classified 62.8% of the high lethality cases, with good clinical utility, building the evidence base for brief suicide assessment. Only resolution of precipitant made a unique statistical significant contribution, indicating that youths who had resolution of precipitants were less likely to make a medically lethal attempt, suggesting that this is an important factor to work on in suicide interventions for youths. This study adds to the current literature on youth suicide, building evidence on the usage of recognized risk and protective factors in brief suicide assessment, and casting light on the relevance of resolution of precipitants in interventions for youth suicides, with the ultimate aim to reduce medical lethality and premature death of youths from suicide.

Acknowledgement

Administrative support for this work was funded by the James Cook University Internal Research Grant CRT/2016/30.

References

- Ackard, D., Neumark-Sztainer, D., Story, M., Perry, C., 2006. Parent-child connectedness and behavioral and emotional health among adolescents. *Am. J. Prev. Med.* 30 (1), 59–66.
- Ang, R., Chia, B., Fung, D., 2006. Gender differences in life stressors associated with child and adolescent suicides in Singapore from 1995 to 2003. *Int. J. Soc. Psychiatry* 52 (6), 561–570.
- Apter, A., Bursztein, C., Bertolote, J.M., Fleishmann, A., Wasserman, D., 2009. Part 12. Young people and suicide. Suicide on all continents in the young. In: Wasserman, D., Wasserman, C. (Eds.), *The Oxford Textbook of Suicidology and Suicide Prevention. A Global Perspective*. Oxford University Press, Oxford.
- Aseltine Jr., R.H., James, A., Schilling, E.A., Glanovsky, J., 2007. Evaluating the SOS suicide prevention program: a replication and extension. *BMC Public Health* 7, 161.
- Baber, K., Bean, G., 2009. Frameworks: a community-based approach to preventing youth suicide. *J. Community Psychol.* 37 (6), 684–696.
- Beautrais, A.L., 1998. Risk factors for serious suicide attempts among young people: a case control study. In: Kosky, R., Eshkeviri, H., Goldney, R., Hassan, R. (Eds.), *Suicide Prevention: The Global Context*. Plenum Press, New York, pp. 267–282.
- Beautrais, A.L., 2000. Risk factors for suicide and attempted suicide among young people. *Aust. N.Z. J. Psychiatry* 34, 420–436.
- Berman, A., Jobs, D., 1994. Treatment of suicidal adolescent. In: Leenaars, A., Maltzberger, J., Neimeyer, R. (Eds.), *Treatment of Suicidal People*. Taylor and Francis, Washington, pp. 89–100.

- Bertolote, J.M., Fleischmann, A., 2002. A global perspective in the epidemiology of suicide. *Suicidologi* 7 (2), 6–8.
- Bhugra, D., 2002. Suicidal behaviour in South Asians in the UK. *Crisis* 23 (3), 108–113.
- Blum, R.W., Nelson-Mmari, K., 2004. The health of young people in a global context. *J. Adolesc. Health* 35, 402–418.
- Boldero, J., Fallon, B., 1995. Adolescent help-seeking: what do they get help for and from whom? *J. Adolesc. Health* 18, 193–209.
- Borowsky, W., Ireland, M., Resnick, M.D., 2001. Adolescent suicide attempts: risks and protectors. *Pediatrics* 107 (3), 485–493.
- Bourke, L., Evans, P., 2000. Youth workers in Sydney: doing a lot with a little. *Youth Stud. Aust.* 19 (1), 38–43.
- Brausch, A.M., Gutierrez, P.M., 2010. Differences in non-suicidal self-injury and suicide attempts in adolescents. *J. Youth Adolesc.* 39 (3), 233–242.
- Brent, D., Moritz, G., Liotus, L., Schweers, J., Balach, L., Roth, C., Perper, J., 1998. Familial risk factors for adolescent suicide: a case control study. In: Kosky, R., Eshkeviri, H., Goldney, R., Hassan, R. (Eds.), *Suicide Prevention: The Global Context*. Plenum Press, New York, pp. 41–50.
- Bridge, J., Goldstein, T., Brent, D., 2006. Adolescent suicide and suicidal behaviour. *J. Child Psychol. Psychiatry* 47, 372–394.
- Budinger, M.C., Cwik, M.F., Riddle, M.A., 2014. Awareness, attitudes, and use of crisis hotline among youth at-risk for suicide. *Suicide Life. Threat. Behav.* 45 (2), 192–198.
- Cantor, C., 2000. Suicide in the Western world. In: Hawton, K., Van Heeringen, K. (Eds.), *The International Handbook of Suicide and Attempted Suicide*. John Wiley & Sons, West Sussex, pp. 9–28.
- Carlton, P.A., Deane, F.P., 2000. Impact of attitudes and suicidal ideation on adolescents' intentions to seek professional psychological help. *J. Adolesc.* 23 (1), 35–45.
- Cash, S.J., Bridge, J.A., 2009. Epidemiology of youth suicide and suicidal behavior. *Curr. Opin. Pediatr.* 21, 613–619.
- Cauce, A.M., Srebnik, D., 2003. Before treatment: adolescent mental health help-seeking. *Prev. Res.* 10, 6–9.
- Chan, D., 1995. Reasons for living among Chinese adolescents in Hong Kong. *Suicide Life. Threat. Behav.* 25 (3), 347–355.
- Chen, J., Choi, Y.C., Mori, K., Sawada, Y., Sugano, S., 2012. Recession, unemployment, and suicide in Japan. *Japan Labor Rev.* 9 (2), 75–92.
- Cheok, C., Ang, Y.G., Chew, W., Tan, H.Y., 2000. Adjusting to military life – servicemen with problems coping and their outcomes. *Singapore Med. J.* 41 (5), 218–220.
- Chia, B., 1981. Suicidal Behavior in Singapore. SEAMIC, Tokyo.
- Chia, B., 1999. Too Young to Die: An Asian Perspective on Youth Suicide. Times, Selangor.
- Chia, B.H., Chia, A., Tai, B.C., 2008. Suicide letters in Singapore. *Arch. Suicide Res.* 12, 74–81.
- Chong, M., Yeh, E., Wen, J., 1992. Suicidal behaviour in Taiwan: orientation to cross-society comparison. In: Kok, L., Tseng, W. (Eds.), *Suicidal Behaviour in the Asia Pacific Region*. Singapore University Press, Kent Ridge, pp. 69–82.
- Chong, S.A., Verma, S., Vaingankar, J.A., Chan, Y.H., Wong, L.Y., Heng, B.H., 2007. Perception of the public towards the mentally ill in developed Asian country. *Soc. Psychiatry Psychiatr. Epidemiol.* 42 (9), 734–739.
- Cole, E.C., Protinsky, H.O., Cross, L.H., 1992. An empirical investigation of adolescent suicidal ideation. *Adolescence* 27 (108), 813–818.
- Curtis, C., 2010. Youth perceptions of suicide and help-seeking: 'They'd think I was weak or "mental"'. *J. Youth Stud.* 13 (6), 699–715.
- Daniel, S.S., Goldston, D.B., 2009. Interventions for suicidal youth: a review of the literature and developmental considerations. *Suicide Life. Threat. Behav.* 39, 252–268.
- Deane, F.P., Wilson, C.J., Ciarrochi, J., 2001. Suicidal ideation and help-negation: not just hopelessness or prior help. *J. Clin. Psychol.* 57 (7), 1–14.
- Dieserud, G., Gerhardsen, R.M., Van den Weghe, H., Corbett, K., 2010. Adolescent suicide attempts in Bærum, Norway, 1984–2006. *Crisis* 31 (5), 255–264.
- Eisenberg, M., Ackard, D., Resnick, M., 2007. Protective factors and suicide risk in adolescents with a history of sexual abuse. *J. Pediatr.* 151, 482–487.
- Eisenberg, M., Resnick, M., 2006. Suicidality among gay, lesbian and bisexual youth: the role of protective factors. *J. Adolesc. Health* 39, 662–668.
- Evans, E., Hawton, K., Rodham, K., 2005. Suicidal phenomena and abuse in adolescents: a review of epidemiological studies. *Child Abuse Negl.* 29 (1), 45–58.
- Fergusson, D.M., Woodward, L.J., Horwood, L.J., 2000. Risk factors and life processes associated with the onset of suicidal behavior during adolescence and early adulthood. *Psychol. Med.* 30 (1), 23–39.
- Frederico, M., Davis, C., 1996. Gatekeeper training and youth suicide prevention. Report for Youth Suicide Prevention Initiative: Education and Training Consultancy. Department of Health and Family Services, Canberra, Australia.
- Gould, M., Fisher, P., Parides, M., Flory, M., Shaffer, D., 1996. Psychosocial risk factors for child and adolescent completed suicides. *Arch. Gen. Psychiatry* 53, 1155–1162.
- Gould, M.S., Greenberg, T., Velting, D.M., Shaffer, D., 2003. Youth suicide risk and preventive interventions: a review of the past 10 years. *J. Am. Acad. Child Adolesc. Psychiatry* 42, 386–405.
- Grunbaum, J.A., Lowry, R., Kann, L., Pateman, B., 2000. Prevalence of health risk behaviors among Asian American/Pacific Islander high school students. *J. Adolesc. Health* 27, 322–330.
- Hawton, K., 1998. Why has suicide increased in young males? *Crisis* 19, 119–124.
- Hawton, K., 2000. Sex and suicide: gender differences in suicidal behavior. *Br. J. Psychiatry* 177, 484–485.
- Ho, B.K.W., Kua, E.H., 1998. Parasuicide: a Singapore perspective. *Ethn. Health* 3, 255–263.
- Ho, B.K.W., Hong, C., Kua, E.H., 1999. Suicidal behavior among young people in Singapore. *Gen. Hosp. Psychiatry* 21, 128–133.
- Hoberman, H., Garkindl, B., 1988. Completed suicide in youth. *Can. J. Psychiatry* 33,

- 494–504.
- Judd, L.L., Schettler, P.J., Akiskal, H., Coryell, W., Fawcett, J., Fiedorowicz, J.G., Solomon, D.A., Keller, M.B., 2012. Prevalence and clinical significance of subsyndromal manic symptoms, including irritability and psychomotor agitation, during bipolar major depressive episodes. *J. Affect. Disord.* 138 (3), 440–448.
- Kalafat, J., 1997. Prevention of youth suicide. In: Weissberg, R.P., Gullotta, T.P. (Eds.), *Healthy Children 2010: Enhancing Children's Wellness. Issues in Children's and Families' Lives*, vol. 8. pp. 175–213.
- Kalafat, J., Elias, M.J., 1995. Suicide prevention in an educational context: broad and narrow foci. *Suicide Life. Threat. Behav.* 25 (1), 125–133.
- Kerfoot, M., Butler, A., 1988. *Problems of Childhood and Adolescence*. Macmillan, Basingstoke.
- Kerr, D.C.R., Preuss, L.J., King, C.A., 2006. Suicidal adolescents' social support from family and peers: gender-specific associations with psychopathology. *J. Abnorm. Child Psychol.* 34 (1), 103–114.
- King, K.A., Smith, J., 2000. Project SOAR: a training program to increase school counselors' knowledge and confidence regarding suicide prevention and intervention. *J. Sch. Health* 70 (10), 402–407.
- Klerman, G., 1988. The current age of youth melancholia: evidence for increase in depression in adolescents and young adults. *Br. J. Psychiatry* 152, 4–14.
- Kok, L., 1988. *Attempted Suicide in Singapore from Drug Overdose*. (Unpublished MD thesis) National University of Singapore, Singapore.
- Kok, L., Tseng, W., 1992. Orientation to cross-society comparison. In: Kok, L., Tseng, W. (Eds.), *Suicidal Behaviour in the Asia Pacific Region*. Singapore University Press, Kent Ridge, pp. 1–14.
- Kyle, J., 2013. Spirituality: its role as a mediating protective factor in youth at risk for suicide. *J. Spiritual. Mental Health* 15 (1), 47–67.
- LeardMann, C.A., Pwll, T.M., Smith, T.C., Bell, M.R., Smith, B., Boyko, E.J., ... Hoge, C.W., 2013. Risk factors associated with suicide in current and former US military personnel. *JAMA* 310 (5), 496–506.
- Lee, A., Tsang, C.K., 2004. Youth risk behavior in a Chinese population: a territory-wide youth risk behavioural surveillance in Hong Kong. *Public Health* 118 (2), 88–95.
- Lee, M., Wong, B., Chow, B., McBride-Chang, C., 2006. Predictors of suicide ideation and depression in Hong Kong adolescents: perceptions of academic and family climates. *Suicide Life. Threat. Behav.* 36, 82–96.
- Lee, S.Y., Hong, J.S., Espelage, D.L., 2010. An ecological understanding of youth suicide in South Korea. *Sch. Psychol. Int.* 31, 531–546.
- Lester, D., 2000. Substance abuse is a factor in many suicides. In: Roleff, T. (Ed.), *Teen Suicide*. Greenhaven, California, pp. 57–62.
- Li, X.Y., Phillips, M.R., Zhang, Y.P., Xu, D., Yang, G.H., 2008. Risk factors for suicide in China's youth: a case-control study. *Psychol. Med.* 38, 397–406.
- Limosin, F., Loze, J., Philippe, A., Casadebaig, F., Rouillon, F., 2007. Ten-year prospective follow-up study of the mortality by suicide in schizophrenic patients. *J. Schizophr. Res.* 94 (1), 23–28.
- Liu, K., Yip, P., 2008. Mainland China. In: Yip, P. (Ed.), *Suicide in Asia: Causes and Prevention*. Hong Kong University Press, pp. 31–48.
- Lo, W., 1992. Suicidal behaviour in Hong Kong. In: Kok, L., Tseng, W. (Eds.), *Suicidal Behaviour in the Asia Pacific Region*. Singapore University Press, Kent Ridge, pp. 83–111.
- Logan, J., Hall, J., Karch, D., 2011. Suicide categories by patterns of known risk factors a latent class analysis. *Arch. Gen. Psychiatry* 68, 935–941.
- Loh, C., Tai, B., Ng, W., Chia, A., Chia, B., 2012. Suicide in young Singaporeans aged 10–24 years between 2000 to 2004. *Arch. Suicide Res.* 16 (2), 174–182.
- Lynch, M., Howard, P., El-Mallakh, E., Matthews, J., 2008. Assessment and management of hospitalized suicidal patients. *J. Psychosoc. Nurs. Ment. Health Serv.* 46 (7), 45–52.
- Males, M., 2000. The extent of teen suicide is exaggerated. In: Roleff, T. (Ed.), *Teen Suicide*. Greenhaven, California, pp. 70–87.
- Malone, K., Quinlivan, K., Grant, T., Kelleher, C., 2013. Ageing towards 21 as a risk factor for Young Adult Suicide in the UK and Ireland: 'many young men of twenty said goodbye'. *Epidemiol. Psychiatr. Sci.* 22 (3), 263–267.
- Maris, R., Berman, A., Silverman, M., 2000. *Comprehensive Textbook of Suicidology*. Guilford Press, New York.
- Mazza, J.J., 2006. Youth suicidal behavior: a crisis in need of attention. In: Villarruel, F.A., Luster, T. (Eds.), *Adolescent Mental Health*. Greenwood Publishing Group, pp. 156–177.
- Mittendorfer-Rutz, E.M., Wasserman, D., 2004. Trends in adolescent suicide mortality in the WHO European Region. *Eur. Child Adolesc. Psychiatry* 13, 321–331.
- Nock, M.K., Kessler, R.C., 2006. Prevalence of and risk factors for suicide attempts versus suicide gestures: analysis of the National Comorbidity Survey. *J. Abnorm. Psychol.* 115, 616–623.
- Offer, D., Howard, K.I., Schonert, K.A., Ostrov, E., 1991. To whom do adolescents turn for help? Differences between disturbed and nondisturbed adolescents. *J. Am. Acad. Child Adolesc.* 30, 623–630.
- Osborn, D., Levy, G., Nazareth, I., King, M., 2008. Suicide and severe mental illnesses. Cohort study within the UK general practice research database. *Schizophr. Res.* 99 (1), 134–138.
- Palmer, B.A., Pankratz, V.S., Bostwick, J.M., 2005. The lifetime risk of suicide in schizophrenia: a reexamination. *Arch. Gen. Psychiatry* 62 (3), 247–253.
- Peng, K., Choo, A., 1990. Suicide and parasuicide in Singapore. *Med. Sci. Law* 30 (3), 225–233.
- Pfeffer, C., 1986. *The Suicidal Child*. Guilford, New York.
- Poteet, D., 1987. Adolescent suicides: a review of 87 cases of completed suicide in Shelby County, Tennessee. *Am. J. Forensic Med. Pathol.* 8, 12–17.
- Prinstein, M.J., Boergers, J., Spirito, A., 2001. Adolescents' and their friends' health-risk behavior: factors that alter or add to peer influence. *J. Pediatr. Psychol.* 26 (5), 287–298.
- Pritchard, C., 1995. *Suicide – The Ultimate Rejection? A Psycho-social Study*. Open University Press, Buckingham.
- Rohaimi, N.A., 2016. Suicide risks among adolescents: a look behind the black curtain. *Int. J. Pharm. Sci. Res.* 7 (6), 2266.
- Rickwood, D.J., Deane, F.P., Wilson, C.J., 2007. When and how do young people seek professional help for mental health problems? *Med. J. Aust.* 187 (7 Suppl), S35–S39.
- Sawyer, M.G., Arney, F.M., Baghurst, P.A., Clark, J.J., Graetz, B.W., Kosky, R.J., ... Zubrick, S.R., 2000. *The Mental Health of Young People in Australia*. Publications Production Unit, Commonwealth Department of Health and Aged Care, Canberra, Australia.
- Schilling, E.A., Aseltine Jr., R.H., Glanovsky, J., James, A., Jacobs, D., 2009. Adolescent alcohol use, suicidal ideation, and suicidal attempts. *J. Adolesc. Health* 44, 335–341.
- Schneidman, E., 2001. *Understanding Suicide*. American Psychological Association, Washington.
- Shaffer, D., Pfeffer, C., 2001. Practice parameter for the assessment and treatment of children and adolescents with suicidal behavior. *J. Am. Acad. Child Adolesc.* 40 (7 Suppl), 24S–51S.
- Shilubane, H.N., Ruiter, R.A., Bos, A.E., Reddy, P.S., van den Borne, B., 2014a. High school students' knowledge and experience with a peer who committed or attempted suicide: a focus group study. *BMC Public Health* 18 (14), 1081.
- Shilubane, H.N., Ruiter, R.A.C., Bos, A.E.R., van den Borne, B., James, S., 2014b. Psychosocial correlates of suicide ideation in South African adolescents. *Child Psychiatry Hum. Dev.* 45 (2), 153–162.
- Shilubane, H.N., Ruiter, R.A.C., van den Borne, B., Sewpaul, R., James, S., Reddy, P.S., 2012. Suicide and related health risk behaviours among school learners in South Africa: results from the 2002 and 2008 national youth risk behavior surveys. *BMC Public Health* 13, 926.
- Strunk, C.M., Sorter, M.T., Ossege, J., King, K.A., 2014. Emotionally troubled teens' help-seeking behaviors: an evaluation of surviving the Teens' suicide prevention and depression awareness program. *J. Sch. Nurs.* 30 (5), 366–375.
- Sung, S.C., Kyoung, H.J., 2012. Risk factors related to suicidal ideation and attempted suicide: comparative study of Korean and American youth. *J. Sch. Nurs.* 28 (6), 448–458.
- Takahashi, Y., 1998. Suicide in Japan: what are the problems? In: Kosky, R., Eshkevari, H., Goldney, R., Hassan, R. (Eds.), *Suicide Prevention: The Global Context*. Plenum Press, New York, pp. 121–130.
- Taliaferro, L.A., Muehlenkamp, J., 2013. Risk and protective factors that distinguish adolescents who attempt suicide from those who only consider suicide in the past year. *Suicide Life. Threat. Behav.* 44 (1), 6–22.
- Teft, L., 2000. Many factors put teens at risk of suicide. In: Roleff, T. (Ed.), *Teen Suicide*. Greenhaven, California, pp. 66–69.
- Tiller, J., Krupinski, J., Burrows, G., Mackenzie, A., Hallenstein, H., Johnstone, G., 1998. Youth suicide: the Victorian coroner's study. In: Kosky, R., Eshkevari, H., Goldney, R., Hassan, R. (Eds.), *Suicide Prevention: The Global Context*. Plenum Press, New York, pp. 87–91.
- Tsoi, W.F., Kok, L.P., 1982. Suicidal behavior in Singapore for the year 1980. *Singapore Med. J.* 23, 299–305.
- Ung, E.K., 2003. Youth suicide and parasuicide in Singapore. *Ann. Acad. Med. Singapore* 32 (1), 8–12.
- Värnik, A., Kõlves, K., Allik, J., Arensman, E., Aromaa, E., van Audenhove, C., ... Hegerl, U., 2009. Gender issues in suicide rates, trends and methods among youths aged 15–24 in 15 European countries. *J. Affect. Disord.* 113 (3), 216–226.
- Wasserman, D., Cheng, Q., Jiang, G.X., 2005. Global suicide rates among young people aged 15–19. *World Psychiatry* 4, 114–120.
- Williams, J.M., Pollock, L.R., 2000. The psychology of suicidal behaviour. In: Hawton, K., Van Heeringen, K. (Eds.), *The International Handbook of Suicide and Attempted Suicide*. John Wiley & Sons, West Sussex, pp. 79–94.
- World Health Organization, 2013. *Mental Health Action Plan 2013–2020*. World Health Organization, Geneva.
- World Health Organization, 2014. *Preventing Suicide: A Global Imperative*. World Health Organization, Geneva.
- Wyman, P.A., Brown, C.H., Inman, J., Cross, W., Schmeelk-Cone, K., Guo, J., Pena, J.B., 2008. Randomized trial of a gatekeeper program for suicide prevention: 1-year impact on secondary school staff. *J. Consult. Clin. Psychol.* 76 (1), 1054–1115.
- Young, R., Sweeting, H., Ellaway, A., 2011. Do schools differ in suicide risk? The influence of school and neighbourhood on attempted suicide, suicidal ideation and self-harm among secondary school pupils. *BMC Public Health* 11, 874–889.
- Zhang, J., Li, N., Tu, X., Xiao, S., Jia, C., 2011. Risk factors for rural young suicide in China: a case-control study. *J. Affect. Disord.* 129, 244–251.