

ResearchOnline@JCU


This is the **Accepted Version** of a paper published in the
journal: Musculoskeletal Science and Practice

Land, Helen, Watt, Kerriane, and Gordon, Susan (2017) *Reply to the letter to the editor regarding 'Clinical assessment of subacromial shoulder impingement – which factors differ from the asymptomatic population?'*. Musculoskeletal Science and Practice, 29. e14-e15.

[10.1016/j.msksp.2017.04.001](https://doi.org/10.1016/j.msksp.2017.04.001)

© 2016. This manuscript version is made available under
the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>



Reply to Musculoskeletal Science and Practice Feedback

Thank you for your comments regarding the recent publication 'Clinical Assessment of Subacromial Shoulder Impingement – Which Factors Differ from the Asymptomatic Population?' Not surprisingly, impingement is a term which does not reflect the underlying cause of all shoulder pain. Hence there is healthy debate, especially amongst physical therapists, regarding alternate terminology (Braman, Zhao, Lawrence, Harrison, & Ludewig, 2013; J. S. Lewis, 2011; McFarland et al., 2013). However, it continues to be a term used throughout the medical literature and in an attempt to embrace this wider audience, until there is agreement about terminology, it was chosen for use in this paper.

The paper did not represent itself as investigating the acromial irritation theory (J. S. Lewis, 2011), as you have suggested, but investigated factors purported in the literature which potentially alter the size of the subacromial space in those with shoulder pain (Land & Gordon, 2017). With current research suggesting rotator cuff tendinopathy is a major factor in the source of pain in this subacromial area (Jeremy Lewis, 2016; Jeremy. Lewis & Ginn, 2015; J. S. Lewis, 2010) and tendinopathy related research suggesting this may be related to a compressive load (Cook & Purdam, 2012) we felt it relevant to explore this relationship. No comment was able to be made about whether compressive loading or irritation of the tendon or both was the cause of the pain in these participants as this was not the aim of the paper.

In preparation for this study, previous literature on clinical tests used to assess external factors in those with subacromial shoulder impingement (SSI) was reviewed (Land & Gordon, 2017). The few included studies that compared symptomatic and asymptomatic subjects did not definitely show which clinical assessments are able to detect a difference in each of these factors with many having methodological limitations with respect to recruitment of subjects, matching of subjects for dominance and comparison of values calculated from both shoulders within each group prior to comparison between groups (Land & Gordon, 2017).

The systematic review 'Is thoracic spine posture associated with shoulder pain, range of motion and function?' cited in the letter included research about non-specific shoulder pain and thoracic spine posture (Barrett, O'Keeffe, O'Sullivan, Lewis, & McCreesh, 2016). It reported that thoracic kyphosis did not appear to be a major contributor to non-specific shoulder pain, based on studies across various populations with the majority of included studies having moderate to high levels of bias (Barrett et al., 2016). The authors concluded as with other systematic reviews that 'higher quality research is warranted' (Barrett et al., 2016; Haik, Albuquerque-Sendin, Moreira, Pires, & Carmargo, 2016; Trampas & Kitsios, 2006). Other references cited in the letter relate specifically to the athletic throwing population (Bach & Goldberg, 2006; Dashottar & Borstad, 2012; Gates, Gupta, McGarry, Tibone, & Lee, 2012; Manske & Ellenbecker, 2013) and cadaver studies ((Bach & Goldberg, 2006; Dashottar & Borstad, 2012; Gates et al., 2012) neither of which should be directly compared to the current study.

People who were regularly participating in strength training were excluded from our study, as it is common to include 'strength training' within a prescribed exercise program for this condition and we wanted to understand the presentation prior to intervention (Carmargo et al., 2015; Holmgren, Bjornsson Hallgren, Oberg, Adolfsson, & Johansson, 2012; Tate, McClure, Young, Salvatori, & Michener, 2010).

Accepted outcome measures used in shoulder intervention studies are predominantly changes in pain scores (VAS, NRS) (Jensen, Karoly, & Braver, 1986) and changes in a functional score (SPADI, DASH) (Hudak, Amadio, & Bombardier, 1996; Roach, Budiman-Mak, Songsiridej, & Lertratanakul,

1991). Changes in these measures could be attributed to a biopsychosocial effect of the intervention experience (Woolf, 2010). The measures provided in this recent publication have provided objective comparison of symptomatic and asymptomatic groups which provides new information to inform clinical management.

It was timely and relevant therefore to study a clearly defined homogenous patient group, which commonly presents to general clinical practice, using well-designed research methodology. We look forward to sharing the outcome of intervention studies in this defined shoulder pain population in the future.

- Bach, H. G., & Goldberg, B. A. (2006). Posterior capsular contracture of the shoulder. *Journal of the American Academy of Orthopaedic Surgeons*, 14(5), 265 -277.
- Barrett, E., O'Keeffe, M., O'Sullivan, K., Lewis, J., & McCreesh, K. (2016). Is thoracic spine posture associated with shoulder pain, range of motion and function? A systematic review. *Manual Therapy*, 26, 38 - 46. doi: 10.1016/j.math.2016.07.008
- Braman, J., P., Zhao, K. D., Lawrence, R. L., Harrison, A. K., & Ludewig, P. M. (2013). Shoulder impingement revisited: evolution of diagnostic understanding in orthopedic surgery and physical therapy. *Medical & Biological Engineering & Computing*. doi: 10.1007/s11517-013-1074-1
- Carmargo, P. R., Alburquerque-Sendin, F., Avila, M. A., Haik, M. N., Vieira, A., & Salvini, T. F. (2015). Effects of Stretching and Strengthening Exercises, With and Without Manual Therapy, on Scapular Kinematics, Function, and Pain in Individuals With Shoulder Impingement: A Randomized Controlled Trial. *Journal of Orthopaedic & Sports Physical Therapy*, 45(12), 984 - 997. doi: 10.2519/jospt.2015.5939
- Cook, J. L., & Purdam, C. (2012). Is compressive load a factor in the development of tendinopathy? *British Journal of Sports Medicine*, 46, 163 - 168. doi: 10.1136/bjsports-2011-090414
- Dashottar, A., & Borstad, J. (2012). Posterior glenohumeral joint capsule contracture. *Shoulder & Elbow*, 4(4), 230 - 236. doi: 10.1111/j.1758-5740.2012.00180
- Gates, J. J., Gupta, A., McGarry, M. H., Tibone, J. E., & Lee, T. Q. (2012). The Effect of Glenohumeral Internal Rotation Deficit Due to Posterior Capsular Contracture on Passive Glenohumeral Joint Motion. *The American Journal of Sports Medicine*, 40(12), 2794 - 2800. doi: 10.1177/0363546512462012
- Haik, M. N., Alburquerque-Sendin, F., Moreira, R. F. C., Pires, E. D., & Carmargo, P. R. (2016). Effectiveness of physical therapy treatment of clearly defined subacromial pain: a systematic review of randomised controlled trials. *British Journal of Sports Medicine*, 0, 1 - 14. doi: 10.1136/bjsports-2015-095771
- Holmgren, T., Bjornsson Hallgren, H., Oberg, B., Adolfsson, L., & Johansson, K. (2012). Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: Randomized Control Study. *British Medical Journal*(BMJ 2012;344:e787), 9. doi: 10.1136/bmj.e787
- Hudak, P. L., Amadio, P. C., & Bombardier, C. (1996). Development of an Upper Extremity Outcome Measure: The DASH (Disabilities of the Arm, Shoulder, and Hand) *American Journal of Industrial Medicine*, 29, 602 - 608.
- Jensen, M. P., Karoly, P., & Braver, S. (1986). The measurement of clinical pain intensity: a comparison of six methods. *Pain*, 27, 117-126.
- Land, H., & Gordon, S. (2017). Clinical Assessment of Factors Associated with Subacromial Shoulder Impingement: A Systematic Review. *Physical Therapy Reviews*.

- Lewis, J. (2016). Rotator cuff related shoulder pain: Assessment, management and uncertainties. *Manual Therapy, 23*, 57 - 68. doi: 10.1016/j.math.2016.03.009
- Lewis, J., & Ginn, K. (2015). Rotator cuff tendinopathy and subacromial pain syndrome. In G. Jull, A. Moore, D. Falla, J. Lewis, C. McCarthy, & M. Sterlin (Eds.), *Grieve's Modern Musculoskeletal Physiotherapy* (Fourth ed., pp. 563-568): Elsevier.
- Lewis, J. S. (2010). Rotator Cuff Tendinopathy: A model for the continuum of pathology and related management. *British Journal of Sports Medicine, 44*, 918-923.
- Lewis, J. S. (2011). Subacromial impingement syndrome: a musculoskeletal condition or a clinical illusion? *Physical Therapy Reviews, 16*(5), 388 - 398. doi: 10.1179/1743288X11Y.0000000027
- Manske, R. C., & Ellenbecker, T. S. (2013). CURRENT CONCEPTS IN SHOULDER EXAMINATION OF THE OVERHEAD ATHLETE. *International Journal of Sports Physical Therapy, 8*(5), 554 - 578.
- McFarland, E. G., Maffulli, N., Del Buono, A., Murrell, G. A. C., Garzon-Muvdi, J., & Petersen, S. A. (2013). Impingement is not impingement: the case for calling it "rotator cuff disease". *Muscles, Ligaments and Tendons Journal, 3*(3), 196 - 200.
- Roach, K. E., Budiman-Mak, E., Songsiridej, N., & Lertratanakul, Y. (1991). Development of a Shoulder Pain and Disability Index. *Arthritis Care and Research, 4*(4), 143 - 149.
- Tate, A. R., McClure, P. W., Young, I. A., Salvatori, R., & Michener, L. A. (2010). Comprehensive Impairment-Based Exercise and Manual Therapy Intervention for Patients With Subacromial Impingement Syndrome: A Case Series. *Journal of Orthopaedic & Sports Physical Therapy, 40*(8), 474 - 493.
- Trampas, A., & Kitsios, A. (2006). Exercise and manual therapy for the treatment of impingement syndrome of the shoulder: A systematic review. *Physical Therapy Reviews, 11*, 125-142.
- Woolf, C. J. (2010). What is this thing called pain? *The Journal of Clinical Investigation, 120*(11), 3742 - 3744. doi: 10.1172/JCI45178