

GSA Annual Meeting in Seattle, Washington, USA - 2017

Paper No. 32-1
Presentation Time: 1:35 PM

COSMOGENIC ¹⁰BE AND ²⁶AL STUDIES OF THE RISING STAR SITE, CRADLE OF HUMANKIND, SOUTH AFRICA: MYSTERY OF THE TRUE DENUDATION RATES

MAKHUBELA, Tebogo V.¹, KRAMERS, J.D.¹, SCHERLER, D.², WITTMANN, H.³, DIRKS, P.H.G.M.⁴ and WINKLER, S.R.⁵, (1)Department of Geology, University of Johannesburg, Auckland Park, Johannesburg, 2006, South Africa, (2)Helmholtz Centre Potsdam, GFZ German Research Center for Geosciences, Potsdam, 14473, Germany; Department of Geological Sciences, Freie University of Berlin, Berlin, 12249, Germany, (3)Helmholtz Centre Potsdam, GFZ German Research Center for Geosciences, Potsdam, 14473, Germany, (4)School of Geosciences, University of the Witwatersrand, Private Bag 3, Johannesburg, 2050, South Africa; Geosciences, College of Science and Engineering, James Cook University, Townsville, 4811, Australia, (5)Accelerator Mass Spectrometry, iThemba LABS, Johannesburg, 2193, South Africa, vmakhubela@uj.ac.za

Assessing the history and rate of landscape change of the Cradle of Humankind (CoH), UNESCO World Heritage Site, South Africa, is important for better understanding of paleo-ecosystems and hominin evolution in southern Africa, as well as fossil deposition and preservation in caves [1]. In the CoH, terrestrial cosmogenic nuclides have previously been used for this [1,2,3], as well as for burial dating of the fossil-bearing cave deposits [4,5,6]. We present new *in situ* cosmogenic ¹⁰Be and ²⁶Al concentrations ([¹⁰Be] and [²⁶Al], respectively) in quartz and chert from soils, chert float and chert horizons in outcropping dolomite, on the peneplain surface above the Rising Star cave system.

Soil samples yield the highest [¹⁰Be] followed by clast samples, with bedrock chert samples yielding the lowest (Table 1). The average apparent erosion rates for the soil and clast samples are similar within 1σ: 3.58 ± 1.91 m/Ma and 4.29 ± 0.69 m/Ma, respectively. The average apparent erosion rate for the bedrock chert is up to three times faster than for soils and clasts (11.28 ± 1.09 m/Ma). All soil samples have ²⁶Al/¹⁰Be ratios significantly lower than the 6.75 surface steady state ratio.

Based on ¹⁰Be denudation rates previously found (3.6 m/Ma, [2]; 3.44 m/Ma, [1]), the landscape across the CoH is considered old and eroding slowly. High erosion rates similar to our results (5.13 - 15.02 m/Ma) for chert bedrock are ascribed to fast river incision or a recent partial collapse event [1,2]. In contrast, we think our high outcrop erosion rates reflect true denudation and the low apparent values from soil samples indicate long retention of quartz on surface, while dolomite is largely removed in solution. The quartz then experiences periods of burial and reworking in caves and river terraces, resulting in low ²⁶Al/¹⁰Be ratios.

Samples	[¹⁰ Be]	[²⁶ Al]
Soils	2.28 – 3.91	13.13 – 19.58
Clasts	1.02 – 2.43	7.79 – 14.04
Bedrock chert	0.457 – 0.904	3.79 – 6.66

- [1] Dirks et al., 2016. J. Hum. Evol., 96, 19-34.
- [2] Dirks et al., 2010. Science, 328, 205-208.
- [3] Dirks and Berger, 2013. J. Afr. Earth Sci. 78, 109-131.
- [4] Partridge et al., 2003. Science, 300, 607-612.
- [5] Gibbon et al., 2014. Quat Geochron., 24, 10-15.
- [6] Granger et al., 2015. Nature, 522, 85-88.

Session No. 32

[D3. The Geochemistry, Biogeochemistry, and Ecology of Cave and Karst Settings](#)
Sunday, 22 October 2017: 1:30 PM-5:30 PM

Room 612 (Washington State Convention Center)

Geological Society of America Abstracts with Programs. Vol. 49, No. 6
doi: 10.1130/abs/2017AM-297563

reproduce and distribute it freely, for noncommercial purposes. Permission is hereby granted to any individual scientist to download a single copy of this electronic file and reproduce up to 20 paper copies for noncommercial purposes advancing science and education, including classroom use, providing all reproductions include the complete content shown here, including the author information. All other forms of reproduction and/or transmittal are prohibited without written permission from GSA Copyright Permissions.

[Back to: D3. The Geochemistry, Biogeochemistry, and Ecology of Cave and Karst Settings](#)

[Previous Abstract](#) | [Next Abstract >>](#)
