

Hybridization between damselfishes *Dascyllus aruanus* and *D. reticulatus* on the Great Barrier Reef




Fig. 1 Anterior and lateral view of *Dascyllus aruanus* (a), the hybrid (b) and *D. reticulatus* (c). Color features that separate this hybrid from both *D. aruanus* and *D. reticulatus* are marked in red

the two lineages distributed evenly within the hybrid genotype. While Pomacentridae hybridization is rarely observed, this record emphasizes the potential for natural hybridization in areas of high abundance and well within geographic distribution limits.

References

- Montanari SR, Hobbs JP, Pratchett MS, van Herwerden L (2016) The importance of ecological and behavioural data in studies of hybridisation among marine fishes. *Reviews in Fish Biology and Fisheries* 26:181–198
- Yaakub SM, Bellwood DR, van Herwerden L, Walsh FM (2006) Hybridization in coral reef fishes: introgression and bi-directional gene exchange in *Thalassoma* (family Labridae). *Mol Phylogenet Evol* 40:84–100

J. L. Johansen (✉)
 Marine Science Institute, University of Texas at Austin, Port Aransas, TX 78373, USA
 e-mail: jacob.johansen@my.jcu.edu.au

S. He · M. K. Pappas · M. L. Berumen
 King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia

G. Frank
 James Cook University, Townsville, QLD 4811, Australia

A. S. Hoey
 ARC Centre of Excellence for Coral Reef Studies, Townsville, QLD 4811, Australia

Received: 13 August 2016 / Accepted: 27 February 2017 / Published online: 6 March 2017
 © Springer-Verlag Berlin Heidelberg 2017

Coral Reefs (2017) 36:717
 DOI 10.1007/s00338-017-1563-z