



Anomalies in T Cell Function Are Associated With Individuals at Risk of *Mycobacterium abscessus* Complex Infection

Viviana P. Lutzky^{1†}, Champa N. Ratnatunga^{1,2,3†}, Daniel J. Smith^{1,4}, Andreas Kupz², Denise L. Doolan^{1,2}, David W. Reid^{1,4}, Rachel M. Thomson^{3,4,5}, Scott C. Bell^{1,3,4†} and John J. Miles^{1,2,3,6*†}

¹QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia, ²Centre for Biodiscovery and Molecular Development of Therapeutics, Centre for Biosecurity and Tropical Infectious Diseases, AIITHM, James Cook University, Cairns, QLD, Australia, ³Faculty of Medicine, University of Queensland, Brisbane, QLD, Australia, ⁴Department of Thoracic Medicine, The Prince Charles Hospital, Brisbane, QLD, Australia, ⁵Gallipoli Medical Research Institute, Brisbane, QLD, Australia, ⁶Institute of Infection and Immunity, Cardiff University School of Medicine, Cardiff, United Kingdom

OPEN ACCESS

Edited by:

Vladimir Brusic,
The University of Nottingham
Ningbo China, China

Reviewed by:

António Gil Castro,
University of Minho, Portugal
Derin Keskin,
Dana-Farber Cancer Institute,
United States

*Correspondence:

John J. Miles
john.miles@jcu.edu.au

[†]These authors have contributed
equally to this work.

Specialty section:

This article was submitted
to T Cell Biology,
a section of the journal
Frontiers in Immunology

Received: 23 November 2017

Accepted: 28 May 2018

Published: 11 June 2018

Citation:

Lutzky VP, Ratnatunga CN, Smith DJ,
Kupz A, Doolan DL, Reid DW,
Thomson RM, Bell SC and Miles JJ
(2018) Anomalies in T Cell Function
Are Associated With Individuals
at Risk of *Mycobacterium*
abscessus Complex Infection.
Front. Immunol. 9:1319.
doi: 10.3389/fimmu.2018.01319

The increasing global incidence and prevalence of non-tuberculous mycobacteria (NTM) infection is of growing concern. New evidence of person-to-person transmission of multidrug-resistant NTM adds to the global concern. The reason why certain individuals are at risk of NTM infections is unknown. Using high definition flow cytometry, we studied the immune profiles of two groups that are at risk of *Mycobacterium abscessus* complex infection and matched controls. The first group was cystic fibrosis (CF) patients and the second group was elderly individuals. CF individuals with active *M. abscessus* complex infection or a history of *M. abscessus* complex infection exhibited a unique surface T cell phenotype with a marked global deficiency in TNF α production during mitogen stimulation. Importantly, immune-based signatures were identified that appeared to predict at baseline the subset of CF individuals who were at risk of *M. abscessus* complex infection. In contrast, elderly individuals with *M. abscessus* complex infection exhibited a separate T cell phenotype underlined by the presence of exhaustion markers and dysregulation in type 1 cytokine release during mitogen stimulation. Collectively, these data suggest an association between T cell signatures and individuals at risk of *M. abscessus* complex infection, however, validation of these immune anomalies as robust biomarkers will require analysis on larger patient cohorts.

Keywords: non-tuberculous mycobacteria, cystic fibrosis, immunoprofiling, pulmonary non-tuberculous mycobacteria infection, T cells

INTRODUCTION

Pulmonary infection caused by non-tuberculous mycobacteria (NTM) is an emerging threat with serious public health consequences. Mortality rates of 10–40% due to lung disease caused by these lesser known “cousins” of *Mycobacterium tuberculosis* (TB) have been increasingly reported in the developed world (1–4). The prolonged treatment regimens lasting months to years and increasing antibiotic resistance to front-line antibiotics make these pathogens difficult and expensive infections to treat. Over 180 species of NTM are known to cause disease in humans of which the *Mycobacterium*

avium complex (MAC) and the *Mycobacterium abscessus* complex (MABS) are of dominant clinical interest (5). These species account for over 80% of NTM disease worldwide and are among the most common causative agents for NTM lung disease (6). The global increase in disease prevalence over the past 10–15 years has led to an increased focus on patient-oriented research (7, 8).

The emergence and spread of human transmissible clones of MABS has been recently reported (9) and is the first evidence of person-to-person transmission of NTM that were, up until to now, considered environmentally acquired by susceptible individuals. MABS infection is associated with rapid decline in lung function and extensive lung damage which can be life threatening, particularly in patients already compromised with respiratory problems such as those with cystic fibrosis (CF). Multi-drug resistance (MDR) of these pathogens contributes to prolonged and difficult treatment regimens and high relapse rates, both of which lead to increased morbidity/mortality and escalating treatment costs in a group of patients who are already highly susceptible to opportunistic infections. The presence of MABS is an absolute contradiction to lung transplantation (10–13).

Non-tuberculous mycobacteria infections are also a growing health concern among the elderly population. Pre-existing lung diseases, such as chronic obstructive pulmonary disease and bronchiectasis are known risk factors for developing NTM infection as are lung malignancies, immune modulatory treatments, and HIV/AIDS (7, 14). The worldwide increase in NTM infections in apparently immunocompetent middle aged to elderly patients, in the setting of an aging population contributes to an increased population of susceptible individuals at-risk of developing NTM infection.

Delineating immune function in NTM infection is of fundamental interest in order to understand how and why these infections: (i) occur in specific at-risk populations; (ii) progress in some patients and; (iii) resolve in others. The importance of Th1-type cell-mediated immunity in anti-mycobacterial immunity is well established. Low production of the Th1 cytokines IFN γ and TNF α and more recently, low production of IL-17 and IL-10 have been associated with NTM infection (15–21).

Cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), programmed cell death protein 1 (PD-1), and T-cell immunoglobulin domain and mucin domain 3 (TIM-3) are negative regulatory check points that are important for T cell tolerance and regulation during the immune response. Widely known for their use as targets in cancer immunotherapy (22), these immune checkpoints have also been shown to play an important role in T cell exhaustion during chronic infections such as TB (23–27). The role of these molecules in NTM infection has yet to be explored. Information on T cell “quality” in terms of cytokine production is also lacking. Polyfunctional T cells are known to determine pathogenesis and disease progression of TB and other infectious and immune-related diseases (26, 28–30).

In order to discover new targets for therapeutic intervention and rational vaccine design, an improved understanding of the molecular and cellular host defense mechanisms that provide protective immunity toward NTM is required. The present study comprehensively characterized the immune profile of NTM patients by performing high-dimensional flow cytometry-based

analysis in two cohorts of NTM patients. The first group was CF patients and the second group was immunocompetent middle aged to elderly patients with MABS infection. We show across both groups abnormalities in global T cell function that associate with individuals at risk of infection.

PATIENTS AND METHODS

Patient Cohorts

Two patient cohorts were studied. The CF patient cohort ($n = 24$) included three groups of patients; (i) CF patients with active pulmonary MABS infection (defined as MABS identified by at least one positive culture within a 12-month period) $n = 6$ (CF^{Act}) and one with MAC infection; (ii) patients who had a previous diagnosis of MABS infection who had undergone treatment and were now in remission as defined by at least six clear sputum samples (smear and culture negative over a 1-year period; CF^{Past} $n = 8$) and; (iii) patients who were confirmed as having chronic *Pseudomonas aeruginosa* (Pa) infection by the Leeds criteria (31), but with no history of or current NTM infection were included as a “within-disease” control group (CF^{Controls} $n = 9$). Five of the six patients with active MABS infection had chronic Pa infection and one had intermittently positive sputum cultures ($\leq 50\%$ of serial sputum cultures positive) for Pa. Sex, age, and demographic-matched healthy controls were recruited from unrelated adult volunteers (HC^A $n = 10$). The second patient cohort was elderly patients diagnosed with active NTM (NTM^{Act} $n = 10$) all of whom had confirmed MABS infection. Healthy controls (HC^B) sex, age, and demographic-matched to the NTM^{Act} patients were recruited from healthy adult volunteers ($n = 10$). All samples were obtained with written consent and all protocols were approved by the Human Research Ethics Committees of the QIMR Berghofer Medical Research Institute, The Prince Charles Hospital, and Greenslopes Private Hospital, Australia.

Blood Samples and Processing

Peripheral blood mononuclear cells (PBMC) were separated from venous blood by Ficoll–Paque PLUS (GE Health) density gradient method and were cryopreserved in R10 medium (RPMI-1640 containing 10% fetal bovine serum (FBS) with 100 U/ml penicillin and 100 μ g/ml streptomycin) supplemented with 10% DMSO (Sigma-Aldrich). Thawed cells were rested overnight in R10 medium at 37°C. Cells were then stained for viability, counted, and aliquoted into three 96-well plate. Cells in plate one were resuspended in staining buffer (PBS with 2% FBS) and stained for direct *ex vivo* flow cytometric analysis. Cells in plate two and three were split into two aliquots and one aliquot was activated with PMA ionomycin (PMA/I) (Ebioscience) at 1 \times final concentration in the presence of Brefeldin A 1 μ g/ml (Ebioscience) and Monensin 0.1 μ g/ μ l (Ebioscience) for 6 h at 37°C. The other aliquot was incubated in R10 without PMA/I as an *ex vivo* baseline control.

Flow Cytometric Analysis

The cells in plate one were stained for surface markers delineating major immune cell lineages and evaluated for TIM-3 expression.

Panel one included surface markers α CD4-FITC (BD), α CD8-Percp-Cy5.5 (Biolegend), α CD16-PECY7 (BD), α CD19-BV421 (BD), α CD14-APC (BD), and α TIM-3-PE (R&D Systems). Cells in plate two were stained with panel 2 which included surface markers α CD4-FITC (BD), α CD8-Percp Cy5.5 (Biolegend), activation marker α CD25-PE (BD), and exhaustion marker α PD-1-BV605 (BD). Staining for intracellular exhaustion marker α CLTA-4-BV421 (BD) and nuclear transcription factor α FOXP3-APC was performed after fixation and permeabilization on ice with FOXP3 Permeabilization kit (Ebioscience) according to manufacturers' instructions. Plate three cells were activated with PMA/I and incubated with α CD107a-FITC (BD) during activation. Cells were then washed and resuspended in staining buffer and surface staining was performed with panel 3; α CD3-PECY7 (BD), α CD4-BV711 (BD), and α CD8-Percp-Cy5.5 (BD) markers. Intracellular cytokines were stained with α INF γ -AlexaFlour700 (BD), α TNF α -APC (BD), and α IL-2-PE (BD) after fixation and permeabilization with Fix/Perm buffer kit (BD) for intracellular staining. Stained samples were run on a BD LSR Fortessa 4 laser cytometer (BD). Sample acquisition was performed on BD FACSDiva 8.0 (BD) and data were analyzed with FlowJo v10 (TreeStar) and Cytobank¹ for viSNE analysis (32).

Statistical Analysis

Statistical analysis was performed with SPSS 22, Graphpad PRISM (v6.05), and Gmine² (33). Comparison of means over multiple groups was performed using one-way ANOVA tests with Tukeys *post-hoc* comparisons, while comparisons between two groups were performed using unpaired *t*-tests or Wilcoxon rank test. Hierarchical clustering and biomarker identification was performed in Gmine. Stepwise regression was performed to identify variables associated with NTM disease. The patient in CF^{Act} group with active MAC infection was excluded from these analyses. However, the patients' data point was included in scatter plots as a blue circle to visually demonstrate how active MAC infection compares to active MABS infection. Polyfunctionality analysis was performed using Pestle and SPICE V5³ (34).

RESULTS

Distinct T Cell Phenotype in CF Patients Susceptible to MABS Infection

We first investigated the phenotypic and functional immune profiles of PBMCs in CF patients to probe for functional deficiencies that could underlie predisposition to NTM infection. Cohorts were categorized as CF with active NTM infection (CF^{Act}), CF with a past history of NTM infection (CF^{Past}), CF with chronic Pa infection (and no history of NTM infection CF^{Control}) and healthy controls (HC^A), all matched in both age (ANOVA $P = 0.350$) and gender distribution (Chi-square $P = 0.445$). Demographic and clinical characteristics of patient groups are shown in Table 1. All patients had either active or past MABS

TABLE 1 | Table shows demographic of cystic fibrosis (CF) cohorts and elderly patient group.

	CF ^{Act}	CF ^{Past}	CF ^{Controls}	NTM ^{Act}
Mean age (SD)	32.6 (13.6)	34.9 (10.4)	33.1 (8.0)	75.6 (9.23)
Male:female	06:01	06:02	06:03	03:07
Non-tuberculous mycobacteria (NTM) infection at time of sample	MABS (6) MAC (1) ^a	None	None	MABS (10)
History of NTM infection				
MABS	0	8	0	0
MAC	3 ^b	0	0	3 ^c
Other infections				
<i>Pseudomonas aeruginosa</i>	5	8	9	4
<i>Aspergillus</i> spp	0	0	0	1
<i>Burkholderia</i> spp ^d	0	1	1	0
Lung function				
>70% FEV1	4	1	3	8
30–70% FEV1	3	6	5	2
<30% FEV1	0	1	1	0
Radiographic features				
Bronchiectasis	7	8	9	10

CF^{Act}-CF patients with active NTM infection. CF^{Past}-CF patients with past NTM infection. Both groups of patients had a history of *P. aeruginosa* infection. CF^{Controls}-control CF patients with chronic *P. aeruginosa* infection. NTM^{Act}-elderly patients with active NTM infection. N/A, not applicable; MABS, *Mycobacterium abscessus* complex; MAC, *Mycobacterium avium* complex.

^aOne patient in CF^{Act} group had active MAC infection. This patient was excluded from ANOVA and biomarker analysis but included in general profiling analysis. Data point is shown as a blue circle in scatter plots for ANOVA analysis in Figures 1A and 2A.

^bOne patient in CF^{Act} group had prior history of MAC infection while two subsequently developed MAC infection.

^cIn NTM^{Act} elderly patient group, one patient had a history of MAC infection prior to MABS infection while two others developed MAC infection after treatment of current episode of MABS infection.

^d*Burkholderia cepacia* complex.

infection with the exception of one patient who had active MAC infection.


Flow cytometric analysis of PBMCs revealed no significant differences in the percentage of B cells (CD19⁺), total CD4⁺ T cells (CD3⁺CD4⁺), total CD8⁺ T cells (CD3⁺CD8⁺), or in TIM-3 expression levels on cell subsets between the CF patients and healthy controls (data not shown). Comparison of Tregs (CD4⁺ CD25⁺ FOXP3⁺) between cohorts showed higher percentages in CF^{Act} and CF^{Past} groups compared to the CF^{Control} group (Figure 1A). Analysis of the individual expression of activation marker CD25 and exhaustion marker CTLA-4 on CD4⁺ T cells revealed significantly higher expression of CD25 and CTLA-4 in CF^{Act} group compared to the CF^{Control} group (Figure 1A). Higher co-expression of CD25 and CTLA-4 was seen on CD4⁺ T cells in CF^{Act} group compared to the CF^{Control} group and higher percentages of CD25 and CTLA-4 double-negative CD4⁺ T cells were observed in the CF^{Control} group and HC^A group (Figure 1A).

We next compared the overall pattern of immune marker expression of the CD4⁺ and CD8⁺ T cell compartments in terms of surface "phenotypic fingerprint." Analysis of triple, double, single, or nil expression of markers CD25, CTLA-4, and PD-1 on CD4⁺ T cells revealed a common fingerprint in CF^{Act} and CF^{Past} groups which was distinct from the CF^{Control} group (Figures 1B,C). A higher number of CLTA-4 single-positive cells were seen in patients with either active or past NTM infection as compared

¹www.Cytobank.org.

²http://cgenome.net/GMine/.

³https://exon.niaid.nih.gov/spice/.



with the healthy control subjects (Figures 1B,C). The T cell fingerprint following PMA/I activation was not significantly different to *ex vivo* T cells (Figures 1B,C). The T cell fingerprint on CD8⁺ T cells was similar between different groups (Figure S1A in Supplementary Material) with significance found in CD8⁺ CD25 single positive T cells *ex vivo* in the CF^{Act} group and in the PMA/I-activated CF^{Past} group (Figure S1A in Supplementary Material). These data reveal a difference in systemic T cell phenotypes in CF patients with active or past NTM disease, particularly in CD4⁺ T cells, compared to CF patients with more common chronic Pa infection. There was no difference between CF patients with active or past NTM disease and healthy controls in terms of T cell fingerprint.

Distinct T Cell Function in CF Patients Susceptible to MABS Infection

Given the differences in surface T cell phenotypes between cohorts we next analyzed cytokine production post-mitogen stimulation. T cell cytokine production after PMA/I stimulation revealed a specific signature associated with NTM disease. TNF α -producing CD4⁺ T cells were significantly lower in both CF^{Act} and CF^{Past} groups compared to the CF^{Control} group (Figure 2A). TNF α production in CD4⁺ T cells was also markedly lower in both CF^{Act} and CF^{Past} compared to HC^A group, though this difference did not reach statistical significance (Bonferroni *post-hoc* test). IFN γ ⁺ CD4⁺ T cells in the CF^{Act} group were higher than the CF^{Control} group (Figure 2A). In the CD8⁺ subset, IFN γ ⁺ T cells from the CF^{Act} and CF^{Past} groups were higher than the CF^{Control} group (Figure 2A).

Polyfunctionality in CD4⁺ and CD8⁺ T cells was next examined. As CD107a represents degranulation and cytolytic activity, the expression of this marker was included in the polyfunctionality profile in addition to TNF α , IFN γ , and IL-2. Both CF^{Act} and CF^{Past} groups showed a unique polyfunctionality profile compared to both CF^{Controls} and HC^A groups (Figure 2B). Both CD4⁺ TNF α -producing single positive T cells (mono-functional) and CD4⁺ TNF α ⁺ IL2⁺ double-positive cells (dual-functional) were seen to be significantly reduced in both CF^{Act} and CF^{Past} groups compared to both control groups (Figures 2B,C). Significantly higher IFN γ mono-functional T cells were seen in the CF^{Past} group compared to the CF^{Control} group though there was no difference compared to the HC^A group. There was also no difference in the number of triple- and quadruple-functional T cells between groups. When CD8⁺ T cell polyfunctionality was compared, significantly higher numbers of IFN γ producing mono-functional cells were seen in both CF^{Act} and CF^{Past} groups compared to both control groups though there was no significant difference in terms of overall polyfunctionality profile (Figure S2 in Supplementary Material). Total TNF α -producing CD8⁺ T cells were significantly higher in the HC^A and CF^{Control} groups compared to CF^{Act} and CF^{Past} groups.

Hierarchical clustering analysis of cytokine production and CD107a expression data showed a grouping of CF^{Act} and CF^{Past} groups, while the CF^{Control} and HC^A groups clustered together (Figure 3A). Based on global cytokine and CD107a expression profiles, patients with NTM disease (either past or present) could be grouped together. There was no clear separation of the active and past NTM infection groups indicating that based on all clustering variables; no global differences were seen between


these two groups. The same pattern was observed in the chronic Pa infection group CF^{Control} and the HC^A group where both groups clustered together.


Given this common hierarchical clustering result between CF^{Act} and CF^{Past} cohorts, we next redefined the cohorts for subsequent data analysis. Patients who had either active or past NTM infection (CF^{Act} and CF^{Past}) were defined as the “NTM disease” cohort and CF patients with chronic Pa infection and healthy controls (CF^{Control} and HC^A) were defined as the “control” cohort (i.e., persons with no history of NTM infection). These two variables were then used as outcomes to analyze data for predictive biomarkers using GMine multivariate analysis software (33). CCA analysis showed significant clustering between these groups (Figure 3B). Biomarker analysis identified 13 significant predictors of “NTM disease” after correction for multiple comparisons (FDR) (Figure 4A). Unsurprisingly, significant predictors included combinations of TNF α and IFN γ production. The stepwise regression model (area under the curve, AUC 100%) identified percentage CD8⁺ IL2⁺ TNF α ⁻ T cells and percentage CD8⁺ IFN γ ⁺ TNF α ⁻ T cells as the best predictors of NTM disease (Figure 4B).

Distinct T Cell Function in Elderly Patients With Active MABS Infection

To determine whether this immune profile would also be found in other independent disease cohorts, we next investigated elderly patients with active NTM infection (NTM^{Act}). In Australia, the rate of notified NTM cases per 100,000 population has increased by approximately 17% per year between 2012 and 2015 (35). The reasons for this increase are unknown, and there is yet no predictor to identify at-risk individuals. To determine if underlying immune dysfunction may be a predictive factor for NTM infection we compared NTM^{Act} patients with elderly healthy controls (HC^B). We found that Tregs were increased in the peripheral blood of elderly NTM^{Act} patients compared to elderly HC^B (Figure 5A). Elevated CD25 and CTLA-4 expression was also seen on CD4⁺ T cells in the NTM^{Act} group alongside increased PD-1 expression on CD4⁺ CD25⁺ T cells. Analysis of the phenotypic fingerprint of CD25, CTLA-4, and PD-1 expression in CD4⁺ T cells showed a specific signature in NTM^{Act} patients with elevated CD25 and CTLA-4 double-positive T cells as well as elevated CD25 single-positive T cells in the NTM^{Act} group (Figures 5A,B). T cells negative for all three markers (triple-negative) were significantly higher in the HC^B group compared to NTM^{Act} patients. Four significant phenotypic differences *ex vivo* and during mitogen stimulation were observed between the NTM^{Act} group and HC^B group (Figure 5C). *Ex vivo* CD8⁺ T cell fingerprint was identical between HC^B and NTM^{Act} cohorts (data not shown).

Mitogen stimulation and polyfunctionality analysis in the elderly cohorts revealed a very different profile to that seen in CF patients. In the NTM^{Act} group, significantly higher TNF α production by CD8⁺ T cells was observed (Figure 6A), while there was no difference in the number of TNF α -producing CD4⁺ T cells between groups (Figure S3A in Supplementary Material). IFN γ single-positive CD8⁺ T cells were similar in patients and controls (Figures 6B,C) as was the overall polyfunctionality profile in CD4⁺ T cells (Figures S3A,B in Supplementary Material).








FIGURE 4 | Top immune biomarker candidates for diagnosis of non-tuberculous mycobacteria (NTM) infection status in cystic fibrosis cohorts. **(A)** Flow cytometric biomarkers for NTM disease state from cytokine secretion profile data (Wilcoxon rank test). Graph shows log odds ratio for each biomarker candidate for diagnosis of NTM disease state (negative versus exposed state) with 95% CI. (P, probability; FDR, false discover rate; AUC, area under the curve). **(B)** Stepwise regression model (forward selection) for NTM infection status. Fitted model contains 18 variables shown in table with a final AIC of -30 and an AUC 1. Plots show contribution of each variable to model and fitted value for each patient/control (green dots) when model is applied showing an AUC = 1.

DISCUSSION

The increased incidence and prevalence of NTM disease in recent years, warrants a more comprehensive understanding of immunity in susceptible individuals (7). Due to increasing antibiotic resistance of NTM strains and poor patient-outcomes in particular MABS, immunomodulatory strategies may emerge as important adjuvants to conventional anti-mycobacterial therapy in NTM disease. Here, we dissect faults in immunity in two cohorts of patients (CF and elderly) with NTM lung infection and matched controls to identify the underlying blood immune signatures of each disease. Blood was examined given: (i) T cells are known to traffic between the blood and lung (36); (ii) NTM-specific T cells have been observed in the

blood (37); and (iii) blood is easy to access for diagnostic tests. The first cohort included CF patients with active NTM disease, CF patients with past NTM disease who had been successfully treated and were now in disease remission, CF patients with chronic Pa infection who had no history of NTM infection and a group of matched healthy controls. The second cohort included elderly patients with NTM infection and matched healthy controls. All control individuals in both cohorts had active or past MABS infection with the exception of one patient who had MAC infection.

The frequency of Tregs was significantly increased in CF patients with active and past NTM infection compared to CF patients with chronic Pa infection. These data contrast with the elderly cohort, where Tregs significantly were higher in elderly




FIGURE 5 | Evidence of CD4⁺ T cell activation and exhaustion in immunocompetent individuals with active non-tuberculous mycobacteria (NTM) infection.

(A) Flow cytometric analysis of *ex vivo* CD4⁺ T cells showed significantly more Tregs in NTM^{Act} group compared to HC^B group ($P = 0.028$). Significantly more CD25⁺ CTLA4⁺ and CD25⁺ CTLA4⁻ CD4⁺ T cells were seen in NTM^{Act} group ($P = 0.002$ and $P = 0.009$, respectively) compared to HC^B. A reciprocal increase in CD25⁻ CTLA4⁻ CD4⁺ T cells was seen in the HC^B group ($P = 0.001$) compared to disease group. Higher numbers of PD1⁺ CD4⁺ T cells ($P = 0.021$) and CD25⁺ PD1⁺ CD4⁺ T cells ($P = 0.011$) were seen in the NTM^{Act} group. CD25⁺ PD1⁻ T cells were significantly higher in NTM^{Act} group ($P < 0.001$) and CD25⁻ and PD1⁻ CD4⁺ T cells were significantly higher in HC^B group ($P < 0.001$). **(B)** Phenotyping and polyfunctionality profiling of CD4⁺ T cells by SPICE showed differences in NTM^{Act} and HC^B groups directly *ex vivo* and post PMA/I stimulation. Phenotype profiles were significantly different between groups both *ex vivo* ($P = 0.0013$) and post stimulation ($P = 0.022$). Significantly more CD25⁺ CTLA4⁺ T cells were observed in NTM^{Act} groups compared to HC^B. **(C)** SPICE dot plots show phenotype profile of all combinations of markers CD25, CTLA4, and PD1 in CD4⁺ T cells both *ex vivo* and post PMA/I stimulation between NTM^{Act} and HC^B groups. Significantly different expression compared to HC^B (Wilcoxon rank test $P < 0.05$) is indicated with # symbol.

NTM patients compared to healthy controls. This was similarly observed by Hector et al., where lower number of Tregs was found in both the airways and peripheral blood of CF patients compared

to healthy controls and a further reduction in Tregs was seen in patients with chronic Pa infection (38). The lack of difference in Treg percentages between CF NTM patients and healthy controls




FIGURE 6 | Divergence of CD8⁺ T cell cytokine profiles in immunocompetent individuals with active non-tuberculous mycobacteria (NTM) infection. **(A)** Flow cytometric analysis of *ex vivo* CD8⁺ T cells showed significantly more IFN γ ⁻ TNF α ⁺ cells ($P = 0.01$) and IL2⁻ TNF α ⁺ cells ($P = 0.048$) in NTM^{Act} compared to HC^B. **(B)** Polyfunctionality profiling of CD8⁺ T cells shown in pie charts were not significantly different between the two groups. However, significantly higher TNF α mono-functional CD8⁺ T cells were observed in NTM patient group ($P = 0.002$). **(C)** SPICE dot plots show polyfunctionality profile of all combinations of cytokine expression in CD8⁺ T cells after PMA/I stimulation between NTM^{Act} and HC^B groups. Significantly different expression compared to HC^B (Wilcoxon rank test $P < 0.05$) is indicated with # symbol.

could be an indication of an increase in Tregs following NTM infection. This rationale would align with our findings of elevated Tregs in the elderly immunocompetent patients with active NTM infection. Of note, increased Treg numbers have been observed in the peripheral blood of TB patients (39–41) indicating overlap between TB and NTM immunopathology.

When examining the T cell fingerprint, CF patients with active and past MABS infection were different to control patients with

chronic Pa infection, though they were similar to the healthy control group. This finding suggests differences in T cell immunity between CF patients susceptible to MABS and CF patients who have no history of NTM infection.

Distinct activation and exhaustion profiles were seen in both patient cohorts with NTM infection compared to corresponding controls. Higher numbers of CD4⁺ T cells co-expressing the activation marker CD25 and the exhaustion marker CTLA-4

were seen in CF patients with both active and past NTM infection. In the elderly cohort, a similar pattern of increased CD25 and CTLA-4 co-expression was observed on CD4⁺ T cells. PD-1 expression was also increased on CD4⁺ T cells, suggesting an exhausted immune phenotype. High PD-1 expression on T cells associates with increased TB disease burden (26). Increased PD-1 expression on T cells, B cells, NK cells, and monocytes has also been reported in patients with MAC infection (42). In mice, PD-1 gene knockout can enhance TB resistance by preventing overproduction of IFN γ (43). However, another study showed that PD-1 gene knockout mice can be more susceptible to NTM infection (44). To date, the immune checkpoint CTLA-4 has not been studied in the context of NTM. Our data present the first finding of elevated CTLA-4 on T cells in NTM infection. The significance of this as well as elevated PD-1 expression on T cells indicates a degree of immune suppression in NTM infection. No PD-1 (or CTLA-4) antibody blockade therapy has been examined *in vivo* on TB or NTM. Future research should focus on the prospect of treating NTM patients using PD-1 and/or CTLA-4 antibody blockade.

TNF α directly activates macrophages to restrict mycobacterial growth and induces apoptosis of infected macrophages leading to bacterial killing (45–47). TNF α is also essential for granuloma formation and disease restriction during mycobacterial pathogenesis highlighting its importance for *in vivo* control of the pathogen (48). Animals deficient in TNF α are highly susceptible to disseminated forms of TB (49). The cytokine polyfunctionality profiles seen in CF cohorts revealed underlying global TNF α deficiencies that could explain susceptibility to NTM infection in CF patients. Low TNF α secretion (mono-functional TNF α -secreting CD4⁺ T cells) was seen in CF patients with both active and past NTM infection, while increased TNF α secreting mono-functional CD8⁺ T cells were seen in elderly patients with NTM infection. Given that CF patients with both active and past infection exhibit this TNF α deficiency in response to mitogen, it is more likely that this phenotype is an underlying predisposition to disease rather than a direct effect of the disease. The contrasting pattern of TNF α secretion in the two disease scenarios is significant in the larger context of mycobacterial pathobiology. Studies of NTM immunity are conflicting and show both low and high levels of TNF α production (12, 42, 50–52), however, these studies did not perform in depth mapping of cell subsets. TNF α production has an important role in host resistance as treatment with anti-TNF α therapies are associated with increased susceptibility to active TB and reactivation of latent TB infection (53) and is correlated with NTM activation in autoimmune diseases, such as rheumatoid arthritis, Crohn's disease, ankylosing spondylitis, and psoriasis (54). Here, we have shown that the TNF α profiles can vary with disease setting. Thus, taking into account the individual patient profile is critical when interpreting these findings. Moreover, we have also shown that TNF α secretion levels vary according to cell subset and it may be specific deficiencies in certain cell subsets that predispose to disease. It is possible that low dose TNF α replacement therapy, *via* aerosol, for example, may aid standard of care in NTM treatment in CF. Indeed, other cytokine replacement therapies, such as IFN γ , IFN α , IL-2, GM-CSF, and IL-12 have shown promise against TB, MDR-TB, MAC, and

MABS [reviewed in Ref. (53)]. Systemically administered IFN γ has shown the most promise for clinical use (55).

The importance of IFN γ in anti-mycobacterial immunity is widely accepted and extensively studied in TB (56). Reduced IFN γ production in NTM infection has been shown in several studies (51, 57, 58) though contradictory results have also been reported (16). Akin to TNF α , we show that IFN γ secretion varies with cell subset and disease scenario. In the CF NTM cohort, increased IFN γ secretion was seen in both CD4⁺ and CD8⁺ T cells compared to healthy controls, whereas in the elderly NTM cohort, there was no significant increase in IFN γ -secreting cells in both CD4⁺ or CD8⁺ T cells. This could indicate an inadequate protective response rather than a deficiency of cytokine. Comparison of MABS infection in these two disease cohorts once again highlights the importance of the clinical context when searching for risk factors. However, one limitation of the study is the profiling of circulating lymphocytes which may differ to those at the site of disease.

The search for environmental and behavioral risk factors for NTM infection in CF patients has found increased acquisition in the tropics and decreased acquisition with macrolide treatment (59). Here, we show that it is also possible to define immune parameters in the circulatory blood that help to identify at-risk individuals. Specifically, we showed that CF patients can stratify CF into two groups based on Treg frequencies, CD4⁺ T cell surface phenotype (CD25 and CTLA-4) and cytokine production by CD4⁺/CD8⁺ T cells (IFN γ , TNF α , CD107a, and IL-2), with at-risk patients exhibiting a distinct deficiency in TNF α . We suggest that this immune signature could be further refined and validated in independent cohort studies, and then ultimately developed for use as a diagnostic tool to identify individuals who are at high risk of developing NTM infection. Targeted behavioral interventions for at-risk patients may subsequently decrease the risk of NTM acquisition from the environment or infected patients.

The role of CD8⁺ T cells in NTM infection in humans remains unclear. Research in TB in primate and mouse models has shown that CD8⁺ T cells are important in controlling experimental infection. However, the precise mechanism by which they contribute to protection is unknown (60). Regulatory CD8⁺ T cells have been identified in human TB, though their role in protection is also not clear (60).

This study specifically focused on MABS infection. It is important to highlight that the CF patient with active MAC infection (shown as a blue circle in **Figures 1A** and **2A**) exhibited a T cell fingerprint and T cell functional profile very similar to patients with MABS infection (note; this patient was not included in statistical analysis). Additionally, it is also important to note the 2 of 6 CF MABS patients and 3 of 10 elderly MABS patients either had a previous history of MAC infection or subsequently developed MAC infection. This tentatively suggests that the susceptibility to NTM infection may occur through a common immune dysfunction pathway, although further antigen-specific immune studies are required to validate this hypothesis. If NTM-specific immune dysfunction is common in the individuals, targeted immunotherapies may help in correcting this deficit.

Limitations of this study include the small sample size due to restrictions in patient recruitment within a confined study

duration. An additional, larger patient cohort will be essential to validate our predictive model. Limitations in sample volumes also restricted the analysis to *ex vivo* phenotyping and mitogen-activation signatures. Thus, antigen-specific immune responses were not conducted in this study. However, further research is warranted given the identification of significant global immune anomalies in this exploratory study. Anomalies in mitogen-triggered activation signatures in the global T cell compartment suggest an underlying immune deficiency in the patients that would likely translate to NTM-specific responses as seen in TB (61, 62).

In summary, our study presents the first data on immune checkpoint expression on T cell subsets in human MABS infection as well as the first comparison of T cell polyfunctionality between CF and non-CF patient groups with MABS infection in response to mitogen stimulation. We show that MABS infection in two different patient groups exhibit specific immune phenotypes and show dysregulation in type 1 cytokine production and a global decrease in T cell "quality." In CF patients, TNF α -mediated immunity may hold the key to understanding the increased risk of MABS infection and guide future therapeutic interventions. In elderly individuals, interference with checkpoint molecules (PD-1 and CTLA-4) may guide future therapeutic interventions. Collectively, the study has revealed many potential associations between T cell phenotype and individuals at risk of MABS infection. The idea of an underlying immune dysfunction which predisposes certain individuals to NTM infection is attractive but very speculative. Larger populations and further functional experiments will be required to validate this hypothesis.

REFERENCES

- Morimoto K, Iwai K, Uchimura K, Okumura M, Yoshiyama T, Yoshimori K, et al. A steady increase in nontuberculous mycobacteriosis mortality and estimated prevalence in Japan. *Ann Am Thorac Soc* (2014) 11(1):1–8. doi:10.1513/AnnalsATS.201303-067OC
- Society TRCotBT. Pulmonary disease caused by *Mycobacterium avium*-intracellulare in HIV-negative patients: five-year follow-up of patients receiving standardised treatment. *Int J Tuberc Lung Dis* (2002) 6(7):628–34.
- Andrejak C, Thomsen VO, Johansen IS, Riis A, Benfield TL, Duhaut P, et al. Nontuberculous pulmonary mycobacteriosis in Denmark: incidence and prognostic factors. *Am J Respir Crit Care Med* (2010) 181(5):514–21. doi:10.1164/rccm.200905-0778OC
- Fleshner M, Olivier KN, Shaw PA, Adjemian J, Strollo S, Claypool RJ, et al. Mortality among patients with pulmonary non-tuberculous mycobacteria disease. *Int J Tuberc Lung Dis* (2016) 20(5):582–7. doi:10.5588/ijtld.15.0807
- Fangous MS, Mougari F, Gouriou S, Calvez E, Raskine L, Cambau E, et al. Classification algorithm for subspecies identification within the *Mycobacterium abscessus* species, based on matrix-assisted laser desorption ionization-time of flight mass spectrometry. *J Clin Microbiol* (2014) 52(9):3362–9. doi:10.1128/JCM.00788-14
- Hoefsloot W, van Ingen J, Andrejak C, Angeby K, Bauriaud R, Bemer P, et al. The geographic diversity of nontuberculous mycobacteria isolated from pulmonary samples: an NTM-NET collaborative study. *Eur Respir J* (2013) 42(6):1604–13. doi:10.1183/09031936.00149212
- Prevots DR, Marras TK. Epidemiology of human pulmonary infection with nontuberculous mycobacteria: a review. *Clin Chest Med* (2015) 36(1):13–34. doi:10.1016/j.ccm.2014.10.002
- Henkle E, Aksamit T, Barker A, Daley CL, Griffith D, Leitman P, et al. Patient-centered research priorities for pulmonary nontuberculous mycobacteria (NTM) infection. An NTM Research Consortium Workshop Report. *Ann Am Thorac Soc* (2016) 13(9):S379–84. doi:10.1513/AnnalsATS.201605-387WS

ETHICS STATEMENT

All samples were obtained with written consent and all protocols were approved by the Human Research Ethics Committees of the QIMR Berghofer Medical Research Institute, The Prince Charles Hospital, and Greenslopes Private Hospital, Australia (QIMR Berghofer HREC P2045).

AUTHOR CONTRIBUTIONS

SB and JM conceptualized the study. VL, CR, DS, AK, DD, DR, RT, and JM performed the experiments and analyzed the data. CR, VL, and JM wrote the manuscript with input from all authors. DR, RT, SB, and JM supervised the study. All authors approved the final manuscript.

ACKNOWLEDGMENTS

JM is supported by a NHMRC CDF Level 2 Fellowship (1131732). DR and SB were supported by Queensland Health Fellowships. This work was supported by a QIMR Berghofer Clinician Research Collaboration Award.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <https://www.frontiersin.org/articles/10.3389/fimmu.2018.01319/full#supplementary-material>.

- Bryant JM, Grogono DM, Rodriguez-Rincon D, Everall I, Brown KP, Moreno P, et al. Emergence and spread of a human-transmissible multidrug-resistant nontuberculous mycobacterium. *Science* (2016) 354(6313):751–7. doi:10.1126/science.aaf8156
- Martiniano SL, Nick JA, Daley CL. Nontuberculous mycobacterial infections in cystic fibrosis. *Clin Chest Med* (2016) 37(1):83–96. doi:10.1016/j.ccm.2015.11.001
- Qvist T, Gilljam M, Jonsson B, Taylor-Robinson D, Jensen-Fangel S, Wang M, et al. Epidemiology of nontuberculous mycobacteria among patients with cystic fibrosis in Scandinavia. *J Cyst Fibros* (2015) 14(1):46–52. doi:10.1016/j.jcf.2014.08.002
- Bar-On O, Mussaffi H, Mei-Zahav M, Prais D, Steuer G, Stafler P, et al. Increasing nontuberculous mycobacteria infection in cystic fibrosis. *J Cyst Fibros* (2015) 14(1):53–62. doi:10.1016/j.jcf.2014.05.008
- Park IK, Olivier KN. Nontuberculous mycobacteria in cystic fibrosis and non-cystic fibrosis bronchiectasis. *Semin Respir Crit Care Med* (2015) 36(2):217–24. doi:10.1055/s-0035-1546751
- Chan ED, Iseman MD. Underlying host risk factors for nontuberculous mycobacterial lung disease. *Semin Respir Crit Care Med* (2013) 34(1):110–23. doi:10.1055/s-0033-1333573
- Chan ED, Bai X, Kartalija M, Orme IM, Ordway DJ. Host immune response to rapidly growing mycobacteria, an emerging cause of chronic lung disease. *Am J Respir Cell Mol Biol* (2010) 43(4):387–93. doi:10.1165/rcmb.2009-0276TR
- Lim A, Allison C, Price P, Waterer G. Susceptibility to pulmonary disease due to *Mycobacterium avium*-intracellulare complex may reflect low IL-17 and high IL-10 responses rather than Th1 deficiency. *Clin Immunol* (2010) 137(2):296–302. doi:10.1016/j.clim.2010.07.011
- Cooper AM, Mayer-Barber KD, Sher A. Role of innate cytokines in mycobacterial infection. *Mucosal Immunol* (2011) 4(3):252–60. doi:10.1038/mi.2011.13
- Kim SY, Koh WJ, Kim YH, Jeong BH, Park HY, Jeon K, et al. Importance of reciprocal balance of T cell immunity in *Mycobacterium abscessus* complex lung disease. *PLoS One* (2014) 9(10):e109941. doi:10.1371/journal.pone.0109941

19. Matsuyama M, Ishii Y, Yageta Y, Ohtsuka S, Ano S, Matsuno Y, et al. Role of Th1/Th17 balance regulated by T-bet in a mouse model of *Mycobacterium avium* complex disease. *J Immunol* (2014) 192(4):1707–17. doi:10.4049/jimmunol.1302258
20. Becker KL, van Ingen J, Ten Oever J, Merkus PJ, Ferwerda G, Netea MG, et al. Deficient interleukin-17 production in response to *Mycobacterium abscessus* in cystic fibrosis. *Eur Respir J* (2016) 47(3):990–3. doi:10.1183/13993003.00446-2015
21. Allie N, Alexopoulou L, Quesniaux VJ, Fick L, Kranidioti K, Kollias G, et al. Protective role of membrane tumour necrosis factor in the host's resistance to mycobacterial infection. *Immunology* (2008) 125(4):522–34. doi:10.1111/j.1365-2567.2008.02865.x
22. Hoos A. Development of immuno-oncology drugs – from CTLA4 to PD1 to the next generations. *Nat Rev Drug Discov* (2016) 15(4):235–47. doi:10.1038/nrd.2015.35
23. Kaufmann DE, Walker BD. PD-1 and CTLA-4 inhibitory cosignaling pathways in HIV infection and the potential for therapeutic intervention. *J Immunol* (2009) 182(10):5891–7. doi:10.4049/jimmunol.0803771
24. Smith P, Walsh CM, Mangan NE, Fallon RE, Sayers JR, McKenzie AN, et al. *Schistosoma mansoni* worms induce anergy of T cells via selective up-regulation of programmed death ligand 1 on macrophages. *J Immunol* (2004) 173(2):1240–8. doi:10.4049/jimmunol.173.2.1240
25. Das S, Suarez G, Beswick EJ, Sierra JC, Graham DY, Reyes VE. Expression of B7-H1 on gastric epithelial cells: its potential role in regulating T cells during *Helicobacter pylori* infection. *J Immunol* (2006) 176(5):3000–9. doi:10.4049/jimmunol.176.5.3000
26. Jurado JO, Alvarez IB, Pasquinelli V, Martinez GJ, Quiroga MF, Abbate E, et al. Programmed death (PD)-1:PD-ligand 1/PD-ligand 2 pathway inhibits T cell effector functions during human tuberculosis. *J Immunol* (2008) 181(1):116–25. doi:10.4049/jimmunol.181.1.116
27. McNab FW, Berry MP, Graham CM, Bloch SA, Oni T, Wilkinson KA, et al. Programmed death ligand 1 is over-expressed by neutrophils in the blood of patients with active tuberculosis. *Eur J Immunol* (2011) 41(7):1941–7. doi:10.1002/eji.201141421
28. Szaba FM, Kummer LW, Duso DK, Koroleva EP, Tumanov AV, Cooper AM, et al. TNF α and IFN γ but not perforin are critical for CD8 T cell-mediated protection against pulmonary *Yersinia pestis* infection. *PLoS Pathog* (2014) 10(5):e1004142. doi:10.1371/journal.ppat.1004142
29. Frey O, Meisel J, Hutloff A, Bonhagen K, Bruns L, Kroczeck RA, et al. Inducible costimulator (ICOS) blockade inhibits accumulation of polyfunctional T helper 1/T helper 17 cells and mitigates autoimmune arthritis. *Ann Rheum Dis* (2010) 69(8):1495–501. doi:10.1136/ard.2009.119164
30. Pizzemolo T, Guggino G, La Manna MP, Di Liberto D, Dieli F, Caccamo N. Functional signatures of human CD4 and CD8 T cell responses to *Mycobacterium tuberculosis*. *Front Immunol* (2014) 5:180. doi:10.3389/fimmu.2014.00180
31. Lee TWR, Brownlee KG, Conway SP, Denton M, Littlewood JM. Evaluation of a new definition for chronic *Pseudomonas aeruginosa* infection in cystic fibrosis patients. *J Cyst Fibros* (2003) 2(1):29–34. doi:10.1016/S1569-1993(02)00141-8
32. Amir el AD, Davis KL, Tadmor MD, Simonds EF, Levine JH, Bendall SC, et al. viSNE enables visualization of high dimensional single-cell data and reveals phenotypic heterogeneity of leukemia. *Nat Biotechnol* (2013) 31(6):545–52. doi:10.1038/nbt.2594
33. Proietti C, Zakrzewski M, Watkins TS, Berger B, Hasan S, Ratnatunga CN, et al. Mining, visualizing and comparing multidimensional biomolecular data using the Genomics Data Miner (GMine) Web-Server. *Sci Rep* (2016) 6(1):38178. doi:10.1038/srep38178
34. Roederer M, Nozzi JL, Nason MC. SPICE: exploration and analysis of post-cytometric complex multivariate datasets. *Cytometry A* (2011) 79(2):167–74. doi:10.1002/cyto.a.21015
35. Thomson R, Donnan E, Konstantinos A. Notification of nontuberculous mycobacteria: an Australian perspective. *Ann Am Thorac Soc* (2017) 14(3):318–23. doi:10.1513/AnnalsATS.201612-994OI
36. Jennrich S, Lee MH, Lynn RC, Dewberry K, Debes GF. Tissue exit: a novel control point in the accumulation of antigen-specific CD8 T cells in the influenza A virus-infected lung. *J Virol* (2012) 86(7):3436–45. doi:10.1128/JVI.07025-11
37. Steindor M, Nkwouano V, Mayatepek E, Mackenzie CR, Schramm D, Jacobsen M. Rapid detection and immune characterization of *Mycobacterium abscessus* infection in cystic fibrosis patients. *PLoS One* (2015) 10(3):e0119737. doi:10.1371/journal.pone.0119737
38. Hector A, Schafer H, Poschel S, Fischer A, Fritzsching B, Ralhan A, et al. Regulatory T-cell impairment in cystic fibrosis patients with chronic *Pseudomonas* infection. *Am J Respir Crit Care Med* (2015) 191(8):914–23. doi:10.1164/rccm.201407-1381OC
39. Guyot-Revoll V, Innes JA, Hackforth S, Hinks T, Lalvani A. Regulatory T cells are expanded in blood and disease sites in patients with tuberculosis. *Am J Respir Crit Care Med* (2006) 173(7):803–10. doi:10.1164/rccm.200508-1294OC
40. Ribeiro-Rodrigues R, Resende Co T, Rojas R, Toossi Z, Dietze R, Boom WH, et al. A role for CD4+CD25+ T cells in regulation of the immune response during human tuberculosis. *Clin Exp Immunol* (2006) 144(1):25–34. doi:10.1111/j.1365-2249.2006.03027.x
41. Chen X, Zhou B, Li M, Deng Q, Wu X, Le X, et al. CD4(+)CD25(+)FoxP3(+) regulatory T cells suppress *Mycobacterium tuberculosis* immunity in patients with active disease. *Clin Immunol* (2007) 123(1):50–9. doi:10.1016/j.clim.2006.11.009
42. Shu CC, Wang JY, Wu MF, Wu CT, Lai HC, Lee LN, et al. Attenuation of lymphocyte immune responses during *Mycobacterium avium* complex-induced lung disease due to increasing expression of programmed death-1 on lymphocytes. *Sci Rep* (2017) 7:42004. doi:10.1038/srep42004
43. Sakai S, Kauffman KD, Sallin MA, Sharpe AH, Young HA, Ganusov VV, et al. CD4 T Cell-Derived IFN- γ plays a minimal role in control of pulmonary *Mycobacterium tuberculosis* infection and must be actively repressed by PD-1 to prevent lethal disease. *PLoS Pathog* (2016) 12(5):e1005667. doi:10.1371/journal.ppat.1005667
44. Barber DL, Mayer-Barber KD, Feng CG, Sharpe AH, Sher A. CD4 T cells promote rather than control tuberculosis in the absence of PD-1-mediated inhibition. *J Immunol* (2011) 186(3):1598–607. doi:10.4049/jimmunol.1003304
45. Jayaraman P, Sada-Ovalle I, Nishimura T, Anderson AC, Kuchroo VK, Remold HG, et al. IL-1 β promotes antimicrobial immunity in macrophages by regulating TNFR signaling and caspase-3 activation. *J Immunol* (2013) 190(8):4196–204. doi:10.4049/jimmunol.1202688
46. Balcewicz-Sablinska MK, Keane J, Kornfeld H, Remold HG. Pathogenic *Mycobacterium tuberculosis* evades apoptosis of host macrophages by release of TNF-R2, resulting in inactivation of TNF- α . *J Immunol* (1998) 161(5):2636–41.
47. Bermudez LE, Young LS. Tumor necrosis factor, alone or in combination with IL-2, but not IFN- γ , is associated with macrophage killing of *Mycobacterium avium* complex. *J Immunol* (1988) 140(9):3006–13.
48. Roach DR, Bean AGD, Demangel C, France MP, Briscoe H, Britton WJ. TNF regulates chemokine induction essential for cell recruitment, granuloma formation, and clearance of mycobacterial infection. *J Immunol* (2002) 168(9):4620–7. doi:10.4049/jimmunol.168.9.4620
49. Flynn JL, Goldstein MM, Chan J, Triebold KJ, Pfeffer K, Lowenstein CJ, et al. Tumor necrosis factor- α is required in the protective immune response against *Mycobacterium tuberculosis* in mice. *Immunity* (1995) 2(6):561–72. doi:10.1016/1074-7613(95)90001-2
50. Greinert U, Schlaak M, Rusch-Gerdes S, Flad HD, Ernst M. Low in vitro production of interferon- γ and tumor necrosis factor- α in HIV-seronegative patients with pulmonary disease caused by nontuberculous mycobacteria. *J Clin Immunol* (2000) 20(6):445–52. doi:10.1023/A:1026407815946
51. Kwon YS, Kim EJ, Lee SH, Suh GY, Chung MP, Kim H, et al. Decreased cytokine production in patients with nontuberculous mycobacterial lung disease. *Lung* (2007) 185(6):337–41. doi:10.1007/s00408-007-9040-z
52. Sampaio EP, Elloumi HZ, Zelazny A, Ding L, Paulson ML, Sher A, et al. *Mycobacterium abscessus* and *M. avium* trigger toll-like receptor 2 and distinct cytokine response in human cells. *Am J Respir Cell Mol Biol* (2008) 39(4):431–9. doi:10.1165/rcmb.2007-0413OC
53. Tomioka H. Adjunctive immunotherapy of mycobacterial infections. *Curr Pharm Des* (2004) 10(26):3297–312. doi:10.2174/1381612043383232
54. Desai AA, Marks DJ. Atypical mycobacteria: showerheads, anti-TNF therapy and Crohn's disease. *Expert Rev Clin Immunol* (2010) 6(5):695–9. doi:10.1586/eci.10.61
55. Reljic R, Paul MJ, Arias MA. Cytokine therapy of tuberculosis at the crossroads. *Expert Rev Respir Med* (2009) 3(1):53–66. doi:10.1586/17476348.3.1.53
56. Kumar P. IFN γ -producing CD4+ T lymphocytes: the double-edged swords in tuberculosis. *Clin Transl Med* (2017) 6(1):21. doi:10.1186/s40169-017-0151-8

57. Vankayalapati R, Wizel B, Samten B, Griffith DE, Shams H, Galland MR, et al. Cytokine profiles in immunocompetent persons infected with *Mycobacterium avium* complex. *J Infect Dis* (2001) 183(3):478–84. doi:10.1086/318087
58. Safdar A, White DA, Stover D, Armstrong D, Murray HW. Profound interferon gamma deficiency in patients with chronic pulmonary nontuberculous mycobacteriosis. *Am J Med* (2002) 112:756–9. doi:10.1016/S0002-9343(02)01313-X
59. Sherrard LJ, Tay GT, Butler CA, Wood ME, Yerkovich S, Ramsay KA, et al. Tropical Australia is a potential reservoir of non-tuberculous mycobacteria in cystic fibrosis. *Eur Respir J* (2017) 49(5):1–4. doi:10.1183/13993003.00046-2017
60. Lin PL, Flynn JL. CD8 T cells and *Mycobacterium tuberculosis* infection. *Semin Immunopathol* (2015) 37(3):239–49. doi:10.1007/s00281-015-0490-8
61. Riou C, Berkowitz N, Goliath R, Burgers WA, Wilkinson RJ. Analysis of the phenotype of *Mycobacterium tuberculosis*-specific CD4+ T cells to discriminate latent from active tuberculosis in HIV-uninfected and HIV-infected individuals. *Front Immunol* (2017) 8:968. doi:10.3389/fimmu.2017.00968
62. Abebe F, Belay M, Legesse M, Mihret A, Franklin KS. Association of ESAT-6/CFP-10-induced IFN-gamma, TNF-alpha and IL-10 with clinical tuberculosis: evidence from cohorts of pulmonary tuberculosis patients, household contacts, and community controls in an endemic setting. *Clin Exp Immunol* (2017) 189(2):241–9. doi:10.1111/cei.12972

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Lutzky, Ratnatunga, Smith, Kupz, Doolan, Reid, Thomson, Bell and Miles. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.