

# SCIENTIFIC REPORTS

OPEN

## Parasite detection in the ornamental fish trade using environmental DNA

A. Trujillo-González<sup>1,5</sup>, R. C. Edmunds<sup>2</sup>, J. A. Becker<sup>3</sup> & K. S. Hutson<sup>1,4</sup>

Effective border control relies on stringent biosecurity protocols to detect and prevent introductions of exotic pests and diseases. Detection of pathogens and parasites in the live ornamental fish trade using environmental DNA (eDNA) techniques has the potential to improve current biosecurity practices. We examined water samples from 11 target consignments (cyprinids susceptible to *Dactylogyrus* spp. infections) and seven non-target fish consignments (non-cyprinids, not susceptible to *Dactylogyrus* spp. infections) imported from Southeast Asia to Australia for the presence of eDNA from five *Dactylogyrus* species (Monogenea: Dactylogyridae). A four-step predictive framework was used to predict putative positive and putative negative detections from quantitative PCR assays. Both target and non-target consignments were positive for *Dactylogyrus* spp. eDNA as confirmed by Sanger sequencing. Positive detections for *Dactylogyrus* spp. eDNA in non-target fish consignments demonstrates the possibility of source water contamination, limiting the applicability of eDNA screening methods at border control. This study suggests that screening for parasite eDNA within ornamental fish consignments should be tested during pre-export quarantine periods to avoid false positive detections at border control. Lastly, the proposed predictive framework has a broad utility for minimizing false positive and false negative eDNA detections of aquatic organisms.

The ornamental fish trade is a known route of exotic pathogen translocations globally<sup>1–5</sup>. Parasites and their infected hosts have been co-introduced to non-native environments with detrimental effects on biodiversity, ecosystems, industries, and dependent local communities<sup>6</sup>. To minimize pathogen translocation through the ornamental fish trade, governments can establish quarantine measures based on scientific risk analyses that consider the origin and history of fish stocks, parasite life cycles, host susceptibility to infection, risk of transmission to native species, and the reliability of detection methods<sup>3,7</sup>. Australia for example, has stringent mandatory pre-export quarantine requirements, biosecurity protocols at border control, and post arrival mandatory quarantine requirements following strict biosecurity import risk assessments of ornamental fish imports<sup>3,8</sup>. Despite current biosecurity protocols<sup>9</sup>, recent surveys of ornamental fish species imported to Australia have shown that a high diversity of parasites were not detected during inspection at border control, highlighting the need for more detection sensitivity<sup>10</sup>. Considering the limitation of gross visual inspection under current biosecurity protocols it is important to explore new and complimentary methods to increase biosecurity rigor and the possible integration of molecular genetic techniques.

Environmental DNA (eDNA) refers to the DNA that is naturally shed by organisms, such as through epidermal sloughing, metabolic waste excretions or post-mortem decay, into their local environment<sup>11</sup>. In the case of microscopic parasites, life stages like eggs, spores, cysts, larvae, juveniles and adults can be present in the water column, in sediment, or in extracellular DNA disassociated from host organisms<sup>12</sup>. As such, parasite genomic (gDNA) and nucleic (nDNA) can be captured with eDNA samples<sup>12</sup>, extracted, and screened for target species using standard molecular genetic techniques like quantitative real-time polymerase chain reaction (qPCR)<sup>11,13,14</sup>.

<sup>1</sup>Centre for Sustainable Tropical Fisheries and Aquaculture, College of Science and Engineering, James Cook University, 1 James Cook Dr, Townsville, QLD, 4811, Australia. <sup>2</sup>Centre for Tropical Water & Aquatic Ecosystem Research (TropWATER), College of Science and Engineering, James Cook University, 1 James Cook Dr, Townsville, QLD, 4811, Australia. <sup>3</sup>School of Life and Environmental Sciences, Faculty of Science, University of Sydney, 425 Werombi road, Camden, NSW, 2570, Australia. <sup>4</sup>Cawthron Institute, 98 Halifax Street East, Nelson, 7010, New Zealand. <sup>5</sup>Institute for Applied Ecology, University of Canberra, 40 Bunda Street, Canberra, ACT, 2601, Australia. Correspondence and requests for materials should be addressed to A.T.-G. (email: [alejandro.trujillogonzalez@canberra.edu.au](mailto:alejandro.trujillogonzalez@canberra.edu.au))

Environmental DNA could enable species-level detection and monitoring in aquatic parasitology with important benefits to human health, animal welfare, freshwater fisheries, coastal aquaculture, conservation, and ecosystem health<sup>12</sup>. Indeed, captured and extracted eDNA from water samples has been shown to accurately detect pathogenic trematodes infecting wild amphibians<sup>15</sup> and to monitor parasites infecting farmed<sup>16–18</sup> and wild fish species<sup>19</sup>. Environmental DNA was recently proposed to be a non-destructive and sensitive detection tool for biosecurity, and was used to determine the presence of ornamental fish species present at low densities within high risk mixed imports<sup>20</sup>. Screening water used to import ornamental fish consignments for the presence of parasites has the potential for biosecurity monitoring advancement; however, there are no studies to date that have specifically tested this utility of eDNA.

False positive and false negative errors are commonly encountered in qPCR analyses<sup>21–23</sup>. From a biosecurity perspective, misinterpreting qPCR data could lead to pathogen-free consignments being considered hazards during quarantine inspection (i.e., false positive error), or high-risk pathogens going undetected in infected consignments (i.e., false negative error). As such, preventative measures must be developed to ensure accurate interpretation of qPCR data<sup>22</sup> and reduce the possibility of false positive and negative results.

The aim of this study was to determine if eDNA screening by qPCR is a plausible detection tool for biosecurity. A four-step predictive framework was designed to minimize the possibility of false positive and false negative qPCR detections for the presence or absence of five ectoparasitic monogenean flukes (*Dactylogyrus anchoratus*, *D. formosus*, *D. intermedius*, *D. vastator* and *D. ostraviensis*) previously detected by necropsies infecting ornamental cyprinid fishes (*Carassius auratus* and *Pethia conchonius*) imported from Southeast Asia to Australia<sup>10</sup>.

## Methods

***Dactylogyrus* spp. eDNA collection.** All water samples analysed for the presence of eDNA from *Dactylogyrus* species in this study were collected during a cross-sectional survey for the presence of nationally listed aquatic pathogens associated with at least one ornamental fish host<sup>8</sup>. Briefly, 37 ornamental fish consignments representing 11 farmed freshwater and seven marine wild caught fish species were imported from Southeast Asia to Australia in 2015 following Australian Biosecurity Import Conditions (BICON) and subjected to Australian quarantine protocols, which involved gross visual inspection and clearance by Australian Quarantine Services<sup>9</sup>. A 'consignment' of fish was defined as a unique fish species within a shipment of fish, identified by an invoice containing details of the numbers and species of fish, date of shipment, origin and destination, accompanied by health certification<sup>3,9</sup>. Following release from quarantine inspection, all consignments were transported by road to an Approved Arrangement Site (AA Site) at the University of Sydney (Camden, Australia).

Freshwater consignments arrived at the AA Site in either one large plastic bag or several medium plastic bags, containing 40 to 200 individuals depending on species and size<sup>8</sup>. Each plastic bag contained approximately 1–5 L of freshwater and was sealed with either rubber bands or metal clasps. All consignments were housed inside large Styrofoam boxes during transit (12–48 hours including export, delivery, inspection, and release to the importer) before water samples were collected from each consignment and preserved. Negative controls (distilled water) were collected prior to collecting triplicate 15 mL samples from each fish consignment. To minimize the risk of eDNA cross contamination, each 15 mL replicate was collected from all plastic bags holding each consignment using a new disposable 20 mL sterile glass pipette attached to an automatic pipette controller (EasyPet, Eppendorf). Water samples were dispensed directly into individual pre-labelled DNA-free 50 mL centrifuge tubes, each with 33.5 mL absolute ethanol and 1.5 mL 3 M sodium acetate for preservation and then stored at room temperature<sup>16</sup>. Following water sample collection, 30 fish from each consignment were randomly selected, euthanized, and examined for the presence of monogenean parasites by necropsy, as described in a separate study<sup>10</sup>. In brief, all 30 fish were sequentially surveyed for external parasites by an experienced parasitologist using a compound microscope to carefully examine gill samples from each fish for the presence or absence of parasites<sup>10</sup>. A sample size of 30 fish per consignment was selected to achieve a minimum detection prevalence of 10% with 95% confidence limits determined by using exact binomial approximation<sup>8</sup>. As such, samples where no parasites were detected by necropsy were considered to have an apparent prevalence of 0%, with a 95% confidence interval (CI) of 0–11.4%, assuming a perfect test<sup>8</sup>. Environmental DNA was extracted using cetyl trimethylammonium bromide (CTAB), which included phenol-chloroform isolation and terminal isopropanol precipitation<sup>16</sup>. All DNA was resuspended in 60 µL 1x Tris-EDTA (TE) buffer and stored at –20 °C until screening for *Dactylogyrus* spp. eDNA by qPCR. Animal ethics, method and sampling approval was obtained from the University of Sydney Animal Ethics Committee (approval number: 720) and all methods were performed in accordance with guidelines and regulations of the University of Sydney Animal Ethics Committee.

**Design of species-specific *Dactylogyrus* primers and assay validation.** Novel species-specific oligonucleotide primers were designed to detect and discriminate between five *Dactylogyrus* species (Monogenea: Capsalidae): *Dactylogyrus anchoratus* (Dujardin, 1845), *Dactylogyrus formosus* Kulwiec, 1927, *Dactylogyrus intermedius* Wegener, 1909, *Dactylogyrus ostraviensis* Rehulka, 1988, and *Dactylogyrus vastator* Nybelin, 1924. All five *Dactylogyrus* spp. are highly specific to cyprinid fish hosts<sup>24,25</sup>. All qPCR assays targeted the internal transcribed spacer 1 (*ITS1*) between base pair 366 and 588. The *ITS1* is a high abundance nuclear gene known to be detectable in eDNA extracted from water samples<sup>26</sup> and to provide species-level resolution for *Dactylogyrus*<sup>10</sup> and other helminths given its low intraspecific yet high interspecific variability<sup>27</sup>. Each *Dactylogyrus*-specific primer was designed to target the *ITS1* region that contained the most mismatches ( $\geq 1$ ) between target and all non-target *Dactylogyrus* species (Table 1). To achieve this, previously accessioned *Dactylogyrus* spp. *ITS1* nucleotide sequences<sup>10</sup> were downloaded from GenBank (NCBI) and aligned using ClustalW ([www.genome.jp/tools/clustalw](http://www.genome.jp/tools/clustalw), version 1.81).

All qPCR assays were tested for specificity *in silico* using the National Center for Biotechnology Information (NCBI) Primer BLAST<sup>28</sup>, Amplify4 ([engels.genetics.wisc.edu/amplify](http://engels.genetics.wisc.edu/amplify)), and Amplifx 1.7.0 (Nicolas Jullien;

Parasite species	Primer	Amplicon (bp)	Annealing (°C)	Primer sequence (5'-3')	qPCR efficiency (%)	R <sup>2</sup>	Limit of detection (ng/μL)
<i>Dactylogyrus anchoratus</i>	D. anchoratus F	185	60	5'-GCCATCCTTGAGGGAATATGCCCA-3'	75.12	0.981	0.00065
	D. anchoratus R			5'-GAGTTTACGTTGACCGCCCGACAT-3'			
<i>Dactylogyrus formosus</i>	D. formosus F	184	65	5'-ATCATCCTTGTGGGAATCTGCCCG-3'	119.55	0.984	0.0079
	D. formosus R			5'-AAGTGTACGTTGACCGCCAGCAG-3'			
<i>Dactylogyrus ostraviensis</i>	D. ostraviensis F	120	65	5'-TCGTGCTGACGACCTTGG-3'	97.3	0.98	0.00092
	D. ostraviensis R			5'-CACATACTGCAGTGACCCCT-3'			
<i>Dactylogyrus vastator</i>	D. vastator F	210	60	5'-GTTGCGGAACCTGAACCTAGCCA-3'	98.99	0.95	0.00009
	D. vastator R			5'-AGACTGCACGACACGTTACCAA-3'			
<i>Dactylogyrus intermedius</i>	D. intermedius F	210	60	5'-TCAGAAATCTGAACCTATCCAATAC-3'	104.6	0.982	1.32E-07
	D. intermedius R			5'-TGCCGCACGACACGTTA-3'			


**Table 1.** Primers for *Dactylogyrus* spp. *ITS1* eDNA assay. The efficiency, R<sup>2</sup> and limit of detection for each quantitative PCR assay is provided. Primer cross-reactivity tests are provided in Support Information 1.

CNRS, Aix-Marseille Université: crn2m.univ-mrs.fr/pub/amplifx-dist). For Amplify4 and Amplifx 1.7.0 *in silico* tests, virtual PCRs were run against *ITS1* nucleotide sequences for all five target *Dactylogyrus* species. All assays demonstrated specificity to the targeted *Dactylogyrus* species across all three *in silico* tests. Primers were synthesized (standard desalting; Sigma-Aldrich, Australia), resuspended in 1x TE at 100 μM, and stored at −20 °C. Lastly, all qPCR assays were tested for species-specificity *in vitro* using both end-point PCR and qPCR with previously extracted genomic DNA (gDNA) from each target *Dactylogyrus* species<sup>10</sup>. All assays demonstrated specificity to the targeted *Dactylogyrus* species across all *in vitro* tests (Table 1; Supplementary Information 1), produced 120–210bp amplicons and performed optimally at assay-specific annealing temperatures (60 °C or 65 °C; Table 1).

Quantitative PCR assays (10 μL or 20 μL) contained 3 or 6 μL gDNA, 0.5 or 1 μL each PCR primer (400 nM), 5 or 10 μL PowerUP® SYBR GreenER qPCR Master Mix (Life Technologies, Australia) and 1 or 2 μL MilliQ® water, respectively, and were performed under the following fast cycling conditions (ramp rate = 2.70 °C/sec): UDG incubation at 50 °C for 2 min, initial denaturation at 95 °C for 2 min, 40 cycles of 95 °C denaturation for 15 sec then 60 or 65 °C primer-specific annealing for 60 sec (Table 1), and terminal dissociation curve generation (60–95 °C at 0.15 °C/sec). Previously extracted *Dactylogyrus* spp. gDNA<sup>10</sup> was quantified on a NanoDrop™ spectrophotometer (Invitrogen Inc.) and then each species-specific gDNA sample was serially diluted 1:10 to generate a five-point standard curve for each target *Dactylogyrus* species ( $1 \times 10^{-2}$ – $1 \times 10^{-6}$  ng/μL). Species-specific gDNA standards were used as template to determine assay amplification efficiency (E; i.e., increase in amplicon per cycle)<sup>29</sup> and limit of detection (LOD; i.e., lowest gDNA standard detected across all technical qPCR replicates) for each corresponding species-specific qPCR assay. All qPCR assays were run on a QuantStudio3™ Real-Time PCR System (ThermoFisher Scientific Inc., Brisbane), and threshold cycle value (C<sub>t</sub>) based on a common fluorescence threshold of 0.2. Melting temperature (T<sub>m</sub>) values were determined for each amplicon using QuantStudio™ Design and Analysis Software (version 1.4.2). All data was exported to Microsoft Excel for comparative analyses.

**Stepwise criteria for eDNA detection and samples tested for *Dactylogyrus* spp.** A four-step conservative predictive framework was developed to minimise the risk false positive and false negative results in qPCR T<sub>m</sub> analysis<sup>21,22,30</sup>. These criteria were selected considering the need to accurately determine absence from disease in biosecurity<sup>31</sup> and future applications of T<sub>m</sub> analysis to ensure accurate and reliable detection. For each qPCR assay the T<sub>m</sub> of each amplicon was compared to the mean T<sub>m</sub> of the corresponding species-specific gDNA, which was calculated from all technical qPCR replicates across the entire standard curve  $\pm 99.7\%$  CI<sup>32</sup>. The absolute difference between the mean T<sub>m</sub> of the species-specific gDNA standard curve and each individual qPCR technical replicate amplicon within a corresponding species-specific assay ( $|\Delta T_m|$ ) was calculated by subtracting the T<sub>m</sub> of each technical replicate amplicon from the mean T<sub>m</sub> of the corresponding species-specific gDNA standard. Calculated  $|\Delta T_m|$  values were then used to categorise each putative positive detection (i.e., amplicon) into one of three confidence levels: CL 1 = high (amplicon expected to be positive for *Dactylogyrus* spp. detection), CL 2 = medium (amplicon suspected to be positive for *Dactylogyrus* spp. detection), and CL 3 = low (amplicon predicted to not be positive for *Dactylogyrus* spp. detection, i.e., false positive) (Fig. 1).

Amplicons were categorized as CL 1 if: (1) amplification curves crossed the common threshold fluorescence within 40-cycles (Criterion 1.1, Fig. 1), (2) T<sub>m</sub> values were within 99.7% CI of the corresponding species-specific mean gDNA standard T<sub>m</sub> (Criterion 2: CL 1, Fig. 1), and (3) agarose gel visualization confirmed length to match that observed and expected for corresponding species-specific gDNA standard (Criterion 3, Fig. 1). Amplicons were categorized as CL 2 if they matched CL 1 criteria (see above) but exhibited a  $|\Delta T_m|$  outside 99.7% CI and  $\leq 1$  °C from mean T<sub>m</sub> of corresponding species-specific standards (Criterion 2: CL 2, Fig. 1). Amplicons were categorized as CL 3 if they matched CL 1 criteria but exhibited  $|\Delta T_m|$  outside 99.7% CI and  $> 1$  °C from mean T<sub>m</sub> of corresponding species-specific standard (Criterion 2: CL 3, Fig. 1). Putative positive CL 1, CL 2, and CL 3 amplicons were Sanger sequenced (Australian Genome Research Facility, Brisbane) for *Dactylogyrus* spp. level confirmation (NCBI BLAST; Criterion 4, Fig. 1). If any given *Dactylogyrus* spp. eDNA assay had  $\geq 2$  putative positive amplicons categorized as CL 1 or CL 2 then two representatives for each CL were chosen for Sanger sequencing (one with lowest and one with highest  $|\Delta T_m|$  value), otherwise one or both putative positive amplicons were sequenced. If any *Dactylogyrus* spp. eDNA assay had  $\geq 2$  putative positive amplicons categorized as CL 3 then the



**Figure 1.** Predictive framework designed to interpret qPCR amplicon data for eDNA detection determination.

amplicons with the lowest and highest  $|\Delta T_m|$  values (i.e., most and least likely to be confirmed as positive detections) were sequenced, otherwise both putative positive amplicons were sequenced.

Amplicons were considered to be putative false negative detections if no amplification curves were produced or failed to cross the common fluorescence threshold within 40 cycles (Criterion 1.2) but exhibited  $|\Delta T_m|$  values within 99.7% CI of mean  $T_m$  of corresponding species-specific standards (false negative, Fig. 1). Amplicons categorized as putative false negatives were re-amplified by qPCR to determine if a  $|\Delta T_m|$  value within 99.7% CI of mean  $T_m$  of corresponding species-specific standards and expected amplicon length were produced when amplified using 1  $\mu$ L of PCR product from initial amplification. Putative false negative amplicons were re-amplified using six replicate 20  $\mu$ L qPCRs containing 1  $\mu$ L of post-PCR product, 1  $\mu$ L of each PCR primer (400 nM), 10  $\mu$ L PowerUP® SYBR GreenER qPCR Master Mix (Life Technologies, Australia) and 8  $\mu$ L MilliQ® water, and were run under the same cycling conditions described above. Any amplicons produced from qPCR re-amplification that met Criteria 1, 2, and 3 (see above; Fig. 1) was Sanger sequenced for confirmation.

If an entire assay did not produce any amplicons that crossed common fluorescence threshold within 40 cycles (Criterion 1.2, Fig. 1) and no amplicons exhibited a discernible  $T_m$  then the entire assay was repeated. An assay was considered negative if neither initial or subsequent qPCR runs produced amplicons that crossed common fluorescence threshold within 40 cycles (Criterion 1.2, Fig. 1) and neither initial or subsequent qPCR runs produced amplicons with detectable  $T_m$  (Criterion 2, Fig. 1).

Species-specific qPCR assays were used to test extracted DNA in water samples from target and non-target fish consignments for the presence of *Dactylogyrus* spp. eDNA (Fig. 2). Imported consignments were considered 'target' or 'non-target' fish consignments based on published records of infection for any of the *Dactylogyrus* spp. targeted in this study ( $n = 5$ )<sup>5,24,25,33</sup>. Based on this criteria, seven goldfish (*Carassius auratus* (Linnaeus, 1758)) consignments were considered targets for *D. anchoratus*, *D. formosus*, *D. intermedius*, and *D. vastator* whereas four rosy barb (*Pethia conchonius* (Hamilton, 1822)) consignments were considered targets for *D. ostraviensis* (Fig. 2). Based on the same criteria, one guppy (*Poecilia reticulata*, Peters 1859), one pearl gourami (*Trichopodus leerii* (Bleeker, 1852)), one three-spot gourami (*Trichopodus trichopterus* (Pallas, 1770)), one green swordtail (*Xiphophorus hellerii* Heckel, 1848), and three platyfish (*Xiphophorus maculatus* (Günther, 1866)) consignments were considered non-target hosts for all five *Dactylogyrus* species. All target and non-target host fish consignments were screened for the presence of eDNA from all five *Dactylogyrus* species using species-specific qPCR assays (Fig. 2) followed by assessment of each produced amplicon based the selection criteria described above (Fig. 1).

## Results

**Positive *Dactylogyrus* spp. eDNA detection in target fish consignments.** *Dactylogyrus* spp. eDNA was detected in all consignments where *Dactylogyrus* spp. were detected by standard necropsies. Specifically, eDNA from *D. formosus* and *D. vastator* was detected in water samples from all *C. auratus* consignments, and eDNA from *D. anchoratus* and *D. intermedius* was detected in all consignments except for consignments 4 and 6, respectively (Fig. 2). *Dactylogyrus anchoratus* was detected by both approaches (eDNA and necropsy) in consignments 6 and 7 while neither approach detected parasites in consignment 4. *Dactylogyrus ostraviensis* eDNA was detected in all target *P. conchonius* consignments, while necropsies did not detect *D. ostraviensis* in consignment 12 (Fig. 2). *Dactylogyrus* spp. eDNA was detected in five *C. auratus* and one *P. conchonius* consignments considered to have *Dactylogyrus* spp. apparent prevalence of 0% (95% CI 0–11.4%) by necropsy<sup>10</sup> (Fig. 2). No eDNA was detected in negative controls.

**Positive *Dactylogyrus* spp. eDNA detections in non-target fish consignments.** A total of 39 amplicons produced across all 58 qPCR tests of non-target fish consignments were confirmed positive for *Dactylogyrus* spp. eDNA (Fig. 2). *Dactylogyrus formosus*, *D. intermedius*, and *D. vastator* eDNA was detected



Fish pop.	Fish species	Exporter	<i>Dactylogyrus anchoratus</i>		<i>Dactylogyrus formosus</i>		<i>Dactylogyrus intermedius</i>		<i>Dactylogyrus vastator</i>		<i>Dactylogyrus ostraviensis</i>	
			Necropsy	eDNA	Necropsy	eDNA	Necropsy	eDNA	Necropsy	eDNA	Necropsy	eDNA
3	<i>Carassius auratus</i>	Singapore 2	-	4/6*	-	4/6*	-	4/6*	-	4/6*	-	6/6
4	<i>Carassius auratus</i>	Singapore 2	-	0/12	-	4/6*	-	6/6*	-	6/6*	-	5/6
5	<i>Carassius auratus</i>	Thailand 1	-	4/6*	-	6/6*	20 (8–39)	5/6	13.3 (2–27)	6/6	-	-
6	<i>Carassius auratus</i>	Thailand 1	26.6 (12–46)	6/6	6.7 (0.82–22)	5/6	43.3 (25–63)	0/12	40 (23–59)	4/6	-	-
7	<i>Carassius auratus</i>	Thailand 1	3.3 (0.1–17.2)	4/6	43.3 (25–63)	4/6	30 (15–49)	6/6	16.6 (2–29)	6/6	-	-
8	<i>Carassius auratus</i>	Malaysia 1	-	1/6*	3.3 (0.1–17.2)	3/3	3.3 (0.1–17.2)	3/3	3.3 (0.1–17.2)	3/3	-	-
9	<i>Carassius auratus</i>	Malaysia 1	-	5/6*	6.6 (0.82–22)	4/6	3.3 (0.1–17.2)	4/6	-	4/6*	-	-
13	<i>Pethia conchonius</i>	Singapore 2	-	-	-	1/6	-	1/6	-	2/6	26.6 (12–46)	6/12
14	<i>Pethia conchonius</i>	Thailand 1	-	-	-	-	-	-	-	-	33.3 (17–53)	4/6
15	<i>Pethia conchonius</i>	Thailand 2	-	-	-	-	-	-	-	1/6	73.3 (54–88)	4/6
16	<i>Pethia conchonius</i>	Malaysia 1	-	-	-	-	-	-	-	1/6	-	2/12*
17	<i>Poecilia reticulata</i>	Sri Lanka 2	-	-	-	2/6	-	1/6	-	-	-	-
18	<i>Trichopodus leerii</i>	Sri Lanka 1	-	-	-	-	-	-	-	1/6	-	5/6
19	<i>Trichopodus trichopterus</i>	Thailand 1	-	-	-	-	-	-	-	-	-	-
22	<i>Xiphophorus helleri</i>	Sri Lanka 2	-	-	-	-	-	-	-	-	-	-
23	<i>Xiphophorus maculatus</i>	Thailand 1	-	-	-	-	-	1/6	-	2/6	-	-
24	<i>Xiphophorus maculatus</i>	Singapore 2	-	-	-	-	-	1/6	-	-	-	5/6
25	<i>Xiphophorus maculatus</i>	Sri Lanka 2	-	-	-	4/6	-	-	-	-	-	-

**Figure 2.** Comparison between necropsies and environmental DNA (eDNA) detection of *Dactylogyrus* species in imported ornamental fish consignments. Detections by necropsy presented as mean apparent prevalence % (95% Confidence Interval, CI)<sup>10</sup> and eDNA detections as confirmed positive amplicons/total number of amplicons. Grey areas indicate qPCR assays of target fish consignments, and asterisks (\*) indicate consignments where *Dactylogyrus* spp. were not detected by necropsies but were detected by eDNA assays. Negative symbols (–) indicate that no parasites were detected in a total of 30 fish and had an apparent prevalence = 0% (95% CI = 0–11.4%)<sup>10</sup>, and that no parasite eDNA was detected from a total of six eDNA sample replicates.


in *P. conchonius* consignment 13 (Singapore 2; Fig. 2). *Dactylogyrus intermedius* and *D. ostraviensis* eDNA was detected in *X. maculatus* consignment 24 (Singapore 2, Fig. 2) while *D. vastator* and *D. intermedius* eDNA was detected in *X. maculatus* consignment 23 (Thailand 1; Fig. 2). Similarly, *D. ostraviensis* eDNA was detected in *C. auratus* consignments 3 and 4 as well as *X. maculatus* consignment 24 (Singapore 2; Fig. 2). Lastly, *D. formosus*, *D. intermedius*, *D. vastator*, and *D. ostraviensis* eDNA was detected by qPCR in *P. reticulata* consignment 17, *T. leerii* consignment 18, and *X. maculatus* consignment 25 (Sri Lanka; Fig. 2). No *Dactylogyrus* spp. were reported in non-target fish consignments by necropsies<sup>10</sup>.

**Accuracy of predictive framework.** All amplicons categorized as high confidence of *Dactylogyrus* detection (CL 1) from all *Dactylogyrus* spp. qPCR assays were confirmed positive by Sanger sequencing (Fig. 1 Criterion 4). All amplicons categorized as moderate confidence (CL 2) from *D. anchoratus*, *D. formosus*, and *D. intermedius* qPCR assays were also confirmed positive by Sanger sequencing (Fig. 1 Criterion 4). Of the amplicons categorized as CL 2 from *D. ostraviensis* and *D. vastator* qPCR assays, 80% and 87.5% ( $n = 4/5$  and  $7/8$ ) were confirmed positive by Sanger sequencing, respectively. These two CL 2 amplicons were unable to be confirmed as positive detections due to poor sequencing quality (i.e., not due to non-target amplification; see Fig. 3D for *D. ostraviensis* and Fig. 4 for *D. vastator*).


No low confidence (CL 3) categorized amplicons from *D. anchoratus*, *D. formosus*, *D. intermedius*, or *D. ostraviensis* qPCR assays were confirmed positive by Sanger sequencing. However, 81.25% ( $n = 13/16$ ) of CL 3 categorized amplicons from *D. vastator* qPCR assays were confirmed positive by Sanger sequencing (Fig. 1, Criterion 4). One *D. vastator* qPCR assay amplicon from *T. trichopterus* consignment 14 was initially considered a putative false negative (Fig. 1 Criterion 2) but was subsequently categorized as CL 1 following qPCR reamplification (Fig. 1) and confirmed positive by Sanger sequencing (Fig. 1 Criterion 4, Fig. 4 “amplicon 19\_4”). All other putative false negative amplicons produced during *Dactylogyrus* spp. eDNA assays were confirmed negative following the selective framework (Fig. 1, Support Information 2).

**Amplicon sequence confirmation.** All confirmed positive *D. anchoratus* amplicons were 100% homologous to *D. anchoratus ITS1* GenBank sequences (AJ564111, AJ490161, MF356241, KY859795, MF662103, MF356243, and MF356242). All confirmed positive *D. formosus* amplicons were 100% homologous to *D. formosus ITS1* GenBank sequences (AJ564135, MF356239, KM525669, KX369215, and KC876018). All confirmed positive *D. intermedius* amplicons were 100% homologous to *D. intermedius ITS1* GenBank sequences (KC876017, KX369220, MF356236, MF356244, KJ854364, MF356237, and MF356240). All confirmed positive *D. ostraviensis* amplicons were 100% homologous to *D. ostraviensis ITS1* GenBank sequences (MF356250 and MF356249; which are the only two sequences available)<sup>10</sup>.

Confirmed positive *D. vastator* amplicons, unlike all other *Dactylogyrus* spp. amplicons, separated into two distinct groups (Fig. 4). *Dactylogyrus vastator* Group 1 amplicons exhibited an average  $T_m \pm SD$  of  $86.64^\circ\text{C} \pm 0.59$  with average  $|\Delta T_m|$  being  $\pm 0.6^\circ\text{C}$  away from  $T_m$  of gDNA standards ( $|\Delta T_m|$ ; Fig. 4), while amplicons in Group 2 exhibited an average  $T_m \pm SD$  of  $85.37^\circ\text{C} \pm 0.47$  with average  $|\Delta T_m|$  being  $\pm 1.97^\circ\text{C}$  away from  $T_m$  of gDNA standards (Fig. 4). The six confirmed positive *D. vastator* amplicons that fell within the 99.7% CI of *D. vastator* gDNA standards (Group 1) were 98–100% homologous to the following *D. vastator ITS1* GenBank sequences: MF356235 (Thailand), KY207446 (Croatia), AJ564159 (Czech Republic), MF806586 (Iran), MF356246 (Thailand), KY201104 (Italy), and KY201092 (Bosnia and Herzegovina). The 11 positive *D. vastator* amplicons that fell outside the 99.7% CI of the same *D. vastator* gDNA standards (Group 2) were 96–100% homologous to



**Figure 3.** Absolute difference in melting temperature ( $|\Delta T_m|$ ) between sequenced amplicons derived from environmental DNA assays for *Dactylogyrus anchoratus* (A), *Dactylogyrus formosus* (B), *Dactylogyrus intermedius* (C) and *Dactylogyrus ostraviensis* (D) and their corresponding genomic DNA standards. Grey and black bars in Panels A–D represent confirmed positive and confirmed negative amplicons, respectively. Horizontal dotted lines in Panels A–D represent the upper 99.7% Confidence interval for  $T_m$  of species-specific standards. \*\*Forward and reverse sequences were low in quality; however, a 72 bp fragment of consensus alignment was found to be 100% similar to *Cyprinus carpio* GenBank sequence LN599613 (i.e. considered as confirmed negative).



**Figure 4.** Absolute difference in melting temperature ( $|\Delta T_m|$ ) between *Dactylogyrus vastator* amplicons derived from environmental DNA (eDNA) assays and genomic DNA (gDNA) standards confirmed by Sanger sequencing. Grey and black bars represent confirmed positive and confirmed negative amplicons, respectively. Horizontal dotted lines represent the upper 99.7% CI for  $T_m$  of serially diluted *D. vastator* gDNA standard, while Group 2 amplicons had 2–18 base pair differences between sequences obtained compared to *D. vastator* gDNA. Asterisk (\*): consensus sequence could not be determined for this amplicon because reverse sequence failed; however, forward sequence had 93.8% similarity to *Contracaecum* sp. [GenBank accession KM463761] and 91% similarity to *Contracaecum rudolphii* Hartwich, 1964 [GenBank accession JQ071409] and thus this amplicon was considered as a confirmed negative detection. ClustalW alignment of all *D. vastator ITS1* amplicon sequences provided in Support Information 3.

the following *D. vastator ITS1* GenBank sequences: KX369223 (China), MF356247 (Thailand), KY201103 (Czech Republic), and KM487695 (China). Groups 1 and 2 *D. vastator* amplicons differed by a total of 16 fixed nucleotide differences (Support Information 3).

## Discussion

The developed qPCR assays detected *Dactylogyrus* spp. eDNA in all consignments where necropsies detected *Dactylogyrus* spp.<sup>10</sup>. Species-specific qPCR assays were able to detect *Dactylogyrus* spp. eDNA in six target fish consignments where necropsies considered *Dactylogyrus* spp. to have an apparent prevalence of 0% (95% CI 0–11.4; Fig. 2). As such, qPCR-based eDNA detection had higher surveillance sensitivity than necropsies, detecting *Dactylogyrus* spp. DNA in triplicate 15 mL water samples and confirming amplicons by Sanger sequencing.

However, *D. intermedius*, which was reported to infect *C. auratus* in consignment 6 by necropsy<sup>10</sup> was not detected by eDNA screening in any qPCR technical replicates ( $n = 12$ ; Fig. 2). Consequently, this was the only false negative eDNA detection observed in this study (1/90 tests; Fig. 2). It is possible that *D. intermedius* present in consignment 6 were genetically distinct from *D. intermedius* infecting consignments 5, 7, 8 and 9 (Fig. 2). The possibility of unique ITS1 genotypes in *D. intermedius* is supported by sequenced data of *D. vastator*, which displayed two ITS1 genotypes observed across screened goldfish consignments (Fig. 4; Support Information 3). Unlike the *D. vastator* assay, the *D. intermedius* assay appears to target an ITS1 region that is sufficiently hyper-variable to prevent primer binding<sup>27,34,35</sup>; however, this was unknown at the time of assay development due to limited nucleotide sequence information available for *D. intermedius* populations. Such a lack of comprehensive nucleotide sequence information has also limited other molecular genetic studies aimed at investigating parasite diversity<sup>36,37</sup>. As such, successful implementation of the four-step predictive framework relied on the comprehensiveness of species-specific gDNA standards, suggesting  $|\Delta T_m|$  analysis requires careful interpretation given the inherent dependence on sequence homology between amplicons and standards for targeted gene(s) that may or may not be known. This study highlights the need for comprehensive nucleotide sequence data and robust corresponding morphological taxonomy to ensure accuracy of designed qPCR assays and corresponding standards for  $|\Delta T_m|$  analyses.

A total of 39 amplicons from non-target fish consignments were confirmed positive for *Dactylogyrus* spp. eDNA (Fig. 2). Considering that all *Dactylogyrus* spp. in this study are highly specific to cyprinid species<sup>24,25,33</sup>, positive detections in water samples from non-target consignments suggest that detected eDNA was not present due to active shedding from live infesting *Dactylogyrus* parasites. This interpretation is further supported by the absence of infection records for the selected *Dactylogyrus* specimens in non-target host fish species<sup>24,25</sup> and non-detection by necropsies (Fig. 2). *Dactylogyrus* spp. are ectoparasites that occur naturally in southeast Asia<sup>5,38</sup> and their environmental stages (i.e., eggs and oncomiracidia) could be present in recirculating aquaculture systems, raceways, or ponds used to rear freshwater species by exporting companies. As such, it is possible that exporters could have used a water source contaminated with *Dactylogyrus* spp. environmental life stages<sup>12</sup> or degraded eDNA to transport exported fish consignments. If exporters do not use clean (e.g., filtered or UV treated) water to export ornamental fish consignments, then the accuracy and interpretability of eDNA assays at border control is limited, given that their applicability would depend greatly in differentiating between live infections and dead or inactive environmental parasite stages in the water column. Furthermore, considering that Australian quarantine officers have limited time to process imported consignments, eDNA-based detection by qPCR may not be applicable or reliable at border control using  $T_m$  analysis to carefully interpret qPCR results within an acceptable timeframe and biosecurity standard.

Screening water samples for parasite eDNA by qPCR could be a valuable detection method during pre-export quarantine periods. Current risk analyses from the Australian Government Department of Agriculture and Water Resources aim to ensure off-shore biosecurity in exporting countries<sup>39</sup> by enforcing strict regulations and health requirements prior to export<sup>9</sup>. For example, all imported goldfish consignments must be certified free of infection from gill flukes *Dactylogyrus extensus* and *D. vastator* prior to export<sup>9</sup>. Both species are reported to cause significant economic losses in Asian cyprinid aquaculture<sup>1,40</sup>, and could pose significant risks to Australian aquarium shops selling cyprinids if live parasite infections go undetected during quarantine<sup>1</sup>.

Detection of eDNA by qPCR assays could be conducted on ornamental fish consignments during the mandatory quarantine period prior to export to support mandatory pre-export health certifications<sup>9</sup>. For instance, qPCR assays could be developed to assess the origin of parasite eDNA based on DNA decay rates by targeting various DNA fragment lengths<sup>41–43</sup>. Abundant long DNA fragments would indicate active shedding from live parasites while abundant short DNA fragments would indicate degrading DNA in the absence of live, shedding organisms<sup>41,42</sup>. Similarly, qPCR assays could also assess cellular activity by targeting environmental RNA (eRNA)<sup>12,43,44</sup>. Environmental RNA is indicative of active gene transcription and is proportionally less abundant in dormant stages than in metabolically active stages<sup>12</sup>. Given that RNA is less able to persist extracellularly and degrades quickly in dead or sloughed-off cells<sup>12</sup>, detection of eRNA by qPCR could be employed to determine the presence of metabolically active parasites infecting fish ready for export. Future research should consider designing qPCR assays to differentiate between active parasite infections and dead or non-active parasite stages and the applicability of eDNA detection during pre-export quarantine periods.

In conclusion, this first attempt at applying eDNA to ornamental fish parasite biosecurity highlights both the utility of incorporating molecular methods into biosecurity protocols as well as the limitations that need to be addressed if future applications and full integration are to be successful. We present a novel and comprehensive four-step predictive framework (Fig. 1) for the accurate interpretation of species-specific eDNA data and reduce false positive and false negative detections generated by Sybr-based qPCR assays. The interpretability and reliability of eDNA detection at border control specifically is limited; however, eDNA screening could prove highly valuable if implemented following pre-export quarantine periods. Further research needs to address limitations encountered in this study and test the viability of eDNA-based detection methods in other stages of quarantine and biosecurity surveillance.

## Data Availability

Data for this study can be accessed as: Trujillo Gonzalez, A. (2018). Parasite detection in the ornamental fish trade using environmental DNA. James Cook University. [Data Files] <https://doi.org/10.25903/5b90c1897397a>.

## References

- Kahn, S. A., Wilson, D. W., Perera, R. P., Hayder, H. & Gerrity, S. E. Import risk analysis on live ornamental finfish. *Australian Quarantine and Inspection Service, Canberra, Australia*, pp 172 (1999).
- Corfield, J. *et al.* Review of the impacts of introduced aquarium fish species that have established wild populations in Australia. Prepared for the Australian Government Department of the Environment and Water Resources, ID number: 49547, pp 278 (2007).
- Whittington, R. & Chong, R. Global trade in ornamental fish from an Australian perspective: the case for revised import risk analysis and management strategies. *Prev. Vet. Med.* **81**, 92–116 (2007).
- Chang, A. L. *et al.* Tackling aquatic invasions: Risks and opportunities for the aquarium fish industry. *Biol. Inv.* **11**, 773–785 (2009).
- Trujillo-González, A., Becker, J. A. & Hutson, K. S. Parasite dispersal in the goldfish trade. *Adv. Parasitol.* **100**, 239–281 (2018).
- Lymbery, A. J., Morine, M., Kanani, H. G., Beatty, S. J. & Morgan, D. L. Coinvaders: the effects of alien parasites on native hosts. *Int. J. Parasitol. Par. Wild.* **3**, 171–177 (2014).
- Hine, P. Problems of applying risk analysis to aquatic organisms. In: Rogers, C. (Ed.), *Risk Analysis in Aquatic Animal Health*. Office International des Epizooties, Paris, pp. 71–82 (2001).
- Becker, J. A., *et al.* Aquatic Animal Health Subprogram: Strategic approaches to identifying pathogens of quarantine concern associated with the importation of ornamental fish. Fisheries Research and Development Corporation, Project No. 2014/001, pp. 83 ISBN 978-1-74210-399-0 (2016).
- [BICON] Biosecurity Import Conditions System (BICON) In: Australian Department of Agriculture and Water Resources. Accessed 13 March 2018, [www.agriculture.gov.au/import/online-services/bicon](http://www.agriculture.gov.au/import/online-services/bicon) (2018).
- Trujillo-González, A., Becker, J. A., Vaughan, D. B. & Hutson, H. S. Monogenean parasites infect ornamental fish imported to Australia. *Parasitol. Res.* **117**, 995–1011 (2018).
- Thomsen, P. & Willerslev, E. Environmental DNA - an emerging tool in conservation for monitoring past and present biodiversity. *Biol. Conserv.* **183**, 4–18 (2015).
- Bass, D., Stentiford, G. D., Littlewood, D. T. J. & Hartikainen, H. Diverse applications of environmental DNA methods in parasitology. *Trends Parasitol.* **31**, 499–513, <https://doi.org/10.1016/j.pt.2015.06.013> (2015).
- Barnes, M. A. & Turner, C. R. The ecology of environmental DNA and implications for conservation genetics. *Conserv. Gene.* **17**, 1–17 (2016).
- Goldberg, C. S. *et al.* Critical considerations for the application of environmental DNA methods to detect aquatic species. *Methods Ecol. Evol.* **7**, 1299–1307 (2016).
- Huwer, J. R., Koprivnikar, J., Johnson, P. T. J. & Whyard, S. Development and application of an eDNA method to detect and quantify a pathogenic parasite in aquatic ecosystems. *Ecol. Appl.* **25**, 991–1002, <https://doi.org/10.1890/14-1530.1> (2015).
- Bastos-Gomes, G. *et al.* Use of environmental DNA (eDNA) and water quality data to predict protozoan parasites outbreaks in fish farms. *Aquaculture* **479**, 467–473 (2017).
- Hallett, S. L. *et al.* Density of the waterborne parasite *Ceratomyxa shasta* and its biological effects on salmon. *Appl. Environ. Microbiol.* **78**, 3724–3731 (2012).
- Agawa, Y., Tani, K., Yamamoto, S., Hirano, C. & Shirakashi, S. Development of a quantitative polymerase chain reaction assay for the detection of skin fluke *Neobenedenia girellae* larvae from environmental water. *Fish. Sci.* **82**, 827–833 (2016).
- Rusch, J. C. *et al.* Catching the fish with the worm: a case study on eDNA detection of the monogenean parasite *Gyrodactylus salaris* and two of its hosts, Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*). *Parasit. Vectors.* **11**, 1–12 (2018).
- Collins, R. A., Armstrong, K. F., Holyoake, A. J. & Keeling, S. Something in the water: biosecurity monitoring of ornamental fish imports using environmental DNA. *Biol. Inv.* **15**, 1209–1215 (2013).
- Schmidt, B. R., Kéry, M., Ursenbacher, S., Hyman, O. J. & Collins, J. P. Site occupancy models in the analysis of environmental DNA presence/absence surveys: a case study of an emerging amphibian pathogen. *Methods Ecol. Evol.* **4**, 646–653 (2013).
- Lahoz-Monfort, J. J., Guillera-Aroita, G. & Tingley, R. Statistical approaches to account for false-positive errors in environmental DNA samples. *Mol. Ecol. Resour.* **16**, 673–685 (2016).
- Ficetola, G. F. *et al.* Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Mol. Ecol. Resour.* **15**, 543–556 (2015).
- Whittington, I. D., Cribb, B. W., Hamwood, T. E. & Halliday, J. E. Host-specificity of monogenean (platyhelminth) parasites: a role for anterior adhesive areas? *Int. J. Parasitol.* **30**, 305–20 (2000).
- Cribb, T. H., Chisholm, L. A. & Bray, R. A. Diversity in the Monogenea and Digenea: does lifestyle matter? *Int. J. Parasitol.* **32**, 321–328 (2002).
- Minamoto, T. *et al.* Nuclear internal transcribed spacer-1 as a sensitive genetic marker for environmental DNA studies in common carp *Cyprinus carpio*. *Mol. Ecol. Resour.* **17**, 324–333 (2017).
- Van Herwerden, L., Blair, D. & Agatsuma, T. Intra- and interindividual variation in ITS1 of *Paragonimus westermani* (Trematoda: Digenea) and related species: Implications for phylogenetic studies. *Mol. Phyl. Evol.* **12**, 67–73 (1999).
- Johnson, M. *et al.* NCBI BLAST: a better web interface. *Nucleic Acids Res.* **36**, W5–W9 (2008).
- Ruijter, J. M. *et al.* Amplification efficiency: Linking baseline and bias in the analysis of quantitative PCR data. *Nucleic Acids Res.* **37**, e45:1–12 (2009).
- Davidson, A. D. & Hewitt, C. L. How often are invasion-induced ecological impacts missed? *Biol. Inv.* **16**, 1165–1173 (2014).
- [OIE] Aquatic Animal Health Code. In: World Organization for Animal Health. Accessed on the 14th of February, <http://www.oie.int/en/standard-setting/aquatic-code/access-online/> (2019).
- Ririe, K. M., Rasmussen, R. P. & Wittwer, C. T. Product differentiation by analysis of DNA melting curves during the polymerase chain reaction. *Analyt. Biochem.* **245**, 154–160 (1997).
- Řehulka, J. *Dactylogyrus ostraviensis* n. sp. (Dactylogyridae: Monogenea) from the gills of *Barbus conchoniensis*. *Syst. Parasitol.* **12**, 77–80, <https://doi.org/10.1007/BF00182030> (1988).
- Van Herwerden, L., Caley, M. J. & Blair, D. Regulatory motifs are present in the ITS1 of some flatworm species. *J. Exp. Zool. Mol. Develop. Evol.* **296B**, 80–86 (2003).
- Warberg, R., Jensen, K. T. & Frydenberg, J. Repetitive sequences in the ITS1 region of ribosomal DNA in congeneric microphallid species (Trematoda: Digenea). *Parasitol. Res.* **97**, 420–423 (2005).
- Gómez, F. Problematic biases in the availability of molecular markers in protists: The example of the dinoflagellates. *Acta Protozool.* **53**, 63–75, <https://doi.org/10.4467/16890027AP.13.0021.1118> (2014).
- Palesse, S., Meadors, W. A., de Buron, I., Roumillat, W. A. & Strand, A. E. Use of molecular tools in identification of philometrid larvae in fishes: Technical limitations parallel our poor assessment of their biodiversity. *Parasitol. Res.* **109**, 1725–1730, <https://doi.org/10.1007/s00436-011-2481-6> (2011).
- Ling, F., Tu, X., Huang, A. & Wang, G. Morphometric and molecular characterization of *Dactylogyrus vastator* and *D. intermedius* in goldfish (*Carassius auratus*). *Parasitol. Res.* **115**, 1755–1765, <https://doi.org/10.1007/s00436-016-4913-9> (2016).



39. Hood, Y., Sadler, J., Poldy, J., Starkey, C. S. & Robinson, A. P. Biosecurity system reforms and the development of a risk-based surveillance and pathway analysis system for ornamental fish imported into Australia. *Prev. Vet. Med.* in press, <https://doi.org/10.1016/j.prevetmed.2018.11.006> (2018).
40. Zhang, X. P. *et al.* The efficacy of four common anthelmintic drugs and traditional Chinese medicinal plant extracts to control *Dactylogyrus vastator* (monogenea). *Aquaculture* **420–421**, 302–307 (2014).
41. Bylemans, J., Furlan, E. M., Gleeson, D. M., Hardy, C. M. & Duncan, R. P. Does size matter? an experimental evaluation of the relative abundance and decay rates of aquatic environmental. *DNA. Environ. Sci. Tech.* **52**, 6408–6416, <https://doi.org/10.1021/acs.est.8b01071> (2018).
42. Robson, H. *et al.* Fine-tuning for the tropics: application of eDNA technology for invasive fish detection in tropical freshwater ecosystems. *Mol. Ecol.* **16**, 922–932 (2016).
43. Pochon, X., Zaiko, A., Fletcher, L., Laroche, O. & Wood, S. Wanted dead or alive? using metabarcoding of environmental DNA and RNA to distinguish living assemblages for biosecurity applications. *Plos One* **12**, e0187636, <https://doi.org/10.1371/journal.pone.0187636> (2017).
44. Zaiko, A., Pochon, X., Garcia-Vazquez, E., Olenin, S. & Wood, S. A. Advantages and limitations of environmental DNA/RNA tools for marine biosecurity: Management and surveillance of non-indigenous species. *Front. Mar. Sci.* **5**, <https://doi.org/10.3389/fmars.2018.00322> (2018).

## Acknowledgements

We thank Dr Terrence Miller, Dr Paul Hick, Alison Tweedie, and Joshua Allas for their assistance during sampling events, and Dr Giana Bastos-Gomes and Julie Goldsbury for their assistance and comments on DNA extraction methods. We thank Dr Lynne van Herwerden from James Cook University for her useful comments. This study was funded by the Australian Government through the Fisheries Research and Development Corporation (FRDC) (Project No. 2014/001), the University of Sydney, and James Cook University.

## Author Contributions

Alejandro Trujillo-González wrote the main manuscript text, prepared all figures, extracted all DNA associated to the study, designed all qPCR assays and ran all qPCR tests and analyses. Dr Richard C. Edmunds contributed to the editing process of the manuscript text, designed the primers used in all qPCR assays and provided advice during qPCR testing and analysis. Dr Joy A. Becker contributed to the editing process of the manuscript text, sample collection, and was the primary researcher of the FRDC grant 2014/001: Aquatic Animal Health Subprogram: Strategic approaches to identifying pathogens of quarantine concern associated with the importation of ornamental fish. This grant funded all materials and equipment needed for sampling and qPCR analysis. Dr Kate S. Hutson is the primary supervisor of Alejandro Trujillo-González, contributed to the editing process of the manuscript text, sample collection, provided advice during qPCR analysis and was the secondary researcher of the FRDC grant 2014/001.

## Additional Information

**Supplementary information** accompanies this paper at <https://doi.org/10.1038/s41598-019-41517-2>.

**Competing Interests:** The authors declare no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019