

Microtopographic refuges enhance recruitment and survival but inhibit growth of propagules of the tropical macroalga *Sargassum swartzii*

Zoe Loffler*, Andrew S. Hoey

ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, QLD 4811, Australia

ABSTRACT: For organisms with discrete life-histories, any mechanism that enhances growth and/or survival at critical life-history transitions may significantly influence adult population size. On coral reefs, structurally complex microhabitats enhance the early post-settlement growth and/or survival of fishes and corals; however, the importance of such microhabitats to the early life stages of tropical macroalgae is largely unknown. Here, we investigate the effect of crevices on the recruitment, growth and survival of propagules of a common coral reef macroalga, *Sargassum swartzii*. We settled *S. swartzii* propagules onto terracotta settlement tiles that had a series of 3 mm deep crevices evenly spaced on their top (exposed) surface. Recruitment of *S. swartzii* was 21 % greater, but propagules were 18 % shorter, 18 d post-settlement within crevices than on adjacent exposed surfaces. Exposing tiles to local fish assemblages for 5 d showed that survival of propagules was 90 % greater in crevices than on exposed areas of the tiles on the reef crest, but not on the reef flat. Underwater video footage revealed that few fishes fed from within the crevices (18 % of all bites), with the majority of feeding being concentrated on the exposed surface of tiles. Interestingly, small-bodied fishes from the family Blenniidae (predominantly *Ecsenius* spp.) accounted for the majority of the feeding activity on the tiles and likely contributed to the mortality of propagules. Structurally complex microhabitats, such as crevices, that shelter vulnerable early post-settlement propagules from herbivory may therefore be important for the persistence of macroalgae on coral reefs.

KEY WORDS: Herbivory · Coral reef · Refuge · Early post-settlement · Blenniidae

Resale or republication not permitted without written consent of the publisher

1. INTRODUCTION

For organisms with discrete life history stages, processes affecting dispersal, settlement and early post-settlement survival can have a large influence on adult population size (Connell 1985, Vadas et al. 1992). Indeed, for many marine organisms, the first few days following settlement have been identified as a critical period shaping adult populations, as recently settled individuals are typically small (Vadas et al. 1992, Hunt & Scheibling 1997), undergo a range of physiological changes (McCormick et al. 2002), and are exposed to a diverse suite of predators for the

first time (Vadas et al. 1992, Almany & Webster 2006, Scheibling & Robinson 2008). For example, mortality in the first days to weeks after settlement has been reported to be as high as 58 to 96.3% for coral recruits (Vermeij & Sandin 2008, Traçon et al. 2013, Doropoulos et al. 2017a), up to 75% for coral reef fishes (Hoey & McCormick 2004, Almany & Webster 2006), >90% for many other marine invertebrates (Gosselin & Qian 1997, Hunt & Scheibling 1997) and up to 99% for temperate marine algae (Vadas et al. 1992, Kendrick & Walker 1994). Factors that mediate rates of mortality at this critical life history transition include the size of an individual (i.e. 'bigger-is-better'

*Corresponding author: zoe.loffler@my.jcu.edu.au

hypothesis; Miller et al. 1988, Bailey & Houde 1989), growth rate (i.e. 'stage duration' hypothesis; Leggett & Deblois 1994), physiological condition (e.g. Hoey & McCormick 2004), inter- and intra-specific competition (Menge & Sutherland 1976, Bonin et al. 2009), and the structural complexity and suitability of the settlement habitat (Diaz-Pulido et al. 2010, Yadav et al. 2016, Doropoulos et al. 2017b, Mallela 2018). Of these, structural complexity of the settlement habitat is thought to be a major driver of rates of survival (Tupper & Boutilier 1997, Nozawa 2008, Brandl & Bellwood 2016, Mallela 2018).

The structural complexity of a habitat (i.e. its 3-dimensional structure, composed of living and dead organisms and abiotic substrata) can mediate rates of mortality through the provision of refugia (Crowder & Cooper 1982, Graham & Nash 2013). The value of structural complexity in providing refugia is dependent on the relative size of available refuges: large enough to allow prey to access, but small enough to exclude predators (Menge & Lubchenco 1981). Therefore, small-scale structural complexity, such as small cracks and crevices, may allow recently settled organisms to avoid the predation pressure of adjacent open or exposed surfaces (Menge & Lubchenco 1981, Brandl et al. 2014, Franco et al. 2015).

On coral reefs, early post-settlement mortality is high (>50%) for both fish and corals (Almany & Webster 2006, Nozawa 2008) and this period is considered critical in the life history of these organisms. Importantly, however, specific microhabitats can enhance the early post-settlement survival of corals and reef fish several-fold, presumably through the provision of refugia (Nozawa 2008, Bonin et al. 2009, Brandl et al. 2014). For example, Nozawa (2008) reported the complete mortality of corals that had settled on exposed surfaces within 4 mo, while up to 12% of corals that had settled in crevices survived the first year post-settlement. Similarly, the early post-settlement survival of coral reef fishes is also enhanced when fishes settle on patch reefs with higher structural complexity (Bonin et al. 2009). Whilst the importance of microhabitats to the settlement and survival of corals and fishes has been relatively well-studied on coral reefs (Nozawa 2008, Bonin et al. 2009, Traçon et al. 2013, Doropoulos et al. 2016), the importance of microhabitats for early life stages of macroalgae has received less attention (Brandl et al. 2014, Poray & Carpenter 2014).

To date, studies investigating spatial and temporal variation in rates of herbivory on coral reef macroalgae have primarily focused on the consumption of adult thalli (Hay 1981, Lewis 1986, Loffler et al.

2015), yet it is often assumed that macroalgal biomass is largely controlled through the grazing of early life stages of macroalgae (Done 1992, Diaz-Pulido & McCook 2003, Bellwood et al. 2004). Spatial and temporal refuges may facilitate the early post-settlement survival of macroalgae by providing a refuge from herbivory through reductions in grazing pressure (Brandl et al. 2014, Brandl & Bellwood 2016), potentially allowing propagules to grow to a size more resistant to grazing (Lubchenco 1983, Stiger et al. 2004, Briggs et al. 2018). In this study, therefore, we investigate how reef microtopography (i.e. crevices) influences the recruitment (defined here as settlement and any mortality during the first 18 d), growth and survival of propagules of the common tropical macroalga *Sargassum swartzii*.

2. MATERIALS AND METHODS

2.1. Study organism

Sargassum is a genus of tall (up to 3 m), leathery, brown macroalgae common on many coastal and/or degraded coral reefs (Done 1992, Chong-Seng et al. 2014). *Sargassum* reproduces through the release of tens of thousands of zoospores per plant (Diaz-Pulido & McCook 2003), with mortality of settled propagules in temperate *Sargassum* near absolute (>99% after 1 yr; Kendrick & Walker 1995). Any reduction in early post-settlement mortality may, therefore, result in a higher rate of survival to adulthood (Vadas et al. 1992). The species *Sargassum swartzii* was used to investigate the value of microtopographic refuges for the recruitment and survival of *Sargassum* propagules, as this species occurs in high abundance on inshore reefs in the northern Great Barrier Reef (GBR) (Wismer et al. 2009).

2.2. Study sites

This study was conducted between October and November 2017 at Lizard Island, a granitic mid-shelf island in the northern GBR, Australia (14° 40' S, 145° 28' E; Fig. S1 in the Supplement at www.int-res.com/articles/suppl/m627p061_supp.pdf). To examine the effect of microtopography on the recruitment, growth and survival of macroalgae propagules, we settled *Sargassum swartzii* propagules to tiles with regular crevices and exposed these tiles to local herbivore assemblages on the exposed reef flat and reef crest (Fig. S1b). These 2 reef

zones were selected as they differ in depth, benthic composition, herbivore abundance and/or biomass and rates of herbivory. Reef flat zones are generally characterised by lower rates of herbivory, lower structural complexity and a higher abundance of macroalgae (especially on inshore reefs) than reef crest zones (Fox & Bellwood 2007, Wismer et al. 2009, Hoey & Bellwood 2010a). Depth differs across the 2 reef zones, with the reef flat experiencing diel tidal changes resulting in water depths between 0.5 m to 2.5 m, and the reef crest experiencing water depths between 1.5 and 3.5 m.

2.3. Collection of adult *Sargassum swartzii* and seeding of propagules

In total, 48 terracotta settlement tiles (110 × 110 × 10 mm), each with 4 crevices (4 mm wide, 3 mm deep, 110 mm long) spaced evenly across the top surface of the tile (Fig. S2), were placed on the bottom of a 1000 l aquarium. Twenty-four hours later, 50 adult, reproductively mature *S. swartzii* thalli were collected by hand from the inshore reefs of the Turtle Group (14° 43' S, 145° 12' E), ~11 km west of Lizard Island (Fig. S1a). Thalli were kept in 80 l plastic aquaria (L × W × H: 60 × 37 × 38 cm) filled with seawater and returned to Lizard Island within 3 h of collection. Thalli were stressed within 5 h of collection using cold-shock to stimulate propagule release. Briefly, thalli were placed in an 80 l plastic aquarium filled with cold seawater (9°C) for 10 min (following Carl et al. 2014), then transferred to the 1000 l aquarium containing the tiles at ambient (26°C) seawater temperature. The *S. swartzii* thalli were then agitated in the 1000 l aquarium by hand twice per day to further encourage release of the propagules and were left in this aquarium with static water and supplemental aeration for 3 d. After 3 d, the adult *S. swartzii* were removed from the tank, and the water remaining in the aquarium was filtered through 20 µm mesh to prevent unattached propagules being released onto the reef surrounding the research station. Propagules were first visible on the tiles 2 d after the adult *Sargassum* were removed from the tanks. The tiles with attached propagules were then cultured in the aquarium with supplemental aeration and unfiltered flow-through seawater for a further 18 d. This period allowed turf algal assemblages to establish on the top surface of the tiles.

To determine if recruitment (i.e. settlement and any mortality during the first 18 d) was higher in crevices than on the top, exposed microhabitat (i.e.

the flat surface of the tiles between crevices; Fig. S2) of the tiles, the density of propagules within crevices and on the exposed microhabitat of the tiles was quantified after 18 d by haphazardly placing a quadrat (L × W: 25 × 4 mm; area: 1 cm²) either over a crevice, matching the width of the quadrat to the width of the crevice (8 quadrats per tile) or on the exposed microhabitat (10 quadrats per tile) of the tile. All measurements were conducted using dissecting microscopes to ensure accurate quantification of the small propagules within the quadrats. The resulting dataset included 864 data points across 48 tiles and 2 treatments (crevice vs. exposed microhabitat) describing the density of propagules on the tiles. To quantify any differences in the growth of propagules that had settled in the crevices versus the exposed microhabitats of the tiles, the height of 10 propagules (per microhabitat per tile), on a subset of 6 haphazardly chosen tiles, were measured to the nearest 0.1 mm with callipers, resulting in 120 data points.

2.4. Herbivore exposure

Tiles seeded with propagules were deployed on the reef for 5 d to determine the susceptibility of propagules within crevices versus those on adjacent exposed microhabitats to herbivory. Each tile was secured to the reef by passing a bolt attached to a stainless-steel base plate through a 5 mm hole in the centre of the tile and fastening with a wingnut; push mounts were used to secure the base plate into drilled holes in the reef (following Mundy 2000). Twelve tiles were deployed at each of 2 reef crest and 2 reef flat sites, with at least 3 m between adjacent tiles within a site. Within each site, 6 tiles were left exposed to local herbivore assemblages, and 6 tiles were placed within exclusion cages (L × W × H: 150 × 150 × 40 mm; 5 mm square plastic mesh) to control for the effects of handling and translocation. After 5 d, the tiles were collected, placed onto a stainless-steel bar with 10 mm plastic spacers between each tile to prevent contact between tiles and transported back to the research station in small aquaria within 30 min of collection. Tiles were placed back into the 1000 l aquarium with flow-through seawater and supplemental aeration, and the density of propagules within crevices and on the exposed microhabitat of each tile was re-quantified as previously described. This resulted in 1728 data points across the 48 tiles, collected before and after deployment, across 2 sites, 2 habitats and the 2 microhabitats per tile.

2.5. Video observations

At each site, 4 of the 6 tiles exposed to herbivores were haphazardly chosen to record feeding activity using stationary underwater video cameras. A small video camera (GoPro HERO3 or HERO4) attached to a dive weight was placed adjacent (30 to 40 cm) to each of the 4 selected tiles at each of the 4 sites (16 cameras per day). Each camera started recording between 08:00 and 09:00 h and recorded continuously for 3 h. This process was repeated each day for the 5 d experimental period, resulting in ~240 h of video footage. The entire video footage was viewed, and for each fish observed taking bites on the tile, the species and total length of each fish, number of bites, and bite location (exposed or crevice) was recorded. Those fishes that were too small to accurately identify to species were recorded to the lowest taxonomic level possible (typically genus). Any bites taken on the sides of the tiles were not included in counts, as propagule densities were only quantified on the top microhabitat of the tiles. To account for differences in the availability of crevices versus adjacent exposed areas, the number of bites recorded on each microhabitat was standardised by the area of that microhabitat (i.e. crevice: 17.6 cm²; exposed microhabitat: 103.4 cm²).

2.6. Statistical analysis

Bayesian linear mixed models were used to analyse differences in the recruitment, growth, and mortality of *S. swartzii* propagules between crevices and exposed surfaces. Analyses were conducted in R version 3.3.2 (R Development Core Team 2016) and the models fitted in STAN with Markov chain Monte Carlo sampling (Stan Development Team 2016b, Carpenter et al. 2017) using the rstanarm package version 2.13.1 (Stan Development Team 2016a). The broom (version 0.4.4; Robinson 2017) and CODA (version 0.19.1; Plummer et al. 2006) packages were used to summarise model outputs using highest posterior density intervals with a probability level of 0.95. Plots were produced using ggplot2 version 2.2.1 (Wickham & Chang 2008).

To analyse differences in propagule density and height between crevices and flat microhabitats of tiles before deployment, the generalized linear mixed models 'stan_glm.nb' and 'stan_glm' were used. The models included microhabitat (i.e. crevice vs. exposed) as a fixed factor and microhabitat nested within tile as a random intercept, to account for the variation between quadrats within each microhabitat and tile. Both models used a Gamma error distribution

with a log-link transformation. Weakly informative priors were used on slope coefficients [density: normal(0,10), height: normal(0,10)], intercept coefficients [density: normal(0,10), height: normal(0,10)], and the error standard deviation [density and height: Cauchy(0,5)], with 6000 iterations, a warmup of 3000, 3 chains and a thinning factor of 4 for both models.

To quantify changes to propagule density after exposure to herbivores, we used the proportion of propagules remaining (i.e. the post-deployment average density divided by the initial average density for the crevice and exposed areas on each tile) as the response variable. This model had the fixed factors of microhabitat (either exposed or crevice), reef zone (either reef flat or reef crest) and treatment (either caged or uncaged). Tile identity was included as a random intercept to account for the non-independence of the flat and crevice microhabitats on the same tile. A Gamma error distribution with a log link transformation was used to analyse the data with the 'stan_glm' function. Weakly informative priors were used on slope coefficients [normal(0,5)], intercept coefficients [normal(0,5)] and error standard deviation [Cauchy(0,5)], with 5000 iterations, a warmup of 2500, 3 chains and a thinning factor of 4.

For all models, diagnostic plots were visually examined to ensure there was convergence of chains and no evidence of heteroscedasticity or autocorrelation. Medians and central intervals from prior and posterior distributions were compared to ensure that the chosen priors were sufficiently wide so as to not dictate any trends, without being flat (non-informative). All \hat{R} values (a measure of sampling efficiency) were <1.05, and the ratio of effective samples to total sample size >0.6.

Planned contrasts comparing the difference in the proportion of propagules remaining between the crevice and exposed microhabitats of each tile were performed for each reef zone and treatment combination. When calculating planned contrasts, the generated model matrix was queried to estimate the probability of the difference by summing the number of slopes >0 if the estimate of the parameter was >0 (and vice versa), dividing this number by the total number of observations, and multiplying by 100 to get a percentage:

$$\text{if } y > 0, \frac{\sum a > 0}{n} \times 100; \text{ if } y < 0, \frac{\sum a < 0}{n} \times 100$$

where y is the estimated value of the slope produced by the model, a is all predicted values of the slope, and n is the total number of estimated values of the slope.

3. RESULTS

3.1. Effect of microhabitat on recruitment and growth of propagules

Prior to deploying the tiles on the reef (i.e. 18 d post-settlement), model estimates indicated that the density of propagules within crevices was 21% higher than the density on the exposed microhabitat of the tiles (mean [95% credible intervals]; crevices: 29.7 [26.9, 32.9] propagules cm^{-2} ; exposed microhabitat: 23.4 [21.0, 25.6] propagules cm^{-2} ; Fig. 1a). Propagules within crevices were, however, 18% shorter than those on the exposed microhabitat (1.05 [0.95, 1.13] vs. 1.27 [1.18, 1.36] mm, respectively; Fig. 1b).

3.2. Effect of microhabitat on propagule survival

There were differences in the survival of propagules between microhabitats on the reef crest, but not on the reef flat (Fig. 2a, Table S1 in the Supplement). Model predictions of propagule survival (i.e. proportion of initial propagules) suggested lower survival of propagules on the exposed surface of uncaged tiles deployed on the reef crest (mean = 0.42; 0.34 lower credible interval, 0.50 upper credible interval), whereas survival in the crevices of the same tiles was 0.80 (0.66, 0.97) (Fig. 2a, Table S1). In contrast, survival of propagules within cages on the reef crest was similar between microhabitats: 0.68 (0.55, 0.82) on the exposed surface vs. 0.68 (0.55, 0.81) in crevices. There was strong evidence (>99% probability) of a difference between the proportion of propagules remaining (i.e. surviving) in crevices compared to the exposed microhabitat of uncaged tiles on the reef crest, but no effect for caged tiles on the reef crest (Fig. 2b, Table S2 in the Supplement).

There was less evidence of an effect of microhabitat on propagule survival for uncaged tiles on the reef flat. Model estimates of survival of propagules on the exposed surface of uncaged tiles deployed on the reef flat was 0.70 (0.58, 0.85), compared to 0.80 (0.65, 0.96) in the crevices. There was an 88.5% probability of having more propagules remaining in the crevices than on the exposed microhabitat of the uncaged tiles on the reef flat (Fig. 2b, Table S2). For caged tiles on the reef flat, survival of propagules was 0.71 (0.57, 0.86) on the exposed surface and 0.63 (0.50, 0.76) in crevices (Fig. 2a).

Overall mean bite rates on the tiles were >150% greater on the reef crest (1.67 ± 0.49 bites $\text{h}^{-1} \text{cm}^{-2}$) compared to the reef flat (1.02 ± 0.45 bites $\text{h}^{-1} \text{cm}^{-2}$). For both reef zones, the majority of bites (standardised by the area of each microhabitat) were taken from the exposed microhabitat of the tile (reef crest: 79%; reef flat: 87%) rather than in crevices. The majority of bites in both reef zones were taken by blennies of the genus *Ecsenius*, averaging 1.19 ± 0.20 bites $\text{h}^{-1} \text{cm}^{-2}$ on the exposed microhabitat and 0.27 ± 0.10 bites $\text{h}^{-1} \text{cm}^{-2}$ in crevices of tiles on the reef crest




Fig. 1. Model estimates of the influence of microhabitat on the recruitment and growth of *Sargassum swartzii* propagules (black symbols and \pm 95% credible intervals), and grey circles showing the distribution of raw data: (a) density of propagules in crevices and exposed surface of experimental tiles 18 d post-settlement; (b) height of propagules in crevices and on flat microhabitat of tiles 18 d post-settlement




Fig. 2. (a) Model estimates of the survival of *Sargassum swartzii* propagules between reef zones (reef crest or reef flat), microhabitat (exposed or crevice) and treatments (caged or uncaged) after 5 d deployment \pm 95% credible intervals. Grey open circles indicate the spread of the raw data. (b) Planned contrasts comparing the survival of *Sargassum swartzii* propagules in crevices vs. exposed microhabitats of tiles, in each reef zone and treatment combination. Lines indicate 95% credible intervals calculated using highest posterior density. Estimates (black diamonds) above the dashed line indicate that there were, on average, more propagules in the crevices than on the exposed microhabitat of the tiles. Black diamonds below the line would indicate that there was a higher proportion of propagules on the exposed microhabitat of the tiles than in the crevices

and 0.61 ± 0.22 bites $\text{h}^{-1} \text{cm}^{-2}$ on the exposed microhabitat and 0.09 ± 0.04 bites $\text{h}^{-1} \text{cm}^{-2}$ in the crevices of tiles on the reef flat (Fig. 3). The only other species that took a substantial number of bites was *Salarias fasciatus* (f. Blenniidae) which took an average of 0.21 ± 0.11 bites $\text{h}^{-1} \text{cm}^{-2}$ on the exposed microhabitat and 0.03 ± 0.02 $\text{h}^{-1} \text{cm}^{-2}$ in crevices on the reef flat, but was not recorded as taking any bites from tiles on the reef crest.

4. DISCUSSION

Our results demonstrate that in areas of high herbivory, microtopographic refuges (crevices) afforded *Sargassum* propagules a survival advantage through the provision of a refuge from herbivory. Such a survival advantage may come at a cost, however, as propagules were 18% shorter in crevices compared to those on the exposed surface of the tiles, indicating that crevices may be less favourable for early post-settlement growth. Furthermore, contrary to previous suggestions that feeding by larger-bodied grazing fishes is a major source of mortality for macroalgal propagules on coral reefs (Diaz-Pulido & McCook 2003, Bellwood et al. 2006), our results suggest that small-bodied blennies, specifically of the genus *Ecsenius*, are likely to have been the primary removers of early post-settlement stage *S. swartzii*.

The refuge afforded to propagules within crevices may be particularly important for the survival of juvenile *S. swartzii*, as *Sargassum* propagules have few chemical or morphological defences (Stiger et




Fig. 3. Average (\pm SE) bite rate by fishes on exposed tiles ($\text{h}^{-1} \text{cm}^{-2}$) between reef zones and microhabitats

al. 2004). On the reef crest, where herbivory is generally high (Fox & Bellwood 2007), survival of *S. swartzii* propagules on the exposed microhabitats of the tiles after 5 d exposure to local herbivore assemblages was only 42%, compared to 80% survival within crevices. Our findings add to the growing literature from a range of ecosystems emphasising the importance of refugia to the early post-settlement survival of organisms. Microtopographic refuges have been identified as important early post-settlement refuges for sessile organisms such as bryozoans (Keough & Downes 1982), encrusting sponges (Maldonado & Uriz 1998) and corals (Nozawa 2008). Indeed, Franco et al. (2015) reported that rates of herbivory on temperate reefs in Portugal structured the distribution of *Laminaria* (kelp) recruits; in areas where herbivory was high, most recruits occurred in crevices, whereas in areas where herbivory was low, the majority of recruits were found in exposed, or open, habitats. This is consistent with our findings that crevices were important for the survival of *Sargassum* propagules in areas where herbivory was high (i.e. the reef crest) but were not as important to survival in areas with lower levels of herbivory (i.e. the reef flat), at least over the 5 d temporal scale of this study.

Microtopographic refuges may be extremely important in mediating benthic community assemblages on coral reefs through the provision of grazing refuges, promoting the survival of benthic taxa otherwise vulnerable to consumption by grazers (Brandl & Bellwood 2016). Microtopographic refuges have been shown to support diverse groups of algal turfs, in contrast to exposed surfaces that are predominantly composed of fast-growing, filamentous green algae (Brandl & Bellwood 2016) and crustose coralline algae (Steneck 1997). The present study provides direct evidence of the lower grazing rates, and hence higher survival of macroalgal propagules, within crevices. Such enhanced survival is likely to result in higher algal species diversity within crevices, as demonstrated for other grazing refugia (Menge & Lubchenco 1981, Hoey & Bellwood 2010b, Brandl & Bellwood 2016). Further, crevices may also enhance diversity of the benthic assemblage by not only reducing overall rates of herbivory but also affecting the components of the algae available to grazers. Grazing fishes generally can only access and crop the apical portions of algae growing within crevices, leaving the holdfast, or basal cells, of the algae intact within the crevice (Hixon & Brostoff 1996). Many algae, including *Sargassum*, can regrow from the holdfast (Loffler et al. 2018), and, conse-

quently, consumption of the apical portions is unlikely to cause mortality of the individual. This reduction in grazing pressure is likely to facilitate the survival of a diverse suite of algae, including *Sargassum*, within crevices in areas of high grazing intensity.

Although settling within crevices appears to confer a survival advantage for *S. swartzii* propagules, it may come at a cost. Eighteen days after settlement, the height of *Sargassum* propagules within crevices was 18% lower than propagules that had settled on the exposed microhabitats of the tiles. This lower growth within crevices could be related to density-dependent factors and/or to different light and water flow regimes within crevices compared to exposed microhabitats. Unlike other algal species (Christie 1973), *Sargassum* propagules are non-motile, and it is therefore likely that the higher initial density in crevices was a result of the negatively buoyant propagules being 'trapped' in a crevice and settling rather than actively choosing to settle in this microhabitat (Norton & Fetter 1981). Kendrick (1994) reported that the growth of temperate *Sargassum* recruits was negatively related to density, indicating that there may be a trade-off between survival and growth within crevices and on exposed surfaces (Amsler et al. 1992, Bergey 1999, 2005, Brandl et al. 2014). Our results suggest that the benefits of settling in a crevice may only outweigh the disadvantages when rates of herbivory are high, by enhancing propagules' survival at a stage characterised by high mortality (Kendrick & Walker 1995, Brandl et al. 2014, Franco et al. 2015). However, the slower growth of propagules within crevices may mean that they are subject to consumption by grazing organisms for longer (i.e. the stage-duration hypothesis; Leggett & Deblois 1994), which could partially or completely negate any protective value of the crevices.

Rapid increases in algal biomass following the exclusion of large herbivores has led to the widely held assumption that large-bodied grazing fishes, such as surgeonfishes and parrotfishes, are primarily responsible for the removal of macroalgal propagules in the epilithic algal matrix (EAM: a conglomerate of turf algae, macroalgal propagules, detritus, sediment and microbes) (Diaz-Pulido & McCook 2003, Bellwood et al. 2004). Our results question this assumption, as few large-bodied grazing fishes fed on the tiles, and none of the fishes previously identified as 'crevice feeders', including *Siganus* spp. and *Zebrosomas* spp. (Fox & Bellwood 2013, Brandl et al. 2015), were observed performing this function within the

crevices of our experimental tiles. This lack of feeding by *Siganus* and *Zebrasoma* may be related to the size of the crevices, with the 3 mm deep, 4 mm wide crevices being too small for these larger-bodied specialised crevice-feeding fishes to access (Fox & Bellwood 2013, Brandl et al. 2014). Small crevices may, therefore, represent an important refuge for the recruitment of benthic marine organisms such as macroalgae.

Small-bodied cryptobenthic blennies *Ecsenius* spp. may play an important role in the early post-settlement mortality of macroalgal propagules, as these fishes took a disproportionately higher number of bites on uncaged tiles than any other grazing fishes. Grazing by the combtooth blenny *Salarias fasciatus* has been demonstrated to reduce the survival of early-post settlement coral recruits (Christiansen et al. 2009, Doropoulos et al. 2012), even though *S. fasciatus* are thought to primarily consume detrital aggregates within the EAM (Wilson 2000). *Ecsenius* spp. may similarly reduce propagule survival through grazing of the EAM even though detrital aggregates may be the primary dietary target (Wilson 2000, Brandl et al. 2018). Although our results suggest that cryptobenthic blennies contribute to the mortality of small *Sargassum* propagules (<2 mm, and up to 25 d post-settlement), it is likely that *Sargassum* and other macroalgal propagules will quickly reach a size refuge from blennies as they grow (Lubchenco 1983, Briggs et al. 2018). It is yet to be determined if propagule removal by blennies at this early post-settlement stage could translate to lower densities of adult macroalgae or if other species of grazing fishes consume larger macroalgal propagules.

Our findings also suggest that the importance of microtopographic refuges for the survival of macroalgal propagules varies among reef zones. On the reef flat, where conditions for *Sargassum* growth are favourable (McCook 1996) and herbivory is generally low (Hay 1981, Fox & Bellwood 2007), the survival of *Sargassum* propagules did not differ between caged and uncaged tiles. This result is consistent with the findings of previous studies that have shown that the density and/or biomass of herbivorous fishes (Lewis & Wainwright 1985, Wismer et al. 2009) and rates of herbivory (Hay 1981, Fox & Bellwood 2007) are lower on the reef flat than the reef crest, and such studies have led to suggestions that the reef flat is a spatial refuge from herbivory (Hay et al. 1983, Lewis & Wainwright 1985). In areas of lower herbivory, refuges may be less critical for the survival of *Sargassum* propagules (Franco et al. 2015).

The results of our study highlight the potential importance of crevices for the early post-settlement survival and growth of *S. swartzii*; however, further research is needed to investigate how the size of refugia and the potential agents of mortality (i.e. herbivores) change with macroalgae size and vary among macroalgal taxa. Whilst crevices were of uniform size in the present study, the size and depth of crevices relative to propagule size is likely to have a large influence on the degree of protection afforded to propagules (Lubchenco 1983, Bergley 1999). Furthermore, the fish species responsible for the removal of propagules are likely to change as the juvenile *Sargassum* grows. We do not know which fishes remove these larger (0.5 to 5 cm) juveniles, whether the phenolic content of *S. swartzii* increases ontogenetically, as in *Sargassum mangarevense* (Stiger et al. 2004), possibly decreasing susceptibility to herbivores with age, or whether the protective value of the crevices for early post-settlement propagules significantly affects rates of survival to maturity.

The results of the present study provide a useful insight into the value of microtopographic refuges for the recruitment, growth and survival of newly settled *Sargassum* propagules and is one of the first to highlight the potential role of blennies in consuming newly settled macroalgal propagules on coral reefs. Microtopographic refuges on coral reefs can provide newly settled propagules a refuge from grazing in this vulnerable early post-settlement stage. Higher early-post settlement survival of propagules within refuges may facilitate the persistence of *Sargassum* communities on coral reefs, particularly in areas where herbivory pressure is high.

Author contributions: Z.L. and A.S.H. conceived and designed the experiments. Z.L. performed the experiments and analyzed the data. Z.L. and A.S.H. wrote the manuscript.

Ethical approval. This research was conducted under JCU Ethics A2459. All applicable institutional and national guidelines for the care and use of animals were followed.

Acknowledgements. We thank E. Aston, A. Graba-Landry, K. Hannan, J. Kidgell, M. Larkin and C. Murphy for field and/or video analysing assistance; the staff at Lizard Island Research Station for invaluable field support; and S. Brandl and 2 anonymous reviewers for helpful comments on the manuscript. Financial support was provided by the Ian Potter Doctoral Fellowship at Lizard Island (Z.L.), the Holsworth Wildlife Research Endowment (Z.L.), the Australian Coral Reef Society (Z.L.) and the Australian Research Council (A.S.H.).

LITERATURE CITED

- Almany GR, Webster MS (2006) The predation gauntlet: early post-settlement mortality in reef fishes. *Coral Reefs* 25:19–22
- Amsler CD, Reed DC, Neushul M (1992) The microclimate inhabited by macroalgal propagules. *Br Phycol J* 27: 253–270
- Bailey K, Houde E (1989) Predation on eggs and larvae of marine fishes and the recruitment problem. *Adv Mar Biol* 25:1–83
- Bellwood DR, Hughes TP, Folke C, Nystrom M (2004) Confronting the coral reef crisis. *Nature* 429:827–833
- Bellwood DR, Hughes TP, Hoey AS (2006) Sleeping functional group drives coral-reef recovery. *Curr Biol* 16: 2434–2439
- Bergey EA (1999) Crevices as refugia for stream diatoms: effect of crevice size on abraded substrates. *Limnol Oceanogr* 44:1522–1529
- Bergey EA (2005) How protective are refuges? Quantifying algal protection in rock crevices. *Freshw Biol* 50: 1163–1177
- Bonin MC, Srinivasan M, Almany GR, Jones GP (2009) Interactive effects of interspecific competition and microhabitat on early post-settlement survival in a coral reef fish. *Coral Reefs* 28:265–274
- Brandl SJ, Bellwood DR (2016) Microtopographic refuges shape consumer-producer dynamics by mediating consumer functional diversity. *Oecologia* 182:203–217
- Brandl SJ, Hoey AS, Bellwood DR (2014) Micro-topography mediates interactions between corals, algae, and herbivorous fishes on coral reefs. *Coral Reefs* 33:421–430
- Brandl SJ, Robbins WD, Bellwood DR (2015) Exploring the nature of ecological specialization in a coral reef fish community: morphology, diet and foraging microhabitat use. *Proc Biol Sci* 282:20151147
- Brandl SJ, Goatley CH, Bellwood DR, Tornabene L (2018) The hidden half: ecology and evolution of cryptobenthic fishes on coral reefs. *Biol Rev Camb Philos Soc* 93: 1846–1873
- Briggs CJ, Adam TC, Holbrook SJ, Schmitt RJ (2018) Macroalgae size refuge from herbivory promotes alternative stable states on coral reefs. *PLOS ONE* 13:e0202273
- Carl C, de Nys R, Lawton RJ, Paul NA (2014) Methods for the induction of reproduction in a tropical species of filamentous ulva. *PLOS ONE* 9:e97396
- Carpenter B, Gelman A, Hoffman MD, Lee D and others (2017) Stan: a probabilistic programming language. *J Stat Softw* 76:1–29
- Chong-Seng KM, Nash KL, Bellwood DR, Graham NAJ (2014) Macroalgal herbivory on recovering versus degrading coral reefs. *Coral Reefs* 33:409–419
- Christiansen N, Ward S, Harii S, Tibbetts I (2009) Grazing by a small fish affects the early stages of a post-settlement stony coral. *Coral Reefs* 28:47–51
- Christie AO (1973) Spore settlement in relation to fouling by *Enteromorpha* In: Acker RF Brown BF, DePalma JR, Iverson WP (ed) *Proc Third Int Congr Marine Corrosion Fouling*. Northwestern University Press, Evanston, IL, p 674–681
- Connell JH (1985) The consequences of variation in initial settlement vs. post-settlement mortality in rocky intertidal communities. *J Exp Mar Biol Ecol* 93:11–45
- Crowder LB, Cooper WE (1982) Habitat structural complexity and the interaction between bluegills and their prey. *Ecology* 63:1802–1813
- Diaz-Pulido G, McCook LJ (2003) Relative roles of herbivory and nutrients in the recruitment of coral-reef seaweeds. *Ecology* 84:2026–2033
- Diaz-Pulido G, Harii S, McCook LJ, Hoegh-Guldberg O (2010) The impact of benthic algae on the settlement of a reef-building coral. *Coral Reefs* 29:203–208
- Done TJ (1992) Phase-shifts in coral reef communities and their ecological significance. *Hydrobiologia* 247: 121–132
- Doropoulos C, Ward S, Marshall A, Diaz-Pulido G, Mumby PJ (2012) Interactions among chronic and acute impacts on coral recruits: the importance of size-escape thresholds. *Ecology* 93:2131–2138
- Doropoulos C, Roff G, Bozec YM, Zupan M, Werninghausen J, Mumby PJ (2016) Characterizing the ecological trade offs throughout the early ontogeny of coral recruitment. *Ecol Monogr* 86:20–44
- Doropoulos C, Evensen NR, Gómez-Lemos LA, Babcock RC (2017a) Density-dependent coral recruitment displays divergent responses during distinct early life-history stages. *R Soc Open Sci* 4:170082
- Doropoulos C, Roff G, Visser MS, Mumby PJ (2017b) Sensitivity of coral recruitment to subtle shifts in early community succession. *Ecology* 98:304–314
- Fox RJ, Bellwood DR (2007) Quantifying herbivory across a coral reef depth gradient. *Mar Ecol Prog Ser* 339: 49–59
- Fox RJ, Bellwood DR (2013) Niche partitioning of feeding microhabitats produces a unique function for herbivorous rabbitfishes (Perciformes, Siganidae) on coral reefs. *Coral Reefs* 32:13–23
- Franco JN, Wernberg T, Bertocci I, Duarte P, Jacinto D, Vasco-Rodrigues N, Tuya F (2015) Herbivory drives kelp recruits into 'hiding' in a warm ocean climate. *Mar Ecol Prog Ser* 536:1–9
- Gosselin LA, Qian PY (1997) Juvenile mortality in benthic marine invertebrates. *Mar Ecol Prog Ser* 146:265–282
- Graham NAJ, Nash KL (2013) The importance of structural complexity in coral reef ecosystems. *Coral Reefs* 32: 315–326
- Hay ME (1981) Spatial patterns of grazing intensity on a Caribbean barrier reef: herbivory and algal distribution. *Aquat Bot* 11:97–109
- Hay ME, Colburn T, Downing D (1983) Spatial and temporal patterns in herbivory on a Caribbean fringing reef: the effects on plant distribution. *Oecologia* 58:299–308
- Hixon MA, Brostoff WN (1996) Succession and herbivory: effects of differential fish grazing on Hawaiian coral reef algae. *Ecol Monogr* 66:67–90
- Hoey AS, Bellwood DR (2010a) Among-habitat variation in herbivory on *Sargassum* spp. on a mid-shelf reef in the northern Great Barrier Reef. *Mar Biol* 157:189–200
- Hoey AS, Bellwood DR (2010b) Damselish territories as a refuge for macroalgae on coral reefs. *Coral Reefs* 29: 107–118
- Hoey AS, McCormick MI (2004) Selective predation for low body condition at the larval-juvenile transition of a coral reef fish. *Oecologia* 139:23–29
- Hunt HL, Scheibling RE (1997) Role of early post-settlement mortality in recruitment of benthic marine invertebrates. *Mar Ecol Prog Ser* 155:269–301
- Kendrick GA (1994) Effects of propagule settlement density and adult canopy on survival of recruits of *Sargassum* spp. (Sargassaceae, Phaeophyta). *Mar Ecol Prog Ser* 103: 129–140

- Kendrick GA, Walker DI (1994) Role of recruitment in structuring beds of *Sargassum* spp. (Phaeophyta) at Rottneest Island, Western Australia. *J Phycol* 30:200–208
- Kendrick GA, Walker DI (1995) Dispersal of propagules of *Sargassum* spp. (Sargassaceae, Phaeophyta)—observations of local patterns of dispersal and consequences for recruitment and population structure. *J Exp Mar Biol Ecol* 192:273–288
- Keough MJ, Downes BJ (1982) Recruitment of marine invertebrates: the role of active larval choices and early mortality. *Oecologia* 54:348–352
- Leggett W, Deblois E (1994) Recruitment in marine fishes: Is it regulated by starvation and predation in the egg and larval stages? *Neth J Sea Res* 32:119–134
- Lewis SM (1986) The role of herbivorous fishes in the organization of a Caribbean reef community. *Ecol Monogr* 56: 183–200
- Lewis SM, Wainwright PC (1985) Herbivore abundance and grazing intensity on a Caribbean coral-reef. *J Exp Mar Biol Ecol* 87:215–228
- Loffler Z, Bellwood DR, Hoey AS (2015) Among-habitat algal selectivity by browsing herbivores on an inshore coral reef. *Coral Reefs* 34:597–605
- Loffler Z, Graba-Landry A, Kidgell JT, McClure EC, Pratchett MS, Hoey AS (2018) Holdfasts of *Sargassum swartzii* are resistant to herbivory and resilient to damage. *Coral Reefs* 37:1075–1084
- Lubchenco J (1983) *Littornia* and *Fucus*: effects of herbivores, substratum heterogeneity, and plant escapes during succession. *Ecology* 64:1116–1123
- Maldonado M, Uriz MJ (1998) Microrefuge exploitation by subtidal encrusting sponges: patterns of settlement and post-settlement survival. *Mar Ecol Prog Ser* 174: 141–150
- Mallela J (2018) The influence of micro-topography and external bioerosion on coral-reef-building organisms: recruitment, community composition and carbonate production over time. *Coral Reefs* 37:227–237
- McCook LJ (1996) Effects of herbivores and water quality on *Sargassum* distribution on the central Great Barrier Reef: cross-shelf transplants. *Mar Ecol Prog Ser* 139:179–192
- McCormick M, Makey L, Dufour V (2002) Comparative study of metamorphosis in tropical reef fishes. *Mar Biol* 141:841–853
- Menge BA, Lubchenco J (1981) Community organization in temperate and tropical rocky intertidal habitats: prey refuges in relation to consumer pressure gradients. *Ecol Monogr* 51:429–450
- Menge BA, Sutherland JP (1976) Species diversity gradients: synthesis of the roles of predation, competition, and temporal heterogeneity. *Am Nat* 110:351–369
- Miller TJ, Crowder LB, Rice JA, Marschall EA (1988) Larval size and recruitment mechanisms in fishes: toward a conceptual framework. *Can J Fish Aquat Sci* 45: 1657–1670
- Mundy C (2000) An appraisal of methods used in coral recruitment studies. *Coral Reefs* 19:124–131
- Norton TA, Fetter R (1981) The settlement of *Sargassum muticum* propagules in stationary and flowing water. *J Mar Biol Assoc UK* 61:929–940
- Nozawa Y (2008) Micro-crevice structure enhances coral spat survivorship. *J Exp Mar Biol Ecol* 367:127–130
- Plummer M, Best N, Cowles K, Vines K (2006) CODA: convergence diagnosis and output analysis for MCMC. *R News* 6:7–11
- Poray A, Carpenter R (2014) Distributions of coral reef macroalgae in a back reef habitat in Moorea, French Polynesia. *Coral Reefs* 33:67–76
- R Development Core Team (2016) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. www.r-project.org
- Robinson D (2017) broom: convert statistical analysis objects into tidy data frames. R package version 04. <https://github.com/tidymodels/broom>
- Scheibling RE, Robinson MC (2008) Settlement behaviour and early post-settlement predation of the sea urchin *Strongylocentrotus droebachiensis*. *J Exp Mar Biol Ecol* 365:59–66
- Stan Development Team (2016a) rstanarm: Bayesian applied regression modeling via Stan. R package version 2.13.1. <https://github.com/stan-dev/rstanarm>
- Stan Development Team (2016b) Stan modeling language users guide and reference manual, version 2.14.0. <https://github.com/stan-dev/rstan>
- Steneck R (1997) Crustose corallines, other algal functional groups, herbivores and sediments: complex interactions along reef productivity gradients. *Proc 8th Int Coral Reef Symp* 1:695–700
- Stiger V, Deslandes E, Payri CE (2004) Phenolic contents of two brown algae, *Turbinaria ornata* and *Sargassum mangarevense* on Tahiti (French Polynesia): interspecific, ontogenic and spatio-temporal variations. *Bot Mar* 47: 402–409
- Trapon ML, Pratchett MS, Hoey AS, Baird AH (2013) Influence of fish grazing and sedimentation on the early post-settlement survival of the tabular coral *Acropora cytherea*. *Coral Reefs* 32:1051–1059
- Tupper M, Boutilier RG (1997) Effects of habitat on settlement, growth, predation risk and survival of a temperate reef fish. *Mar Ecol Prog Ser* 151:225–236
- Vadas RL, Johnson S, Norton TA (1992) Recruitment and mortality of early post-settlement stages of benthic algae. *Br Phycol J* 27:331–351
- Vermeij MJA, Sandin SA (2008) Density-dependent settlement and mortality structure the earliest life phases of a coral population. *Ecology* 89:1994–2004
- Wickham H, Chang W (2008) ggplot2: an implementation of the Grammar of Graphics. R package version 07, <http://CRANR-project.org/package=ggplot2>
- Wilson S (2000) Trophic status and feeding selectivity of blennies (Blenniidae: Salariini). *Mar Biol* 136:431–437
- Wismer S, Hoey AS, Bellwood DR (2009) Cross-shelf benthic community structure on the Great Barrier Reef: relationships between macroalgal cover and herbivore biomass. *Mar Ecol Prog Ser* 376:45–54
- Yadav S, Rathod P, Alcoverro T, Arthur R (2016) ‘Choice’ and destiny: the substrate composition and mechanical stability of settlement structures can mediate coral recruit fate in post-bleached reefs. *Coral Reefs* 35: 211–222