

ORIGINAL RESEARCH

Early childhood anaemia more than doubles the risk of developmental vulnerability at school-age among Aboriginal and Torres Strait Islander children of remote Far North Queensland: Findings of a retrospective cohort study

Dympna Leonard MPH, APD, PhD Candidate, Adjunct Senior Lecturer¹  |
Petra Buettner PhD, Adjunct Professor^{1,2} |
Fintan Thompson MEpi, Data Analyst/Manager¹ |
Maria Makrides PhD, Professor^{3,4} | Robyn McDermott PhD, Professor¹

¹Australian Institute of Tropical Health and Medicine, James Cook University, Cairns, Queensland, Australia

²Tropical Health Solutions Pty Ltd, Townsville, Queensland, Australia

³Healthy Mothers, Babies and Children, South Australian Health and Medical Research Institute, Adelaide, South Australia, Australia

⁴Discipline of Paediatrics, School of Medicine, The University of Adelaide, Adelaide, South Australia, Australia

Correspondence

Dympna Leonard, Australian Institute of Tropical Health and Medicine, James Cook University, PO Box, 6811, Cairns QLD 4870, Australia.
Email: dympna.leonard@jcu.edu.au

Funding information

National Health and Medical Research Council, Grant/Award Number: APP1092732

Abstract

Aims: Early childhood anaemia, usually attributed to iron deficiency, is associated with persistent detrimental effects on child development. This study investigates the association of anaemia between age six and 23 months with indicators of childhood development at school-age among children of remote Aboriginal and Torres Strait Islander communities of Far North Queensland.

Methods: The triennial Australian Early Development Census (AEDC) encompasses five domains of early childhood development—physical health and wellbeing, social competence, emotional maturity, language and cognitive skills (school-based), communication skills and general knowledge. AEDC 2012 and 2015 assessments were linked with health information for children and their mothers from remote Aboriginal and Torres Strait Islander communities of Far North Queensland.

Results: AEDC assessments were available for 250 children who had measurements of haemoglobin recorded at age 6 to 23 months. More children who had had early childhood anaemia ($n = 66/143$, 46.2%, [37.9%, 54.4%]) were developmentally vulnerable on two or more domains compared to those who had not been anaemic ($n = 25/107$, 23.4% [15.2%, 31.5%], $P < .001$). Multivariable analysis confirmed that early childhood anaemia more than doubled the risk of developmental vulnerability (OR 2.2 [1.1, 4.3] $P = .020$) at school age.

Conclusions: Early childhood anaemia is a risk factor for developmental vulnerability at school-age in this setting. Interventions combining nutrition promotion and multi-micronutrient food fortification, are effective in prevention of early childhood anaemia. Such interventions could also improve early childhood development and subsequent educational achievement.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *Nutrition & Dietetics* published by John Wiley & Sons Australia, Ltd on behalf of Dietitians Association of Australia

1 | INTRODUCTION

Good nutrition in the first thousand days of life—from conception, through pregnancy to around age 2 years—supports the rapid neurodevelopment of early life that provides the scaffolding for subsequent child development.¹ Neurodevelopment is “the dynamic interrelationship between environment, genes and the brain whereby the brain develops across time to establish sensory, motor, cognitive, socioemotional, cultural and behavioral adaptive functions.”² Iron is required for specific neurodevelopment processes before birth and during the first years of life, including neuronal proliferation and growth of axons and dendrites, synapse formation and myelination.^{1,3} Iron deficiency during critical phases of neurodevelopment is associated with persistent deficits in cognitive and behavioural performance.^{4,5} Anaemia is a late stage of iron deficiency; the detrimental effects of iron deficiency impact on the developing brain before anaemia develops.^{5,6}

Anaemia—defined as low haemoglobin—among Aboriginal and Torres Strait Islander women in pregnancy and their children in early life, is prevalent in Far North Queensland and elsewhere in remote northern Australia.^{7–11} There are various causes of anaemia including nutrition-related causes (deficiency of iron and/or folate and/or vitamin B12), chronic infections and genetic conditions.¹²

Anaemia in the first thousand days is usually caused by iron deficiency due to high iron requirements for the rapidly increasing blood volume and tissue growth of pregnancy and early childhood.^{1,5}

The Australian Early Development Census (AEDC) is a national census of early childhood development, conducted every 3 years since 2009.¹³ Each child in the first year of full time school is assessed by his/her teacher on five domains of early childhood development; physical health and wellbeing, social competence, emotional maturity, language and cognitive skills (school-based), communication skills and general knowledge.¹³ These AEDC domains are predictive of outcomes in health, well-being and academic success in later life.¹³ In Australia, AEDC results have been shown to predict subsequent National Assessment Program – Literacy and Numeracy (NAPLAN) scores for numeracy and reading.¹⁴

The 2015 AEDC census with 302 003 child participants found that 22.0% were developmentally vulnerable—scoring below the tenth centile—on one or more domains (DV1) and 11.1% developmentally vulnerable on two or more domains (DV2). More boys were developmentally vulnerable than girls.¹³ Studies in South Australia and the Northern Territory identified perinatal factors associated with developmental vulnerability at school entry; smoking in

pregnancy; anaemia of mothers; low birth weight; prematurity among Aboriginal and non-Aboriginal children.^{15,16}

Here, we report on the association of early childhood anaemia, defined as a child ever having anaemia between age 6 and 23 months, with AEDC assessment results among Aboriginal and Torres Strait Islander children of remote Far North Queensland.

2 | METHODS

This retrospective cohort study used linked information for two cohorts of children and their mothers—the Cape York cohort and the 2009 to 2010 cohort. The Cape York cohort included children born between 2006 and 2008, participants in a previous unpublished study of child growth in remote Cape York communities. The 2009 to 2010 cohort included children born to Aboriginal and/or Torres Strait Islander mothers in 2009 or 2010 in Far North Queensland. Only the first child born to each mother between 2006 and 2010 was included to ensure independence of events for statistical analysis (Figure 1).

Information was sourced from three Queensland Health data collections; (a) Queensland Pathology Services data collection (Auslab) (b) Queensland Perinatal Data Collection (PDC) (c) the Queensland Health community health electronic record system—Ferret—used in Far North Queensland remote Aboriginal and Torres Strait Islander communities (Map 1). Information recorded on Ferret includes results of routine child haemoglobin measurements from age six months. Health information for individual children and their mothers was linked to the child's assessment from AEDC 2012 or AEDC 2015. The final linked de-identified dataset was provided to researchers in May 2017. A full description of this linkage process is described elsewhere.¹⁷ Table S1 provides more information on these data collections.

Ethics approval, granted by Queensland Health Far North Queensland Human Research Ethics Committee (HREC/15/QCH/50-980) in June 2015. In February 2016, approval under the Queensland Public Health Act 2005 was granted by the Director-General of Queensland Health. In September 2016, approval was granted by the Australian Government Department of Education and Training Australian to access the Australian Early Development Census data collection. The complete linked de-identified data were provided to the research group in May 2017.

2.1 | Study variables and definitions

Information on the outcome variables—developmental vulnerability—was sourced from the AEDC data





FIGURE 1 Flow diagram—association of early childhood anaemia and child development at school age among two cohorts of Aboriginal and Torres Strait Islander children and their mothers in Far North Queensland; data available and exclusions

collection.¹³ AEDC has developed categories of developmental status (on track, developmentally at risk, developmentally vulnerable) based on the first AEDC in 2009. Children scoring below the AEDC 2009 tenth centile for each domain are categorised as developmentally

vulnerable for that domain. Children with scores below the tenth centile for one or more of the five AEDC domains are categorised as DV1. Children with scores below the tenth centile for two or more domains are categorised as DV2.¹³ (Table S2).



MAP 1 Far North Queensland—location and year of rollout of the Ferret electronic record system. Based on information provided by the Queensland Health Ferret Support Team, May 2017. Reproduced with permission of the Australian New Zealand Journal of Public Health

Information on the characteristics of children and mothers were sourced from the health service data collections. Anaemia in early childhood was defined using the Queensland Health age-specific criteria; at least one haemoglobin level recorded below 105 g/L from 6 to 11 months, and/or below 110 g/L from 12 to 23 months.¹⁸

Some characteristics are as recorded on health service data collections: mother's usual residence, ethnicity, parity, smoking in pregnancy; child's sex, gestational age at birth, birth weight, method of birth. Criteria to define derived variables including maternal age, body mass index (BMI), anaemia before and during pregnancy, insufficient red cell folate levels (folate status less than optimal for women of reproductive age to prevent neural tube defects)^{19,20} and prematurity and birth weight category of the children are detailed in Table S2. Birth weight z-scores for babies with gestational age of 33 weeks or more, were calculated using the INTERGROWTH-21ST Neonatal Size Calculator.^{21,22}

Australian Bureau of Statistics ranks Statistical Local Areas by deciles of relative socio-economic advantage and disadvantage.²³ A ranking of "1" indicates greatest relative disadvantage while a ranking of "10" indicates greatest relative advantage. Mothers were allocated a Socio-Economic Index for Areas (SEIFA 2011) rank according to usual place of residence.

2.2 | Statistical analysis

Analysis was conducted using Stata version 13 (StataCorp, Lakeway Drive, College Station, Texas). Categorical variables were described using absolute and relative frequencies. The distributions of numerical variables were assessed; symmetrically distributed numerical characteristics were described using mean values and SDs; numerical values with a skewed distribution were described using median and inter-quartile ranges (IQR).

The main outcome variables for bivariate analysis were; (a) developmentally vulnerable for one or more of the five AEDC domains—DV1; and (b) developmentally vulnerable for two or more of the five AEDC domains—DV2. The main outcome variables were presented with 95% confidence interval (95% CI).

The following characteristics of children and their mothers were considered during bi- and multivariable analyses: child sex; birthing method (non-instrumental vaginal, instrumental vaginal, caesarean section); gestational age at birth; premature or not; birth weight; feeding method to age 4 months (only breast milk, only infant formula, both breast milk and formula); child had anaemia at age 6 to 23 months or not; ethnicity of mother (Aboriginal, Torres Strait Islander, both); region of residence of mother; SEIFA category for

residence of mother; maternal age; BMI category of mother (underweight, normal weight, overweight, obese); categories of parity (0-2, ≥ 3); smoking during pregnancy; diabetes during pregnancy; mother had insufficient red cell folate (RCF) level before or during pregnancy; mother anaemic both before and during pregnancy.

Developmental vulnerability was compared by these characteristics of the children and their mothers, using bivariate logistic regression analyses. In addition, the association of developmental vulnerability for each domain was assessed for; mother anaemic both before and during pregnancy; smoking during pregnancy; early childhood anaemia, using Pearson's χ^2 tests.

Multivariable logistic regression analyses were conducted to identify independent risk factors for children considered developmentally vulnerable on two or more domains (DV2) for the complete case analysis. Backward and forward stepwise modelling procedures were initially conducted to establish a basic multivariable model. Characteristics that were not part of the basic model were assessed for potential confounding effects. A confounder was assumed to be a variable that changed estimates of characteristics in the basic model by 10% or more.²⁴

Univariate multiple imputation was conducted using Stata's MI commands for missing values for one characteristic: mother anaemic both before and during pregnancy ($n = 29$). Other characteristics with missing values, including feeding method, parity, BMI of mother, diabetes in pregnancy, and insufficient RCF levels of mother were not imputed because these variables had shown no statistically significant influence on the main outcome variables during bi- and multivariable analyses.

The occurrence of missing values of mother anaemic both before and during pregnancy ($n = 29$) was found unrelated to DV2 using Pearson's χ^2 test. The pattern of missing values was assessed and judged to be "missing at random".²⁵ Logistic regression was used for imputation. Imputation models were based on variables with nil missing data; birth weight of baby; smoking during pregnancy; early childhood anaemia; sex of baby; gestational age of baby; AEDC developmental assessment; birthing method; ethnicity of mother, age of mother, SEIFA index; residential region; and cohort. Twenty imputed data sets were created. Multivariable logistic regression analyses were conducted to identify independent risk factors for DV2 for imputed data.

Results of multivariable models for complete case and imputed data analyses are presented as odds ratios (OR) and 95% confidence intervals. P values of less than .05 were considered statistically significant. The authors used the STROBE checklist for cohort studies to guide the preparation of this paper.²⁶

3 | RESULTS

Ferret longitudinal health records were available for 1155 children of whom 957 were the first child born to his/her mother in the cohort years (Figure 1). Of these 957 children, 708 children had a record of haemoglobin measurement at age 6 to 23 months. Just over one-third (35.3%, $n = 250$) of the 708 children also had a record of AEDC assessment. More than half (58.0%; 95% CI 51.6%, 64.2%) of these 250 children were assessed as DV1 and approximately one third (36.4%; 95% CI 30.4%, 42.7%) as DV2 (Table 1).

Bivariate analysis showed developmental vulnerability was significantly more prevalent among children who had had early childhood anaemia (DV2 $n = 66/143$, 46.2% [95% CI 37.9%, 54.4%]) compared to those who had not been anaemic (DV2 $n = 25/107$, 23.4% [95% CI 15.2%, 31.5%] $P < .001$). Developmental vulnerability was significantly more prevalent among boys compared to girls; children of mothers who were anaemic both before and during pregnancy; children of mothers who smoked in pregnancy. These effects were seen for both the DV1 and the DV2 categories (Tables 1 and 2). Children of Torres Strait Islander mothers were significantly less likely to be categorised as either DV1 or DV2 compared to the children of other Indigenous mothers (Table 2).

Early childhood anaemia was significantly associated with developmental vulnerability for four out of five domains of early childhood development: physical health and wellbeing, social competence, language and cognitive skills (school-based), communication skills and general knowledge. Smoking in pregnancy was significantly associated with developmental vulnerability on three domains: physical health and wellbeing, social competence, communication skills and general knowledge. Anaemia of mothers before and during pregnancy was significantly associated with developmental vulnerability on two domains: physical health and wellbeing, communication skills and general knowledge (Table S3).

Multivariable complete case analyses found that children who had had early childhood anaemia (OR 2.2 [95% CI 1.1, 4.3] $P = .020$) were at significantly higher risk of developmental vulnerability compared to children who had not been anaemic, as were boys compared to girls (OR 2.8 [95% CI 1.5, 5.3] $P = .001$). Characteristics of mothers significantly associated with developmental vulnerability in their children included smoking in pregnancy (OR 2.0 [95% CI 1.02, 3.9] $P = .045$) and anaemia both before and during pregnancy (OR 2.1 [95% CI 1.1, 4.1] $P = .021$). Children of Aboriginal mothers were at higher risk of developmental vulnerability (OR 2.5 [95% CI 1.3, 5.0] $P = .009$) compared to children of Torres

TABLE 1 Developmental vulnerability (in lower 10th percentile of AEDC assessment score at first year of full-time school) in one or more domains (DV1) and in two or more domains (DV2) by characteristics of the child—bivariate analysis (logistic regression)

Part A							
Characteristics of children	Number	AEDC—DV1 (Y)		P-values (DV1 logistic regression)		P values (DV2 logistic regression)	
		n = 145 (58.0%)		AEDC—DV2 (Y)n = 91 (36.4%)			
Sex n = 250							
Boys	132	88 (66.7%) [58.5%, 74.8%]	P = .004*	62 (47.0%) [38.3%, 55.6%]		P < .001*	
Girls	118	57 (48.3%) [39.2%, 57.5%]		29 (24.6%) [16.7%, 32.5%]			
Birth weight category n = 250							
Low birth weight ^a	31	19 (61.3%) [43.1%, 79.5%]	P = .713	n = <15		P = .401	
Normal birth weight	199	115 (57.8%) [50.9%, 64.7%]	Base	68 (34.2%) [27.5%, 40.8%]		Base	
Macrosomic (>4000 g) ^a	20	n ≤ 15	P = .810	n ≤ 15		P = .164	
Premature birth n = 250							
Child—premature birth ^a	32	20 (62.5%) [44.8%, 80.2%]	P = .581	n ≤ 15		P = .356	
Child—full term birth	218	125 (57.3%) [50.7%, 64.0%]		77 (35.3%) [28.9%, 41.7%]			
Birth method n = 250							
Vaginal	172	99 (57.6%) [50.1%, 65.0%]	Base	67 (39.0%) [32.0%, 46.3%]		Base	
Vaginal instrumental ^a	n ≤ 15	n ≤ 15	P = .845	n ≤ 15		P = .186	
Caesarian	67	40 (59.7%) [47.6%, 71.8%]	P = .763	22 (32.8%) [21.3%, 44.4%]		P = .380	
Feeding method—birth to age 4 months n = 189							
Breast milk only	83	48 (57.8%) [47.0%, 68.7%]	Base	34 (41.0%) [30.2%, 51.8%]		Base	
Infant formula only	25	17 (68.0%) [48.3%, 87.7%]	P = .365	n ≤ 15		P = .657	
Both breast milk and infant formula	81	48 (59.3%) [48.3%, 70.2%]	P = .853	30 (37.0%) [26.3%, 47.8%]		P = .606	
Early childhood anaemia (age 6–23 months) n = 250							
Yes	143	92 (64.3%) [56.4%, 72.3%]	P = .019*	66 (46.2%) [37.9%, 54.4%]		P < .001*	
No	107	53 (49.5%) [39.9%, 59.2%]		25 (23.4%) [15.2%, 31.5%]			
Part B							
Characteristics of children	DV1 = No n = 105	DV1 = Yes n = 145	P-values (logistic regression)	DV2 = No n = 159	DV2 = Yes n = 91	P-values (logistic regression)	
Birth weight g mean (SD) [95% CI]	3234 (654) [3108, 3361]	3161 (615) [3060, 3262]	P = .365	3181 (614) [3085, 3277]	3210 (664) [3072, 3348]	P = .725	
Z-score birthweight adjusted for sex and gestational age mean (SD) [95%CI] {n}	+0.236 (1.1) [+0.2, +0.45] {n = 102}	+0.236 (1.2) [−2.4, +2.9] {n = 143}	P = .500	+0.13 (1.1) (−0.043, +0.30] {n = 156}	+0.27 (1.2) [+0.03, +0.52] {n = 89}	P = .381	
Gestational age at birth weeks median (IQR) [95%CI]	39 (38,40) [39,40]	39 (37,40) [29, 38]	P = .166	39 (38,40) [39, 39.4]	39 (37,40) [38,39]	P = .140	

Note: Categorical variables are shown as absolute and relative frequencies (percentages) with [95% confidence intervals]—child sex, method of birth, birth weight category, premature birth, feeding method, early childhood anaemia. Symmetrically distributed numerical variables are shown as mean (SD) and [95% confidence intervals]—birth weight, z-score birthweight. Numerical values with a skewed distribution are shown as median and inter-quartile range (IQR) and [95% confidence intervals] - gestational age at birth.

^aAEDC require that results relating to cell sizes of 15 or less are not provided to protect confidentiality.

*P < .05.

Strait Islander mothers or both Aboriginal and Torres Strait Islander mothers. The results of analysis following multiple imputation of missing values (n = 29) for

mothers who were anaemic both before and during pregnancy were consistent with the results of complete case analysis (Table 3).

TABLE 2 Developmental vulnerability (in lower 10th percentile of AEDC assessment score at first year of full-time school) in one or more domains (DV1) and in two or more domains (DV2) by characteristics of the mothers—bivariate analysis (logistic regression)

Part A					
Characteristics of mothers n = 250 unless otherwise specified	Number with that characteristic	AEDC—DV1 (Y) n = 145 (58.0%)	P-values (DV1—logistic regression)	AEDC—DV2 (Y) n = 91 (36.4%)	P-values (DV2—logistic regression)
Ethnicity of mother^a					
Aboriginal	132	91 (68.9%) [60.9%, 76.9%]	Base	62 (47.0%) [38.3%, 55.6%]	Base
Torres Strait Islander	97	42 (43.3%) [33.3%, 53.3%]	<i>P</i> < .001*	25 (25.8%) [16.9%, 34.6%]	<i>P</i> = .001*
Aboriginal and Torres Strait Islander	21	n ≤ 15	<i>P</i> = .288	n ≤ 15	<i>P</i> = .023*
Mother's region of residence^a					
Cairns and Hinterland	19	n ≤ 15	<i>P</i> = .218	n ≤ 15	<i>P</i> = .135
Cape York	134	90 (67.2%) [59.1%, 75.2%]	Base	60 (44.8%) [36.2%, 53.3%]	Base
Torres Strait	97	45 (46.4%) [36.3%, 56.5%]	<i>P</i> = .002*	26 (26.8%) [17.8%, 35.8%]	<i>P</i> = .006*
Mother's age when child born—quartiles					
Teenager— <i>younger</i> than 20 years	52	33 (63.5%) [49.9%, 77.0%]	<i>P</i> = .125	17 (32.7%) [19.5%, 45.9%]	<i>P</i> = .824
20-23 years	65	32 (49.2%) [36.7%, 61.7%]	Base	20 (30.8%) [19.2%, 42.9%]	Base
24-30 years	69	39 (56.5%) [44.5%, 68.5%]	<i>P</i> = .398	25 (36.2%) [24.6%, 47.9%]	<i>P</i> = .504
>30 years	64	41 (64.1%) [52.0%, 76.1%]	<i>P</i> = .908	29 (45.3%) [32.8%, 57.8%]	<i>P</i> = .090
SEIFA—mother's usual locality^a					
Mother lives in SEIFA 1 locality	219	128 (58.5%) [51.9%, 65.0%]	<i>P</i> = .703	80 (36.5%) [30.1%, 43.0%]	<i>P</i> = .910
Mother does NOT live in SEIFA 1 locality	31	17 (54.8%) [36.3%, 73.4%]		n ≤ 15	
Mother's parity—cohort pregnancy n = 173					
Parity 0-2	92	58 (63.0%) [53.0%, 73.1%]	<i>P</i> = .859	38 (41.3%) [31.1%, 51.6%]	<i>P</i> = .810
Parity 3 or more	81	50 (61.7%) [50.9%, 72.5%]		32 (39.5%) [28.6%, 50.4%]	

(Continues)

TABLE 2 (Continued)

Mother's body mass index n = 161									
Under or healthy weight	57	36 (63.2%) [50.2%, 76.1%]	Base P = .401	23 (40.4%) [27.2%, 53.5%]	Base P = .338				
Overweight	42	23 (54.8%) [39.1%, 70.5%]	P = .107	n = <15 [16.4, 45.5%]	P = .465				
Obese	62	30 (48.4%) [35.6%, 61.2%]		21 (33.9%) [21.8%, 46.0%]					
Smoking in pregnancy									
Yes	153	101 (66.0%) [58.4%, 73.6%]	P = .001*	65 (42.5%) [34.7%, 50.4%]	P = .013*				
No	97	44 (45.4%) [35.3%, 55.4%]		26 (26.8%) [17.8%, 35.8%]					
Diabetes in pregnancy (gestational diabetes OR pre-existing diabetes) n = 168									
Yes	41	24 (58.5%) [42.8%, 66.2%]	P = .905	16 (39.0%) [23.4%, 54.6%]	P = .746				
No	127	73 (57.5%) [48.8%, 66.2%]		46 (36.2%) [27.7%, 44.7%]					
Anaemia before and during pregnancy n = 221									
Yes	72	53 (73.6%) [63.2%, 84.0%]	P = .004*	35 (48.6%) [36.8%, 60.4%]	P = .019*				
No	149	79 (53.0%) [44.9%, 61.1%]		48 (32.2%) [24.6%, 39.8%]					
Red cell folate level insufficient before OR during pregnancy n = 61 ^a									
Yes	46	25 (54.4%) [39.4%, 69.3%]	P = .405	n = <15	P = .358				
No	n ≤ 15	n ≤ 15		n ≤ 15					
Part B									
Characteristics of mothers	DV1 = No n = 105	DV1 = Yes n = 145	P-values (logistic regression)	DV2 = No n = 159	DV2 = Yes n = 91	P-values (logistic regression)			
Mothers' age in years at baby's birth median (IQR) [95%CI]	24 (20.5, 29.5) [22, 25]	24 (20,31) [23, 26]	P = .570	23 (20,29) [22, 24]	26 (21,32) [23, 28.4]	P = .101			

Note: Categorical variables are shown as absolute and relative frequencies (percentages) with [95% confidence intervals]—mother's ethnicity, region of residence, age group, SEIFA, parity, Body Mass Index, smoking in pregnancy, diabetes in pregnancy, anaemia before and during pregnancy, folate levels insufficient. Numerical values with a skewed distribution are shown as median and inter-quartile range (IQR) and [95%CI]—mother's age in years at birth of baby.

^aAEDC require that results relating to cell sizes of 15 or less are not provided to protect confidentiality.

* P < .05.

TABLE 3 Risk factors for child (n = 250) developmental vulnerability (in lower 10th percentile of AEDC assessment score) in two or more domains (DV2): multi-variable analyses—complete case analyses and analysis with imputed data

Characteristic	Complete case analysis 1 N = 221				Analysis with imputed data ^a N = 250			
	DV2 (yes) 83 (37.6%)	DV2 (No) 138 (62.4%)	Odds-ratio (95% CI)	P-value	DV2 (yes) 91 (35.4%)	DV2 (no) 159 (63.6%)	Odds-ratio (95% CI)	P-value
Child anaemic when aged between 6 and 23 months								
No	24 (28.9%)	72 (52.2%)	1		25 (27.5%)	82 (51.6%)	1	
Yes	59 (71.1%)	66 (47.8%)	2.2 (1.1, 4.3)	P = 0.020	66 (72.5%)	77 (48.4%)	2.2 (1.2, 4.1)	P = .016
Mother anaemic before and during pregnancy								
No	48 (57.8%)	101 (73.2%)	1		53 (58.2%)	117 (73.6%)	1	
Yes	35 (42.2%)	37 (26.8%)	2.1 (1.1, 4.1)	P = 0.021	38 (41.8%)	42 (26.4%)	2.1 (1.1, 4.1)	P = .027
Mother smoked during pregnancy								
No	25 (30.1%)	60 (43.5%)	1		26 (28.6%)	71 (44.7%)	1	
Yes	58 (69.9%)	78 (56.5%)	2.0 (1.02, 3.9)	P = .045	65 (71.4%)	88 (55.4%)	2.2 (1.2, 4.1)	P = .016
Sex of child								
Female	26 (31.3%)	76 (55.1%)	1		29 (31.9%)	89 (56.0%)	1	
Male	57 (68.7%)	62 (44.9%)	2.8 (1.5, 5.3)	P = .001	62 (68.1%)	70 (44.0%)	2.8 (1.6, 5.1)	P = .001
Ethnicity of mother								
Torres Strait Islander	24 (28.9%)	67 (48.6%)	1		25 (27.5%)	70 (44.0%)	1	
Aboriginal	55 (66.3%)	59 (42.8%)	2.5 (1.3, 5.0)	P = .009	62 (68.1%)	72 (45.3%)	2.4 (1.3, 4.7)	P = .008
Both ^a	n ≤ 15	n ≤ 15	1.0 (0.3, 3.8)	P = .990	n ≤ 15	n ≤ 15	0.7 (0.2, 2.6)	P = .623
Age of mother at birth of child								
≤30 years	55 (66.3%)	106 (76.8%)	1		62 (68.1%)	124 (78.0%)	1	
>30 years	28 (33.7%)	32 (23.2%)	2.1 (1.04, 4.2)	P = .039	29 (31.9%)	35 (22.0%)	2.0 (1.03, 4.0)	P = .041

Note: The following characteristics of children and their mothers were considered during multivariable analyses: sex; birthing method (non-instrumental vaginal, instrumental vaginal, caesarean section); gestational age at birth; premature or not; birth weight; feeding method to age 4 months (only breast milk, only infant formula, both breast milk and formula); ethnicity of mother (Aboriginal, Torres Strait Islander, both); region of residence of mother; SEIFA category for residence of mother; maternal age; BMI category of mother (underweight, normal weight, overweight, obese); categories of parity (0–2, ≥3); smoking during pregnancy; diabetes during pregnancy; mother had insufficient red cell folate (RCF) level before or during pregnancy; mother anaemic both before and during pregnancy. Models include all variables shown in Table 3 adjusted for the confounding effect of birth method (no missing values imputed for birth methods). Imputed data are averages of 20 imputations.

Abbreviation: 95% CI, 95% confidence interval.

^aAEDC require that results relating to cell sizes of 15 or less are not provided to protect confidentiality.

4 | DISCUSSION

To the best of our knowledge, this is the first study to demonstrate the association of early childhood anaemia with developmental vulnerability at school age in Australia. Children who had anaemia between age six and 23 months had more than twice the risk of developmental vulnerability at school age compared to those who had not been anaemic. Early childhood anaemia was associated with developmental vulnerability in four of the five domains of early childhood development. Anaemia of mothers also doubled the risk of developmental vulnerability of their children at school age. These findings are consistent with the persistent detrimental effects of iron deficiency during critical phases of neurological development before birth and in early childhood.¹

One limitation of this study is the small numbers of participants. However, the number of children with a Ferret record available (n = 1155) is close to census population information (n = 1147) for the remote communities where the Ferret system was used (Table S4). While the proportion of children with a haemoglobin measured (74.0%) is lower than expected, this is similar (85.0% and 72.1%) to two reports from elsewhere in northern Australia.^{7,11} The proportion of children with a haemoglobin measurement for whom AEDC assessment results were available—250/708 (35.3%)—is consistent with the triennial schedule of the AEDC. Our findings in respect of child's sex, anaemia of mothers and smoking in pregnancy concur with the findings of much larger studies in South Australia and the Northern Territory.^{15,16} Nevertheless, additional studies would be of value to determine whether early childhood

anaemia is associated with developmental vulnerability in other similar settings.

Another limitation is the absence of information on the causes of the early childhood anaemia among these children. Iron deficiency is the usual cause of anaemia in early life.^{5,12} Two Northern Territory studies found iron deficiency was the main cause of anaemia in children there; among 74 pre-school aged children with anaemia 62 (84%) were iron deficient; among 66 school-aged anaemic children, 55 (83%) responded to iron therapy.^{27,28} Comparable studies are needed for Far North Queensland.

The information on child development used here, is derived from the triennial nation-wide AEDC. Initially developed in Canada and adapted subsequently for use in Australia and elsewhere, concerns have been raised about the appropriateness of the AEDC instrument in Australian Aboriginal and Torres Strait Islander settings.^{29,30} Research undertaken in 2008 to 2009 in urban, regional and remote Western Australia provided the basis of an adaptation of the AEDC methodology so that teachers are supported by Indigenous cultural consultants during the assessment of childhood development of Aboriginal and Torres Strait Islander children.²⁹ One strength of the AEDC information is that the information provided encompasses all children in Australia from diverse cultural and linguistic groups including Aboriginal and Torres Strait Islander children.¹³

Our findings show a higher risk of developmental vulnerability among children of Aboriginal mothers compared to children of Torres Strait Islander mothers. This may reflect different historical experiences as Aboriginal people of mainland Queensland were subject to extensive forced removals compared to the Torres Strait.³¹ Loss of access to nutrient-dense traditional food was only one of many grim consequences of this policy.³¹

In settings where the prevalence of early childhood anaemia exceeds 20%, such as Far North Queensland and elsewhere in northern Australia,^{7,10} the World Health Organization recommends interventions that combine nutrition promotion with provision of multi-micronutrient preparations that include iron, for fortification of complementary food.^{12,32} These interventions have been shown to be safe and effective.³² Improvements in haemoglobin levels resulting from such interventions protect against early childhood anaemia, and are also protective against the development deficits associated with early childhood anaemia.³³ Potential mechanisms for this improvement include increased oxygen delivery to tissues and brain facilitating increased interaction of the child with his/her environment.³³ Other possible mechanisms are improvements in iron availability for myelination, synaptogenesis and neurotransmission.³³ One such intervention has been successfully piloted in six remote communities across northern

Australia.³⁴ Operational experience of implementation of these nutrition promotion and anaemia prevention interventions in 65 low- and middle-income countries has been recently compiled to assist program managers and researchers manage the challenges entailed in implementation at scale.^{35,36}

Interventions to prevent anaemia in early childhood will be strengthened by improved food security, and by prevention and treatment anaemia in pregnancy.^{8,37} Evaluation is essential, in particular to assess if prevention of maternal and early childhood anaemia translates into improved developmental indicators for children at school-age.^{1,33}

The Australian government is committed to Closing the Gap between Aboriginal and Torres Strait Islander peoples and other Australians.³⁸ Anaemia prevention and better nutrition provide an opportunity to improve early childhood development, educational attainment and contribute to Closing the Gap between Aboriginal and Torres Strait Islander people and other Australians.

CONFLICT OF INTEREST

The authors affirm that they have no conflict of interest to declare.

AUTHOR CONTRIBUTIONS

D.L. conceived the research, obtained the necessary approvals to secure the data required, conducted preliminary statistical analysis and prepared the first draft of this manuscript. F.T. assisted with data management and preparation, and contributed to statistical analysis. P.B. contributed to study design and guided, supervised and contributed to statistical analysis. R.M. and M.M. contributed to study design, and manuscript development and preparation.

D.L. was supported by a National Health and Medical Research Council post-graduate scholarship APP1092732. The funding body had no role in this research and the researchers were independent from the funder. Other agencies provided non-financial support to this research. We would like to acknowledge and thank the Aboriginal and Torres Strait Islander leaders of the key community-controlled health service organisations in Far North Queensland who considered and endorsed the research proposal. In addition, we acknowledge and thank the Data Custodians for the Queensland Health data collections and the Data Custodian for The Australian Government Department of Education and Training Australian Early Development Census data collection, their research and data management staff for their assistance and support for this work. These agencies extracted, linked and de-identified the information used in this research and provided the linked de-identified dataset to the research

team. Other than this, these agencies had no role in the research.

This research uses data from the Australian Early Development Census (AEDC). The AEDC is funded by the Australian Government Department of Education and Training. The findings and views reported are those of the authors and should not be attributed to the Department or the Australian Government.

ORCID

Dympna Leonard  <https://orcid.org/0000-0002-5081-6070>

REFERENCES

- Georgieff MK, Ramel SE, Cusick SE. Nutritional influences on brain development. *Acta Paediatr.* 2018;107(8):1310-1321.
- Bhutta ZA, Guerrant RL, Nelson CA 3rd. Neurodevelopment, Nutrition, and Inflammation: The Evolving Global Child Health Landscape. *Pediatrics.* 2017;139(Suppl 1):S12-S22.
- Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutr Rev.* 2014;72(4):267-284.
- Larson LM, Phiri KS, Iron PSR. Cognitive Development: What Is the Evidence? *Ann Nutr Metab.* 2017;71(Suppl 3):25-38.
- Domellöf M. Iron requirements in infancy. *Ann Nutr Metab.* 2011;59(1):59-63.
- Georgieff MK. Iron assessment to protect the developing brain. *Am J Clin Nutr.* 2017;106:1588S-1593S.
- Aquino D, Leonard D, Hadgraft N, Marley J. High prevalence of early onset anaemia amongst Aboriginal and Torres Strait Islander infants in remote northern Australia. *Australian J Rur Health.* 2018;26:245-250.
- Leonard D, Buettner P, Thompson F, Makrides M, McDermott R. Anaemia in pregnancy among Aboriginal and Torres Strait Islander women of Far North Queensland: a retrospective cohort study. *Nutrition and Dietetics.* 2018;75:457-467.
- Bar-Zeev S, Barclay L, Kruske S, Kildea S. Factors affecting the quality of antenatal care provided to remote dwelling Aboriginal women in northern Australia. *Midwifery.* 2014;30(3):289-296.
- Leonard D, Buettner P, Thompson F, Makrides M, McDermott R. Anaemia in early childhood among Aboriginal and Torres Strait Islander children of Far North Queensland: A retrospective cohort study. *ANZJPH.* 2019;43(4):319-327.
- Bar-Zeev SJ, Kruske SG, Barclay LM, Bar-Zeev N, Kildea SV. Adherence to management guidelines for growth faltering and anaemia in remote dwelling Australian Aboriginal infants and barriers to health service delivery. *BMC Health Serv Res.* 2013;13:250.
- World Health Organization. *Nutritional Anaemias: Tools for Effective Prevention and Control Geneva.* Geneva: World Health Organization; 2017.
- Australian Early Development Census. *Australian Early Development Census National Report 2015.* Canberra: Australian Government Department of Education; 2016 Accessed December 13, 2018. <https://www.aedc.gov.au/resources/detail/about-the-aedc-data-collection>.
- Brinkman S, Gregory T, Harris J, Hart B, Blackmore S, Janus M. Associations between the Early Development Instrument at Age 5 and Reading and Numeracy Skills at Ages 8, 10 and 12: a Prospective Linked Data Study. *Child Ind Res.* 2013;6:695-708.
- Guthridge S, Lia L, Silburn S, Lia SQ, McKenzie J, Lynch J. Early influences on developmental outcomes among children, at age 5, in Australia's Northern Territory. *Early Child Res Q.* 2016;35:124-134.
- Chittleborough CR, Searlea AK, Smithers LG, Brinkmana S, Lynch JW. How well can poor child development be predicted from early life characteristics? A whole-of-population data linkage study. *Early Child Res Q.* 2016;35:19-30.
- Leonard D, Buettner P, Thompson F, Makrides M, McDermott R. Linking 'data silos' to investigate anaemia among Aboriginal and Torres Strait Islander mothers and children in Far North Queensland. *ANZJPH.* 2018;42(5):256-262.
- Queensland Health. *Primary Clinical Care Manual Paediatrics.* 9th ed. Brisbane: Queensland Government; 2017. <https://www.publications.qld.gov.au/>
- World Health Organization. *Optimal serum and red blood cell folate concentrations in women of reproductive age for prevention of neural tube defects.* Geneva: World Health Organization; 2015.
- Rogers LM, Cordero AM, Pfeiffer CM, et al. Global folate status in women of reproductive age: a systematic review with emphasis on methodological issues. *Ann N Y Acad Sci.* 2018;1431(1):35-57.
- Villar J, Ismail LC, Victora CG, et al. International standards for newborn weight, length and head circumference by gestational age and sex: the Newborn Cross-Sectional Study of the INTERGROWTH-21st Project. *Lancet.* 2014;384:857-868.
- Intergrowth-21st Network. (INTERGROWTH-21ST version 1.3.5); 2014. <https://intergrowth21.tghn.org/newborn-size-birth/#c4> Accessed April 2018.
- Australian Bureau of Statistics. Australian Census of Population and Housing. Socio-Economic Indexes for Areas (SEIFA). *Canberra.* 2011;2033.0.55.001:2013.
- Kleinbaum DG, Kupper LL, Morgenstern H. *Epidemiologic research. Principles and quantitative methods.* New York, New York: Van Nostrand Reinhold; 1982.
- Sterne JAC, White IR, Carlin JB, et al. Multiple imputation for missing data in epidemiological and clinical research: potential and pitfalls. *BMJ.* 2009;338:b2393.
- STROBE. Strengthening the reporting of observational studies in epidemiology—checklist. https://www.strobe-statement.org/fileadmin/Strobe/uploads/checklists/STROBE_checklist_v4_cohort.pdf2007. Accessed June 2019.
- Kruske SG, Ruben AR, BD R. An iron treatment trial in an Aboriginal community: improving non-adherence. *J Paediatr Child Health.* 1999;35:153-158.
- Udovicich C, Perera K, Leahy C. Anaemia in school-aged children in an Australian Indigenous community. *Aust J Rural Health.* 2017;25(5):285-289.
- Silburn S, Brinkman S, Ferguson-Hill S, Styles I, Walker R, Shepherd C. *The Australian Early Development Index (AEDI) Indigenous Adaptation Study.* Perth: Menzies School of Health Research; 2009 http://ccde.menzies.edu.au/sites/default/files/resources/AEDI_Indigenous_Adaptation_Study_Report_Nov_2009.pdf.

30. Janus M, Reid-Westoby, C. *Monitoring the Development of all Children: the Early Development Instrument*. Hamilton, ON: McMaster University; 2016. https://bernardvanleer.org/app/uploads/2016/07/Early-Childhood-Matters-2016_7-1.pdf
31. Australian Law Reform Commission. *Pathways to Justice—An Inquiry into the Incarceration Rate of Aboriginal and Torres Strait Islander Peoples*. Sydney: Australian Law Reform Commission; 2017 <http://apo.org.au/system/files/138661/apo-nid138661-685316.pdf>
32. Dewey KG, Yang ZY, Boy E. Systematic review and meta-analysis of home fortification of complementary foods. *Matern Child Nutr*. 2009;5(4):283-321.
33. Larson LM, Kubes JN, Ramirez-Luzuriaga MJ, Khishen S, A HS, Prado EL. Effects of increased hemoglobin on child growth, development, and disease: a systematic review and meta-analysis. *Ann N Y Acad Sci*. 2019;1450:83-104.
34. Aquino D, Marley J, Senior K et al. The Early Childhood Nutrition and Anaemia Prevention Project—Summary Report. <https://healthinonet.ecu.edu.au/key-resources/publications/25699/?title=Early%20Childhood%20Nutrition%20and%20Anaemia%20Prevention%20Project%20%3A%20summary%20report%3A%20The%20Fred%20Hollows%20Foundation> 2013.
35. Nyhus Dhillon C, Sarkar D, Klemm RD, et al. Executive summary for the micronutrient powders consultation: lessons learned for operational guidance. *Matern Child Nutr*. 2017;13 (Suppl 1):1-9.
36. Pelletier D, DePee S. Micronutrient powder programs: New findings and future directions for implementation science. *Matern Child Nutr*. 2019;15(S5):e12802.
37. Moradi S, Arghavani H, Issah A, Mohammadi H, Mirzaei K. Food insecurity and anaemia risk: a systematic review and meta-analysis. *Public Health Nutr*. 2018;18:1-13.
38. Department of the Prime Minister and Cabinet. Closing the Gap Report 2019. <https://ctgreport.niaa.gov.au/sites/default/files/ctg-report-20193872.pdf?a=1:%202019>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Leonard D, Buettner P, Thompson F, Makrides M, McDermott R. Early childhood anaemia more than doubles the risk of developmental vulnerability at school-age among Aboriginal and Torres Strait Islander children of remote Far North Queensland: Findings of a retrospective cohort study. *Nutrition & Dietetics*. 2020;77:298-309. <https://doi.org/10.1111/1747-0080.12602>