

# Endemic Infection of the Amphibian Chytrid Fungus in a Frog Community Post-Decline

Richard W. R. Retallick<sup>1,2,3</sup>, Hamish McCallum<sup>2\*</sup>, Rick Speare<sup>4</sup>

**1** School of Tropical Biology, James Cook University, Townsville, Queensland, Australia, **2** Department of Zoology and Entomology, University of Queensland, St Lucia, Queensland, Australia, **3** School of Life Sciences—Biology, Arizona State University, Tempe, Arizona, United States of America, **4** Amphibian Diseases Group, School of Public Health and Tropical Medicine, James Cook University, Townsville, Queensland, Australia

**The chytrid fungus *Batrachochytrium dendrobatidis* has been implicated in the decline and extinction of numerous frog species worldwide. In Queensland, Australia, it has been proposed as the cause of the decline or apparent extinction of at least 14 high-elevation rainforest frog species. One of these, *Taudactylus eungellensis*, disappeared from rainforest streams in Eungella National Park in 1985–1986, but a few remnant populations were subsequently discovered. Here, we report the analysis of *B. dendrobatidis* infections in toe tips of *T. eungellensis* and sympatric species collected in a mark-recapture study between 1994 and 1998. This longitudinal study of the fungus in individually marked frogs sheds new light on the effect of this threatening infectious process in field, as distinct from laboratory, conditions. We found a seasonal peak of infection in the cooler months, with no evidence of interannual variation. The overall prevalence of infection was 18% in *T. eungellensis* and 28% in *Litoria wilcoxii/jungguy*, a sympatric frog that appeared not to decline in 1985–1986. No infection was found in any of the other sympatric species. Most importantly, we found no consistent evidence of lower survival in *T. eungellensis* that were infected at the time of first capture, compared with uninfected individuals. These results refute the hypothesis that remnant populations of *T. eungellensis* recovered after a *B. dendrobatidis* epidemic because the pathogen had disappeared. They show that populations of *T. eungellensis* now persist with stable, endemic infections of *B. dendrobatidis*.**

Citation: Retallick RWR, McCallum H, Speare R (2004) Endemic infection of the amphibian chytrid fungus in a frog community post-decline. PLoS Biol 2(11): e351.

## Introduction

Increasingly, the amphibian chytrid fungus (*Batrachochytrium dendrobatidis*) has been implicated as a major contributor to global catastrophic declines in frog populations (Berger et al. 1998; Daszak et al. 1999, 2003). It has been found on frogs in areas where catastrophic declines were reported, it has been shown in the laboratory to be highly pathogenic to some species, and there is pathological evidence to link this fungal parasite to host mortality (Berger et al. 1998). The pathogen may therefore be capable of producing the extremely high mortality observed during declines. However, little information is available on the impact of the fungus on individuals in the field, rather than the laboratory. Furthermore, little has been published on the prevalence of infection among frog populations as a whole, as distinct from the prevalence among morbid frogs only. In addition to data on the prevalence of the pathogen among morbid animals, information on the prevalence of a putative pathogen in the population in general is important to determine the potential effect of the pathogen on the host population (McCallum and Dobson 1995).

In Queensland, Australia, there have been extinctions or major declines of at least 14 frog species in undisturbed, high-elevation rainforest streams, commencing in 1979–1981 in the Conondale and Blackall Ranges (26°50' S, 152°41' E), followed in 1985–1986 in the Eungella region of the Clarke Range (21°07' S, 148°29' E), and, in 1990–1995, in the Wet Tropics bioregion (17°22' S, 145°49' E) (Laurance et al. 1996; McDonald and Alford 1999). Laurance et al. (1996, 1997)

suggested that an epidemic disease was responsible for all these declines, without proposing an agent. Despite the presence of *B. dendrobatidis* in ill and dead frogs collected from the Big Tableland in the Wet Tropics in 1993, it was not recognised as a pathogenic organism until 1998 (Berger et al. 1998) and was described as a new species of fungus in 1999 (Longcore et al., 1999). *B. dendrobatidis* has been suggested as the causative agent of many of the east coast Australian declines (Berger et al. 1999a), although only the decline at Big Tableland (McDonald and Alford 1999) had direct evidence of an association with the presence of *B. dendrobatidis*.

In this paper, we report the retrospective analysis of *B. dendrobatidis* infection on toe tips collected between 1994 and 1998 from six species of frogs at Eungella National Park in east-central Queensland, Australia (Figure 1). Declines at this location were particularly catastrophic. Between 1985 and 1986, the Eungella Gastric-Brooding Frog (*Rheobatrachus vitellinus*) disappeared suddenly from relatively undisturbed


Received March 26, 2004; Accepted August 12, 2004; Published October 5, 2004  
DOI: 10.1371/journal.pbio.0020351

Copyright: © 2004 Retallick et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abbreviations: AIC, Akaike Information Criterion;  $\Delta dev$ , change in deviance; se, standard error

Academic Editor: Georgina M. Mace, Institute of Zoology, Zoological Society of London

\*To whom correspondence should be addressed. E-mail: hmccallum@zen.uq.edu.au



**Figure 1.** Location of Study Sites

1, Mount David Creek; 2, Mount William Creek; 3, Dooloomai Falls; 4, Rawson Creek; 5, Picnic Ground Creek; and 6, Tree Fern Creek.  
DOI: 10.1371/journal.pbio.0020351.g001

rainforest streams, and it is now considered to be extinct (McDonald 1990; Campbell 1999; Department of Environment and Heritage 2003). During the same period, the Eungella Torrent Frog (*Taudactylus eungellensis*) also disappeared, but it was later found to have persisted in a few small populations (McDonald 1990; Couper 1992; McNellie and Hero 1994; Retallick et al. 1997), which are the subject of this paper. A suite of other species that coexisted with *R. vitellinus* and *T. eungellensis* showed no evidence of decline. Unfortunately, the extent of infection with *B. dendrobatidis* in the frog community at Eungella during the period of decline is unknown. The first record of *B. dendrobatidis* at Eungella was from a moribund frog collected in 1995 (Berger et al. 1999a).

## Results

We detected *B. dendrobatidis* on 71 (15.0%) of the 474 toes assessed. Four species showed no sign of infection, while we detected infections on 58 (18.4%) of 316 toes of *T. eungellensis* and 13 (34%) of 47 toes of frogs identified at the time as *Litoria lesueuri* (Table 1). Recently, the taxonomy of *L. lesueuri* has been revised (Donnellan and Mahoney, 2004), with North Queensland members of the complex being either *L. jungguy* or *L. wilcoxii*. From the reported distributions of these species, our “*L. lesueuri*” may have been either one of the species or even hybrids (Donnellan and Mahoney, 2004; Michael Mahoney, personal communication) These species can be distinguished only with genetic information. We therefore describe them as *L. wilcoxii/jungguy* for the remainder of this paper.

Using a logistic model with season (summer, autumn, winter, and spring), species, and site as predictor variables, we found significant effects of each variable on infection, corrected for the effects of the other variables (for season, the change in deviance [ $\Delta dev$ ] = 15.25,  $df = 3$ ,  $p = 0.0061$ ; for site,  $\Delta dev = 13.56$ ,  $df = 5$ ,  $p = 0.02$ ; for species,  $\Delta dev = 17.32$ ,  $df = 5$ ,  $p = 0.004$ ). The most parsimonious model (i.e., one that minimizes the Akaike Information Criterion [AIC]) included each of these predictors but no interaction terms.

Further analysis was concentrated on the two species on which *B. dendrobatidis* was detected, namely *T. eungellensis* and *L. wilcoxii/jungguy*.

### Infection in *T. eungellensis*

The proportions of frogs that were infected in the three largest populations (at Rawson Creek, Dooloomai Falls, and Tree Fern Creek [Figure 1]) differed significantly among those sites ( $\Delta dev = 12.84$ ,  $df = 2$ ,  $p = 0.001$ ). A significantly smaller proportion was infected at Rawson Creek (10.5%) than at Dooloomai Falls (26.7%) or Tree Fern Creek (25.0%). There was a marginally significant difference overall in the infection levels among males, females, and subadults ( $\Delta dev = 5.9$ ,  $df = 2$ ,  $p = 0.052$ ). However, when site was included in the model, any suggestion of a difference in infection level between males, females, or subadults disappeared. When we separated the main *T. eungellensis* populations, it was apparent that the difference in prevalence between those categories was influenced by the population at Tree Fern Creek, where 46% of males ( $n = 13$ ) and no females or subadults were

**Table 1.** Prevalence of Infection of *B. dendrobatidis* in All Frog Toes Examined

	Dooloomai Falls	Mt. David	Mt. William	Picnic Grnd	Rawson Creek	Tree Fern Creek
<i>L. chloris</i>	0 (26)	0 (1)		0 (1)	0 (14)	
<i>L. wilcoxii/jungguy</i>	0.167 (6)			0.286 (35)		0.333 (6)
<i>L. revelata</i>		0 (18)	0 (32)			
<i>M. fasciolatus</i>		0 (4)	0 (1)			
<i>T. eungellensis</i>	0.271 (129)	0 (2)		0.25 (4)	0.102 (157)	0.25 (24)
<i>T. liemi</i>	0 (2)	0 (5)	0 (7)			

Sample sizes are in parentheses. A small number of cases with uncertain diagnoses have been omitted.  
DOI: 10.1371/journal.pbio.0020351.t001

**Table 2.** Prevalence of Infection of *B. dendrobatidis* in *T. eungellensis* by Age/Sex Class and Site, Pooled over Sampling Times

Site	Females	Males	Subadults
Dooloomai Falls	0.286 (21)	0.293 (41)	0.241 (54)
Rawson Creek	0.045 (22)	0.136 (22)	0.110 (109)
Tree Fern Creek	0.000 (6)	0.462 (13)	0.000 (5)

Sample sizes are shown in parentheses. A small number of metamorphlings and adult frogs, the sex of which was not recorded, have been excluded.  
DOI: 10.1371/journal.pbio.0020351.t002

infected (Table 2). When we grouped all sites and times, the estimated overall prevalence of infection was 18.1%.

At these three sites, levels of infection among *T. eungellensis* varied significantly among seasons ( $\Delta dev = 14.605$ ,  $df = 3$ ,  $p = 0.002$ ), but not among years ( $\Delta dev = 3.433$ ,  $df = 3$ ,  $p = 0.26$ ). Further, there was no evidence that the pattern of seasonal variation in infection changed among years ( $\Delta dev = 5.561$ ,  $df = 6$ ,  $p = 0.49$ ). Infection was most prevalent (37.8%) during the winter months (1 June to 31 August) and least prevalent (11.3%) during the summer months (1 December to 28/29 February). Comparing the two sites with the largest sample sizes, Dooloomai Falls and Rawson Creek, there was no evidence that they had differing seasonal patterns of infection ( $\Delta dev = 5.32$ ,  $df = 3$ ,  $p = 0.150$ ), although the level of infection overall was much higher at Dooloomai Falls (log odds ratio = 1.1612, standard error [se] = 0.3586,  $p = 0.0012$ ). Infection levels were much higher in winter and spring combined than in summer and autumn (log odds ratio = 1.360, se = 0.3589,  $p = 0.00015$ ), and there was no evidence of infection levels differing between winter and spring or between summer and autumn ( $\Delta dev = 1.107$ ,  $df = 2$ ,  $p = 0.5$ ). The seasonal changes in infection at the two sites are shown in Figure 2.

### Infection in *L. wilcoxii/jungguy*

*B. dendrobatidis* was detected in 13 of the 47 *L. wilcoxii/jungguy* examined, giving an estimated prevalence of 27.7%, with a 95% confidence interval ranging from 15.6% to 42.6%. There was no evidence that prevalence of infection differed among males, females, and subadults ( $\Delta dev = 2.32$ ,  $df = 2$ ,  $p = 0.31$ ); among sites ( $\Delta dev = 0.175$ ,  $df = 2$ ,  $p = 0.91$ ); or among seasons ( $\Delta dev = 1.44$ ,  $df = 1$ ,  $p = 0.7$ ) (each of the above  $\Delta dev$  terms is corrected for the other terms in the model).


Comparing the prevalence of infection between *L. wilcoxii/jungguy* and *T. eungellensis* was hampered by the fact that prevalence of infection on *T. eungellensis* differed between sites, and that the sampled populations of the two frog species had largely disjunct distributions. When the data were pooled over all sites, there was insufficient evidence to reject a null hypothesis of equal prevalence in the two species, whether the effects of season were allowed for ( $\Delta dev = 0.15$ ,  $df = 1$ ,  $p = 0.7$ ) or not ( $\Delta dev = 2.09$ ,  $df = 1$ ,  $p = 0.15$ ).

### Influence of Infection on Recapture and Survival

Table 3 shows the numbers of *L. wilcoxii/jungguy* and *T. eungellensis* recaptured at any stage later in the study, grouped by their infection status at first capture. A logistic model with recapture as the response and species and infection status as predictors produced no evidence that the effect of *B. dendrobatidis* infection on recapture probability differed

between the species ( $\Delta dev = 0.002$ ,  $df = 1$ ,  $p = 0.96$ ). In both species, the probability of recapture was significantly lower for infected frogs than for uninfected frogs ( $\Delta dev = 5.34$ ,  $df = 1$ ,  $p = 0.02$ ; log odds ratio = -0.6464, se = 0.2856), correcting for the substantially higher overall recapture rate of *T. eungellensis* relative to *L. wilcoxii/jungguy*.

This simple analysis suggests that *B. dendrobatidis* may affect survival, but confounds the probability of recapturing a frog that is present at a site with the probability that a frog is still present at the site. The more sophisticated analysis that follows separates these two components, although it cannot distinguish between death and permanent emigration from



**Figure 2.** Seasonal Patterns of Prevalence of *B. dendrobatidis* in *T. eungellensis*

Solid circles show the observed prevalence, with binomial 95% confidence limits, in frogs pooled over years and age/sex class. The dashed line shows the prevalence predicted from the best-fitting logistic model. Numbers in brackets above each error bar are the sample sizes.

DOI: 10.1371/journal.pbio.0020351.g002

**Table 3.** Recapture and Survival of Infected and Uninfected Frogs

Species	Frogs Marked	Result	Frogs Recaptured (Proportion)	Recapture Period (Days)					Range (Days)
				<30	31–90	91–180	181–370	>371	
<i>L. wilcoxiijungguy</i>	13	+ve	2 (15.4%)	1		1			27–94
	34	–ve	9 (26.5%)	1	2	2	4	1	29–428
<i>T. eungellensis</i>	58	+ve	20 (34.5%)	2	2	2	11	3	25–1,089
	258	–ve	129 (50.0%)	8	23	20	28	50	21–1,449

Shown are numbers of marked *L. wilcoxiijungguy* and *T. eungellensis* frogs that were infected (+ve) and uninfected (–ve) and recaptured during the 4-y monitoring study, and maximum periods of time over which those frogs were known to survive after being toe-clipped.

DOI: 10.1371/journal.pbio.0020351.t003

the site. For brevity, we refer to continued presence at the site as “survival.”

Investigating mark–recapture data for both Rawson Creek and Dooloomai Falls, we found that the best model (on the basis of minimizing the AIC) for which there were sufficient data to estimate almost all parameters had the probability of recapture varying with time but not group (i.e., infected and uninfected males, females, and subadults), and the survival probability differing between the groups but constant through time. However, further analysis in both cases showed that there was no evidence that survival differed between infected and uninfected frogs: At both sites, the group effect was due to survival being highest in females, intermediate in males, and lowest in subadults (Figure 3; Table 4). In the case of Tree Fern Creek, which had a smaller total sample size than the other two sites, and in which only males were infected, the best model had recapture probabilities constant with both time and group, and survival varying with group. At this site, there was some evidence that infected males had lower survival than uninfected males (Figure 3; Table 4).


## Discussion

Our results show unequivocally that remnant populations of *T. eungellensis*, a rainforest frog that almost disappeared as a result of major die-offs in the Eungella area in the mid 1980s, now persist with stable infections of *B. dendrobatidis*. This does not imply that this pathogen cannot have been responsible for the decline. One hypothesis that can be discarded, however, is that *B. dendrobatidis* drove frog populations to low levels, consequently became locally extinct itself, and that frog populations subsequently recovered or stabilised in the absence of the pathogen. Two other hypotheses are consistent with our observed results.

It may be that *B. dendrobatidis* was not responsible for the initial decline of *T. eungellensis* populations. These populations declined more than a decade before this chytrid was formally identified, and the declines occurred over a very short period (McDonald 1990). Because of the rapidity of the declines, no samples were collected during the period of declines, so it is not possible to examine for the presence of *B. dendrobatidis*. Although it is not possible to eliminate all environmental factors as being responsible for the decline of *T. eungellensis* and the disappearance of *R. vitellinus* from Eungella in 1985, the failure to detect substantial climatic

anomalies in the area at that time (McDonald 1990) means that this explanation is unlikely to account for the sudden declines.

Another possibility consistent with our results is that *B.*



**Figure 3.** Estimated Quarterly Survival of Age/Sex Classes of *Taudactylus eungellensis* at Three Sites

Survival for uninfected frogs is shown with circles, and survival for infected frogs is shown with squares; 95% confidence limits around each point are also shown. Where there is no survival estimate for infected frogs, there were insufficient data to estimate the parameter. Sample sizes are given in Table 2.

DOI: 10.1371/journal.pbio.0020351.g003

**Table 4.** Summary of Results of Mark-Recapture Modelling on *T. eungellensis*

Site	Survival Probability Function	Recapture Probability Function	AICc	$\Delta$ AICc	Model Likelihood	Number of Parameters Estimated	Deviance
Dooloomai Falls	Male/female/subadult	Time	427.216	0	1	13	253.105
	Male/female/subadult, infection	Time	430.818	3.6	0.1651	16	249.76
	Male/female/subadult, infection	Time + Male/female/subadult, infection	430.947	3.73	0.1549	21	237.827
	Adult/subadult	Time	432.236	5.02	0.0813	12	260.394
Rawson Creek	Male/female/subadult	Time	710.292	0	1	12	267.793
	Adult/subadult	Time	711.028	0.74	0.6921	11	270.698
	Male/female/subadult, infection status	Time	713.701	3.41	0.1819	15	264.602
Tree Fern Creek	Female/subadult/infected male/uninfected male	Constant	152.294	0	1	5	123.644
	Constant	Female/subadult/infected male/uninfected male	154.37	2.08	0.3541	5	125.721
	Female/subadult/infected male/uninfected male	Female/subadult/infected male/uninfected male	155.442	3.15	0.2073	8	119.319
	Male/female/subadult	Constant	159.062	6.77	0.0339	4	132.745
	Female/subadult/infected male/uninfected male	Time	167.132	14.84	0.0006	17	103.21

“AICc” is the corrected Akaike Information Criterion; “ $\Delta$ AICc” is the change in AICc from the “best” model; and “Model Likelihood” is the likelihood of the model relative to the best model. A parametric bootstrap analysis using the most complex model that could be parameterised from the capture histories (a model with additive effects of group and time on both recapture and survival) indicated that in each case the goodness of fit was adequate.

DOI: 10.1371/journal.pbio.0020351.t004

*dendrobatidis* was a pathogen novel to the ecosystem in 1985, and that it was indeed responsible for the declines. Available molecular evidence (Morehouse et al. 2003), although not conclusive, is consistent with the hypothesis that *B. dendrobatidis* is a recently emerged disease agent. Populations of *T. eungellensis* may have recovered or stabilised following evolution of resistance to the pathogen in the frogs, or evolution of less-pathogenic strains of the fungus. Rapid coevolutionary changes in both host and pathogen following the introduction of a novel pathogen are to be expected, with the best-known example being the coevolution of myxomatosis and its rabbit hosts within a few years of the introduction of the virus to Australia in 1953 (May and Anderson 1983; Fenner and Fantini 1999). Our study provides no direct evidence for such coevolution in frogs and *B. dendrobatidis*, but it identifies this as an important area for future research. If it could be shown that susceptible frog populations were able to develop resistance to the amphibian chytrid, then captive breeding and artificial selection for resistance would provide an avenue for management of this threat to critically endangered species.

It is intriguing that we found similar seasonal fluctuations in infection levels within each year, but no evidence of

variation among the 4 y of the study. This suggests strongly that *B. dendrobatidis* has become endemic and relatively stable in prevalence in these populations. Given that frog numbers and diversity remained broadly similar over this period, it suggests that some form of host–pathogen equilibrium has become established, in contrast to the epizootic that may have occurred 10 y previously.

There are some obvious methodological limitations in our study that we must acknowledge. Infection status was determined by histological examination of toe tips at the time of first capture. Toe tips are often examined for *B. dendrobatidis* because the feet of frogs are a particularly favoured body location for infection by the fungus (Berger 2001). However, in light infections, *B. dendrobatidis* can occur in microscopic clusters, which can potentially be missed in histological sections. Hence, a proportion of frogs found to be negative for *B. dendrobatidis* using histology at first capture may actually have been infected. This possibility of false negatives (Berger 2001; Berger et al. 2002) suggests that the infection prevalences found in this study may underestimate the true prevalence of *B. dendrobatidis* within these frog populations. For individuals classed as infected at the time of first capture, and that subsequently survived for extended

periods, it is not known whether their infection persisted or was cleared after first capture. There is evidence that some frog species can clear *B. dendrobatidis* infection: 50% of experimentally infected *Mixophyes fasciolatus* held at 27 °C with confirmed infection subsequently cleared the infection (Berger 2001; Berger et al. 2004). The prevalences we report are therefore estimated from the prevalence of infection in frogs captured for the first time in the period under consideration.

In this study, infection levels of *B. dendrobatidis* were significantly higher during winter and spring than during summer and autumn. A survey of ill and dead frogs from eastern Australia from 1995 to 1999 showed a similar seasonal prevalence of chytridiomycosis (Berger et al. 2004). In the laboratory, the growth of *B. dendrobatidis* has been shown to peak at about 23 °C, with death or arrested growth occurring in vitro at temperatures above 30 °C (Longcore et al. 1999; Johnson et al. 2003; Piotrowski et al. 2004). Further, infection in some frog species can be cleared in the laboratory by exposing them to temperatures in excess of 37 °C (Woodhams et al. 2003). At Eungella, 23 °C is a typical daytime maximum temperature for winter, with water and air temperatures along streams generally remaining at or below 23 °C for the entire 24-h daylength period. In summer, however, daytime air temperatures along Eungella's streams regularly exceed 23 °C, and at sites such as Rawson Creek can reach as high as 37 °C (R. Retallick, unpublished data).

The different infection levels among sites for *T. eungellensis* reported in Table 2 may be correlated with the degree of sunlight and warmth that reaches the streams. The streams at Dooloomai Falls and Tree Fern Creek are well shaded and remain relatively cool and damp for much of the year. At those sites, the average canopy gap above the centre of the streams (a coarse measure of how much sunlight reaches the stream) are 0.35 m and 1.90 m respectively (R. Retallick, unpublished data). Rawson Creek is considerably wider (average canopy gap = 5.10 m) and thus receives more sunlight. With access to sunny microhabitats, where surface temperatures can exceed ambient air temperatures, frogs at "warmer" streams such as Rawson Creek may be less subject to infection at any time of the year, or may be able to reduce infection through thermoregulation. The idea that frogs may avoid, control, or eliminate infection by differential use of their environment warrants considerable and immediate attention. Such a process may prove to be critical to the relationship between *B. dendrobatidis* and frog populations in the wild.

Our mark-recapture analysis did not find consistent evidence that *T. eungellensis* infected with *B. dendrobatidis* at the time of first capture had a lower survival rate than uninfected *T. eungellensis*. Failure to reject the null hypothesis of no effect of infection on survival cannot, of course, be used as evidence in favor of the null hypothesis, and it is worth noting that the point estimate of survival for infected males was lower than that for uninfected males at each site. Only at Tree Fern Creek, however, did the best-supported model include an infection term. There was also evidence that infection influenced the proportion of frogs that were recaptured (Table 3). Together with the observation that some infected *T. eungellensis* survived for extended periods (a maximum of 1,089 d), our results show that infection with *B. dendrobatidis* did not inevitably lead to rapid death in *T. eungellensis*.

Both epidemiological theory and observations suggest that where a pathogen drives a host species to extinction, there is likely to be a reservoir host within which the pathogen has a reduced effect and is therefore maintained at a higher prevalence (McCallum and Dobson 1995, 2002). *L. wilcoxii/jungguy* appears not to have declined at the same time as *T. eungellensis* and *R. vitellinus* and, in our data, it has a high prevalence of infection. It therefore is a candidate reservoir host. Whether the prevalence of *B. dendrobatidis* differs between the sibling species *L. wilcoxii* and *L. jungguy* at sites where they are sympatric would be an interesting question, but cannot be answered from our study. The *L. lesueurii* complex has a widespread distribution throughout streams on the east coast of Australia (Barker et al. 1995; Donnellan and Mahony 2004) and could therefore play a substantial role in the maintenance and spread of chytrid infection throughout that region. If a species is acting as a reservoir, the prevalence of infection in that species should be higher than in species that are declining as a result of infection (McCallum and Dobson 1995). The prevalence of infection we observed in *L. wilcoxii/jungguy* exceeded that observed in *T. eungellensis*, but we do not have clear evidence that this represents a higher prevalence in the populations as a whole. It is also possible that *B. dendrobatidis* exists as a saprophyte in the environment independent of amphibians (Longcore et al. 1999; Johnson and Speare 2003), or may use tadpoles (which seem relatively little affected by the pathogen) as a reservoir (Daszak et al. 1999).

We did not detect infection in any of the 50 individuals of *Litoria revelata* or the 42 individuals of *L. chloris* we examined, which demonstrates that those species had a lower prevalence of chytridiomycosis than *T. eungellensis* and *L. wilcoxii/jungguy*. We did not record any infected frogs of any species from the two sites at which *L. revelata* was collected. *Litoria chloris*, however, has been shown to carry infection in other areas (Speare and Berger 2004) and in laboratory experiments (Woodhams et al. 2003). Sample sizes in our study for these other frog species may be too small to draw any reliable conclusions about whether they become infected with *B. dendrobatidis* in the wild.

## Materials and Methods

Between 1994 and 1998, 36 excursions to Eungella National Park were made as part of a mark-recapture study of populations of frogs associated with streams. Frogs were captured from six study sites (see Figure 1), which were visited monthly from March 1994 to June 1996, in February and September 1997, and in February and May 1998. When caught for the first time, frogs were toe-clipped using the numbering system devised by Hero (1989) and then released alive. Amputated toe tips were preserved and individually stored in vials filled with 70% ethanol or 10% formalin. To minimize effects on the animals, no further toes were taken from subsequent recaptures of marked frogs; recaptured frogs were identified and released alive.

All frog species encountered were monitored, with a special focus on *T. eungellensis* because of its precarious conservation status and history of decline.

In 1997, toes from six species (278 individual frogs) were histologically prepared in transverse sections stained with Ehrlich's haematoxylin for skeletochronological assessment (see Castanet et al. [1996] for a description of the technique). We subsequently reexamined these sections for *B. dendrobatidis* (Berger et al. 1999b; Pessier et al. 1999). On average, 160 sections per individual were examined. Despite the sections not being stained with eosin, which highlights the keratinised layer of the epidermis where *B. dendrobatidis* occurs (Berger et al. 1999b; Pessier et al. 1999), most diagnoses were unambiguous. A small number, however, were not, and those samples

were excluded from the analysis. For this assessment, samples were labelled “positive” only when the examiners were convinced that the sample was infected with *B. dendrobatidis*. Samples in which no infection was found were labelled “negative.”

For this study, a further 196 archived toes of *T. eungellensis* were analysed by the Australian Animal Health Laboratory in Geelong, Australia, for *B. dendrobatidis* infection using histopathology with immunoperoxidase staining (Berger et al. 2002). The immunoperoxidase stain improves the sensitivity of diagnosis by highlighting structures that are equivocal with haematoxylin and eosin, and that would otherwise be diagnosed as negative.

With the software MARK (Cooch and White 2002), we used mark-recapture modelling to investigate the survival of *T. eungellensis* at different sites. Frogs were grouped into six categories on the basis of their status at first capture (infected and uninfected males, females, and subadults). For both mark-recapture and analysis of prevalence, we grouped the data into 3-mo austral seasons (in Australia, summer runs from 1 December to the end of February, etc.). Mark-recapture modelling of survival requires that several assumptions be satisfied. Most critically, every marked individual present in the population at a given time has the same probability of recapture as all other members of its group, and the same probability of surviving to the next time interval. We used a parametric bootstrap (Cooch and White 2002) to test the goodness of fit of our capture history sets to these assumptions. We analysed prevalence data using logistic models implemented in R (Version 1.6.2) (R Development Core Team 2003). Since the mark-recapture study was performed with no knowledge of the infection status of individual animals, and the examination of histology was carried out with no knowledge of details of the fate of

individual animals in the field, the study was double-blinded in design.

## Acknowledgments

We are grateful to Diana Mendez, Alex Hyatt, and the diagnostic staff at the Australian Animal Health Laboratory for diagnosis of chytrid infection, and the School of Public Health and Tropical Medicine, James Cook University, for work space and technical support. Funding for the original fieldwork was supplied by a grant from Australian Nature Conservation Agency (now Environment Australia) to J-M Hero, RA Alford, and KR McDonald, and an Australian Post-Graduate Research Award to R Retallick. This study was completed under multiple Queensland Parks and Wildlife Service permits, and ethics approval was obtained from the Experimentation Ethics Review Committee (Biological Sciences, James Cook University). Financial support for the retrospective analysis was provided by the Australian Research Council and the US National Science Foundation Integrated Research Challenges in Environmental Biology scheme. Some of the analysis was carried out by H McCallum while he was a visiting scientist at the Centre for Population Biology, Silwood Park.

**Conflicts of interest.** The authors have declared that no conflicts of interest exist.

**Author contributions.** RWRR and RS conceived and designed the experiments. RWRR performed the experiments. HIM analyzed the data. HIM wrote the paper. ■

## References

- Barker J, Grigg G, Tyler M (1995) A field guide to Australian frogs. Chipping Norton: Surrey Beatty and Sons. 407 p.
- Berger L (2001) Diseases in Australian frogs [dissertation]. Townsville, Queensland: James Cook University. 331 p. Available from: James Cook University library (<http://www.jcu.edu>).
- Berger L, Speare R, Hyatt A (1999a) Chytrid fungus and amphibian declines: Overview, implications and future directions. In: Campbell A, editor. Declines and disappearances of Australian frogs. Canberra: Environment Australia. pp. 23–33.
- Berger L, Speare R, Kent A (1999b) Diagnosis of chytridiomycosis in amphibians by histologic examination. *Zoos' Print J* 15: 184–190.
- Berger L, Speare R, Daszak P, Green DE, Cunningham AA, et al. (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proc Natl Acad Sci U S A* 95: 9031–9036.
- Berger L, Hyatt AD, Olsen V, Hengstberger SG, Boyle D, et al. (2002) Production of polyclonal antibodies to *Batrachochytrium dendrobatidis* and their use in an immunoperoxidase test for chytridiomycosis in amphibians. *Dis Aquat Organ* 48: 213–220.
- Berger L, Speare R, Hines HB, Marantelli G, Hyatt AD, et al. (2004). Effect of season and temperature on mortality in amphibians due to chytridiomycosis. *Aust Vet J*. 82: 434–439
- Campbell A (1999) Declines and disappearances of Australian frogs. Canberra: Environment Australia. 234 p.
- Castanet J, Francillon-Vieillot H, Bruce RC (1996) Age estimation in desmognathine salamanders assessed by skeletochronology. *Herpetologica* 52: 160–171.
- Cooch E, White G (2002) Using MARK—A gentle introduction. <http://www.phidot.org/software/mark/docs/book/>. Accessed 25 August 2004.
- Couper PJ (1992) Hope for our missing frogs. *Wildl Aust* 29: 11–12.
- Daszak P, Berger L, Cunningham AA, Hyatt AD, Green DE, et al. (1999) Emerging infectious diseases and amphibian population declines. *Emerg Infect Dis* 5: 735–748.
- Daszak P, Cunningham AA, Hyatt AD (2003) Infectious disease and amphibian population declines. *Divers Distrib* 9: 141–150.
- Department of the Environment and Heritage (2003) National list of threatened fauna Available: <<http://www.deh.gov.au/cgi-bin/sprat/public/publicthreatenedlist.pl?wanted=fauna#ENDANGERED%20FROGS>>. Accessed 21 August 2004.
- Donnellan SC, Mahony MJ (2004) Allozyme, chromosomal and morphological variability in the *Litoria lesueuri* species group (Anura: Hylidae) including a description of a new species. *Aust J Zool* 52: 1–28.
- Fenner F, Fantini B (1999) Biological control of vertebrate pests: The history of myxomatosis—An experiment in evolution. Wallingford: CABI Publishing. 352 p.
- Hero JM (1989) A simple code for toe clipping anurans. *Herpetol Rev* 20: 66–67.
- Johnson ML, Speare R (2003) Survival of *Batrachochytrium dendrobatidis* in water: Quarantine and disease control implications. *Emerg Infect Dis* 9: 922–925.
- Johnson M, Berger L, Philips L, Speare R (2003) Fungicidal effects of chemical disinfectants, UV light, desiccation and heat on the amphibian chytrid, *Batrachochytrium dendrobatidis*. *Dis Aquat Organ* 57: 255–260.
- Laurance WF, McDonald KR, Speare R (1996) Epidemic disease and the catastrophic decline of Australian rain forest frogs. *Conserv Biol* 10: 1–9.
- Laurance WF, McDonald KR, Speare R (1997) In defense of the epidemic disease hypothesis. *Conserv Biol* 11: 1030–1034.
- Longcore JE, Pessier AP, Nichols DK (1999) *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid pathogenic to amphibians. *Mycologia* 91: 219–227.
- May RM, Anderson RM (1983) Epidemiology and genetics in the coevolution of parasites and hosts. *Proc R Soc Lond B Biol Sci* 219: 281–313.
- McCallum HI, Dobson AP (1995) Detecting disease and parasite threats to endangered species and ecosystems. *Trends Ecol Evol* 10: 190–194.
- McCallum H, Dobson A (2002) Disease, habitat fragmentation and conservation. *Proc R Soc Lond B Biol Sci* 269: 2041–2049.
- McDonald K, Alford R (1999) A review of declining frogs in Northern Queensland. In: Campbell A, editor. Declines and disappearances of Australian frogs. Canberra: Environment Australia. pp. 14–22.
- McDonald KR (1990) *Rheobatrachus* Liem and *Taudactylus* Straughan and Lee (Anura: Leptodactylidae) in Eungella National Park, Queensland: Distribution and decline. *Trans R Soc South Aust* 114: 187–194.
- McNellie M, Hero J-M (1994) Mission Amphibian: The search for the missing rainforest frogs of Eungella. *Wildl Aust* 31: 22–23.
- Morehouse EA, James TY, Ganley ARD, Vilgalys R, Berger L, et al. (2003) Multilocus sequence typing suggests that the chytrid pathogen of amphibians is a recently emerged clone. *Mol Ecol* 12: 395–403.
- Pessier AP, Nichols DK, Longcore JE, Fuller MS (1999) Cutaneous chytridiomycosis in poison dart frogs (*Dendrobates* spp.) and White's tree frogs (*Litoria caerulea*). *J Vet Diagn Invest* 11: 194–199.
- Piotrowski JS, Annis SL, Longcore JE (2004) Physiology of *Batrachochytrium dendrobatidis*, a chytrid pathogen of amphibians. *Mycologia* 96: 9–15.
- R Development Core Team (2004) R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available: <http://www.R-project.org>. Accessed 22 August 2004.
- Retallick R, Hero JM, Alford R (1997) Adult population monitoring and larval ecology of the stream-dwelling frogs at Eungella National Park: ANCA/QDOE. 63 p.
- Speare R, Berger L (2004) Chytridiomycosis in amphibians in Australia. Available: <http://www.jcu.edu.au/school/phtm/PHTM/frogs/chyspec.htm>. Accessed 26 Feb 2004.
- Woodhams DC, Alford RA, Marantelli G (2003) Emerging disease of amphibians cured by elevated body temperature. *Dis Aquat Organ* 55: 65–67.