

Article

Synthesis and Investigation of Flavanone Derivatives as Potential New Anti-Inflammatory Agents

Cynthia Sinyeue ^{1,2} , Mariko Matsui ^{1,3} , Michael Oelgemöller ^{4,5} , Frédérique Bregier ² , Vincent Chaleix ² , Vincent Sol ²  and Nicolas Lebouvier ^{1,*} 

- ¹ Institut des Sciences Exactes et Appliquées (ISEA) EA7484, Campus de Nouville, Université de la Nouvelle Calédonie, Noumea 98851, New Caledonia; cynthia.sinyeue@etudiant.unc.nc (C.S.); mmatsui@pasteur.nc (M.M.)
- ² Laboratoire PEIREINE UR 22722, Faculté des Sciences et Techniques, 87060 Limoges, France; frederique.bregier@unilim.fr (F.B.); vincent.chaleix@unilim.fr (V.C.); vincent.sol@unilim.fr (V.S.)
- ³ Group Immunity and Inflammation (GIMIN), Institut Pasteur of New Caledonia, Pasteur Network, Noumea 98845, New Caledonia
- ⁴ College of Science and Engineering, Discipline of Chemistry, James Cook University, Townsville, QLD 4811, Australia; michael.oelgemoller@hs-fresenius.de
- ⁵ Faculty of Chemistry and Biology, Hochschule Fresenius gGmbH—University of Applied Science, 65510 Idstein, Germany
- * Correspondence: nicolas.lebouvier@unc.nc; Tel.: +687-290-251

Abstract: Flavonoids are polyphenols with broad known pharmacological properties. A series of 2,3-dihydroflavanone derivatives were thus synthesized and investigated for their anti-inflammatory activities. The target flavanones were prepared through cyclization of 2'-hydroxychalcone derivatives, the later obtained by Claisen–Schmidt condensation. Since nitric oxide (NO) represents an important inflammatory mediator, the effects of various flavanones on the NO production in the LPS-induced RAW 264.7 macrophage were assessed in vitro using the Griess test. The most active compounds were flavanone (**4G**), 2'-carboxy-5,7-dimethoxy-flavanone (**4F**), 4'-bromo-5,7-dimethoxy-flavanone (**4D**), and 2'-carboxyflavanone (**4J**), with IC₅₀ values of 0.603, 0.906, 1.030, and 1.830 µg/mL, respectively. In comparison, pinocembrin achieved an IC₅₀ value of 203.60 µg/mL. Thus, the derivatives synthesized in this work had a higher NO inhibition capacity compared to pinocembrin, demonstrating the importance of pharmacomodulation to improve the biological potential of natural molecules. SARs suggested that the use of a carboxyl-group in the *meta*-position of the B-ring increases biological activity, whereas compounds carrying halogen substituents in the *para*-position were less active. The addition of methoxy-groups in the *meta*-position of the A-ring somewhat decreased the activity. This study successfully identified new bioactive flavanones as promising candidates for the development of new anti-inflammatory agents.

Keywords: flavanone; anti-inflammatory activity; structure–activity relationship (SAR); RAW264.7; pinocembrin



Citation: Sinyeue, C.; Matsui, M.; Oelgemöller, M.; Bregier, F.; Chaleix, V.; Sol, V.; Lebouvier, N. Synthesis and Investigation of Flavanone Derivatives as Potential New Anti-Inflammatory Agents. *Molecules* **2022**, *27*, 1781. <https://doi.org/10.3390/molecules27061781>

Academic Editors: Heiko Lange and Margherita Brindisi

Received: 31 January 2022

Accepted: 2 March 2022

Published: 9 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Inflammation is involved in many diseases, such as infectious diseases, chronic inflammation, asthma, diabetes, neurodegenerative diseases, coronaviruses, and cancer [1,2]. Anti-inflammatory treatments are mostly based on corticoids or non-steroidal anti-inflammatory drugs (NSAIDs), such as aspirin (2-acetyloxybenzoic acid) and ibuprofen (2-[4-(2-methylpropyl)phenyl]propanoic acid). However, long-term therapy involving these common pharmaceuticals leads to severe side effects such as gastrointestinal ulceration and bleeding, osteoporosis, hypertension, and glaucoma. Consequently, there is a need for new anti-inflammatory compounds with fewer or no side effect [3]. Macrophages are essential in the inflammatory process, and some bacterial endotoxins, such as lipopolysaccharide (LPS), allow their activation [4]. Nitric oxide (NO) is an inflammatory mediator

that influences various biological processes [5]. At high levels, NO can exhibit cytotoxicity and tissue damage. Therefore, inhibition of this mediator can provide therapeutic effects and allow measurement of the degree of inflammation. Natural products such as the sesquiterpene yomogin isolated from *Artemisia princeps* or *Ginkgo biloba* extract are known to inhibit NO production [6,7].

Flavonoids are widely present in plants. A nutrition study revealed that diets rich in fruits and vegetables can help to prevent inflammatory diseases. These beneficial properties have been linked to synergetic effects of bioactive compounds, including flavonoids [8,9]. The latter compounds act as regulators of metabolic processes and have thus been investigated for the treatment of diseases. Consequently, flavonoids have been widely studied for their beneficial effects on human health, including antiallergenic, anti-inflammatory, vasodilating, anti-COVID-19, and antitumor capacities [1,2,10,11]. Flavonoids can furthermore act on important mechanisms in inflammatory processes by controlling regulatory enzymes and transcription factors [12]. Flavanones are a subgroup of flavonoids characterized by two aromatic rings (A and B) linked by a dihydropyrone ring C (Figure 1).




Figure 1. General structure of flavanone.

Pinocembrin (5,7-dihydroxyflavanone) is a natural flavanone present in fruits, spices, propolis, tea, and red wine and has shown beneficial properties, including anti-inflammatory potential [13,14]. A recent *in vivo* study proved the ability of pinocembrin to inhibit the LPS-stimulated inflammatory response in macrophages and to regulate the TLR4-NF- κ B signaling pathway [15]. Furthermore, pinocembrin is also found in pine species that are exploited for their wood [16]. The easy availability of biomass and low cost of co-products from these species may thus provide a renewable access to pinocembrin and its derivatives.

In order to obtain pinocembrin analogues, a two-step synthesis strategy was followed in this study. The chalcones obtained through initial Claisen–Schmidt condensation were subjected to subsequent cyclization to their corresponding flavanones [17,18]. The flavanone derivatives obtained were subsequently examined as regulators for the NO production on the LPS-induced murine macrophage. The aim of this work was to investigate the structural requirements on the A- and B-rings of flavanones for anti-inflammatory activity. Simple structure–activity relationship (SAR) analysis consequently determined the most effective candidates for future optimization studies.

2. Results and Discussion

2.1. Synthesis

A series of 2'-hydroxy chalcones (3A–3L) were synthesized by Claisen–Schmidt condensation from appropriately substituted 2'-hydroxyacetophenones and benzaldehydes. Various experimental conditions, such as acid catalysis [19], base catalysis [20], heat treatment [21], and light activation [22,23], have been reported to achieve the subsequent cyclization to flavanones. Two cyclization procedures were chosen in this work: basic and photochemical activation (Figure 2). The photochemical route was investigated as a milder approach [24]; however, the efficiency of photocyclization is known to depend on the irradiation wavelength [25]. Commercial pinocembrin (PC) was selected as a reference molecule.




Figure 2. Synthesis of chalcone (3A–3L) and flavanone (4A–4L) derivatives. Reaction conditions: (a) excess of NaOH in 25 mL of MeOH; (b) basic activation: 5 eq NaOAc in MeOH (20 mL) under reflux; (c) photochemical method: in EtOH at 419 nm, 30 °C for 7 days.

Following these synthetic approaches, a series of flavanones was prepared with or without methoxy-groups attached to the A-ring, while the B-ring carried various substituents in different positions (Table 1). The isolated yields of chalcones 3 ranged from 23–92%, while those of flavanones 4 varied widely between 7 and 74%, indicating a need for further optimization. After prolonged irradiation for 7 days, the photochemical cyclization gave a low isolated yield of just 7% for 4A, despite a near-complete conversion of 3A of 98%, suggesting significant losses during repeated purification by column chromatography. All other derivatives 4B–F were obtained by base-catalyzed cyclization instead. For the flavanones containing two methoxy-groups on the A-ring (A to F), the highest yield of flavanones overall was obtained with the carboxyl-group (4F), followed by the methoxy substituent (4B). For the series without substitution on the A-ring (G to K), the highest flavanone yield was achieved with the carboxyl-group (4J), followed by the parent flavanone 4G. Thus, when the corresponding benzaldehyde was substituted with an electron-withdrawing group at the 2-position (as in 4F and 4J), the formation of flavanones was effective. In contrast, the addition of electron-donating groups on the A-ring, as for derivatives 4A, G, and L, gave the desired flavanones in only low yields.

Table 1. Synthesis of chalcones 3A–3L and flavanone derivatives 4A–4L.

| Compound | R | R' | Yield (%) | |
|----------|---------------------------------------|-----------------------------|-------------|--------------|
| | | | Chalcones 3 | Flavanones 4 |
| A | 5, 7-(OCH ₃) ₂ | H | 27 | 7* |
| B | 5, 7-(OCH ₃) ₂ | 3'-OCH ₃ | 82 | 74 |
| C | 5, 7-(OCH ₃) ₂ | 4'-OCH ₃ | 92 | 66 |
| D | 5, 7-(OCH ₃) ₂ | 4'-Br | 52 | 69 |
| E | 5, 7-(OCH ₃) ₂ | 4'-Cl | 23 | 66 |
| F | 5, 7-(OCH ₃) ₂ | 2'-COOH | 36 | 72 |
| G | H | H | 62 | 59 |
| H | H | 4'-Cl | 86 | 37 |
| I | H | 4'-OCH ₃ | 56 | 37 |
| J | H | 2'-COOH | 32 | 81 |
| K | H | 2'-OCH ₃ , 5'-Br | 84 | 56 |
| L | 5-OCH ₃ | H | 30 | 9 |

Note: * 98% conversion after irradiation with 419 nm light for 7 days.

Both cyclization methods were found to have limitations. The photochemical method did not require the addition of base and proceeded cleanly, but it demanded an exhaustive irradiation time to achieve a high conversion. The basic activation was found to operate faster but led to by-products that required subsequent purification steps.

2.2. Evaluation of Biological Activities

2.2.1. Biological Activities of Commercial Pinocembrin

Pinocembrin (PC) inhibitory activity was previously investigated on nitrite produced by LPS-induced RAW264.7 [26], and an IC₅₀ of 203.60 µg/mL was determined. The anti-inflammatory bioactivity of commercially available PC was thus studied (Figure 3A),

and the results showed a significant inhibitory response at 200 $\mu\text{g}/\text{mL}$ ($79.36 \pm 7.30\%$) on LPS-dependent NO production, hence confirming the anti-inflammatory effect of PC in this concentration range. No inhibitory effect of PC was observed at 2 or 20 $\mu\text{g}/\text{mL}$ (Figure 3A). The cytotoxicity was also analyzed (Figure 3B), and the results revealed a significant increase in percentage of cytotoxicity for PC at 200 $\mu\text{g}/\text{mL}$ ($94.23 \pm 3.72\%$) compared to LPS-related cytotoxicity ($32.28 \pm 10.01\%$), while no significant difference in cytotoxicity was observed at 2 ($25.76 \pm 14.41\%$) and 20 $\mu\text{g}/\text{mL}$ ($34.45 \pm 7.25\%$). A cytotoxic concentration of 200 $\mu\text{g}/\text{mL}$ could possibly affect the determination of a reliable inhibitory response on NO production, as lower quantities of viable cells could result in inherently lower NO levels. These results thus highlight the importance of evaluating the cytotoxicity of natural compounds and their derivatives.




Figure 3. (A) Anti-inflammatory activity. Nitrite representative of NO production was quantified using Griess reagent, and percent of the inhibitory response was calculated compared to the level of LPS-dependent nitrite production; (B) cytotoxicity of pinocembrin (PC) on LPS-induced RAW264.7. Murine macrophages RAW264.7 (150,000 cells/well in P96) were treated with LPS (*E. coli* 0111:B4) at 1 $\mu\text{g}/\text{mL}$ and pinocembrin at 2, 20, and 200 $\mu\text{g}/\text{mL}$ for 24 h. Cytotoxicity was measured by quantification of LDH production. Results are means \pm SD (at least $n = 3$ well replicates). Mann-Whitney test was used to compare LPS and molecule treatments. ns, nonsignificant; **, $p < 0.01$.

2.2.2. Cytotoxicity of Flavanone Derivatives

Due to the cytotoxicity of PC, the cytotoxicity to LPS-induced RAW264.7 at 2 $\mu\text{g}/\text{mL}$ was evaluated for all flavanone derivatives (Figure 4). Initial cytotoxicity of untreated cells was calculated at $32.41 \pm 3.4\%$, which is consistent with previous observations [27]. No difference with LPS-related cytotoxicity was determined at $33.26 \pm 10.61\%$, and no significant increase was observed for any flavanone at 2 $\mu\text{g}/\text{mL}$, with percentages of cytotoxicity ranging between 20 and 40%. Thus, the potential inhibitory effect of these compounds was investigated at this concentration and compared to PC.

2.2.3. Inhibitory Activity on LPS-Induced NO Production

The anti-inflammatory responses of all synthesized flavanone compounds at 2 $\mu\text{g}/\text{mL}$ were examined by evaluating their inhibitory effect on NO induction produced by murine macrophage RAW264.7 treated with bacterial LPS (Figure 5 and Table 2). The inhibitory response of the reference anti-inflammatory molecule dexamethasone at 100 nM was confirmed ($100.4 \pm 16.14\%$). As noted above, PC did not show an effective response at 2 $\mu\text{g}/\text{mL}$. In contrast, various inhibitory activities were determined for flavanones 4A–L at the same concentration.




Figure 4. Cytotoxicity of all derivatives on LPS-induced RAW264.7. Murine macrophages RAW264.7 (150,000 cells/well in P96) were treated with LPS (*E. coli* 0111:B4) at 1 $\mu\text{g}/\text{mL}$, analogues and pinocembrin (PC) at 2 $\mu\text{g}/\text{mL}$ (in DMSO 0.1%), or anti-inflammatory dexamethasone (DEX) at 100 nM for 24 h. Cytotoxicity was measured by quantification of LDH production. Results are means \pm SD of 4 independent experiments (with $n = 3$ well replicates/experiment). Mann–Whitney test was used to compare LPS and molecule treatments. ns, nonsignificant.




Figure 5. Inhibitory effect of analogues on LPS-induced NO production by RAW264.7. Murine macrophages RAW264.7 (150,000 cells/well in P96) were treated with LPS (*E. coli* 0111:B4) at 1 $\mu\text{g}/\text{mL}$, analogues and pinocembrin (PC) at 2 $\mu\text{g}/\text{mL}$ (in DMSO 0.1%), or anti-inflammatory dexamethasone (DEX) at 100 nM for 24 h. Nitrite representative of NO production was quantified using Griess reagent, and percent of the inhibitory response was calculated compared to the level of LPS-dependent nitrite production. Bars represent mean \pm SD of 4 independent experiments (with $n = 3$ well replicates/experiment). Mann–Whitney test was used to compare LPS and molecule treatments. ns, nonsignificant; *, $p < 0.05$; ***, $p < 0.0005$; ****, $p < 0.0001$.

Table 2. Inhibitory effect of analogues 4A–L on LPS-induced NO production by RAW264.7.

| Compound | Inhibitory Response (%) ^a | p Value ^b |
|------------------|--------------------------------------|-------------------------|
| Pinocembrin (PC) | −21.81 ± 37.42 | 0.1732 ^{ns} |
| Dexamethasone | 100.4 ± 16.14 | <0.0001 ^{****} |
| 4A | 58.99 ± 45.19 | 0.0009 ^{***} |
| 4B | 40.45 ± 49.71 | 0.0272 [*] |
| 4C | 19.14 ± 33.83 | 0.1470 ^{ns} |
| 4D | 61.10 ± 25.50 | <0.0001 ^{****} |
| 4E | 26.85 ± 24.43 | 0.0318 [*] |
| 4F | 78.65 ± 24.73 | <0.0001 ^{****} |
| 4G | 75.65 ± 46.88 | <0.0001 ^{****} |
| 4H | 3.128 ± 34.64 | 0.5605 ^{ns} |
| 4I | −0.87 ± 37.54 | 0.8116 ^{ns} |
| 4J | 72.56 ± 22.70 | <0.0001 ^{****} |
| 4K | 73.29 ± 86.10 | 0.0105 [*] |
| 4L | 64.97 ± 42.37 | 0.0003 ^{***} |

Note: ^a Percentage of inhibitory response of compounds used at 2 µg/mL compared to the level of LPS-dependent nitrite production in RAW264.7 cells expressed in mean ± SD; ^b statistical analyses were conducted. Mann–Whitney test was used to compare LPS and molecule treatments. ^{ns}, nonsignificant; ^{*}, $p < 0.05$; ^{***}, $p < 0.0005$; ^{****}, $p < 0.0001$.

The most active compounds were 4F (78.65 ± 24.73%), 4G (75.65 ± 46.88%), 4J (72.56 ± 22.70%) and 4L (64.97 ± 42.37%), respectively, followed by 4D (61.10 ± 25.50%), and 4A (58.99 ± 45.19%). Due to their large SD variation, the following compounds were considered less effective but still showed a significant inhibitory response: 4K (73.29 ± 86.10%), 4B (40.45 ± 49.71%), and 4E (26.85 ± 24.43%). 4C (19.14 ± 33.83%), 4H (3.12 ± 34.64%), and 4I (−0.87 ± 37.54%) were not significantly bioactive (Table 2).

The IC₅₀ values for the most effective molecules 4F, 4G, 4J, and 4D were subsequently calculated (Table 3). Dexamethasone presented an IC₅₀ of 0.005 µg/mL (95% CI: 0.003–0.008). The most active flavanones were 4G, with the lowest IC₅₀ of 0.603 µg/mL (95% CI: 0.366–1.003) and 4F with an IC₅₀ of 0.906 µg/mL (95% CI: 0.550–1.765), while 4D and 4J furnished lower inhibitory activities with higher IC₅₀ values of 1.030 µg/mL (95% CI: 0.675–1.382) and 1.830 µg/mL (95% CI: 1.467–2.677), respectively.

Table 3. Inhibitory response of PC and selected derivatives on NO produced by LPS-induced RAW264.7.

| Compound | IC ₅₀ (µg/mL) ^a | 95% CI ^b |
|-------------------------------|---------------------------------------|----------------------------|
| Pinocembrin (PC) ^c | 203.60 ^c | 101.30–569.31 ^c |
| Dexamethasone | 0.005 | 0.003–0.008 |
| 4D | 1.030 | 0.675–1.382 |
| 4F | 0.906 | 0.550–1.765 |
| 4G | 0.603 | 0.366–1.003 |
| 4J | 1.830 | 1.467–2.677 |

Note: ^a Concentrations of molecules leading to 50% of inhibitory activity calculated using a logarithm-transformed (inhibitor) vs. normalized response equation; ^b 95% of Confidence Interval (CI); ^c [26].

2.3. Structure–Activity Relationship Study

The biological evaluation clearly indicated that the percentage of inhibition depended on the substitution pattern of flavanones 4A–L; Figure 6 summarizes the pharmacomodulation obtained from this study.




Figure 6. Correlations between structures and anti-inflammatory activity.

For the first series with two methoxy-groups at the A-ring, the most active molecules were **4F** (2'-carboxy-5,7-dimethoxy-flavanone) and **4D** (4'-bromo-5,7-dimethoxy-flavanone). Hence, the presence of a carboxyl-group in the *ortho*-position or a bromo substituent in the *para*-position of the B-ring increased the anti-inflammatory activity. In addition, **4D** (4'-bromo-5,7-dimethoxy-flavanone) was more active than the related **4E** (4'-chloro-5,7-dimethoxy-flavanone). For the second series without substitution on the A-ring, the most active compounds were **4G** (flavanone), **4J** (2'-carboxyflavanone), and **4K** (5'-bromo-2'-methoxy-flavanone). Overall, the presence of methoxy-groups on the flavanone skeleton impacted the NO inhibitory activity. Indeed, flavanones with a methoxy-group in the 5-position demonstrated strong inhibitory activity on NO production from LPS-stimulated macrophage cells [28]. For the functionalization on the A-ring, the comparison of **4G** (flavanone), **4L** (5-methoxyflavanone), and **4A** (5,7-dimethoxyflavanone) showed that the addition of a methoxy-group contributes to a drop in activity. Additionally, the flavanone derivative with the methoxy-group in the 3-position (**4B**) was found more active than the corresponding analogue with the methoxy-substituent in the 4-position (**4C**). Pinocembrin (5,7-dihydroxyflavanone) containing two hydroxyl-groups on the A-ring showed no significant effect at the same concentration if compared to the flavanone derivatives synthesized. These results are in agreement with Kim's work, which demonstrated that naringenin (5,7,4'-trihydroxyflavanone) was inactive up to 100 μM [29]. It may thus be suggested that the presence of a hydroxy-group on the A-ring only weakly affects the NO inhibition.

This SARs clearly revealed that a 2'-carboxy-group is beneficial to achieve effective inhibition of NO production. This finding may be due to the electron-withdrawing and highly polar nature of the carboxyl-group. According to Shin's study, a bulky and/or hydrophobic substituent in the *meta*-position of the B-ring results in the most active structure to inhibit NF- κ B activation [30]. This study identified **4B** (3'-methoxyflavanone) as more active than **4C** (4'-methoxyflavanone) and revealed the importance of a carboxyl-group in the *ortho*-position of the B-ring instead, as in **4F** and **4J**.

The predictive analysis of the drug-like absorption of the most active compounds (Table 4) revealed that all compounds meet Lipinski's rule of five [31]: hydrogen bond donors ≤ 5 , hydrogen bond acceptors ≤ 10 , molecular mass ≤ 500 daltons, an octanol-water partition coefficient ($\log p$) ≤ 5 , and a polar surface area $\leq 140 \text{ \AA}^2$. Subsequent future pharmacomodulations, however, may lead to compounds with further improved activity.

Table 4. Predictive analysis of drug-like absorption.

| Compound | IC ₅₀ ($\mu\text{g/mL}$) | M _w (Da) | Log <i>p</i> | Hydrogen Bond Acceptors | Hydrogen Bond Donors | Polar Surface Area (\AA^2) |
|-----------|--|---------------------|-----------------|-------------------------------|----------------------------|--|
| 4D | 1.030 | 363.21 | 4.32 \pm 0.41 | 4 | 0 | 44.77 |
| 4F | 0.906 | 328.32 | 3.23 \pm 0.38 | 6 | 1 | 82.07 |
| 4G | 0.603 | 224.26 | 3.62 \pm 0.26 | 2 | 0 | 26.30 |
| 4J | 1.830 | 268.27 | 3.29 \pm 0.27 | 4 | 1 | 63.60 |

3. Materials and Methods

3.1. Syntheses

3.1.1. General Information

All chemicals were purchased from Sigma–Aldrich (Saint-Louis, MO, USA) and were used as received. HPLC grade solvents were bought from Fischer Chemicals (Leicestershire, UK) and were used without further purification. Reactions were monitored by thin-layer chromatography (TLC) using silica gel-precoated aluminum sheets (60 F254, Merck KGaA, Darmstadt, Germany) with different solvent systems (cyclohexane/EtOAc or DCM/MeOH). Products were visualized with UV irradiation at 365 and 254 nm or by treatment with sulfuric acid-vanillin. Selected compounds were purified by column chromatography (CC) using silica gel 60 (0.015–0.040 mm, Merck) as a stationary phase or by LH-20 Sephadex column chromatography. Electrothermal and Gallenkamp melting point apparatuses (MP, OC, uncorrected) were used to determine the melting points in open capillaries. Nuclear magnetic resonance (NMR) spectra were recorded on a Bruker x400 (^1H nMR: 400 MHz, ^{13}C nMR: 100 MHz) spectrometer or Bruker DPX 500 nMR spectrometer (^1H nMR: 500 MHz, ^{13}C nMR: 125 MHz). The solvents used were CDCl_3 or dimethyl sulfoxide ($\text{DMSO}-d_6$). Chemical shifts (δ) were recorded in parts per million (ppm) relative to residual solvent peaks (CDCl_3 : ^1H nMR: $\delta = 7.26$ ppm and ^{13}C nMR: $\delta = 77.00$ ppm; $\text{DMSO}-d_6$: ^1H nMR: $\delta = 2.50$ ppm and ^{13}C nMR: $\delta = 39.52$ ppm). Coupling constants were reported in hertz (Hz). Multiplicities were reported as s (singlet), br s (broad singlet), d (doublet), t (triplet), br (broad), m (multiplet), dd (doublet of doublets), dt (doublet of triplets), ddt (doublet of doublet of triplet), ddd (doublet of doublets of doublets), app d (apparent doublet), and app t (apparent triplet). High resolution electrospray ionization mass spectrometry (HR ESI-MS) was performed at the ICOA/CBM platform (Orléans University) on a Bruker Q-TOF maXis mass spectrometer, coupled to an Ultimate 3000 RSLC chain (Dionex).

3.1.2. Synthetic Methods

For characterization of the molecular structures, the numbering of atoms shown in Figure 7 was followed.




Figure 7. Structures of chalcone and flavanone derivatives.

- General Procedure for the synthesis of chalcones (3A–L).

A solution of MeOH (25 mL), the corresponding acetophenones 1A–C (1 eq), the appropriate benzaldehyde 2A–G (1 eq), and excess of NaOH were stirred at room temperature, and the progress of the reaction was monitored by TLC. Upon completion, the excess of NaOH was neutralized by addition of HCl (1 M) with pH control. The solvent was evaporated, the residue was taken up in EtOAc, and the resulting solution was washed with distilled water. The organic layer was dried with MgSO_4 and filtered, and the solvent was evaporated under reduced pressure. The crude product was purified by crystallization from methanol (MeOH) or by column chromatography.

(E)-1-(2'-hydroxy-4,6-dimethoxyphenyl)-3-phenylprop-2-en-1-one or 2'-Hydroxy-4',6'-dimethoxy-chalcone or flavokavain B (3A)

Prepared following the general procedure starting from 2-hydroxy-4,6-dimethoxyacetophenone (**1A**, 0.1458 g, 0.74 mmol) and benzaldehyde (**2A**, 0.078 g, 0.74 mmol) with 10 eq of NaOH. Yellow crystals (0.057 g, 0.20 mmol), 27% yield, m.p. (Gallenkamp apparatus): 89 °C (85–86 °C, [32]).

¹H-NMR (400 MHz, CDCl₃): δ 14.20 (s, 1H, 2'-OH), 7.83 (d, *J* = 15.7 Hz, 1H, β-H), 7.72 (d, *J* = 15.7 Hz, 1H, α-H), 7.54 (dd, *J* = 7.5 Hz, 2.1 Hz, 2H, 2-H, 6-H), 7.30 (m, 3H, 3-H, 4-H, 5-H), 6.05 (d, *J* = 2.4 Hz, 1H, 3'-H), 5.90 (d, *J* = 2.4 Hz, 1H, 5'-H), 3.86 (s, 3H, 4'-OCH₃), 3.77 (s, 3H, 6'-OCH₃). ¹³C-NMR (100 MHz, CDCl₃): δ 166.5 (4'-C), 162.6 (2'-C), 142.3 (β-C), 130 (1-C), 128.89 (3-C, 6-C), 128.4 (5-C), 128.3 (2-C), 127.6 (4-C), 102.7 (1'-C), 91.4 (3'-C), 91.1 (5'-C), 55.7 (4'-OCH₃), 55.5 (6'-OCH₃). ¹H and ¹³C spectral data were consistent with the literature [32].

(E)-1-(2'-hydroxy-4',6'-dimethoxyphenyl)-3-(3-methoxyphenyl)prop-2-en-1-one or 2'-hydroxy-3,4',6'-trimethoxy-chalcone (**3B**)

Prepared following the general procedure starting from 2-hydroxy-4,6-dimethoxyacetophenone (**1A**, 0.50 g, 2.55 mmol) and 3-methoxy-benzaldehyde (**2B**, 0.346 g, 2.55 mmol) with 10 eq of NaOH. Yellow orange crystals (0.655 g, 2.09 mmol), 82% yield, m.p. (Electrothermal apparatus): 100 °C (100 °C, [33]).

¹H-NMR (500 MHz, CDCl₃): δ 14.25 (s, 1H, 2'-OH), 7.87 (d, *J* = 15.7 Hz, 1H, β-H), 7.73 (d, *J* = 15.7 Hz, 1H, α-H), 7.32 (t, *J* = 7.9 Hz, 1H, 5-H), 7.21 (d, *J* = 7.6 Hz, 1H, 6-H), 7.12 (d, *J* = 1.2 Hz, 1H, 2-H), 6.93 (dd, *J* = 8.2 Hz, 1H, 4-H), 6.11 (d, *J* = 2.4 Hz, 1H, 3'-H), 5.96 (d, *J* = 2.4 Hz, 1H, 5'-H), 3.91 (s, 3H, 4'-OCH₃), 3.85 (s, 3H, 6'-OCH₃), 3.83 (s, 3H, 3-OCH₃). ¹³C-NMR (125 MHz, CDCl₃): δ 192.6 (CO), 168.4 (4'-C), 166.3 (6'-C), 162.5 (2'-C), 159.9 (3-C), 142.2 (β-C), 137.0 (1-C), 129.8 (5-C), 127.9 (6-C), 120.9 (α-C), 115.6 (4-C), 113.7 (2-C), 106.4 (1'-C), 93.8 (3'-C), 91.3 (5'-C), 55.8 (3-OCH₃), 55.6 (5-OCH₃), 55.3 (7-OCH₃). ¹H and ¹³C spectral data were consistent with the literature [33,34].

(E)-1-(2'-hydroxy-4',6'-dimethoxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one or 2'-hydroxy-4,4',6'-trimethoxy-chalcone (**3C**) has been previously synthesized by our research group [33].

(E)-3-(4-bromophenyl)-1-(2'-hydroxy-4',6'-dimethoxyphenyl)prop-2-en-1-one or 2'-hydroxy-4-bromo-4',6'-dimethoxy-chalcone (**3D**)

Prepared following the general procedure starting from 2-hydroxy-4,6-dimethoxyacetophenone (**1A**, 0.216 g, 1.10 mmol) and 4-bromo-benzaldehyde (**2D**, 0.204 g, 1.10 mmol) with 2.5 eq of NaOH. Yellow crystals (0.206 g, 0.57 mmol), 52% yield, m.p. (Gallenkamp apparatus): 166 °C (165 °C, [33], 150–151 °C, [32]).

¹H nMR (400 MHz, CDCl₃): δ 14.23 (s, 1H, 2'-OH), 7.90 (d, *J* = 15.6 Hz, 1H, β-H), 7.72 (d, *J* = 15.6 Hz, 1H, α-H), 7.58–7.53 (m, 2H, 3-H, 5-H), 7.51–7.46 (m, 2H, 2-H, 6-H), 6.14 (d, *J* = 2.4 Hz, 1H, 3'-H), 5.99 (d, *J* = 2.4 Hz, 1H, 5'-H), 3.94 (s, 3H, 4'-OCH₃), 3.87 (s, 3H, 6'-OCH₃). ¹³C nMR (100 MHz, CDCl₃): δ 192.5 (CO), 166.3 (4'-C), 162.9 (6'-C), 162.6 (2'-C), 140.9 (β-C), 132.13 (5-C, 3-C) 131.6 (1-C), 129.69 (2-C, 6-C), 128.2 (4-C), 124.2 (α-C), 106.3 (1'-C), 93.8 (3'-C), 91.4 (5'-C), 55.9 (4'-OCH₃), 55.6 (6'-OCH₃). ¹H and ¹³C spectra were consistent with the literature [32].

(E)-3-(4-chlorophenyl)-1-(2'-hydroxy-4',6'-dimethoxyphenyl)prop-2-en-1-one or 4-chloro-2'-hydroxy-4',6'-dimethoxy-chalcone (**3E**)

Prepared following the general procedure starting from 2-hydroxy-4,6-dimethoxyacetophenone (**1A**, 0.50 g, 2.55 mmol) and 4-chloro-benzaldehyde (**2E**, 0.358 g, 2.55 mmol) with 5 eq of NaOH. Yellow crystals (0.189 g, 0.60 mmol), 23% yield, m.p. (Gallenkamp apparatus): 164 °C (173–175 °C, [32], 2006 and 158–168 °C, [33]).

¹H nMR (400 MHz, CDCl₃): δ 14.24 (s, 1H, 2'-OH), 7.88 (d, *J* = 15.6 Hz, 1H, β-H), 7.74 (d, *J* = 15.6 Hz, 1H, α-H), 7.55 (d, *J* = 8.6 Hz, 2H, 2-H, 6-H), 7.44–7.36 (m, 2H, 3-H, 5-H), 6.14 (d, *J* = 2.4 Hz, 1H, 3'-H), 5.99 (d, *J* = 2.4 Hz, 1H, 5'-H), 3.94 (s, 3H, 4'-OCH₃), 3.87 (s, 3H, 6'-OCH₃). ¹H spectral data were consistent with the literature [32].

(E)-2-(3-(2'-hydroxy-4',6'-dimethoxyphenyl)-3-oxoprop-1-en-1-yl)benzoic acid or 2'-hydroxy-2-carboxy 4',6'-dimethoxychalcone (**3F**)

Prepared following the general procedure starting from 2-hydroxy-4,6-dimethoxyacetophenone (**1A**, 0.369 g, 1.89 mmol) and 2-carboxy-benzaldehyde (**2F**, 0.283 g, 1.89 mmol) with 10 eq of NaOH. Yellow solid (0.220 g, 0.67 mmol), 36% yield, m.p. (Gallenkamp apparatus): 161 °C (160–161 °C, [32]).

¹H nMR (400 MHz, CDCl₃): δ 14.04 (s, 1H, 2'-OH), 13.74 (s, 1H, 2-COOH), 8.57 (d, *J* = 15.4 Hz, 1H, β-H), 8.10 (dd, *J* = 7.1 Hz, 1.0 Hz, 1H, 3-H), 7.91 (d, *J* = 15.4 Hz, 1H, α-H), 7.78 (m, 1H, 5-H), 7.66 (m, 1H, 6-H), 7.56 (td, *J* = 8.1 Hz, 1.5 Hz, 1H, 4-H), 6.15 (d, *J* = 2.4 Hz, 1H, 3'-H), 6.12 (d, *J* = 2.4 Hz, 1H, 5'-H), 4.00 (s, 3H, 4'-OCH₃), 3.90 (s, 3H, 6'-OCH₃). ¹H spectral data were consistent with the literature [32]. HRMS (ESI): Calc. for C₁₈H₁₆O₆, [M + H]⁺ *m/z*: 329.10196, found: 329.10204, [M + Na]⁺ *m/z*: 345.07334, found: 345.07320.

(E)-1-(2'-hydroxyphenyl)-3-phenylprop-2-en-1-one or 2'-hydroxy-chalcone (**3G**)

Prepared following the general procedure starting from 2-hydroxyacetophenone (**1B**, 1 g, 7.34 mmol) and benzaldehyde (**2A**, 0.78 g, 7.34 mmol) with 10 eq of NaOH. Yellow solid (1.013 g, 4.52 mmol), 62% yield, m.p. (Gallenkamp apparatus): 77 °C (44–150 °C CAS: 1214-47-7 Aldrich).

¹H nMR (400 MHz, CDCl₃): δ 12.82 (s, 1H, 2'-OH), 7.96 (d, *J* = 15.6 Hz, 1H, β-H), 7.97–7.94 (dd, *J* = 8.1 Hz, 1.6 Hz, 1H, 6'-H), 7.72–7.67 (m, 3H, 2-H, 3-H, 5-H), 7.69 (d, *J* = 15.6 Hz, 1H, α-H), 7.53 (td, *J* = 8.6 Hz, 1.6 Hz, 1H, 5'-H), 7.49–7.45 (m, 4H, 2'-H, 3'-H, 4'-H, 4-H), 7.06 (dd, *J* = 8.5 Hz, 0.9 Hz, 1H, 6-H), 6.98 (ddd, *J* = 8.3 Hz, 7.4 Hz, 1.0 Hz, 1H, 4'-H). ¹³C nMR (100 MHz, CDCl₃): δ 193.7 (CO), 163.6 (2'-C), 145.5 (β-C), 136.1 (1-C), 130.9 (4'-C), 129.6 (5'-C), 129.08 (3-C), 129.07 (5-C), 128.68 (2-C), 128.67 (6-C), 127.7 (4-C), 120.2 (α-C), 118.9 (3'-C), 118.7 (1'-C). HRMS (ESI): Calc. for C₁₅H₁₂O₂, [M + H]⁺ *m/z*: 225.09101, found: 225.09112, [M + Na]⁺ *m/z*: 247.07295, found: 247.07318.

(E)-3-(4-chlorophenyl)-1-(2'-hydroxyphenyl) prop-2-en-1-one or 4-chloro-2'-hydroxy-chalcone (**3H**)

Prepared following the general procedure starting from 2-hydroxyacetophenone (**1B**, 2 g, 14.69 mmol) and 4-chloro-benzaldehyde (**2E**, 2.06 g, 14.69 mmol) with 5 eq of NaOH. Yellow solid (3.265 g, 12.62 mmol), 86% yield, m.p. (Gallenkamp apparatus): 135–140 °C (149–150 °C, [35]).

¹H nMR (400 MHz, CDCl₃): δ 12.76 (s, 1H, 2'-OH), 7.93 (dd, *J* = 8.1 Hz, 1.6 Hz, 1H, 6'-H), 7.89 (d, *J* = 15.6 Hz, 1H, β-H), 7.65 (d, *J* = 15.6 Hz, 1H, α-H), 7.65–7.60 (m, 2H, 3-H, 5-H), 7.54 (ddd, *J* = 8.8 Hz, 7.3 Hz, 1.6 Hz, 1H, 5'-H), 7.47–7.41 (m, 2H, 3'-H, 6'-H), 7.06 (dd, *J* = 8.4 Hz, 0.9 Hz, 1H, 6-H), 6.98 (ddd, *J* = 8.5 Hz, 7.2 Hz, 1.1 Hz, 1H, 4'-H). ¹³C nMR (100 MHz, CDCl₃): δ 193.5 (CO), 163.7 (2'-C), 143.9 (β-C), 136.6 (4-C), 133.1 (1-C), 129.79 (3-C), 129.78 (5-C), 129.6 (4'-C), 129.38 (2-C), 129.37 (6-C), 120.7 (1'-C), 119.9 (α-C), 118.9 (3'-C), 118.7 (5'-C) [34]. HRMS (ESI): Calc. for C₁₅H₁₂ClO₂, [M + H]⁺ *m/z*: 259.05203, found: 259.05249.

(E)-1-(2'-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one or 2'-hydroxy-4-methoxy-chalcone (**3I**)

Prepared following the general procedure starting from 2-hydroxyacetophenone (**1B**, 2 g, 14.69 mmol) and 4-methoxy-benzaldehyde (**2C**, 2 g, 14.69 mmol) with 5 eq of NaOH. Orange solid (2.088 g, 8.21 mmol), 56% yield, m.p. (Gallenkamp apparatus): 84 °C (84–86 °C, [35]).

¹H nMR (400 MHz, CDCl₃): δ 12.95 (s, 1H, 2'-OH), 7.95 (dd, *J* = 8.1 Hz, 1.6 Hz, 1H, 6'-H), 7.93 (d, *J* = 15.4 Hz, 1H, β-H), 7.68–7.63 (m, 2H, 3-H, 5-H), 7.57 (d, *J* = 15.4 Hz, 1H, α-H), 7.51 (ddd, *J* = 8.6 Hz, 7.3 Hz, 1.6 Hz, 1H, 5'-H), 7.05 (dd, *J* = 8.4 Hz, 0.9 Hz, 1H, 6-H), 7.00–6.96 (m, 2H, 2-H, 3'-H), 6.99–6.93 (m, 1H, 4'-H), 3.89 (s, 3H, 4-OCH₃). ¹³C nMR (100 MHz, CDCl₃): δ 193.7 (CO), 163.6 (2'-C), 162.1 (4-C), 145.4 (β-C), 136.1 (6'-C), 130.55 (1-C), 130.54 (3-C), 130.53 (5-C), 129.55 (4'-C), 129.54 (2-C), 127.4 (6-C), 120.2 (1'-C), 118.8 (α-C), 118.6 (5'-C),

117.6 (3'-C), 55.5 (4-OCH₃). HRMS (ESI): Calc. for C₁₆H₁₄O₃, [M + H]⁺ *m/z*: 255.10157, found: 255.10208, [M + Na]⁺ *m/z*: 277.08352 found: 277.08381.

(E)-2-(3-(2'-hydroxyphenyl)-3-oxoprop-1-en-1-yl)benzoic acid or 2'-hydroxy-2-carboxy-chalcone (**3J**)

Prepared following the general procedure starting from 2-hydroxy-acetophenone (**1B**, 2.00 g, 14.69 mmol) and 2-carboxy-benzaldehyde (**2F**, 2.20 g, 14.69 mmol) with 5 eq of NaOH. Yellow solid (1.276 g, 4.76 mmol), 32% yield, m.p. (Gallenkamp apparatus): 144 °C.

¹H nMR (400 MHz, CDCl₃): δ 12.03 (s, 1H, 2'-OH), 7.96 (dd, *J* = 7.8 Hz, 2.3 Hz, 1H, 6'-H), 7.71 (ddd, *J* = 8.4 Hz, 7.4 Hz, 0.9 Hz, 1H, 5-H), 7.67 (dd, *J* = 8.1 Hz, 1.5 Hz, 1H, 6-H), 7.62–7.57 (m, 2H, 3-H, 4-H), 7.54 (ddd, *J* = 8.6 Hz, 7.3 Hz, 1.5 Hz, 1H, 5'-H), 7.05 (dd, *J* = 8.5 Hz, 0.7 Hz, 1H, 3'-H), 6.93 (ddd, *J* = 8.1 Hz, 7.1 Hz, 0.9 Hz, 1H, 4'-H), 6.19 (t, *J* = 6.4 Hz, 1H), 3.81 (dd, *J* = 17.6 Hz, 6.2 Hz, 1H, β-H), 3.46 (dd, *J* = 17.6 Hz, 6.8 Hz, 1H, α-H). ¹³C nMR (100 MHz, CDCl₃): δ 201.7 (CO), 169.9 (2-C, COOH), 162.6 (2'-C), 149.3 (β-C), 137.2 (1-C), 129.84 (3-C), 129.83 (4'-C), 129.82 (5'-C), 129.63 (4-C), 129.62 (5-C), 125.93 (6'-C), 125.92 (6-C), 122.6 (2-C), 119.3 (1'-C), 119.1 (α-C), 118.8 (3'-C). HRMS (ESI): Calc. for C₁₆H₁₂O₄, [M + H]⁺ *m/z*: 269.08084, found: 269.08139, [M + Na]⁺ *m/z*: 291.06278, found: 291.06338.

(E)-3-(5-bromo-2-methoxyphenyl)-1-(2'-hydroxyphenyl) prop-2-en-1-one or 5-bromo-2'-hydroxy-2-methoxy-chalcone (**3K**)

Prepared following the general procedure starting from 2-hydroxy-acetophenone (**1B**, 1.00 g, 7.34 mmol) and 5-bromo-2-methoxy-benzaldehyde (**2G**, 1.58 g, 7.34 mmol) with 10 eq of NaOH. Yellow solid (2.056 g, 6.17 mmol), 84% yield, m.p. (Gallenkamp apparatus): 121 °C.

¹H nMR (400 MHz, CDCl₃): δ 12.84 (s, 1H, 2'-OH), 8.16 (d, *J* = 15.7 Hz, 1H, β-H), 7.95 (dd, *J* = 8.1 Hz, 1.5 Hz, 1H, 6'-H), 7.78 (d, *J* = 2.4 Hz, 1H, 6-H), 7.74 (d, *J* = 15.7 Hz, 1H, α-H), 7.56–7.48 (m, 2H, 5'-H, 3'-H), 7.06 (dd, *J* = 8.4 Hz, 0.9 Hz, 1H, 4-H), 6.98 (td, *J* = 8.2 Hz, 7.2 Hz, 1.2 Hz, 1H, 4'-H), 6.87 (d, *J* = 8.9 Hz, 1H, 3-H), 3.94 (s, 3H, 2-OCH₃). ¹³C nMR (100 MHz, CDCl₃): δ 193.9 (CO), 163.6 (2'-C), 157.9 (2-C), 139.2 (β-C), 136.4 (1-C), 134.4 (4-C), 131.4 (6-C), 129.75 (4'-C), 129.74 (5'-C), 125.7 (6'-C), 121.7 (α-C), 120.1 (1'-C), 118.9 (5-C), 118.6 (3'-C), 113.1 (3-C), 55.9 (2-OCH₃). HRMS (ESI): Calc. for C₁₆H₁₃BrO₃, [M + H]⁺ *m/z*: 333.01208, found: 333.01247, [M + Na]⁺ *m/z*: 354.99403, found: 354.99467.

(E)-1-(2'-hydroxy-6-methoxyphenyl)-3-phenylprop-2-en-1-one or 2'-hydroxy-6'-methoxy-chalcone (**3L**)

Prepared following the general procedure starting from 2-hydroxy-6-methoxy-acetophenone (**1C**, 0.5 g, 3.0 mmol) and benzaldehyde (**2A**, 0.32 g, 3.0 mmol) with 10 eq of NaOH. Yellow crystals (0.657 g, 2.58 mmol), 30% yield, m.p. (Electrothermal apparatus): 100 °C.

¹H nMR (500 MHz, CDCl₃): δ 13.13 (s, 1H, 2'-OH), 7.86 (d, *J* = 15.6 Hz, 1H, β-H), 7.80 (d, *J* = 15.6 Hz, 1H, α-H), 7.62–7.58 (m, 2H, 3-H, 5-H), 7.43–7.36 (m, 3H, 2-H, 4-H, 6-H), 7.34 (t, *J* = 8.1 Hz, 1H, 4'-H), 6.60 (d, *J* = 8.2 Hz, 1H, 5'-H), 6.41 (d, *J* = 8.3 Hz, 1H, 3'-H), 3.93 (s, 3H, 6'-OCH₃). ¹³C nMR (125 MHz, CDCl₃): δ 194.5 (4-C, CO), 164.9 (2'-C), 161.0 (6'-C), 142.9 (β-C), 136.0 (4'-C), 135.3 (1-C), 130.3 (4'-C), 128.45 (3-C, 5-C), 127.6 (1'-C), 126.8 (2-C, 6-C), 112.1 (3'-C), 112.0 (5'-C), 101.6 (α-C), 79.4 (β-H), 55.9 (6'-OCH₃).

- Flavanone **3A** synthesis by photochemical activation.

Chalcone **3A** (85 mg, 1 eq) was dissolved in 70 mL of ethanol in a Pyrex vessel. The solution was purged with purified nitrogen gas for 10 min before the reaction vessel was sealed. Irradiation was conducted in an RPR-200 photochemical reactor equipped with 16 fluorescent tubes (8 W each, 419 ± 25 nM) at approximately 30 °C [36]. The progress of the reaction was followed by TLC analysis, and after 7 days, almost complete conversion to the desired cyclized product was observed. The solution was evaporated to dryness, and the residue was purified on a chromatographic column.

5,7-dimethoxy-2-phenyl-2,3-dihydro-4H-benzopyran-4-one or 5,7-dimethoxy-2-phenylchroman-4-one or 5,7-dimethoxy-flavanone (**4A**)

This product was purified by repeated column chromatography (C₆H₁₂/EtOAc 90:10, 70:30 and 50:50). White solid (0.006 g, 0.02 mmol), 7% yield, m.p. (Gallenkamp apparatus): 144–146 °C.

¹H nMR (400 MHz, CDCl₃): δ 7.41–7.26 (m, 5H, 2'-H, 3'-H, 4'-H, 5'-H, 6'-H), 6.10 (d, *J* = 2.3 Hz, 1H, 6-H), 6.03 (d, *J* = 2.3 Hz, 1H, 8-H), 5.35 (dd, *J* = 13.1 Hz, 3.0 Hz, 1H, 2-H), 3.83 (s, 3H, 5-OCH₃), 3.76 (s, 3H, 7-OCH₃), 2.96 (dd, *J* = 16.6 Hz, 13.1 Hz, 1H, 3ax-H), 2.74 (dd, *J* = 16.6 Hz, 3.1 Hz, 1H, 3eq-H). ¹³C nMR (100 MHz, CDCl₃): δ 192.7 (4-C, CO), 166.26 (7-C), 165.9 (5-C), 162.5 (8a-C), 135.6 (1'-C), 128.45 (3'-C, 5'-C), 127.7 (4'-C), 126.85 (2'-C, 6'-C), 106.4 (4a-C), 93.8 (8-C), 93.7 (6-C), 79.5 (2-C), 56.8 (7-OCH₃), 55.9 (4-OCH₃), 38.9 (3-C). Calc. for C₁₇H₁₆O₄, [M + H]⁺ *m/z*: 285.11214 Da, found 285.11258 Da, [M + Na]⁺ *m/z*: 307.09408 found : 307.09413.

- General Procedure for flavanones synthesis by base activation (**4B–L**).

To a solution of chalcones **3B–L** (1 eq) in methanol (20 mL) was added 5 eq of sodium acetate. The mixture was heated to reflux for 24 h and was monitored by TLC. The solvent was evaporated, EtOAc (20 mL) was added, and the mixture was washed with distilled water (3 × 20 mL). The solution was dried over anhydrous MgSO₄ and filtered. The solvent was evaporated under reduced pressure, and the residue was purified by column chromatography (CC) using silica gel 60, 0.2–0.5 mm C35-70mesh ASTH CAS: 7631-86-9 (Scharlau Spain) as stationary phase to give flavanones **4B–L**.

5,7-dimethoxy-2-(3-methoxyphenyl)-2,3-dihydro-4H-benzopyran-4-one or 5,7,3'-trime-thoxy-flavanone (**4B**)

Prepared following the general procedure starting from chalcone **3B** (0.603 g, 1.92 mmol). This product was purified by column chromatography (C₆H₁₂/EtOAc 90:10, 70:30, 50:50). Pale yellow solid (0.445 g, 1.41 mmol), 74% yield, m.p. (Electrothermal apparatus): 89–91 °C.

¹H nMR (500 MHz, CDCl₃): δ 7.32 (t, *J* = 7.9 Hz, 1H, 5'-H), 7.02 (m, 2H, 2'-H, 6'-H), 6.90 (dd, *J* = 8,3 Hz, 2.1 Hz, 1H, 4'-H), 6.16 (d, *J* = 2.3 Hz, 1H, 6-H), 6.09 (d, *J* = 2.3 Hz, 1H, 8-H), 5.38 (dd, *J* = 13.3 Hz, 2.8 Hz, 1H, 2-H), 3.89 (s, 3H, 7-OCH₃), 3.83 (s, 3H, 5-OCH₃), 3.82 (s, 3H, 3'-OCH₃), 3.00 (dd, *J* = 16.5 Hz, 13.3 Hz, 1H, 3eq-H), 2.79 (dd, *J* = 16.50 Hz, 2.80 Hz, 1H, 3ax-H), ¹³C nMR (125 MHz, CDCl₃): δ 189.2 (4-C, CO), 166.0 (7-C), 164.9 (5-C), 162.3 (8a-C), 159.9 (3'-C), 129.89 (1'-C), 129.88 (5'-C), 118.3 (6'-C), 114.0 (4'-C), 111.8 (4a-C), 106.1 (2'-C), 93.6 (6-C), 93.2 (8-C), 79.1 (2-C), 56.2 (7-OCH₃), 55.6 (5-OCH₃), 55.3 (3'-OCH₃), 45.6 (3-C). HRMS (ESI): Calc. for C₁₈H₁₈O₅, [M + H]⁺ *m/z*: 315.12270 Da, found 315.12292, [M + Na]⁺ *m/z*: 337.10464, found: 337.10455.

5,7-dimethoxy-2-(4-methoxyphenyl)-2,3-dihydro-4H-benzopyran-4-one or 5,7,4'-trime-thoxy-flavanone (**4C**)

Prepared following the general procedure starting from chalcone **3C** (0.32 g, 1.08 mmol). This product was purified by column chromatography (C₆H₁₂/EtOAc 70:30). Pale yellow solid (0.213 g, 0.68 mmol), 66% yield, m.p. (Electrothermal apparatus): 60–70 °C.

¹H-NMR (400MHz, CDCl₃): δ 7.44–7.34 (m, 2H, 2'-H, 6'-H), 7.01–6.91 (m, 2H, 3'-H, 5'-H), 6.16 (d, *J* = 2.3 Hz, 1H, 6-H), 6.11 (d, *J* = 2.3 Hz, 1H, 8-H), 5.38 (dd, *J* = 13.1 Hz, 2.9 Hz, 1H, 2-H), 3.92 (s, 3H, 7-OCH₃), 3.85 (s, 3H, 5-OCH₃), 3.84 (s, 3H, 4'-OCH₃), 3.06 (dd, *J* = 16.5 Hz, 13.1 Hz, 1H, 3eq-H), 2.79 (dd, *J* = 16.5 Hz, 2.9 Hz, 1H, 3ax-H). ¹³C nMR (100 MHz, CDCl₃): δ 189.5 (4-C, CO), 165.9 (7-C), 165.1 (5-C), 162.3 (8a-C), 159.9 (4'-C), 130.8 (1'-C), 127.71 (2'-C), 127.70 (6'-C), 114.17 (3'-C), 114.16 (5'-C), 106.0 (4a-C), 93.6 (6-C), 93.1 (8-C), 79.0 (2-C), 56.2 (7-OCH₃), 55.6 (5-OCH₃), 55.4 (4'-OCH₃), 45.4 (3-C). HRMS (ESI): Calc. for C₁₈H₁₈O₅, [M + H]⁺ *m/z*: 315.12270 Da, found: 315.12282, [M + Na]⁺ *m/z*: 337.10464, found: 337.10436.

2-(4-bromophenyl)-5,7-dimethoxy-)-2,3-dihydro-4H-benzopyran-4-one or 4'-bromo-5,7-dimethoxy-flavanone (**4D**)

Prepared following the general procedure starting from chalcone **3D** (0.1033 g, 0.28 mmol). This product was purified by column chromatography (C₆H₁₂/EtOAc 70:30). Pale yellow solid (0.0719 g, 0.20 mmol), 69% yield, m.p. (Electrothermal apparatus): 60–70 °C.

^1H nMR (500 MHz, CDCl_3): δ 7.62–7.54 (m, 2H, 2'-H, 6'-H), 7.38–7.33 (d, $J = 8.3$ Hz, 2H, 3'-H, 5'-H), 6.17 (d, $J = 2.3$ Hz, 1H, 6-H), 6.13 (d, $J = 2.3$ Hz, 1H, 8-H), 5.40 (dd, $J = 12.8$ Hz, 3.2 Hz, 1H, 2-H), 3.92 (s, 3H, 5-OCH₃), 3.86 (s, 3H, 7-OCH₃), 2.99 (dd, $J = 16.5$ Hz, 12.8 Hz, 1H, 3ax-H), 2.81 (dd, $J = 16.5$ Hz, 3.2 Hz, 1H, 3eq-H). ^{13}C nMR (125 MHz, CDCl_3): δ 192.3 (4-C, CO), 168.5 (7-C), 166.4 (5-C), 162.5 (8a-C), 134.5 (1'-C), 132.11 (3'-C), 132.10 (5'-C), 129.69 (2'-C), 129.68 (6'-C), 124.2 (4'-C), 106.3 (4a-C), 93.8 (8-C), 91.3 (6-C), 77.3 (2-C), 55.9 (7-OCH₃), 55.6 (5-OCH₃). Calc. for $\text{C}_{17}\text{H}_{15}\text{BrO}_4$, $[\text{M} + \text{H}]^+$ m/z : 363.02265 Da, found 363.02329, $[\text{M} + \text{Na}]^+$ m/z : 385.00459 found: 385.00514.

2-(4-chlorophenyl)-5,7-dimethoxy-)-2,3-dihydro-4H-benzopyran-4-one or 4'-chloro-5,7-dimethoxy-flavanone (**4E**)

Prepared following the general procedure starting from chalcone **3E** (0.127 g, 0.40 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 90:10, 70:30, 50:50). Colorless solid (0.084g, 0.26 mmol), 66% yield, m.p. (Gallenkamp apparatus): 120 °C.

^1H nMR (400 MHz, DMSO): δ 7.57–7.53 (m, 2H, 2'-H, 6'-H), 7.52–7.47 (m, 2H, 3'-H, 5'-H), 6.25 (d, $J = 2.3$ Hz, 1H, 6-H), 6.22 (d, $J = 2.3$ Hz, 1H, 8-H), 5.57 (dd, $J = 12.5$ Hz, 3.0 Hz, 1H, 2-H), 3.82 (s, 3H, 7-OCH₃), 3.79 (s, 3H, 5-OCH₃), 3.02 (dd, $J = 16.3$ Hz, 12.5 Hz, 1H, 3eq-H), 2.67 (dd, $J = 16.3$ Hz, 3.0 Hz, 1H, 3ax-H). ^{13}C nMR (100 MHz, CDCl_3): δ 187.9 (4-C, CO), 165.9 (7-C), 164.5 (5-C), 162.3 (8a-C), 138.5 (1'-C), 133.4 (4'-C), 128.99 (3'-C), 128.98 (5'-C), 128.86 (2'-C), 128.85 (6'-C), 105.9 (4a-C), 94.2 (6-C), 93.5 (8-C), 77.9 (2-C), 56.2 (7-OCH₃), 56.4 (5-OCH₃), 45.1 (3-C). HRMS (ESI): Calc. for $\text{C}_{17}\text{H}_{15}\text{ClO}_4$, $[\text{M} + \text{H}]^+$ m/z : 319.07316 Da, found 319.07330, $[\text{M} + \text{Na}]^+$ m/z : 341.05511, found: 341.05529.

2-(5,7-dimethoxy-4-oxo-3,4-dihydro-2H-benzopyran-2-yl)benzoic acid or 2'-carboxy-5,7-dimethoxy-flavanone (**4F**)

Prepared following the general procedure starting from chalcone **3F** (0.220g, 0.67 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 70:30). White solid (0.159 g, 0.48 mmol), 72% yield, m.p. (Gallenkamp apparatus): 168 °C.

^1H nMR (400 MHz, CDCl_3): δ 13.75 (s, 1H, 2'-OH), 7.93 (d, $J = 7.7$ Hz, 1H, 3'-H), 7.70–7.65 (m, 1H, 4'-H), 7.63–7.53 (m, 2H, 5'-H, 6'-H), 6.18 (dd, $J = 7.9$ Hz, 5.4 Hz, 1H, 2-H), 6.13 (d, $J = 2.3$ Hz, 1H, 6-H), 5.94 (d, $J = 2.3$ Hz, 1H, 8-H), 3.89 (dd, $J = 18.5$ Hz, 5.3 Hz, 1H, 3ax-H), 3.43 (dd, $J = 18.5$ Hz, 7.9 Hz, 1H, 3eq-H). ^{13}C nMR (125 MHz, CDCl_3): δ 200.2 (4-C, CO), 170.5 (2'-COOH), 167.8 (7-C), 166.7 (5-C), 162.7 (8a-C), 150.3 (2'-C), 134.2 (4'-C), 129.2 (3'-C), 126.0 (1'-C), 125.6 (6'-C), 123.1 (5'-C), 105.7 (4a-C), 93.8 (6-C), 91.1 (8-C), 77.48 (2-C), 55.68 (7-OCH₃), 55.67 (5-OCH₃), 48.94 (3-C). Calc. for $\text{C}_{18}\text{H}_{16}\text{O}_6$, $[\text{M} + \text{H}]^+$ m/z : 329.10197 Da, found: 329.10205, $[\text{M} + \text{Na}]^+$ m/z : 351.08391, found: 351.08389.

2,3-dihydro-2-phenyl-4H-1-benzopyran-4-one or flavanone (**4G**)

Prepared following the general procedure starting from chalcone **3G** (0.5g, 2.23 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 90:10, 70:30, 50:50). Colorless solid (0.294 g, 1.31 mmol), 59% yield, m.p. (Gallenkamp apparatus): 70 °C (75–76 °C, [37]).

^1H nMR (400 MHz, CDCl_3): δ 7.97 (dd, $J = 8.1$ Hz, 1.7 Hz, 1H, 5-H), 7.58–7.37 (m, 6H, 2'-H, 3'-H, 4'-H, 5'-H, 6'-H, 6-H), 7.11–7.05 (m, 2H, 7-H, 8-H), 5.52 (dd, $J = 13.3$ Hz, 2.9 Hz, 1H, 2-H), 3.13 (dd, $J = 16.9$ Hz, 13.3 Hz, 1H, 3ax-H), 2.93 (dd, $J = 16.9$ Hz, 2.9 Hz, 1H, 3eq-H). ^{13}C nMR (125 MHz, CDCl_3): δ 191.9 (4-C, CO), 161.6 (8a-C), 138.8 (1'-C), 136.2 (7-C), 128.86 (3'-C, 5'-C), 128.8 (4'-C), 127.1 (5-C), 126.16 (2'-C, 6'-C), 121.6 (6-C), 120.9 (4a-C), 118.1 (8-C), 79.6 (2-C), 44.7 (3-C) [37].

2-(4-chlorophenyl)-2,3-dihydro-4H-benzopyran-4-one or 4'-chloro-flavanone (**4H**)

Prepared following the general procedure starting from chalcone **3H** (1.5 g, 5.80 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 70:30). Colorless solid (0.556 g, 2.15 mmol), 37% yield, m.p. (Gallenkamp apparatus): 80–90 °C (93–95 °C, [37]).

^1H nMR (400 MHz, CDCl_3): δ 7.96 (ddd, $J = 7.8$ Hz, 1.6 Hz, 0.4 Hz, 1H, 5-H), 7.55 (ddd, $J = 8.7$ Hz, 7.2 Hz, 1.8 Hz, 1H, 2'-H), 7.48–7.41 (m, 4H, 3'-H, 5'-H, 6'-H, 6-H), 7.12–7.04 (m, 2H, 7-H, 8-H), 5.50 (dd, $J = 13.1$ Hz, 2.9 Hz, 1H, 2-H), 3.07 (dd, $J = 16.8$ Hz, 13.1 Hz, 1H, 3ax-H), 2.91 (dd, $J = 16.8$ Hz, 2.9 Hz, 1H, 3eq-H). ^{13}C nMR (100 MHz, CDCl_3): δ 191.5 (4-C, CO), 161.6 (8a-C), 137.3 (1'-C), 136.3 (4'-C), 133.86 (6-C), 133.85 (7-C), 129.1 (3'-C, 5'-C), 127.52 (2'-C), 127.51 (6'-C), 127.1 (5-C), 121.8 (4a-C), 118.1 (8-C), 78.8 (2-C), 44.6 (3-C) ([37]).

2,3-dihydro-2-(4-methoxyphenyl)-4H-1-benzopyran-4-one or 4'-methoxy-flavanone (**4I**)

Prepared following the general procedure starting from chalcone **3I** (1 g, 3.93 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 90:10, 70:30, 50:50). Pale yellow solid (0.374 g, 1.47 mmol), 37% yield, m.p. (Gallenkamp apparatus): 85 °C (87–88 °C, [37]).

^1H nMR (400 MHz, CDCl_3): δ 7.95 (dd, $J = 8.0$ Hz, 1.5 Hz, 1H, 5-H), 7.55–7.50 (m, 1H, 2'-H), 7.46–7.41 (m, 2H, 3'-H, 5'-H), 7.01–6.95 (m, 2H, 6'-H, 6-H), 7.10–7.03 (m, 2H, 7-H, 8-H), 5.46 (dd, $J = 13.3$ Hz, 2.9 Hz, 1H, 2-H), 3.86 (s, 3H, 4'-OCH₃), 3.13 (dd, $J = 16.8$ Hz, 13.3 Hz, 1H, 3ax-H), 2.89 (dd, $J = 16.7$ Hz, 2.9 Hz, 1H, 3eq-H). ^{13}C nMR (100 MHz, CDCl_3): δ 162.6 (4'-C), 137.2 (1'-C), 129.84 (6-C), 129.83 (7-C), 129.65 (3'-C), 129.64 (5'-C), 125.9 (5-C), 125.95 (2'-C), 125.94 (6'-C), 122.7 (4a-C), 118.8 (8-C), 76.6 (2-C), 43.3 (3-C) [37].

2-(2'-Carboxyphenyl)benzopyran-4-one or 2'-carboxyflavanone (**4J**)

Prepared following the general procedure starting from chalcone **3J** (0.6 g, 2.24 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 70:30). Colorless solid (0.486 g, 1.81 mmol), 81% yield, m.p. (Gallenkamp apparatus): 130 °C.

^1H nMR (500 MHz, CDCl_3): δ 12.01 (br s, 2'-COOH) 7.94 (d, $J = 7.8$ Hz, 1H, 5-H), 7.69 (ddd, $J = 7.5$ Hz, 7.5 Hz, 0.8 Hz, 1H, 3'-H), 7.65 (dd, $J = 8.1$ Hz, 1.4 Hz, 1H, 6'-H), 7.60–7.55 (m, 2H, 4'-H, 5'-H), 7.51 (dd, $J = 8.1$ Hz, 1.4 Hz, 1H, 6-H), 7.06 (dd, $J = 8.4$ Hz, 0.7 Hz, 1H, 7-H), 6.91 (ddd, $J = 7.9$ Hz, 7.3 Hz, 0.9 Hz, 1H, 8-H), 6.16 (app t, $J = 6.5$ Hz, 1H, 2-H), 3.80 (dd, $J = 17.6$ Hz, 6.2 Hz, 1H, 3ax-H), 3.47 (dd, $J = 17.6$ Hz, 6.8 Hz, 1H, 3eq-H). ^{13}C nMR (100 MHz, CDCl_3): δ 201.6 (4-C, CO), 169.9 (2'-COOH), 162.6 (8a-C), 134.4 (1'-C), 129.6 (4'-C), 129.8 (6-C), 129.6 (7-C), 129.81 (3'-C), 129.80 (5'-C), 125.9 (5-C), 122.68 (2'-C), 122.67 (6'-C), 119.3 (4a-C), 118.4 (8-C), 76.6 (2-C), 43.3 (3-C). Calc. for $\text{C}_{16}\text{H}_{12}\text{O}_4$, $[\text{M} + \text{H}]^+$ m/z : 269.08084 Da, found: 269.08049, $[\text{M} + \text{Na}]^+$ m/z : 291.06278, found: 291.06275.

2-(5-bromo-2-methoxyphenyl)-2,3-dihydro-4H-benzopyran-4-one or 5'-bromo-2'-methoxy-flavanone (**4K**)

Prepared following the general procedure starting from chalcone **3K** (1.5 g, 4.50 mmol). This product was purified by column chromatography (hexane/dichloromethane 90:10, 70:30, 50:50, 70:30 then ethyl acetate and methanol). White solid (0.846 g, 0.25 mmol), 56% yield, m.p. (Gallenkamp apparatus): 130–135 °C.

^1H nMR (400 MHz, CDCl_3): δ 7.97 (dd, $J = 7.8$ Hz, 1.8 Hz, 1H, 5-H), 7.80 (dd, $J = 2.5$ Hz, 0.6 Hz, 1H, 6'-H), 7.54 (ddd, $J = 8.9$ Hz, 7.1 Hz, 1.7 Hz, 1H, 8-H), 7.46 (dd, $J = 8.9$ Hz, 2.5 Hz, 1H, 4'-H), 7.13–7.06 (m, 2H, 6-H, 7-H), 6.82 (d, $J = 8.8$ Hz, 1H, 3'-H), 5.80 (dd, $J = 13.4$ Hz, 2.8 Hz, 1H, 2-H), 3.85 (s, 3H, 2'-OCH₃), 3.00 (dd, $J = 16.8$ Hz, 13.4 Hz, 1H, 3ax-H), 2.85 (dd, $J = 16.8$ Hz, 2.8 Hz, 1H, 3eq-H). ^{13}C nMR (100 MHz, CDCl_3): δ 192.2 (4-C, CO), 161.7 (8a-C), 154.7 (2'-C), 136.1 (4'-C), 131.9 (6'-C), 129.8 (6-C), 129.3 (7-C), 127.1 (5-C), 121.7 (4a-C), 120.9 (1'-C), 118.1 (5'-C), 112.3 (3'-C, 8-C), 74.2 (2-C), 55.6 (2'-OCH₃), 43.7 (3-C). Calc. for $\text{C}_{16}\text{H}_{13}\text{BrO}_3$, $[\text{M} + \text{H}]^+$ m/z : 333.01208 Da, found: 333.01236, $[\text{M} + \text{Na}]^+$ m/z : 354.99403, found: 354.99396.

5-methoxy-2-phenyl-2,3-dihydro-4H-benzopyran-4-one or 5-methoxy-flavanone (**4L**)

Prepared following the general procedure starting from chalcone **3L** (0.657 g, 2.58 mmol). This product was purified by column chromatography ($\text{C}_6\text{H}_{12}/\text{EtOAc}$ 90:10, 70:30 and 50:50). Colorless solid (0.056 g, 0.2 mmol), 9% yield, m.p. (Electrothermal apparatus): 95–98.7 °C.

^1H nMR (500 MHz, CDCl_3): δ 7.49–7.34 (m, 6H, 2'-H, 3'-H, 4'-H, 5'-H, 6'-H, 7-H), 6.66 (dd, $J = 8.4$ Hz, 0.6 Hz, 1H, 8-H), 6.55 (d, $J = 8.3$ Hz, 1H, 6-H), 5.44 (dd, $J = 13.2$ Hz, 2.9 Hz, 1H,

2-H), 3.93 (s, 3H, 5-OCH₃), 3.07 (dd, *J* = 16.4 Hz, 13.2 Hz, 1H, 3ax-H), 2.86 (dd, *J* = 16.4 Hz, 2.9 Hz, 1H, 3eq-H). ¹³C NMR (125 MHz, CDCl₃): δ 190.7 (4-C, CO), 163.2 (5-C), 160.8 (8a-C), 138.7 (1'-C), 136.1 (7-C), 128.81 (3'-C, 5'-C), 128.68 (4'-C), 126.11 (2'-C, 6'-C), 111.4 (4a-C), 110.2 (8-C), 104.1 (6-C), 78.9 (2-C), 56.2 (5-OCH₃), 45.9 (3-C). Calc. for C₁₆H₁₄O₃, [M + H]⁺ *m/z*: 255.10157 Da, found: 255.10199 Da, [M + Na]⁺ *m/z*: 277.08352, found: 277.08392.

3.2. Biological Assays

3.2.1. Cell Culture and Treatments

The murine macrophage RAW 264.7 cell line (ATCC, TIB-71) was used as an *in vitro* model for studying the immunomodulatory properties of the molecules. Cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM) GlutaMAX™ (Gibco; 10566016) containing 4.5 g/L D-glucose, HEPES buffer at 25 mM, and supplemented with 10% fetal bovine serum (FBS; Gibco), antibiotics, and antifungal solution (penicillin, 10,000 U; streptomycin, 10 mg/mL; amphotericin B, 25 µg/mL). Cell cultures were maintained by cyclic resuspension in 75 cm² culture flasks in a humidified atmosphere with 5% CO₂ at 37 °C. For biological assays, cells were seeded at 150,000 cells/well in 96-well plates and left to recover for 24 h before treatments. The cells were incubated for an additional 24 h with LPS at 1 µg/mL and/or pinocembrin (Sigma–Aldrich, USA) or molecules in dimethyl sulfoxide (DMSO; Sigma Aldrich) at 0.1% and at the final concentration, as indicated in Figures 3–5. Control cells were incubated under the same conditions with or without DMSO at 0.1%, without LPS or molecules. Dexamethasone (Sigma–Aldrich; D4902-100MG) at 100 nM was used as the reference anti-inflammatory molecule inhibiting NO production ([38]). Cell culture supernatants were collected for nitrite quantification and cytotoxicity analyses.

3.2.2. Determination of Cell Mortality

Cytotoxicity was evaluated by quantifying the release of lactate dehydrogenase (LDH) in the culture supernatant that correlates with the amount of cell death and membrane damage, providing an accurate measurement of cellular toxicity [39]. LDH was quantified using the commercial CytoTox 96® Non-Radioactive Cytotoxicity Assay (Promega; G1780) following the manufacturer's specifications. Absorbance at 450 nm (A₄₅₀) was read using a microplate spectrophotometer (Multiskan™ FC, Thermo Fisher Scientific). LDH in the supernatants was normalized against absorbance obtained for total lysed cells, and results were expressed as percent of cytotoxicity, as recommended for LDH-based assays [40].

3.2.3. Quantification of Nitric Oxide (NO)

Nitrite (NO₂TM) production was determined as an indicator of Nitric Oxide (NO) synthesis in cell culture supernatant, as previously described [41], using the Griess Reagent System (Promega; G2930) and following the supplier's recommendations. Briefly, 50 µL of Griess reagent A (1% sulfanilamide in 5% phosphoric acid) was added to 50 µL of cellular supernatant and incubated for 5–10 min at room temperature, protected from light, before additional dispensing of Griess reagent B (0.1% N-1-naphthylethylenediamine dihydrochloride in water). Absorbance at 570 nm (A₅₇₀) was read using a microplate spectrophotometer (Multiskan™ FC, Thermo Fisher Scientific). Standard calibration curves were prepared using the provided sodium nitrite and after serial dilutions (0–100 µM). Due to the good correlation factor of this linear regression (*R*² = 0.98), the nitrite concentrations accumulated in the cellular supernatants was determined. The reduction or increase in nitrite levels was assessed, and results for molecules are expressed as the percentage of inhibitory response of nitrite production in the target sample relative to the LPS-induced nitrite production level, calculated as the inhibition of NO produced according to the following equation:

$$\% \text{ Inhibitory response} = 100 \times [1 - ([\text{NO}]_{\text{M+LPS}} - [\text{NO}]_{\text{Blank}}) / ([\text{NO}]_{\text{LPS}} - [\text{NO}]_{\text{Blank}})]$$

where $[\text{NO}]_{\text{Blank}}$, $[\text{NO}]_{\text{LPS}}$, and $[\text{NO}]_{\text{M+LPS}}$ are the concentrations of NO quantified when cells were not treated, induced with LPS, and treated with the molecules and LPS, respectively.

3.2.4. Predictive Analysis of Drug-Like Absorption

Log P predictions were performed using the software ACD/ChemSketch (ACD/Labs). Molecular properties were determined using the Molinspiration Cheminformatics website (<http://www.molinspiration.com/cgi-bin/properties>, accessed on 26 February 2022).

3.2.5. Statistical Analysis

Results are provided as mean \pm standard deviation (SD). Statistical analyses were performed using the software statistical package Prism 9.0 (GraphPad Software LLC, USA). Cytotoxicity and anti-inflammatory inhibitory response were evaluated using a Mann–Whitney nonparametric test to compare distribution between treatments and LPS induction. p values ≤ 0.05 were considered significant. The concentrations of molecules leading to 50% of inhibitory activity (IC_{50}) were also calculated using the software Prism 9.0 from a logarithm-transformed (inhibitor) vs. normalized response equation.

4. Conclusions

Twelve flavanone derivatives were successfully synthesized and were evaluated for their ability to inhibit NO production against commercial pinocembrin. Six of these derivatives were found to be highly active, with percentage inhibition responses greater than 50%. The most active flavanone derivatives were characterized by the presence of a carboxyl group in the *ortho*-position or a bromo-group in the *para*-position of the B-ring. The importance of the presence of methoxy-groups on the inhibition of NO production was noted, in line with findings by Lee in 2015. In particular, the incorporation of a methoxy group on ring A decreased the biological activity. In contrast, the least active molecules were 4'-chloro flavanone, 4'-methoxyflavanone, and pinocembrin. SAR identified flavanone 4F carrying a carboxyl group in the *ortho*-position of the B-ring as a potential new lead compound for anti-inflammatory activity.

Author Contributions: Conceptualization, C.S., V.S. and N.L.; methodology, C.S., V.S., N.L., M.O. and M.M.; validation, V.S., N.L. and M.O.; formal analysis, C.S., M.M., M.O., F.B. and V.C.; investigation, C.S., M.M. and N.L.; resources, N.L., M.O., M.M. and V.S.; writing—original draft preparation, C.S.; writing—review and editing, N.L., V.S., M.O. and M.M.; visualization, C.S. and M.M.; supervision, M.O., V.S. and N.L.; project administration, N.L.; funding acquisition, N.L. and M.O. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the foundation of University of New Caledonia (Project NORAS), the French Ministry of Higher Education, Research and Innovation (MESRI), the Institut Pasteur of New Caledonia, and the Pasteur Network. S.C., and M.M. thanks the Government of New Caledonia for scholarships.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: This work was supported by the foundation of University of New Caledonia (Project NORAS) and by the Government of New Caledonia for the PhD thesis of S.C. and the position of M.M. The authors are thankful to the College of Science and Engineering, Discipline of Chemistry, at James Cook University in Townsville for technical support, providing analysis equipment, and NMR experiments. The cell culture facility and analytical platform at the Institut Pasteur of New Caledonia are supported by the French Ministry of Higher Education, Research and Innovation (MESRI) and the Pasteur Network. We are thankful for the use of the analysis equipment of PEIRENE Laboratory (Limoges University, France), BISCEM Platform (Unité Mixte de Services,

Inserm 042, CNRS 2015, CHU of Limoges), for nMR experiments and grateful to Cyril Colas (ICOA, Orléans University, France) for HRMS experiments.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are not available from the authors.

Abbreviations

The following abbreviations are used in this manuscript.

| | |
|--------------------------------|---------------------------------------|
| NMR. | Nuclear magnetic resonance |
| ¹ H | Proton nuclear magnetic resonance |
| ¹³ C | Carbon nuclear magnetic resonance |
| NO | Nitric oxide |
| NOS | Nitric oxide synthase |
| LPS | Lipopolysaccharide |
| MeOH | Methanol |
| EtOAc | Ethyl acetate |
| EtOH | Ethanol |
| TLC | Thin Layer chromatography |
| PC | Pinocembrin |
| IC ₅₀ | Half maximal inhibitory concentration |
| LDH | Lactate dehydrogenase |
| SD | Standard deviation |
| DMSO | Dimethyl sulfoxide |
| DEX | Dexamethasone |
| DCM | Dichloromethane |
| CC | Column chromatography |
| m.p. | Melting point |
| Hz | Hertz |
| HRMS | High-resolution mass spectrometry |
| ESI | Electrospray ionization |
| M | Mass |
| CAS | Chemical abstracts service |
| Da | Dalton |
| W | Watt |
| C ₆ H ₁₂ | Cyclohexane |
| SAR | Structure–activity relationship |
| CI | Confidence Interval |

References

1. Maleki, S.J.; Crespo, J.F.; Cabanillas, B. Anti-inflammatory Effects of Flavonoids. *Food Chem.* **2019**, *299*, 125124. [[CrossRef](#)] [[PubMed](#)]
2. Tutunchi, H.; Naeini, E.; Ostadrahimi, A.; Hosseinzadeh-Attar, M.J. Naringenin, a Flavanone with Antiviral and Anti-Inflammatory Effects: A Promising Treatment Strategy against COVID-19. *Phytother. Res.* **2020**, *34*, 3137–3147. [[CrossRef](#)] [[PubMed](#)]
3. Gautam, R.; Jachak, S.M. Recent Developments in Anti-Inflammatory Natural Products. *Med. Res. Rev.* **2009**, *29*, 767–820. [[CrossRef](#)]
4. Ialenti, A.; Moncada, S.; Di Rosa, M. Modulation of Adjuvant Arthritis by Endogenous Nitric Oxide. *Br. J. Pharmacol.* **1993**, *110*, 701–706. [[CrossRef](#)] [[PubMed](#)]
5. Stichtenoth, D.O.; Frolin, J.C. Nitric Oxide and Inflammatory Joint Diseases. *Br. J. Rheumatol.* **1998**, *37*, 246–257. [[CrossRef](#)] [[PubMed](#)]
6. Ryu, J.H.; Lee, H.J.; Jeong, Y.S.; Ryu, S.Y.; Han, Y.N. Yomogin, an Inhibitor of Nitric Oxide Production in LPS-Activated Macrophages. *Arch. Pharmacol. Res.* **1998**, *21*, 481–484. [[CrossRef](#)]
7. Kobuchi, H.; Droy-lefaix, M.T.; Christen, Y.; Packer, L. Ginkgo Biloba Extract (Egb 761): Inhibitory Effect on Nitric Oxide Production in the Macrophage Cell Line RAW 264.7. *Biochem. Pharm.* **1997**, *53*, 897–903. [[CrossRef](#)]
8. Schwingshackl, L.; Schwedhelm, C.; Hoffmann, G.; Knüppel, S.; Preterre, A.; Iqbal, K.; Bechthold, A. Food Groups and Risk of Colorectal Cancer. *Int. J. Cancer* **2018**, *142*, 1748–1758. [[CrossRef](#)]
9. Van Breda, S.G.; De Kok, T.M.C.M. Smart Combinations of Bioactive Compounds in Fruits and Vegetables May Guide New Strategies for Personalized Prevention of Chronic Diseases. *Mol. Nutr. Food Res.* **2018**, *62*, 1700597. [[CrossRef](#)]
10. Pietta, P.G. Flavonoids as Antioxidants. *J. Nat. Prod.* **2000**, *63*, 1035–1042. [[CrossRef](#)]

11. Kanadaswami, C.; Lee, L.T.; Lee, P.H.; Hwang, J.J.; Ke, F.C.; Huang, Y.T.; Lee, M.T. The Antitumor Activities of Flavonoids. *Vivo* **2005**, *19*, 895–910.
12. Lolli, G.; Cozza, G.; Mazzorana, M.; Tibaldi, E.; Cesaro, L.; Donella-Deana, A.; Meggio, F. Inhibition of Protein Kinase CK2 by Flavonoids and Tyrphostins. A Structural Insight. *Biochemistry* **2012**, *51*, 6097–6107. [[CrossRef](#)] [[PubMed](#)]
13. Shen, X.; Liu, Y.; Luo, X.; Yang, Z. Advances in Biosynthesis, Pharmacology, and Pharmacokinetics of Pinocembrin, a Promising Natural Small-Molecule Drug. *Molecules* **2019**, *24*, 2323. [[CrossRef](#)] [[PubMed](#)]
14. Soromou, L.W.; Chu, X.; Jiang, L.; Wei, M.; Huo, M.; Chen, N.; Guan, S.; Yang, X.; Chen, C.; Feng, H.; et al. In Vitro and in Vivo Protection Provided by Pinocembrin against Lipopolysaccharide-Induced Inflammatory Responses. *Int. Immunopharmacol.* **2012**, *14*, 66–74. [[CrossRef](#)] [[PubMed](#)]
15. Gan, W.; Li, X.; Cui, Y.; Xiao, T.; Liu, R.; Wang, M.; Wei, Y. Pinocembrin Relieves Lipopolysaccharide and Bleomycin Induced Lung Inflammation via Inhibiting TLR4-NF-KB-NLRP3 Inflammasome Signaling Pathway. *Int. Immunopharmacol.* **2021**, *90*, 107230. [[CrossRef](#)]
16. Gabaston, J.; Richard, T.; Cluzet, S.; Pinto, A.P.; Dufour, M.; Corio-Costet, M.; Mérillon, J. Pinus Pinaster Knot: A Source of Polyphenols against *Plasmopara Viticola*. *J. Agric. Food Chem.* **2017**, *65*, 8884–8891. [[CrossRef](#)]
17. French, D.; Schifano, P.; Cortés-Concepcion, J.; Hargrove-Leak, S. Li-Al Layered Double Hydroxides as Catalysts for the Synthesis of Flavanone. *Catal. Commun.* **2010**, *12*, 92–94. [[CrossRef](#)]
18. Shi, L.; Feng, X.E.; Cui, J.R.; Fang, L.H.; Du, G.H.; Li, Q.S. Synthesis and Biological Activity of Flavanone Derivatives. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 5466–5468. [[CrossRef](#)]
19. Brennan, C.M.; Hunt, I.; Jarvis, T.C.; Johnson, D.; McDonnell, P.D. Stereoelectronic effects in ring closure reactions: The 2'-hydroxychalcone—flavanone equilibrium, and related systems. *Can. J. Chem.* **1990**, *1*, 1780–1785. [[CrossRef](#)]
20. Keane, D.D.; Marathe, K.G.; O'Sullivan, W.I.; Philbin, E.M.; Simons, R.M.; Teague, P.C. Configuration and Conformation of 3-Arylidene flavanones. *J. Org. Chem.* **1970**, *35*, 2286–2290. [[CrossRef](#)]
21. Hoshino, Y.; Takeno, N. Thermal Isomerization Equilibrium between 2'-Hydroxychalcones and Flavanones. *Bull. Chem. Soc. Jpn.* **1986**, *59*, 2903–2904. [[CrossRef](#)]
22. Pandey, G.; Krishna, A.; Kumaraswamy, G. Photosensitized (Set) Conversion of 2'-Hydroxychalcones to Flavonoids a Probable Biogenetic Pathway. *Tetrahedron Lett.* **1987**, *28*, 4615–4616. [[CrossRef](#)]
23. Geresh, S.; Levy, O.; Markovits, Y.; Shani, A. On the Mechanism of Intramolecular Photocycloaddition of Substituted O-Allylphenols to Cyclic Ethers. *Tetrahedron* **1975**, *31*, 2803–2807. [[CrossRef](#)]
24. Oelgemöller, M.; Hoffmann, N. Studies in Organic and Physical Photochemistry—an Interdisciplinary Approach. *Org. Biomol. Chem.* **2016**, *14*, 7392–7442. [[CrossRef](#)]
25. Matsushima, R.; Hiroyuki, K. Photochemical Cyclization of 2'-Hydroxychalcones. *J. Chem. Soc. Perkin Trans. 2* **1985**, *6*, 743–748. [[CrossRef](#)]
26. Tundis, R.L.; Frattaruolo, G.C.; Armentano, B.; Badolato, M.; Loizzo, M.; Aiello, F.; Cappello, A. An ancient remedial repurposing: Synthesis of new pinocembrin fatty acid acyl derivatives as potential antimicrobial/anti-inflammatory agents. *Nat. Prod. Res.* **2018**, *33*, 1–7. [[CrossRef](#)]
27. Forest, V.; Figarol, A.; Boudard, D.; Cottier, M.; Grosseau, P.; Pourchez, J. Adsorption of lactate dehydrogenase enzyme on carbon nanotubes: How to get accurate results about the cytotoxicity of these nanomaterials. *Langmuir* **2015**, *31*, 3635–3643. [[CrossRef](#)]
28. Lee, J.H. In-Vitro Evaluation for Antioxidant and Anti-Inflammatory Property of Flavanone Derivatives. *Food Biosci.* **2015**, *11*, 1–7. [[CrossRef](#)]
29. Kim, H.K.; Cheon, B.S.; Kim, Y.H.; Kim, S.Y.; Kim, H.P. Effects of Naturally Occurring Flavonoids on Nitric Oxide Production in the Macrophage Cell Line RAW 264.7 and Their Structure-Activity Relationships. *Biochem. Pharm.* **1999**, *58*, 759–765. [[CrossRef](#)]
30. Shin, S.Y.; Woo, Y.; Hyun, J.; Yong, Y.; Koh, D.; Lee, Y.H.; Lim, Y. Relationship between the Structures of Flavonoids and Their NF-KB-Dependent Transcriptional Activities. *Bioorganic Med. Chem. Lett.* **2011**, *21*, 6036–6041. [[CrossRef](#)]
31. Lipinski, C.A.; Lombardo, F.; Dominy, B.W.; Feeney, P.J. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv. Drug Deliv. Rev.* **2001**, *46*, 3–26. [[CrossRef](#)]
32. Boeck, P.; Bandeira Falcão, C.A.; Leal, P.C.; Yunes, R.A.; Filho, V.C.; Torres-Santos, E.C.; Rossi-Bergmann, B. Synthesis of Chalcone Analogues with Increased Antileishmanial Activity. *Bioorganic Med. Chem.* **2006**, *14*, 1538–1545. [[CrossRef](#)] [[PubMed](#)]
33. Thieury, C.; Lebouvier, N.; Le Guével, R.; Barguil, Y.; Herbette, G.; Anthaume, C.; Hnawia, E.; Asakawa, Y.; Nour, M.; Guillaudeux, T. Mechanisms of Action and Structure-Activity Relationships of Cytotoxic Flavokawain Derivatives. *Bioorganic Med. Chem.* **2017**, *25*, 1817–1829. [[CrossRef](#)] [[PubMed](#)]
34. Detsi, A.; Majdalani, M.; Kontogiorgis, C.A.; Hadjipavlou-Litina, D.; Kefalas, P. Natural and synthetic 2'-hydroxy-chalcones and aurones: Synthesis, characterization and evaluation of the antioxidant and soybean lipoxygenase inhibitory activity. *Bioorganic Med. Chem.* **2009**, *17*, 8073–8085. [[CrossRef](#)] [[PubMed](#)]
35. Qin, C.X.; Chen, X.; Hughes, R.A.; Williams, S.J.; Woodman, O.L. Understanding the Cardioprotective Effects of Flavonols: Discovery of Relaxant Flavonols without Antioxidant Activity. *J. Med. Chem.* **2008**, *51*, 1874–1884. [[CrossRef](#)]
36. Krüger, K.; Lüdke, V.; Pettinger, J.; Ashton, L.; Bonnet, L.; Motti, C.A.; Lex, J.; Oelgemöller, M. Photochemical synthesis of cyclic peptide models from phthalimido acetamides and phthaloyl dipeptide esters. *Tetrahedron Lett.* **2018**, *59*, 1427–1430. [[CrossRef](#)]
37. Kavala, V.; Lin, C.; Kuo, C.W.; Fang, H.; Yao, C.F. Iodine Catalyzed One-Pot Synthesis of Flavanone and Tetrahydropyrimidine Derivatives via Mannich Type Reaction. *Tetrahedron* **2012**, *68*, 1321–1329. [[CrossRef](#)]

38. Mazzi, E.A.; Li, N.; Bauer, D.; Mendonca, P.; Taka, E.; Darb, M.; Thomas, L.; Williams, H.; Soliman, K.F.A. Natural product HTP screening for antibacterial (*E.coli* 0157:H7) and anti-inflammatory agents in (LPS from *E. coli* O111:B4) activated macrophages and microglial cells; focus on sepsis. *BMC Complementary Altern. Med.* **2016**, *16*, 467. [[CrossRef](#)]
39. Haslam, G.; Wyatt, D.; Kitos, P.A. Estimating the Number of Viable Animal Cells in Multi-Well Cultures Based on Their Lactate Dehydrogenase Activities. *Cytotechnology* **2000**, *32*, 63–75. [[CrossRef](#)]
40. Riss, T.; Niles, A.; Moravec, R.; Karassina, N.; Vidugiriene, J. Cytotoxicity Assays: In Vitro methods to measure dead cells. In *Assay Guidance Manual*; Markossian, S., Grossman, A., Brimacombe, K., Arkin, M., Auld, D., Austin, C.P., Baell, J., Chung, T.D.Y., Coussens, N.P., Dahlin, J.L., et al., Eds.; Eli Lilly & Company and the National Center for Advancing Translational Sciences: Bethesda, MD, USA, 2004.
41. Kumar-Roiné, S.; Matsui, M.; Reybier, K.; Darius, H.T.; Chinain, M.; Pauillac, S.; Laurent, D. Ability of Certain Plant Extracts Traditionally Used to Treat Ciguatera Fish Poisoning to Inhibit Nitric Oxide Production in RAW 264.7 Macrophages. *J. Ethnopharmacol.* **2009**, *123*, 369–377. [[CrossRef](#)]