



Avant, leading protection and support when it matters most

Award-winning defence

Our over 300-strong* medico-legal defence team provides you with on-the-ground support across Australia.

Comprehensive support

Access our medico-legal experts, 24/7 in emergencies, and risk advisers. As well as a comprehensive library of online medico-legal resources.

Australia's most popular medical indemnity insurer, covering over 50% of doctors

Dr Mark Green
Avant member



Join today

avant.org.au/practitioners
1800 128 268



Avant
By doctors for doctors

IMPORTANT: Professional indemnity insurance products are issued by Avant Insurance Limited, ABN 82 003 707 471, AFSL 238 765. Please read the relevant Product Disclosure Statement or policy wording, available at www.avant.org.au before deciding whether to acquire, or continue to hold the product. *Accurate as at 30/06/2022. MJN1033 03/23 (DT-2889)

Self-discharge as a marker of surgical cultural competency and cultural safety for Aboriginal and/or Torres Strait Islander patients

Introduction

Australian Aboriginal and/or Torres Strait Islander patients experience disparities in surgical access^{1–4} and outcomes.^{5–11} Several factors impact the provision of surgical care to Aboriginal and/or Torres Strait Islander people including cultural factors, social determinants of health (social, political and economic factors), historical experiences (loss of land and culture, trans-generational trauma, grief and loss), racism and social exclusion.¹² Suboptimal surgical cultural competency and cultural safety may be a contributing factor to these disparities.

Cultural competency and cultural safety

In the healthcare setting cultural competency is a broad concept describing strategies to improve the accessibility and effectiveness of healthcare for people from ethnic/racial minority groups.¹³ This includes the ability of healthcare providers, institutions and systems, to incorporate the individual values, beliefs and behaviours of a population into the delivery and structure of healthcare provision.¹³ Cultural safety encompasses critical consciousness where healthcare professionals and organizations hold themselves accountable, recognizing their own stereotypes, prejudices and biases and working to ensure that they are providing culturally safe care.^{14,15} Increasing attention is being paid to healthcare cultural competency or safety and the potential correlation with disparities in surgical care.¹³

Monitoring cultural competency and cultural safety in the healthcare system is difficult.¹⁶ Cultural competency can be measured directly, for example, patient satisfaction or indirectly, for example, self-discharge.^{16,17} Self-discharge occurs when an inpatient chooses to leave the hospital before discharge is recommended by their clinicians.¹⁸ It reflects a failure to reach consensus between the healthcare staff and patient regarding the need for continued inpatient care. Thus, self-discharge can be utilized as a proxy for measuring healthcare cultural competency and cultural safety.

Self-discharge

Aboriginal and/or Torres Strait Islander status is the most significant variable contributing to self-discharge.¹⁹ Nationally Aboriginal and/or Torres Strait Islander hospitalized patients are six times more likely to self-discharge.¹⁹ Rates of self-discharge for all

hospitalisations are 3.1% for Aboriginal and/or Torres Strait Islander patients and 0.5% for other Australians.¹⁹ Alarming there has been an increasing rate of self-discharge over time, from 11 per 1000 in 2004–2005 to 16 per 1000 in 2016–2017.¹⁹ This may indicate a lack of hospital responsiveness to Aboriginal and/or Torres Strait Islander patient's needs.²⁰ One study from the state of Victoria found that 30% of Aboriginal and/or Torres Strait Islander people have experienced racism in health settings.¹⁷ When asked about their experiences with general practitioners in non-remote areas Aboriginal and/or Torres Strait Islander Australians reported that their doctor rarely or never: showed respect for what was said (15%), listened to them (20%) or spent enough time with them (21%).¹⁶ Potential markers of cultural competency and cultural safety like self-discharge following a surgical admission have not previously been examined for patients of Aboriginal and/or Torres Strait Islander descent in Australia.

Self-discharge is related to increased rates of postoperative complications, mortality,²¹ readmission^{21,22} and increased healthcare expenditure.^{22–24} These adverse events may be preventable.²² Healthcare staff are commonly concerned that these patients may be susceptible to serious health consequences as the result of not receiving the necessary care.¹⁸

The surgical care journey is complex and many different healthcare workers may be involved in patient care. Self-discharge measures the cultural competency and cultural safety along this journey.

Reasons for self-discharge

The reasons for self-discharge are diverse and complex. Qualitative Australian studies have identified several factors related to self-discharge in Aboriginal and/or Torres Strait Islander patients. Patients noted institutionalized racism²⁴ and a distrust of the healthcare system^{22,24} including anxiety, anger and negative associations with hospital.²⁵ Family and social obligations,^{24–26} including financial responsibilities especially on payday,²² pressure from partners and family to return home²² and transport cost and availability (may self-discharge when transport options become available).²² Some patients felt unwelcome or misunderstood²⁵ and that there was a lack of cultural safety.²⁴ Other factors included medical staff that lacked appropriate cultural training,²⁵ behaved in an inappropriate, insensitive or racist manner²² and that there was an absence of Aboriginal and/or Torres Strait Islander staff.²² Social isolation

and loneliness were also identified.^{24,26} Communication, language and cultural barriers between staff and patients were identified^{22,24,26} leading to miscommunication with hospital staff.^{24,26} Patients felt that they lacked an understanding of the care they were receiving,²⁴ felt that the treatment had finished²⁴ or felt that it is okay to leave.²⁶

The patient is always in a vulnerable position in relation to the health care system. Hence it should be an important task for a health care system to take responsibility and adopt to the cultural needs of their patients. However, current rates of self-discharge imply ineffective surgical cultural competency and cultural safety.

Quality metrics for cultural competency and cultural safety

Quality metrics (standardized measurable indicators) are widely used to assess health system performance and identify areas where quality improvement is needed. Efforts to reduce disparities require metrics that provide the foundation for focused and tailored interventions. Quality metrics are also used to monitor the impacts of a healthcare quality improvement initiatives. There is a lack of validated metrics of surgical cultural competency and cultural safety.

Strategies that aim to increase cultural competency and cultural safety may reduce healthcare disparities.¹³ Intervention strategies include: education and/or training of health professionals or students, culturally specific health programs or resources for Aboriginal and/or Torres Strait Islander people and increasing the number of Aboriginal and/or Torres Strait Islander health workers.¹² Increased Aboriginal and/or Torres Strait Islander health workers or liaison officers may be useful in decreasing self-discharge.^{25,27} However the care of Aboriginal and/or Torres Strait Islander patients should incorporate the entire health workforce. It is therefore imperative and urgent that cultural competency and cultural safety for the entire workforce should be improved. There is a lack of evidence from rigorous evaluations on the effectiveness of interventions for improving cultural competency and cultural safety in health care for global Indigenous populations.^{12,28} Previous studies examining cultural competency improvement strategies in Indigenous populations have not utilized self-discharge as a study end point. The Royal Australian College of Surgeons (RACS) has recently incorporated training on cultural competency and safety into the Surgical Education and Training program and added cultural competency and cultural safety to the RACS Framework. It will be important to assess the impact of this initiative on cultural competency and cultural safety.

Self-discharge data is readily available in administrative datasets and could be used as a metric to monitor, benchmark and improve, the cultural competency and cultural safety of a surgical department and possibly also in other specialties.

Conclusion

Nationally Aboriginal and/or Torres Strait Islander patients are significantly more likely, than other Australians, to self-discharge. This may indicate suboptimal healthcare, including surgical cultural competency and cultural safety. Self-discharge data is readily

available in administrative datasets and could be used as a metric to monitor, benchmark and improve, the cultural competency and cultural safety of a surgical department and most likely also other departments.

Author contributions

Elzerie de Jager: Conceptualization; funding acquisition; investigation; project administration; resources; validation; writing – original draft; writing – review and editing. **Ronny Gunnarsson:** Supervision; validation; writing – review and editing. **Yik-Hong Ho:** Resources; supervision; validation; writing – review and editing.

Acknowledgements

We acknowledge Prof Yvonne Cadet-James for proofing this manuscript as an Aboriginal and Torres Strait Islander cultural mentor to ensure cultural sensitivity. We gratefully acknowledge the Traditional Owners of the Townsville Region; the Wulgurukaba of Gurambilbarra and Yunbenun, Bindal, Gugu Badhun and Nywaigi people. We pay our respects to their cultures, their ancestors and their Elders, past and present – and all future generations. Open access publishing facilitated by James Cook University, as part of the Wiley - James Cook University agreement via the Council of Australian University Librarians.

Disclosures

Elzerie de Jager was a Doctor of Philosophy candidate at an Australian university; this research degree was supported by an Australian Government Research Training Program (RTP) Scholarship. This work was also supported by the Avant Doctor in Training Research Scholarship.

References

1. Australian Institute of Health and Welfare. Admitted Patient Care 2017–18: Australian Hospital Statistics. Canberra; 2019.
2. Cheng WYC, Blum P, Spain B. Barriers to effective perioperative communication in indigenous Australians: an audit of Progress since 1996. *Anaesth. Intensive Care* 2004; **32**: 542–7.
3. Ishak M. Differences in the surgical admissions to hospital between the indigenous and non-indigenous populations of Australia. *ANZ J. Surg.* 2003; **73**: 200–4.
4. Prin M, Guglielminotti J, Mtalimanja O, Li G, Charles A. Emergency-to-elective surgery ratio: a global indicator of access to surgical care. *World J. Surg.* 2018; **42**: 1971–80.
5. Alizzi AM, Knight JL, Tully PJ. Surgical challenges in rheumatic heart disease in the Australian indigenous population. *Heart Lung Circ.* 2010; **19**: 295–8.
6. Wiemers P, Mamey L, Muller R *et al.* Cardiac surgery in Indigenous Australians—how wide is “the gap”? *Heart Lung Circ.* 2014; **23**: 265–72.
7. Prabhu A, Tully PJ, Bennetts JS, Tuble SC, Baker RA. The morbidity and mortality outcomes of indigenous Australian peoples after isolated

- coronary artery bypass graft surgery: the influence of geographic remoteness. *Heart Lung Circ.* 2013; **22**: 599–605.
8. Anonymous. Injury in indigenous communities: a call for enlightened action. *ANZ J. Surg.* 2007; **77**: 930–1.
 9. Henman K, Gordon CL, Gardiner T *et al.* Surgical site infections following caesarean section at Royal Darwin Hospital, Northern Territory. *Healthcare Infect.* 2012; **17**: 47–51.
 10. Thong DW, Kim J, Dobson B, Cheung H, Arthur T. Variation in antimicrobial prescription and complications post emergency appendectomy in Australia: do we follow recommended guidelines? *ANZ J. Surg.* 2019; **90**(3): 251–256.
 11. Singh TP, Moxon J V, Healy GN, Cadet-James Y, Gollidge J. Presentation and outcomes of indigenous Australians with peripheral artery disease; 2018.
 12. Clifford A, McCalman J, Bainbridge R, Tsey K. Interventions to improve cultural competency in health care for indigenous peoples of Australia, New Zealand, Canada and the USA: a systematic review. *International J. Qual. Health Care* 2015; **27**: 89–98.
 13. Thackrah RD, Thompson SC. Refining the concept of cultural competence: building on decades of progress. *Med. J. Aust.* 2013; **199**: 35–8.
 14. Ronald MM, Aramoana JKA, Hill AG. Surgeons and cultural safety and cultural competency: the road to transformation. *ANZ J. Surg.* 2020; **90**: 2563–6.
 15. Curtis E, Jones R, Tipene-Leach D *et al.* Why cultural safety rather than cultural competency is required to achieve health equity: a literature review and recommended definition. *Int. J. Equity Health* 2019; **18**: 174.
 16. Council Australian Health Ministers' Advisory. Aboriginal and Torres Strait Islander health performance framework 2017 Report. Canberra; 2017.
 17. Kelaher MA, Ferdinand AS, Paradies Y. Experiencing racism in health care: the mental health impacts for Victorian Aboriginal communities. *Med. J. Aust.* 2014; **201**: 44–7.
 18. Yong TY, Fok JS, Hakendorf P, Ben-Tovim D, Thompson CH, Li JY. Characteristics and outcomes of discharges against medical advice among hospitalised patients. *Intern. Med. J.* 2013; **43**: 798–802.
 19. Australian Institute of Health and Welfare. Aboriginal and Torres Strait Islander health performance framework (HPF)—AIHW Indigenous HPF; 2020.
 20. Bridging the Health Equity Gap A Submission from the Cooperative Research Centre for Aboriginal Health to the National Health and Hospitals Reform Commission; 2008.
 21. Glasgow JM, Vaughn-Sarrazin M, Kaboli PJ. Leaving against medical advice (AMA): risk of 30-day mortality and hospital readmission. *J. Gen. Intern. Med.* 2010; **25**: 926–9.
 22. Henry B, Dunbar T, Barclay L, Thompson R. Self-discharge against medical advice from Northern Territory hospitals. Darwin; 2007.
 23. Aliyu ZY. Discharge against medical advice: sociodemographic, clinical and financial perspectives. *Int. J. Clin. Pract.* 2002; **56**: 325–7.
 24. Shaw C. An Evidence-Based Approach to Reducing Discharge against Medical Advice amongst Aboriginal and Torres Strait Islander Patients. Deeble Institute for Health Policy Research & Australian Healthcare and Hospitals Association. Deakin, ACT; 2016.
 25. Katzenellenbogen JM, Sanfilippo FM, Hobbs MS *et al.* Voting with their feet - predictors of discharge against medical advice in aboriginal and non-aboriginal ischaemic heart disease inpatients in Western Australia: an analytic study using data linkage. *BMC Health Serv. Res.* 2013; **13**(330). doi: 10.1186/1472-6963-13-330.
 26. Franks C, Beckmann K. A qualitative analysis of patients taking their own leave from Alice Springs Hospital in 1998—version details—trove. *Aboriginal Islander Health Worker J* 2002; **26**: 3–8.
 27. Einsiedel LJ, Van Iersel E, MacNamara R *et al.* Self-discharge by adult aboriginal patients at Alice Springs hospital, Central Australia: insights from a prospective cohort study. *Aust. Health Rev.* 2013; **37**: 239–45.
 28. Truong M, Paradies Y, Priest N. Interventions to Improve Cultural Competency in Healthcare: A Systematic Review of Reviews. Vol. 14, BMC Health Services Research. BioMed Central, p. 99; 2014.

Elzerie de Jager,* MBBS (Hons), PhD 

Ronny Gunnarsson,†‡§ MD, PhD

Yik-Hong Ho,*¶ MD, FRACS

*College of Medicine and Dentistry, The James Cook University, Townsville, Queensland, Australia, †General Practice/Family Medicine, School of Public Health and Community Medicine, Institute of Medicine, the Sahlgrenska Academy, University of Gothenburg, Goteborg, Sweden, ‡Research, Education, Development & Innovation, Primary Health Care, Västra Götaland Region, Sweden, §Primary Health Care Center for Homeless People, Närhälsan, Västra Götaland Region, Sweden and ¶Townsville Clinical School, The Townsville Hospital, Townsville, Queensland, Australia

doi: 10.1111/ans.18238