



Draft Genome Sequence of a Novel Adenovirus Recovered from the Metagenome of Agile Wallabies

God'spower Richard Okoh,^a Ellen Ariel,^a David Whitmore,^a Paul F. Horwood^a

^aJames Cook University, College of Public Health, Medical and Veterinary Sciences, Townsville, Queensland, Australia

ABSTRACT Here, we report the draft genome sequence of a novel agile wallaby adenovirus that was detected in the fecal metagenome of agile wallabies. The genome is 31,512 bp long, with a G+C content of 34.4%. Currently, the pathogenic and zoonotic potential of this novel virus is unknown.

Adenoviruses are nonenveloped, icosahedral viruses with linear, unsegmented double-stranded DNA genomes (1). The genomes of adenoviruses range in size from 26 to 48 kb and commonly contain between 22 and 40 genes (1, 2). The family *Adenoviridae* is divided into six genera, namely, *Mastadenovirus*, *Aviadenovirus*, *Atadenovirus*, *Siadenovirus*, *Ichtadenovirus*, and *Testadenovirus* (1, 3). Adenoviral infections do not always result in disease, although they have been associated with both single and multipathogen disease processes (4–8).

The novel Agile wallaby atadenovirus 1 (AwAdv-1) described in this report belongs to the genus *Atadenovirus* and was originally identified in the metagenome of free-ranging agile wallabies (*Notamacropus agilis*) (G. R. Okoh, E. Ariel, D. Whitmore, P. F. Horwood, submitted for publication). Briefly, five fresh fecal samples were collected from the ground at grazing sites around James Cook University and Townsville University Hospital (Townsville, Australia) in 2021. The samples were homogenized, pooled, and then virally enriched by filtration (0.25 μm), ultracentrifugation (100,000 × g), and digestion with DNase I (20 U/mL). Viral DNA was then extracted using QIAamp MinElute virus kit (Qiagen). Library preparation using the Nextera DNA XT kit and Illumina sequencing (NovaSeq 6000) were performed at Macrogen (Seoul, South Korea) in paired-end 151-bp format. For this report, the sequencing reads (71,170,820) were trimmed (Trimmomatic v0.39) (9) to remove low-quality reads, normalized using BBNorm v39.01 (<https://sourceforge.net/projects/bbmap/>), and *de novo* assembled using SPAdes v3.15.5 in “careful” mode (10). The resulting contigs were searched using Diamond BLASTX (11) against the NCBI nonredundant (nr) protein database to identify the contigs corresponding to adenovirus. To assemble the genome, the reads were mapped to the identified adenoviral contig using Geneious v11.1.5 (<https://www.geneious.com>). Prediction of open reading frames (ORFs) was performed using Glimmer3 in Geneious v11.1.5, and ORF annotations were determined by conducting a BLASTX search against the NCBI nr protein databases (12). All tools were run with default parameters unless otherwise specified.

The assembled genome of AwAdv-1 was found to be 31,512 bp long, with a coverage depth of 22× and 34.4% G+C content. The genome was predicted to contain 32 ORFs with an orientation typical of atadenoviruses. Of the 32 ORFs, 26 were annotated as having various similarities to the coding genes of other atadenoviruses (Tables 1). The IVa2, penton base protein, PX, and hexon genes showed the highest amino acid identity (71% to 87%) to the reference mammalian atadenoviruses (Table 1). The AwAdv-1 genome possesses multiple insertions and deletions in most of the genes except IVa2, pX, pVI, pVIII, and U-exon. Two fiber genes, namely, fiber and IV-1 (homologous to the fiber 2 gene in lizard adenovirus 2), were present in the genome of AWAdv-1, instead of the single long fiber gene in mammalian atadenoviruses. Phylogenetic analysis based on the full amino

Editor Jelle Matthijnssens, Katholieke Universiteit Leuven

Copyright © 2023 Okoh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to God'spower Richard Okoh, godspower.okoh@jcu.edu.au, or Paul F. Horwood, paul.horwood@jcu.edu.au.

The authors declare no conflict of interest.

Received 15 February 2023

Accepted 26 April 2023

Published 18 May 2023

TABLE 1 Sequence comparison with the genomes of mammalian adenoviruses

Data for virus (GenBank accession no.): ^a														
Agile wallaby adenovirus 1		Ovine adenovirus (NC_004037)			Bovine adenovirus D (NC_002685)			Bovine adenovirus E (NC_020074)			Odocoileus adenovirus 1 (NC_035619)			
ORF	Description (putative)	Length (nt)	Length (aa)	Length (nt)	Length (aa)	Amino acid identity (%)	Length (nt)	Length (aa)	Amino acid identity (%)	Length (nt)	Length (aa)	Amino acid identity (%)	Length (nt)	Amino acid identity (%)
orf0001	Hypothetical protein	255	85	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0002	Hypothetical protein	402	134	861	286	39	819	272	45	891	296	40	930	309
orf0003	p32K	1,032	344	363	120	29	378	125	29	378	125	26	393	130
orf0004	LH1	396	132	1149	382	43	1,161	386	42	1,152	383	45	1,146	381
orf0005	E1B 55K	1,293	431	984	327	75	966	321	74	966	321	72	1,209	402
orf0006	IVa2	903	301	3,216	1,071	58	3,222	1,073	58	3,222	1,073	57	3,228	1,075
orf0007	Pol	3,243	1,081	591	1,788	50	1,803	600	51	1,803	600	51	1,800	599
orf0008	pTP	1,773	52K	987	329	1,008	335	62	1,059	343	62	1,035	344	60
orf0009	pIIA	1,767	589	1,707	568	53	1,722	573	54	1,551	516	54	1,014	337
orf0010	Penton base protein	1,347	449	1,359	452	68	1,353	450	67	1,359	452	68	1,353	450
orf0011	pVII	345	115	336	111	54	360	119	57	357	118	56	354	117
orf0012	px	102	34	216	71	87	216	71	81	219	72	0	No data	No data
orf0013	pVI	669	223	666	221	54	603	200	57	612	203	56	678	225
orf0014	Hexon	2,730	910	2,736	911	74	2,733	910	72	2,733	910	76	2,733	910
orf0015	23K endopeptidase	606	202	606	201	59	606	201	60	606	201	61	606	201
orf0016	DNA binding protein	999	333	1,149	382	56	1,143	380	58	1,140	379	56	1,158	385
orf0017	100K	2,055	685	1,878	625	54	1,887	628	56	1,887	628	54	1,914	637
orf0018	33K	512	170	402	133	39	405	134	65	408	135	39	414	137
orf0019	pVIII	780	260	654	217	44	669	222	44	672	223	45	681	226
orf0020	U-exon	165	55	177	58	35	165	54	54	165	54	54	165	54
orf0021	Fiber	891	297	1,632	543	36	1,608	535	36	1,332	443	43	1,422	473
orf0022	IV-1	1,554	518	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0023	E4.3	600	200	714	237	34	654	217	38	657	218	38	705	234
orf0024	E4.2	678	226	663	220	40	678	219	42	660	219	42	678	219
orf0025	RH0	441	147	429	142	36	429	142	37	429	142	38	429	142
orf0026	RH5	381	127	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0027	Hypothetical protein	606	202	597	198	26	624	207	27	651	216	24	624	207
orf0028	Hypothetical protein	372	124	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0029	Hypothetical protein	201	67	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0030	Hypothetical protein	366	122	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0031	Hypothetical protein	372	124	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0032	Hypothetical protein	100a	67	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0033	Hypothetical protein	366	122	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data
orf0034	Hypothetical protein	372	124	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data	No data

^a aa, amino acids.




FIG 1 Phylogenetic analysis of Agile wallaby atadenovirus 1 (shown in red), based on the amino acid sequence of penton-based protein. All the sequences used in this analysis belong to the genus *Atadenovirus*. Following multiple sequence alignment of amino acid sequences using the Muscle program in Geneious v11.1.5, a maximum likelihood phylogenetic tree was constructed using MEGA X (13) with the best model of amino acid substitution (LG + G) and 1,000 bootstrap replications.

acid sequence of penton base protein showed that AwAdV-1 belongs to the genus *Atadenovirus* and forms a distinct cluster with another marsupial adenovirus known as possum adenovirus 1 (Fig. 1).

The pathogenic potential of AwAdV-1 is unclear; however, it could be a suitable candidate for future research in vaccinology, diagnostics, and therapeutics.

Data availability. The raw sequence reads for this study have been deposited in the NCBI SRA database under BioProject accession number [PRJNA907146](#) and BioSample accession number [SAMN31952915](#). The novel genome sequence has been deposited at GenBank under the accession number [OQ792214](#).

ACKNOWLEDGMENTS

We thank the Australian Wildlife Society and College of Public Health, Medical and Veterinary Sciences, James Cook University, for funding this research.

REFERENCES

1. Madarame H, Uchiyama J, Tamukai K, Katayama Y, Osawa N, Suzuki K, Mizutani T, Ochiai H. 2019. Complete genome sequence of an adenovirus-1 isolate from an African pygmy hedgehog (*Atelerix albiventris*) exhibiting respiratory symptoms in Japan. *Microbiol Resour Announc* 8:e00695-19. <https://doi.org/10.1128/MRA.00695-19>.
2. Davison AJ, Benkő M, Harrach B. 2003. Genetic content and evolution of adenoviruses. *J Gen Virol* 84:2895–2908. <https://doi.org/10.1099/vir.0.19497-0>.
3. Böszörményi KP, Podgorski Ii, Vidovszky MZ, Sós E, Benkő M, Harrach B. 2020. Full genome sequence analysis of a novel adenovirus from a captive polar bear (*Ursus maritimus*). *Virus Res* 277:197846. <https://doi.org/10.1016/j.virusres.2019.197846>.
4. Jesse ST, Ciurkiewicz M, Siesenop U, Spitzbarth I, Osterhaus ADME, Baumgärtner W, Ludlov M. 2022. Molecular characterization of a bovine adenovirus type 7 (bovine atadenovirus F) strain isolated from a systemically infected calf in Germany. *Virol J* 19:89. <https://doi.org/10.1186/s12985-022-01817-y>.
5. Fent GM, Fulton RW, Saliki JT, Caseltine SL, Lehmkuhl HD, Confer AW, Purdy CW, Briggs RE, Loan RW, Duff GC. 2002. Bovine adenovirus serotype 7 infections in postweaning calves. *Am J Vet Res* 63:976–978. <https://doi.org/10.2460/ajvr.2002.63.976>.
6. Thomson D, Meers J, Harrach B. 2002. Molecular confirmation of an adenovirus in brushtail possums (*Trichosurus vulpecula*). *Virus Res* 83:189–195. [https://doi.org/10.1016/s0168-1702\(01\)00437-3](https://doi.org/10.1016/s0168-1702(01)00437-3).
7. Marschang RE. 2019. Emerging reptile viruses, p 267–273. In Miller RE, Lamberski N, Calle PP (ed), *Fowler's zoo and wild animal medicine: current therapy*, vol 9. Elsevier, St. Louis, MO.
8. Chen EC, Yagi S, Kelly KR, Mendoza SP, Tarara RP, Canfield DR, Maninger N, Rosenthal A, Spinner A, Bales KL, Schnurr DP, Lerche NW, Chiu CY. 2011. Cross-species transmission of a novel adenovirus associated with a fulminant pneumonia outbreak in a new world monkey colony. *PLoS Pathog* 7:e1002155. <https://doi.org/10.1371/journal.ppat.1002155>.
9. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pshkin AV, Sirotnik AV, Vyazhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
11. Buchfink B, Reuter K, Drost HG. 2021. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* 18:366–368. <https://doi.org/10.1038/s41592-021-01101-x>.
12. Sayers EW, Beck J, Bolton EE, Bourexis D, Brister JR, Canese K, Comeau DC, Funk K, Kim S, Klimke W, Marchler-Bauer A, Landrum M, Lathrop S, Lu Z, Madden TL, O'Leary N, Phan L, Rangwala SH, Schneider VA, Skripchenko Y, Wang J, Ye J, Trawick BW, Pruitt KD, Sherry ST. 2021. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 49:D10–D17. <https://doi.org/10.1093/nar/gkaa892>.
13. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Mol Biol Evol* 35:1547–1549. <https://doi.org/10.1093/molbev/msy096>.