

Grasses of Townsville

2nd edition

Nanette B. Hooker

Photographs by Greg Calvert, Chris Gardiner and Russell Cumming



Published by James Cook University Townsville QLD 4811 Australia
©2023 Nanette Hooker.
First edition 2016
Second edition 2023

Author: Hooker, N.B. (Nanette Barbara)
Photographers: Calvert G.A., Gardiner C.P., Cumming R.J.
Scanned specimens Australian Virtual Herbarium
Title: Grasses of Townsville 2nd edition
ISBN: 978-0-6457422-7-5
Subjects: Poaceae - Grasses - Townsville - Identification
Author's email: nanette.hooker@jcu.edu.au



Front cover photographs
Cymbopogon ambiguus by Russell Cumming
Chionachne cyathopoda by Russell Cumming

This is an open access book distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<https://researchonline.jcu.edu.au/81169/>

Grasses of Townsville

Contents

Grasses of the Townsville area.....	1
Grass genera summary	3
Key to grass genera in Townsville	4
Key to the species with thickened rachis and a single raceme or spike	16
Subfamilies and Tribes of Grasses (excluding bamboos) in Townsville	17
Grass genera and species descriptions.....	26
Townsville grasses species list	361
Townsville grasses common names list	365
References	369


Grasses of the Townsville area

Welcome to the grasses of the Townsville area. The genera covered in this publication are those found in the lowland areas around Townsville as far north as Bluewater, south to Alligator Creek and west to the base of Hervey's Range.

This book provides a means of identifying the native and naturalised grasses of the Townsville region. The aim of this book is to help students, researchers, conservationists and anyone else who is interested in the Poaceae family. This book provides a description and photographs of the genera and species.

The grasses belong to a very widespread and large family called the Poaceae. The original family name, Gramineae, is used in some publications, in Australia the preferred family name is Poaceae. It is one of the largest flowering plant families of the world, comprising more than 700 genera, and more than 10,000 species. In Australia there are over 1300 species including non-native grasses. There are 175 natives, 65 non-natives and 4 cultivated grasses found in the Townsville area, and I have described grass species, subspecies and varieties.

The grasses have highly modified flowers arranged in a variety of ways. Because they are highly modified and specialized, there are also many terms used to describe the various features. Hence there is a lot of terminology that chiefly applies to grasses, but some terms are used also in the sedge family. The basic unit of the grass inflorescence (the flowering part) is the spikelet. The spikelet consists of 1-2 basal glumes (bracts at the base) that subtend 1-many florets or flowers. Each basic floret consists of lemma and palea; these enclose the male and female organs.



There are many basic variations on this basic pattern, sometimes the palea may be missing for instance or the floret may be sterile or neuter, i.e. there are no male or female organs inside. Some spikelets are bisexual (both stamens and ovary present) or unisexual (only male or female organs present). Sometimes male and female flowers are in different spikelets, then the plant is said to be monoecious, if male on one plant and female on another as for the Beach Spinifex then the plant is dioecious.

Because of the difficulties in understanding the various terms, in this book illustrations are used as much as possible. To find your way to a genus, there are two routes you can follow, the first method is 'flick till you find'. The second is to use what is called a 'key'. This is based on pairs of contrasting statements or couplets. Both statements must be read to find out which one 'fits' your plant and then you go to the corresponding number and continue like this till you find the description that fits. Diagrams have been used as much as possible to supplement the words.

Several useful references are:

Jacobs SWL, Whalley RDB, Wheeler DJB (2008), *Grasses of New South Wales* (Fourth Edition). University of New England, Armidale

Mallett K, Orchard AE (Eds) (2002), *Flora of Australia Volume 43, Poaceae 1: Introduction and Atlas*. ABRIS/CSIRO Publishing, Melbourne

Wheeler DJB, Jacobs SWL, Whalley RDB (2002), The grass plant. Extract from pages 6-23 of *Grasses of New South Wales* (Third Edition). University of New England, Armidale
http://www.fog.org.au/grasses_of_nsw/grasses_of_nsw.htm

There are several groups of plants which are sometimes mistaken for grasses.

Families which have grass-like species or could be mistaken for grasses are sedges (Cyperaceae) which are found in moist habitats.

The following table provides a comparison of these families, all of whom do not have typical petal-like flowers.

	Poaceae (grasses)	Cyperaceae (sedges)	Lomandra	Dianella
Stems	Hollow or solid Round	Solid Triangular or round	Round	Round
Leaves	2-ranked	3-ranked	Spiral	2-ranked
Ligules	Mostly present	Absent	Absent	Absent
Flowers	Commonly bisexual, enclosed by a lemma and palea and subtended by sterile glumes	Bisexual or unisexual and monoecious, enclosed by a single bract (glume)	Unisexual with petaloid perianth	Bisexual petaloid perianth
Floral formula	P0 A3 G(2)	P0 A3 G(2-3)	P3+3 A6 <u>G</u> 3	P3+3 A6 <u>G</u> 3
Fruit	Caryopsis	Achene	Capsule	Capsule

Grass Genera summary

The 75 genera of grass which have been found in Townsville area are listed here.

<i>Alloteropsis</i>	<i>Hymenachne</i>
<i>Ancistrachne</i>	<i>Hyparrhenia</i>
<i>Andropogon</i>	<i>Imperata</i>
<i>Aristida</i>	<i>Ischaemum</i>
<i>Arthragrostis</i>	<i>Leersia</i>
<i>Arundinella</i>	<i>Lepturus</i>
<i>Arundo</i>	<i>Megathyrsus</i>
<i>Axonopus</i>	<i>Melinis</i>
<i>Bothriochloa</i>	<i>Mnesithea</i>
<i>Brachyachne</i>	<i>Ophiuros</i>
<i>Capillipedium</i>	<i>Oplismenus</i>
<i>Cenchrus</i>	<i>Oryza</i>
<i>Chionachne</i>	<i>Oxychloris</i>
<i>Chloris</i>	<i>Panicum</i>
<i>Chrysopogon</i>	<i>Paspalidium</i>
<i>Cleistochloa</i>	<i>Paspalum</i>
<i>Cymbopogon</i>	<i>Perotis</i>
<i>Cynodon</i>	<i>Phragmites</i>
<i>Dactyloctenium</i>	<i>Pseudopogonatherum</i>
<i>Dichanthium</i>	<i>Pseudoraphis</i>
<i>Digitaria</i>	<i>Sacciolepis</i>
<i>Dinebra</i>	<i>Sarga</i>
<i>Diplachne</i>	<i>Schizachyrium</i>
<i>Echinochloa</i>	<i>Sehima</i>
<i>Ectrosia</i>	<i>Setaria</i>
<i>Eleusine</i>	<i>Sorghum</i>
<i>Elionurus</i>	<i>Spinifex</i>
<i>Elytrophorus</i>	<i>Sporobolus</i>
<i>Enneapogon</i>	<i># Stenotaphrum</i>
<i>Enteropogon</i>	<i>Themeda</i>
<i>Eragrostis</i>	<i>Thuarea</i>
<i>Eremochloa</i>	<i>Triodia</i>
<i>Eriachne</i>	<i>Tripogonella</i>
<i>Eriochloa</i>	<i>Urochloa</i>
<i>Eulalia</i>	<i>Vacoparis</i>
<i>Hackelochloa</i>	<i>Whiteochloa</i>
<i>Hemarthria</i>	<i># Zoysia</i>
<i>Heteropogon</i>	

Some garden (cultivated) plants are included and are indicated by #.




The species numbers in the genera description includes species, subspecies and varieties.

The world numbers are Accepted Names from the World Flora Online (<https://www.worldfloraonline.org/>).





Australian number includes species that have not given a species name. The number is taken from AusGrass 2 (<https://ausgrass2.myspecies.info/>) and Australian Virtual Herbarium (<https://avh.ala.org.au/>).

Key to the grass Genera in Townsville




1. Spikelets morphologically or functionally unisexualgo to 2
Spikelets bisexualgo to 4
2. Plants dioecious i.e. with male and female spikelets on different plants, grows on sandy seashores *Spinifex*
Plants monoecious, i.e. with male and female spikelets on the same plant, grows on sandy seashores or creek banks.....go to 3

		
Dioecious plant - female flowers	Dioecious plant - male flowers	Monoecious plant- male and female flowers



3. Plant tufted, more than 1 m tall, grows on creek banks..... *Chionachne*
Plant prostrate, grows on sandy seashores..... *Thuarea*

			
Spikelet with one floret	Spikelet with many florets	Panicoid spikelet Paniceae tribe	Panicoid spikelet pair Andropogoneae tribe





4. Spikelets with one-to-many florets, if two-flowered both florets or lower one bisexualgo to 5
Spikelets two flowered, lower floret male or barren, upper bisexual or female go to 31
5. Inflorescence a solitary, bilateral spike (almost cylindrical); spikelets solitary and partially embedded in rachis; lower glume absent or obscure; a coastal grass *Lepturus*
Grasses without the above combination of charactersgo to 6
6. Fertile lemmas distinctly nine-lobed..... *Enneapogon*
Fertile lemmas entire, dentate or slightly lobed.....go to 7

		
Florets		
Fertile lemmas distinctly 9-lobed	Fertile lemmas entire	Fertile lemmas dentate or slightly lobed


7. Leaf blades hard, woody, needle-like..... *Triodia*
 Leaf blades not hard, woody, needle-like.....go to 8
8. Plants tall and reed like, usually more than 2 m tallgo to 9
 Plants not tall and reed like go to 10
9. Lemmas not hairy (glabrous); rachilla (axis of spikelet) hairy *Phragmites*
 Lemmas hairy; rachilla not hairy *Arundo*

	
Florets	Florets
Glumes	Glumes
Lemma	Lemma
Lemmas hairy = <i>Arundo</i>	Rachilla hairy = <i>Phragmites</i>


10. Spikelets with two or more bisexual florets, or if only one bisexual floret with sterile florets (empty lemmas) above it go to 11
 Spikelets with one bisexual floret, with no sterile florets above it go to 22

			
Florets	Sterile florets	Floret	Floret
Glumes	Fertile florets	Glumes	Upper glume
	Glumes		Rachis
Spikelets with two or more bisexual florets, or if one bisexual floret with sterile florets above it		Spikelets with one bisexual floret	




11. Inflorescence digitate or subdigitate..... go to 12
 Inflorescence a raceme, spike or panicle go to 16






12. Florets unawned or shortly awned..... go to 13
 Florets with distinct awns..... go to 14





13. Axis of each inflorescence branch ending in a bristle; spikelet-bearing axis disarticulating *Dactyloctenium*
 Axis of each inflorescence branch ending in a spikelet; spikelet-bearing axis not disarticulating *Eleusine*
14. Lowest lemma dorsally compressed (lying on front or back when placed on a flat surface) *Enteropogon*
 Lowest lemma laterally compressed (lying on the side when placed on a flat surface) go to 15
15. Lemmas very broad, wing-like *Oxychloris*
 Lemmas not very broad, not wing-like..... *Chloris*
16. Spikelets distinctly awned go to 17
 Spikelets unawned or mucronate or shortly awned..... go to 19
17. Inflorescence an interrupted spicate panicle *Elytrophorus*
 Inflorescence an open or contracted panicle go to 18
18. Inflorescence a contracted panicle; spikelets with one or two basal bisexual flowers, with male or empty lemmas above them, the upper lemmas reduced to awns..... *Ectrosia*
 Inflorescence an open panicle, spikelets with two bisexual florets only *Eriachne*
19. Inflorescence a spike or raceme *Tripogonella*
 Inflorescence a once-branched open panicle go to 20
 Inflorescence a panicle with secondary branches..... go to 21

		
Inflorescence a spike or raceme	Inflorescence once branched	Inflorescence with secondary branches


20. Spikelets 1-6 mm long, with 2-9 florets, fertile florets 2.5-6 mm long *Dinebra decipiens*
Spikelets 5-18 mm long, with 4-14 florets, fertile florets 5-18 mm long *Diplachne fusca*
21. Spikelets with two bisexual florets *Eriachne*
Spikelets with three or more bisexual florets *Eragrostis*
22. Spikelets awned go to 23
Spikelets unawned go to 26
23. Inflorescence unbranched - a spike or raceme *Perotis*
Inflorescence branched - an open or contracted panicle go to 24
24. Glumes absent or rudimentary *Oryza*
Glumes well-developed go to 25

		
Glumes absent or rudimentary	Glumes well-developed	


25. Lemma with a 3-branched awn (very rarely 1-branched); spikelets with one floret *Aristida*
Lemma with a single awn; spikelets with 2 florets (very rarely 1) *Eriachne*
26. Inflorescence digitate go to 27
Inflorescence a spike or raceme or an open or contracted panicle go to 28

	
Inflorescence a digitate or subdigitate	Inflorescence a raceme, spike or panicle


27. Glumes shorter than floret *Cynodon*
 Glumes longer than floret *Brachyachne*




28. Inflorescence a single raceme; mat-forming plant used as an ornamental grass *Zoysia*
 Inflorescence an open or contracted panicle; tufted grass..... go to 29
29. Glumes absent or rudimentary *Leersia*
 Glumes well developed..... go to 30




30. Inflorescence a one-branched panicle *Dinebra neesii*
 Inflorescence a panicle with secondary branches, open or contracted..... *Sporobolus*



31. Inflorescence a raceme or spike (unbranched) go to 32
 Inflorescence with branches (panicle) go to 44



32. Leaves when crushed are lemon-scented; lower glume of sessile spikelet 2 toothed.....*Elionurus*
 Leaves when crushed, NOT lemon scented; lower glume of sessile spikelet entire or notched .. go to 33
33. Spikelets sessile distinctly awned..... go to 34
 All spikelets not distinctly awned..... go to 37
34. Awns more than 4.5 cm long, intertwined with other awns at maturity *Heteropogon*
 Awns less than 4.5 cm long, not intertwined with other awns at maturity go to 35
35. Annual; inflorescence usually subtended by a leaf-like bract; spikelets usually covered with long, silky white hairs*Schizachyrium*
 Perennial; inflorescence not subtended by a leaf-like bract; spikelets glabrous or covered with short white hairs go to 36
36. Lower glume of pedicelled spikelet slightly asymmetrical and strongly nerved, pedicels and internodes of the inflorescence densely bearded with white hairs..... *Sehima*
 Lower glume of pedicelled spikelet not asymmetrical and faintly nerved; pedicels and internodes of the inflorescence not bearded with white hairs*Dichanthium*
37. Creeping or prostrate grass go to 38
 Tufted erect or semi-erect grass go to 39
38. Raceme subtended by a spathe (leaf-like bract), growing on sandy beaches *Thuarea*
 Raceme not subtended by a spathe, growing on a range of soils.....*Stenotaphrum*
39. Raceme or spike with 2-6 spikelets; with a single cleistogamous spikelet in the axils *Cleistochloa*
 Raceme or spike with more than 6 spikelets, no axillary spikelets present..... go to 40




Axillary spikelets




Cleistogamous spikelet

40. Sessile spikelets with curved spines on the lower margin *Eremochloa*
 Spikelets with no spines on the margin go to 41
41. Raceme or spike not subtended by a leaf-like bract, spikelets fully exposed go to 42
 Raceme or spike partly or wholly enclosed by a leaf-like bract at the base go to 43
42. Spikelets solitary; inflorescence branches ending in a bristle..... *Paspalidium*
 Spikelets in pairs; inflorescence branch ending in a spikelet *Hemarthria*
43. Lower glume has prominent sculpturing *Hackelochloa*
 Lower glume smooth..... *Mnesithea*




44. Inflorescence a spatheate panicle (leaf-like bracts arranged throughout the panicle) go to 45
 Inflorescence not a spatheate panicle go to 51




45. Leaves lemon-scented when crushed *Cymbopogon*
 Leaves not lemon-scented when crushed go to 46
46. Fertile spikelets within an involucre of four male or barren spikelets..... *Themeda*
 Fertile spikelets without an involucre of spikelets..... go to 47
47. Sessile spikelet awned..... go to 48
 All spikelets awnless go to 49
48. Spikelets are covered with red or brown hairs *Hyparrhenia*
 Spikelets are covered with white hairs *Andropogon*
49. Spikelets in opposite neat rows on axis; spikelets often stick out at maturity *Ophiuros*
 Spikelets on one side of axis; spikelets erect at maturity go to 50
50. Lower glume has prominent sculpturing *Hackelochloa*
 Lower glume smooth..... *Mnesithea*


51. Inflorescence a digitate or subdigitate panicle go to 52
 Inflorescence an open or contracted panicle go to 60
52. Pedicelled spikelet distinctly awned (awn more than 3 mm long) go to 53
 All spikelets awnless or shortly awned (awn 3 mm or less long) go to 57
53. Racemes 2, appressed and interlocking and only separating at maturity; pedicels swollen ...*Ischaemum*
 Racemes 2 or more always separate; pedicels not swollen..... go to 54
54. Delicate annual; awns intertwining at maturity; all spikelets very small, 1.5-2.5 mm long
 *Pseudopogonatherum*
 Perennial; awns not intertwining at maturity; sessile spikelets usually more than 3 mm long ... go to 55
55. Pedicels of spikelets and glumes with silky brown hairs *Eulalia*
 Pedicels of spikelets and glumes glabrous or with white hairs..... go to 56
56. Rachis joints and pedicels with a translucent furrow between thickened margins *Bothriochloa*
 Rachis joints and pedicels without a translucent mid-line *Dichanthium*




57. Lower glume 40-70% of spikelet length; spikelet shortly awned.....*Alloteropsis*
 Lower glume less than 40% of spikelet length or absent; spikelet awnless go to 58




58. Lower floret adjacent to the inflorescence axis *Axonopus*
 Lower floret positioned away from the inflorescence axis go to 59




59. Palea of upper floret nearly completely clasped by lemma *Digitaria*
 Palea of upper floret not completely clasped by lemma *Paspalum*



60. Inflorescence spicate panicle, panicle branches short and often difficult to see go to 61
 Inflorescence open or contracted, panicle branches obvious, sometimes appressed to main axis go to 65




61. Spikelets covered in long, white hairs giving the inflorescence a fluffy appearance *Imperata*
 Spikelets not covered in long, white hairs; inflorescence not fluffy go to 62
62. Spikelets subtended by hairs, bristles or spines go to 63
 Spikelets not subtended by hairs, bristles or spines go to 64




63. Spikelets falling at maturity without any bristles or spines attached *Setaria*
 Spikelets falling at maturity with bristles or spines attached *Cenchrus*
64. Annual 10-60 cm tall; inflorescence 1-13 cm long; spikelets hump-backed *Sacciolepis*
 Aquatic perennial 50-350 cm tall; inflorescence 8-50 cm long; spikelets not hump-backed *Hymenachne*


65. Upper glume and lower lemma with hooked hairs *Ancistrachne*
 Upper glume and lower lemma without hooked hairs go to 66
66. Inflorescence a once-branched panicle go to 67
 Inflorescence with secondary branches go to 76




67. Inflorescence terminating in a bristle (inspect carefully since it looks similar to an awn) go to 68
 Inflorescence terminating in a spikelet go to 69




68. Upper lemma covered with wrinkles, as long as or longer than upper glume *Paspalidium*
 Upper lemma smooth, shorter than upper glume *Pseudoraphis*
69. Spikelet with a bead-like swelling at base (lower glume) *Eriochloa*
 Spikelet without a bead-like swelling at base go to 70







70. Spikelets distinctly awned go to 71
Spikelets awnless or mucronate..... go to 72
71. Glumes more or less equal and similar; decumbent grass of shaded habitats *Optismenus*
Glumes very unequal and dissimilar; erect grass of wet habitats *Echinochloa*
72. Ligule absent; glumes very unequal; palea tip reflexed *Echinochloa*
Ligule present..... go to 73
73. Palea of upper floret completely clasped by lemma (or nearly so) *Digitaria*
Palea of upper floret not completely clasped by lemma go to 74
74. Lower glume absent or very much reduced *Paspalum*
Lower glume present go to 75
75. Spikelets laterally compressed *Whiteochloa*
Spikelets dorsally compressed *Urochloa*



76. Inflorescence purple or red, fading to white; spikelets with awns arising from between apical lobes of upper glume and lower lemma, sometimes not obvious if spikelet covered in long hairs..... *Melinis*
Inflorescence various colours; spikelets awnless or with awn arising terminally from lemmas or glumes go to 77
77. Spikelets in pairs or triplets, sessile spikelet awned go to 78
Spikelets solitary or paired, awnless go to 83
78. Inflorescence with sweet, spicy smell when crushed; pedicels with a translucent furrow between thickened margins *Capillipedium*
Inflorescence without distinctive smell; pedicels without a translucent furrow go to 79







79. Spikelets laterally compressed go to 80
Spikelets dorsally compressed go to 81

 		
Spikelets laterally compressed (glumes and lemmas folded or rounded)	Spikelets dorsally compressed (glumes and lemmas flat)	





80. Spikelets all alike; spikelets breaking above glumes *Arundinella*
Pedicelled spikelets different to sessile spikelets; spikelets breaking up in spikelet pairs or triplets
..... *Chrysopogon*
81. Pedicelled spikelets reduced to narrow, linear glumes *Vacoparis*
Pedicelled spikelets usually well-developed go to 82
82. Lemma awns of sessile spikelet absent or 1-1.5 cm long if present *Sorghum*
Lemma awns of sessile spikelet 2-9 cm long *Sarga*
83. Spikelets dorsally compressed go to 84
Spikelets laterally compressed go to 86
84. Panicle contracted, spikelets very closely packed *Sorghum*
Panicle open, spikelets widely spaced go to 85
85. Fertile lemma smooth *Panicum*
Fertile lemma rugose (wrinkled) *Megathyrsus*
86. Robust perennial 1.5-2 m tall; inflorescence purplish (Vetiver Grass) *Chrysopogon*
Annual or weak perennial 15-110 cm tall; inflorescence green go to 87
87. Plant 55-110 cm tall; grows on sandy, alluvial soils; inflorescence open or contracted, branches
persistent *Whiteochloa*
Plant 17-60 cm; grows on rocky hillsides; inflorescence open; branches deciduous stalk
..... *Arthrargrostis*

Key to species with thickened rachis and a single raceme or spike, not awned

1. Spikelets with 1 to many florets, if 2-flowered both florets or lower one bisexual *Lepturus repens*
 Spikelets 2-flowered, lower floret male or barren, upper bisexual or female 2
2. Spikelets often dissimilar and usually paired, with one sessile and the other pedicelled (occasionally both pedicelled, but unequally, or in 3's with 2 sessile and 1 pedicelled) or rarely with the pedicelled spikelet reduced to the pedicel; upper lemma of sessile spikelet usually geniculate awned; glumes firm in texture 3
 Spikelets all alike and usually solitary, rarely paired; upper lemma not geniculate awned; glumes usually membranous 6
3. Lower glume of sessile spikelet 2-toothed *Elionurus citreus*
 Lower glume of sessile spikelet not 2-toothed 4

Lepturus repens	Elionurus citreus
	
	

4. Spikelets of each pair similar, both bisexual *Hemarthria uncinata*
 Spikelets of each pair dissimilar, pedicelled male or sterile or spikelets borne singly *Eremochloa bimaculata*

Hemarthria uncinata	Eremochloa bimaculata
	
	


Subfamilies and Tribes of grasses in Townsville

There are 13 subfamilies in Poaceae. The grasses found in Townsville are represented in 6 subfamilies, the largest of which is the subfamily Panicoideae. The spikelet drawings refer to species in Townsville.

Ehrhartoideae

Tribe Oryzeae

1. Spikelets without glumes or glumes rudimentary.
 2. Spikelets one-flowered or three-flowered with the two lower florets reduced to sterile lemmas
 3. Aquatic or wetland grasses
 4. Disarticulation at the base of the lemmas
- Townsville Genera: *Leersia* (A=6 or 1-3), *Oryza* (A=6)




Arundinoideae

1. Common members of these taxa are robust, "reed-like" grasses with plumose panicles
2. Stems usually hollow
3. Disarticulation above the glumes and between the florets

Tribe Arundineae


Townsville Genera: *Arundo*, *Phragmites*



Aristidoideae

Tribe Aristideae


1. Spikelets with a single floret
 2. Lemma with a 3-branched awn at the apex (though the 2 lateral awns may be reduced)
 3. Callus of lemma well developed
 4. Ligule a fringed-membrane or a fringe of hairs
 5. Disarticulation above the glumes
- Townsville Genera: *Aristida*



Danthonioideae

1. Spikelets with multiple florets
2. Lemmas bifid at apex, an awn emerging from between the lobes
3. Glume usually \pm equal and as long as column of florets
4. Ligule of hairs
5. Disarticulation above the glumes and between the florets

Townsville Genera: *Elytrophorus*




Chloridoideae

Tribe Pappophoreae

1. Lemmas with 5-13 veins, all of which extend into apical awns
2. Ligule a fringe of hairs
3. Spikelets with 3 or more florets
4. Disarticulation above the glumes but not between the florets


Townsville Genera: *Enneapogon*



Tribe Triodieae

1. Spikelets with one or more florets
2. Lemmas rounded on back, with 3-9 nerves
3. Leaf blades rigid, needle-like


Townsville Genera: *Triodia*



Tribe Eriachneae

1. Spikelets usually with two bisexual florets, awned or unawned





Townsville Genera: *Eriachne*











Tribe Cynodonteae





Most members of the tribe possess 2 or more of the following characteristics:


1. Laterally compressed spikelets
 2. Lemmas with 1-3 veins or 7-13 veins
 3. Spike-like branches of the inflorescence
 4. Coarse hairs near the junction of the sheath and blade
 5. Disarticulation variable (e.g. beneath fertile florets, beneath the glumes, at base of branches)
- Townsville Genera: *Brachyachne*, *Chloris*, *Cynodon*, *Dactyloctenium*, *Dinebra*, *Diplachne*, *Ectrosia*, *Eleusine*, *Enteropogon*, *Eragrostis*, *Lepturus*, *Oxychloris*, *Perotis*, *Sporobolus*, *Tripogonella*, *Zoysia*.

Cynodonteae spikelets			
 <p>Floret Glumes</p>	 <p>Floret Glumes</p>	 <p>Grain Fertile floret Glumes</p>	 <p>Floret Upper glume</p>
<i>Brachyachne</i>	<i>Cynodon</i>	<i>Sporobolus</i>	<i>Zoysia</i>

 <p>Awns Glumes</p>	 <p>Floret Upper glume</p>	 <p>Florets Glumes</p>	 <p>Fertile floret Glumes</p>
<i>Perotis</i>	<i>Lepturus</i>	<i>Dinebra</i>	

 <p>Florets Glume</p>	 <p>Florets Glumes</p>	 <p>Florets Glumes</p>	 <p>Florets Glumes</p>
<i>Diplachne</i>	<i>Eleusine</i>	<i>Eragrostis</i>	<i>Tripogonella</i>

 <p>Awns Florets Glumes</p>	 <p>Awns Florets</p>	 <p>Awns Florets Glumes</p>	 <p>Florets Glumes</p>
<i>Chloris</i>		<i>Oxychloris</i>	<i>Enteropogon</i>



Panicoideae


The special feature of this subfamily is the two-flowered spikelet with the lower floret male or barren, and the upper floret bisexual or female.

Of the 74 genera found in Townsville, 49 belong in this subfamily. The three tribes represented are Arundinelleae 1 genus, Paniceae 24 genera and Andropogoneae 24 genera.

Tribe Arundinelleae


1. Spikelets 2-flowered, the lower floret sterile or male
2. Spikelets usually not paired, all alike
3. Disarticulation above the glumes, persistent
4. Glumes usually unequal, the upper exceeding the florets
5. Upper floret awned from sinus (a notch or depression in the apex)
6. Inflorescence a panicle (branched)

Townsville Genera: *Arundinella*



















Tribe Paniceae




1. Spikelets 2-flowered, the lower floret sterile or male
2. Spikelets usually all alike
3. Disarticulation below the glumes
4. The lower glume is usually small or absent
5. The upper glume usually resembles the lemma of the lower floret
6. Upper lemma and palea of spikelet usually indurate (hardened)
7. Inflorescence usually a panicle (branched)



Townsville Genera: *Alloteropsis*, *Ancistrachne*, *Arthrargrostis*, *Axonopus*, *Cenchrus* (*Pennisetum*), *Cleistochloa*, *Digitaria*, *Echinochloa*, *Eriochloa*, *Hymenachne*, *Megathyrsus*, *Melinis*, *Oplismenus*, *Panicum*, *Paspalidium*, *Paspalum*, *Pseudoraphis*, *Sacciolepis*, *Setaria*, *Spinifex*, *Stenotaphrum*, *Thuarea*, *Urochloa*, *Whiteochloa*.




Panicoideae spikelets		
 <p>Awn Lower floret Upper glume Lower glume</p> <p><i>Alloteropsis</i></p>	 <p>Lower floret Upper glume Lower glume</p> <p><i>Ancistrachne</i></p>	 <p>Upper glume Lower floret Lower glume Stalk Fertile floret</p> <p><i>Arthragrostis</i></p>
 <p>Lower floret Upper glume Rachis</p> <p><i>Axonopus</i></p>	 <p>Spikelets</p> <p><i>Cenchrus</i></p>	 <p>Spines Spikelets</p>
 <p>Lower floret Upper floret Lower glume Upper glume</p> <p><i>Cleistochloa</i></p>	 <p>Lower lemma Lower glume Lemma Palea</p> <p><i>Digitaria</i></p>	 <p>Lower floret Upper glume Lower floret Lower glume Upper glume</p> <p><i>Echinochloa</i></p>
 <p>Upper glume Lower floret Lower glume</p> <p><i>Eriochloa</i></p>	 <p>Upper floret Lower floret Upper glume Lower glume</p> <p><i>Hymenachne</i></p>	 <p>Upper glume Lower floret Lower glume</p> <p><i>Megathyrsus</i></p>






 <p>Male florets Female florets</p>	 <p>Upper glume Lower floret Lower glume</p>	 <p>Upper floret Lower floret Upper glume Lower glume</p>
<i>Thuarea</i>	<i>Urochloa</i>	<i>Whiteochloa</i>





Tribe Andropogoneae





1. Spikelets 2-flowered, the lower floret sterile or male
2. Spikelets usually paired, one sessile the other pedicelled, usually dissimilar
3. Disarticulation usually in the branch axes beneath the sessile or short-pedicelled spikelet
4. Both glumes usually concealing the florets
5. Upper lemma and palea of spikelet usually of weak texture
6. Inflorescence a collection of rames (a series of paired spikelets, one sessile and one pedicellate)

Andropogoneae spikelets		
 <p>Awn Pedicelled spikelet Sessile spikelet Pedicel</p>	 <p>Lower glume Upper glume Lower lemma Upper lemma</p> <p>Pedicelled spikelet (this example does not have any paleas)</p>	 <p>Awn Lower glume Upper glume Lower lemma Upper lemma palea</p> <p>Sessile spikelet (this example does not have a lower palea)</p>
<p>Spikelet pair, many of the genera have a triplet of spikelets (2 pedicelled and 1 sessile) at the end of the branches</p>	<p>Drawings from Tothill & Hacker (1983)</p>	

Townsville Genera: *Andropogon*, *Bothriochloa*, *Capillipedium*, *Chionachne*, *Chrysopogon*, *Cymbopogon*, *Dichanthium*, *Elionurus*, *Eremochloa*, *Eulalia*, *Hackelochloa*, *Hemarthria*, *Heteropogon*, *Hyparrhenia*, *Imperata*, *Ischaemum*, *Mnesithea*, *Ophiuros*, *Pseudopogonatherum*, *Sarga*, *Schizachyrium*, *Sehima*, *Sorghum*, *Themeda*, *Vacoparis*.

Andropogoneae spikelets		
 <p>Pedicelled spikelet Awns Sessile spikelet</p>	 <p>Pedicelled spikelet Awn Pedicel Sessile spikelet</p>	 <p>Awn Pedicelled spikelets Sessile spikelet</p>
<i>Andropogon</i>	<i>Bothriochloa</i>	<i>Capillipedium</i>





Grass Genera and species descriptions

Alloteropsis


Cockatoo Grasses

Alloteropsis is from the Greek *allotrios* (belonging to another) and *opsis* (appearance), the spikelets and inflorescences somewhat resemble another genus of grass.

Tufted perennials or annuals, culms erect or decumbent. The inflorescence is digitate, and the spikelets are usually paired and usually awned. The glumes are unequal, the lower glume 50-75% of the spikelet, the upper glume as long as the spikelet and densely hairy along the marginal nerves. The lower glume is shorter than the spikelet.

Subfamily: Panicoideae; Tribe: Paniceae

World = 8, Australia = 2



Townsville species

Alloteropsis cimicina

Annual Cockatoo Grass

Alloteropsis semialata

Cockatoo Grass

Key to the species of *Alloteropsis*

Tussocky, perennial; spikelets 4-7.5 mm long *Alloteropsis semialata*
Annual, 1-3 culms; spikelets 3.5-5.5 mm long *Alloteropsis cimicina*

***Alloteropsis cimicina* - Annual Cockatoo Grass**

Derivation

cimicina - from the Latin *cimex* (bug) and *-ina* (resemblance), mature spikelets bear a fanciful resemblance to a small bug.

Habit

An annual, erect grass, the culms are 15-100 cm tall.

Inflorescence

The panicle digitate with 4-11 racemes, 5-10 cm long, and the spikelets are 3.5-5.5 mm long. The raceme rachis is glabrous.

Habitat

This species occurs on woodlands and grasslands.



Spikelets



***Alloteropsis semialata* - Cockatoo Grass**

Derivation

semialata - from the Latin *semi* (half) and *ala* (wing) and *-ata* (possessing), refers to the winged margins of the upper glume.

Habit

A perennial, tufted grass, the culms are 20-150 cm tall.

Inflorescence

The panicle digitate with 2-4(28) racemes, 2-22 cm long, and the spikelets are 4-7.5 mm long. The raceme rachis is pilose.

Habitat

This species occurs on woodlands and grasslands.



Spikelet




Ancistrachne

From Greek *ankistron* (fishhook) and *achne* (glume), referring to the often-hooked hairs on the spikelet.

Tufted or scrambling perennial with wiry culms, leaves cauline (growing on stems) and usually less than 10 cm long. The inflorescence is a depauperate panicle or a single raceme. The spikelets (upper glume and lower lemma) are covered with hooked or curved spines or hairs.

Subfamily: Panicoideae; Tribe: Paniceae
World = 4, Australia = 2



Spikelets

Townsville species

Ancistrachne uncinulata Hooky Grass



***Ancistrachne uncinulata* - Hooky Grass**

Derivation

uncinulata - from the Latin *uncinulus* (small hook) and *-ata* (possessing), referring to the hooked hairs on the upper glume and lower lemma.

Habit

A perennial, shrubby, erect or spreading grass, the culms is 30-200 cm tall.

Inflorescence

Inflorescence is a narrow panicle 2-14 cm long with few spikelets. The rigid, hooked spines on the spikelets make it a distinctive species.

Habitat

This species usually grows on sandstone soils.



Andropogon

From the Greek words *aner* (man) and *pogon* (beard), alluding to the awns or to the long hairs on the raceme internodes and pedicels.


Tufted perennials or annuals. The inflorescence of spicate main branches or paniculate, usually with paired or digitate racemes. The spikelets are in pairs, one sessile and the other pedicelled.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 534, Australia = 4

Townsville species

* *Andropogon gayanus* Gamba Grass

It could be confused with Grader Grass (*Themeda quadrivalvis*), Giant Spear Grass (*Heteropogon triticeus*) and Thatch Grass (*Hyparrhenia rufa* subsp. *rufa*).



Key to the species of *Andropogon* and similar genera

1. Inflorescence is a raceme *Heteropogon triticeus*
.....
Inflorescence with branches 2
2. Fertile spikelets within an involucre of four male or barren spikelets *Themeda quadrivalvis*
Fertile spikelets without an involucre of spikelets 3
3. Spikelets covered with red or brown hairs *Hyparrhenia rufa* subsp. *rufa*
Spikelets covered with white hairs *Andropogon gayanus*



Spikelets covered
with brown or red hairs
Hyparrhenia rufa subsp. *rufa*



Spikelets covered
with white hairs
Andropogon gayanus

***Andropogon gayanus* - Gamba Grass**

Derivation

gayanus in honor of Claude Gay (1800-1873) French and - *ana*, indicating connection.

Habit

A perennial, tufted grass, the culms are 150-400 cm tall.

Inflorescence

The panicle is subtended by a spatheole. The rames are paired and are 4-9 cm long. The sessile spikelet is 5-8 mm long. Both sessile and pedicelled spikelets are often with awns.

Habitat

This non-native species was imported into Queensland as a pasture grass, it is now a weed, it is often spread by vehicles and machinery along roads. It grows mainly along roadsides and in disturbed areas.



Spikelets



Aristida


Wiregrasses or Kerosene or Three Awn Grasses

From Latin *arista* (awn or beard of a grain), alluding to the awned lemma.

Tufted, annuals or perennials, usually with slender wiry stems. The inflorescence is a contracted or open panicle. The spikelets solitary. The genus is easily recognised by the lemma awns which are usually 3-branched. The callus (the hard, usually pointed base of the spikelet) is usually sharp and can become embedded into clothing. The glumes remain on the inflorescence after the mature seed falls.

Australian species of *Aristida* usually grow on poor sandy and red soils, although there are a few that prefer black soils and cracking clays.

Subfamily: Aristidoideae; Tribe: Aristideae
World = 395, Australia = 68




Townsville species

Aristida acuta
Aristida calycina
Aristida gracilipes
Aristida holathera
Aristida jerichoensis var. *subspinulifera*
Aristida latifolia
Aristida pernicioso
Aristida queenslandica
Aristida spuria
Aristida superpendens
Aristida utilis
Aristida warburgii

Dark Wiregrass, Branched Wiregrass

Erect Kerosene Grass
 Jericho Wiregrass, Feathertop Threeawn
 Feathertop Wiregrass




Queensland Wiregrass






Parts of *Aristida* spikelet
 Some species does not have a column

Identifying *Aristida* species usually requires looking at the spikelets under magnification. The involute or convolute condition of the lemma, the presence or absence of a lemma awn column and the lengths of the lateral awn branches compared to the median awn branch are diagnostic features of *Aristida* species.



Aristida can be easily identify by the single floret and mostly with 3 awns. They might be confused with some *Chloris* spikelets and with *Arundinella setosa*.

Spikelet with a single floret	Spikelets with 3 or more florets	Spikelets with 2 florets, the lower one sterile or male
Subfamily Aristoideae	Subfamily Chloridoidea	Subfamily Panicoideae
<i>Aristida</i>	<i>Chloris</i>	<i>Arundinella setosa</i>
		

Lateral lemma awns			
Equal or subequal to median awn		Shorter than median awn 1/3 to 2/3 as long as median awn	Absent or very short
			
<i>Aristida acuta</i> <i>Aristida calycina</i> <i>Aristida holathera</i> <i>Aristida jerichoensis</i>		<i>Aristida gracilipes</i> <i>Aristida queenslandica</i>	<i>Aristida spuria</i> <i>Aristida utilis</i>
<i>Aristida latifolia</i> <i>Aristida perniciosa</i> <i>Aristida superpendens</i> <i>Aristida warburgii</i>			

Key to the species of *Aristida*

1. Lateral awns less than 1/3 as long as median awn or absent2
Lateral awns as long as median awn or more than 1/3 its length3
2. Lower internodes glabrous; leaf blades setaceous *Aristida spuria*
Lower internodes hairy; leaf blades flat..... *Aristida utilis*
3. Lemma involute, with a furrow on ventral surface of caryopsis4
Lemma convolute, without a furrow on ventral surface of caryopsis9

Involute lemma: with the margins rolled inwards on the upper surface but not overlapping, and with a furrow on the ventral surface of the caryopsis	Convolute lemma: rolled longitudinally with one edge inside the other
	

4. Lemma column present *Aristida perniciososa*
Lemma column absent5
5. Awns unequal, laterals 1/2-2/3 as long as median6
Awns \pm equal or laterals more than 2/3 as long as median.....7
6. Culm internodes with pilose hairs, especially at base.....*Aristida queenslandica* var. *queenslandica*
Culm internodes glabrous*Aristida queenslandica* var. *dissimilis*
7. Margins of lemma furrow with one to several rows of tubercles
..... *Aristida jerichoensis* var. *subspinulifera*
Margins of lemma furrow smooth or lemma muricate towards apex.....8
8. Glumes acuminate to awned; inflorescence branches usually naked in lower 1/2 *Aristida acuta*
Glumes acute to obtuse; inflorescence branches usually bearing spikelets from base
.....*Aristida calycina*
9. Lemma without a column *Aristida gracilipes*
Lemma with a twisted column (sometimes reduced to a 1/2 spiral) 10
10. Lower glume 3-7-nerved *Aristida warburgii*
Lower glume 1-nerved; 11
11. Column 1-9 mm long *Aristida latifolia*
Column 11-60 mm long 12
12. Awns \pm similar and erect, laterals as long as median or shorter by 2-5 mm*Aristida holathera*
Awns dissimilar, median thicker and usually recurved, laterals slender usually erect and up to 2 cm
shorter *Aristida superpendens*

Aristida acuta

Derivation

acuta - from the Latin *acuo* (sharpen), alluding to the sharply pointed glumes.

Habit


A perennial grass with culms 40-125 cm tall.

Inflorescence

The panicle is a linear or elliptic, 15-30 cm long, 1-10 cm wide. The lemma margin is involute, and the awns do not have a column. The lateral awns are subequal (about 70-85%) to the median awn. The glumes are acuminate and awned.

Habitat

It is found in *Eucalyptus* woodlands and forests on poor soil.



Spikelet



***Aristida calycina* - Feathertop Dark Wiregrass**

Derivation

calycina - from the Greek *kalyx* (cup) and *-ina* (belonging to), the subtending glumes are as long as or longer than the lemma thereby resembling a cup.

Habit



A loosely tufted perennial grass, culms 70-150 cm tall, branched.

Inflorescence

The panicle is open or contracted, linear to elliptic, 12-31.5 cm long, 2-20 cm wide. The spikelets are purple or brown. The lemma margin is involute, and the awns do not have a column. The lateral awns are equal to subequal (about 75-100%) to the median awn. Glumes are acute to obtuse.

Habitat

It grows on red earths, sands and alluvial soils.



Aristida gracilipes

Derivation

gracilipes - from the Latin *gracilis* (slender) and *pes* (foot), alluding to the fine and bushy habit.

Habit

Loosely tufted perennial with culms 60-115 cm tall. It has many branches giving a bushy appearance. It has narrow leaves, 0.5 mm wide.

Inflorescence

The panicle is open to linear, 12-21 cm long, 1-3 cm wide. The lemma margin is convolute, and the awns do not have a column. The lateral awns are unequal (about 60-75%) to the median awn.

Habitat

This species grows in *Acacia*, brigalow, *Eucalyptus* communities, and rainforest fringes, on clays, loams and sandy soils.



Spikelet



***Aristida holathera* - Erect Kerosene Grass**

Derivation

holathera - from Greek *holos*, wholly; *ather*, spike or ear of wheat; awn not disarticulating from base of lemma at maturity.

Habit



An annual or perennial grass, with culms 28-128 cm tall.

Inflorescence

The panicle is open and elliptic, 8-41 cm long, 1-10 cm wide. The lemma margin is convolute, and the awns have a twisted distinct column (15-60 mm long). The lemma has a distinct articulation. The lateral awns are equal to subequal (about 75-100%) to the median awn.

Habitat

It grows on sandy soils.



***Aristida jerichoensis* var. *subspinulifera* - Jericho Wiregrass; Feathertop Threeawn**

Derivation

jerichoensis - denoting origin, from Jericho.

var. *subspinulifera* - subspecies with small spines (within lemma).

Habit


A perennial, tufted grass, the culms are 30-90 cm tall. Leaves and stems are glaucous (of a dull greyish-green or blue colour).

Inflorescence

The panicle is linear or lanceolate, 4-25 cm long, 1-2 cm wide. The lemma margin is involute without a column. The lateral awns are subequal (about 85-90%) to the median awn. The margins of lemma furrow with one to several rows of tubercles.

Habitat

This species grows on red earths or clays.



Spikelet



***Aristida latifolia* - Feathertop Wiregrass**

Derivation

latifolia - from Latin *latus* (broad) and *folium* (leaf), leaf blades broad or relatively broad compared to related species.

Habit



A loosely tufted perennial grass, culms 20-130 cm tall.

Inflorescence

The panicle is contracted and linear, 8-70 cm long 3-3.5 cm wide. The lemma margin is convolute and the awns with 1-9 spirals in the column 1-9 mm long. The lateral awns are equal to subequal (about 85-100%) to the median awn.

Habitat

Grows on clays, red earths, sands and alluvial soils.



Aristida pernicioso

Derivation

pernicioso - from the Latin *per* (very), *noxius* (harmful) and *-osa* (abundance); callus very sharp and readily entangling in wool and clothing.

Habit

Compactly tufted coarse perennial grass with culms 60-140 cm tall.

Inflorescence


The panicle is contracted or spiciform, linear, 15-39 cm long, 1-1.5 cm wide. The lower glume is 5-nerved. The lemma margin is involute, and the awns do not have a column. The lateral awns are equal to subequal (about 75-100%) to the median awn.

Habitat

Occurs in *Eucalyptus* and *Melaleuca* communities on sands and loams.



Spikelet



Aristida queenslandica - Queensland Wiregrass

Derivation

queenslandica - named for the distribution of the species, found mainly in Queensland.

Habit


A perennial, tufted grass with culms erect, 25-109 cm tall.

Inflorescence

The panicle is open and elliptic or loose, 7-64 cm long, 2-9 cm wide. The lemma margin is involute, and the awns do not have a column. The lateral awns are distinctly unequal (about 30-65%) to the median awn.

Habitat

Usually grows on *Eucalyptus* communities on rocky areas and sandstone and granite hills.



Spikelet

Two varieties are recognised

Culm internodes with pilose hairs, especially at base..... var. *queenslandica*
Culm internode glabrous var. *dissimilis*



Aristida spuria

Derivation

spuria - from the Latin *spurius* (false), alluding to the lateral awns being very short or absent.

Habit


A perennial grass with culms decumbent, 20-91 cm tall. It is similar to *Aristida utilis*, but lower internodes glabrous and the leaf blades are involute.

Inflorescence

The panicle is contracted and linear, 11-29 cm long and 0.9-1 cm wide. The lemma margin is involute and the awns with a twisted column. The lateral lemma awns absent or present, 0-4 mm long, shorter than median awn, 0-20% of length of the median awn.

Habitat

Usually grows on *Eucalyptus* woodlands, sometimes in elevated areas and offshore islands. Sandy and loamy soils.



Spikelet



Floret and awn

Aristida superpendens

Derivation

superpendens - from the Latin *super* (above) and *pendeo* (hang), derived from the pendulous spikelets in the upper part of the inflorescence.

Habit


A perennial, tufted grass with culms erect, 60-120 cm tall.

Inflorescence

The panicle is open and elliptic, 14-34 cm long, 5-8 cm wide. The lemma margin is convolute and the awns with a distinct column. The lemma has a distinct articulation. The lateral awns are equal to subequal (about 75-100%)

Habitat

Grows in *Eucalyptus* and *Melaleuca* woodlands on granite sand and alluvial soils.



Spikelet



Aristida utilis

Derivation

utilis - from Latin *utilis* (useful). Peduncles used for the manufacture of hats.

Habit


A perennial and tufted grass with culms 53-130 cm tall. It is similar to *Aristida spuria*, but lower internodes hairy and the leaf blades are flat.

Inflorescence

The panicle is open and lanceolate, 15-48 cm long, 1-6 cm wide. The lemma margin is involute and the awns with a column. The lateral lemma awns absent or present, 1-7 mm long, shorter than median awn, 0-40% of the length of principal.

Habitat

Grows in *Eucalyptus* woodlands in granite and sand country.



Spikelet



Floret and awn



Aristida warburgii

Derivation

warburgii - in honour of Otto Warburg (1859-1938), German botanist.

Habit


It is perennial and tufted; the culms are 30-90 cm tall.

Inflorescence

The panicle is open and elliptic, 11-15 cm long and 2-7 cm wide. The lower glume has 3-7 nerves. The lemma is convolute and the awns with a distinct column. The lemmas awns are equal to subequal (about 70-100%), mature spikelets with central awn strongly recurved and thicker than lateral awn.

Habitat

Grows in *Eucalyptus* and *Melaleuca* communities on sandy soils.



Spikelet



Arthrargrostis

From Greek *arthron* (joint) and *agrostis* (grass), alluding to the disarticulation of the panicle into component parts.

Arthrargrostis was segregated from the genus *Panicum* on the basis of several distinct morphological features. One of its diagnostic characters is the stalked fertile floret.


Tufted or decumbent annual grasses of open habitats. The inflorescence is an open panicle and spikelets solitary.

Subfamily: Panicoideae; Tribe: Paniceae.
World = 5, Australia = 5

This species could be confused with *Panicum* and *Whiteochloa*.

Townsville species


Arthrargrostis deschampsoides




Spikelets

Key to the species of *Arthrargrostis* and related genera

1. Spikelets dorsally compressed.....*Panicum*
- Spikelets laterally compressed2



Spikelets dorsally compressed



Spikelets laterally compressed

2. Plant 55-110 cm tall; grows on sandy, alluvial soils; inflorescence open or contracted, branches persistent *Whiteochloa airoides*
- Plant 17-60 cm; grows on rocky hillsides; inflorescence open; branches deciduous *Arthrargrostis deschampsoides*

Arthrargrostis deschampsioides

Derivation

deschampsioides - name alluding to the similarity to the genus *Deschampsia*.

Habit

A slender, annual grass, the culms are 17-60 cm tall.

Inflorescence

The panicle is open, 5-25 cm long and 1-5 cm wide. The spikelet is laterally compressed; the fertile floret is dorsally compressed and borne on a slender stalk.

Habitat

This species grows on rocky hillsides e.g. Castle Hill and Many Peaks Range.



Spikelet with glumes removed



Arundinella

Reed Grasses

From the Latin *arundo* (a reed) and *-ella* (diminutive suffix).

Tufted annuals and perennials, usually with erect culms, and usually growing in marshy places and along riverbanks. The inflorescence is an open or contracted panicle. The spikelets are solitary or paired, and all are alike. The upper fertile lemma is geniculately (bent like a knee) awned.



Subfamily: Panicoideae; Tribe: Arundinelleae
World = 69, Australia = 4

Townsville species

<i>Arundinella nepalensis</i>	Reed Grass
<i>Arundinella setosa</i>	Reed Grass

Key to the species of *Arundinella*

Grows in rocky ground, dry hills; fertile lemma 3-awned
..... *Arundinella setosa*
Grows along creek beds and swamps; fertile lemma 1-awned
..... *Arundinella nepalensis*



Arundinella setosa



Arundinella nepalensis



Arundinella setosa (left),
Arundinella nepalensis (right)

***Arundinella nepalensis* - Reed Grass**

Derivation

nepalensis - from the Latin *-ense*, denoting origin, from Nepal.

Habit

A perennial, erect grass, the culms are 60-180 cm tall. The ligule is an eciliate membrane.

Inflorescence

The panicle is compound, open; it is 5-40 cm long. The spikelet is 4-7 mm long. Lower glume is muticous; the lemma is 1-awned.

Habitat

This species grows along creek beds and swamps, in Eucalypt forest, Melaleuca swamp, and various types of woodland and in grassland.



Spikelet



***Arundinella setosa* - Reed Grass**

Derivation

setosa - from the Latin *seta* (bristle) and *-osa* (abundance), each raceme subtended by a stout bristle.

Habit


A perennial, tufted grass, the culms are 50-120 cm tall. The ligule is a ciliate membrane.

Inflorescence

The panicle is compound, open; it is 5-30 cm long. The spikelet is 5-7 mm long. The lower glume is awned; the lemma is 3-awned.

Habitat

This species grows in rocky ground, dry hills, in open forests, forest margins and grasslands.



Fertile floret showing three awns



Arundo

Reed Grasses


From Latin *arundo* (reed).

Tall perennials with thick, knotty rhizomes. Culms woody and persistent, 2 to 8 m tall. It grows in very large clumps and resembles bamboo. The inflorescence is an open panicle. The spikelets are solitary, and the lemmas covered with long, soft hairs.

Subfamily: Arundinoideae; Tribe: Arundineae
World = 103, Australia = 1

Townsville species


* *Arundo donax* Giant Reed



Key to the species of *Arundo* and similar species

Lemmas hairy; rachilla not hairy *Arundo donax*

Lemmas not hairy (glabrous); rachilla (axis of spikelet) hairy *Phragmites*



***Arundo donax* - Giant Reed**

Derivation

donax - the Greek word for a type of reed in classical literature, alluding to the easy dispersal of diaspores by the wind.

Habit

A large, tufted perennial, bamboo-like, 2-6 m tall, usually unbranched, with leaves arising along the stem, which is erect or bent over. Closely related and similar to *Phragmites*.

Inflorescence

The inflorescence is a large feathery panicle, 30-60 cm long and 12 cm wide. The spikelets are solitary and comprising of numerous (2-7) florets; the rachilla is glabrous and the lemmas are covered with long silky hairs.

Habitat

This non-native species reproduces vegetatively from rhizomes (underground stems) and can invade nearby bushland. It is now recognised as a weed in most states in Australia; it grows on vacant land, margins of rubbish tips or on roadsides where garden refuse has been tipped.



Axonopus

Carpet Grasses

From the Greek *axon* (axis) and *pous* (foot), alluding to the racemes arising from a common point (digitate).

Stoloniferous or tufted perennials. The inflorescence is digitate or sub-digitate. The spikelets are solitary, and on one side of the rachis (the axis or branch of the inflorescence). The upper glume is abaxial (the side facing away from the rachis). The lower glume is absent or obscure.

Axonopus species are similar to some species of *Paspalum*, however this genus has the upper glume adaxial (the side facing to the rachis).

Subfamily: Panicoideae; Tribe: Paniceae
World = 99, Australia = 2



Townsville species

* *Axonopus compressus* Buffalo Grass, Broadleaf Carpet Grass



Key to the species of *Axonopus* and related genera

1. Lower floret adjacent to the inflorescence axis *Axonopus*
Lower floret positioned away from the inflorescence axis *Paspalum*



***Axonopus compressus* - Buffalo Grass, Broadleaf Carpet Grass**

Derivation

compressus - from the Latin *comprimo* (squeeze together), referring to the flattened culms.

Habit

A perennial, stoloniferous, mat-forming grass, the culms are 15-60 cm tall.

Inflorescence

The panicle is digitate or subdigitate, the spikes are 2-3(-5), 3-10 cm long. The spikelets are 2-2.8 mm long; the lower glume is absent or obscure.

Habitat

This non-native species is used as a popular lawn grass in Townsville.



Spikelet



Bothriochloa



Bluegrasses

From the Greek *bothros* (trench or pit), and *chloa*, (grass), alluding either to the groove in the pedicels or to the pit in the lower glumes of some species.

Tufted, decumbent or stoloniferous perennials, often with branched culms. The inflorescence is a digitate or subdigitate, or an open panicle; and usually has a spicy smell when crushed. The spikelets are in pairs (with terminal triplets), one sessile and one pedicelled. The lower glume of the sessile spikelet of some species has a pit (circular depression). The sessile spikelet is awned and bisexual; the pedicelled spikelet can be bisexual, male or sterile.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 56, Australia = 11

Bothriochloa is closely related to *Dichanthium*, the difference between these two genera requires careful dissection under a microscope. The pedicels of *Bothriochloa* species have a longitudinal translucent furrow, often purple coloured; in cross section the pedicels are dumbbell shaped. The pedicels of *Dichanthium* species are rounded.



Bothriochloa bladhii subsp. *bladhii* is closely related and similar to the genus *Capillipedium*.

Townsville species

<i>Bothriochloa bladhii</i> subsp. <i>bladhii</i>	Forest Bluegrass
<i>Bothriochloa decipiens</i>	Pitted Grass
<i>Bothriochloa ewartiana</i>	Desert Bluegrass
* <i>Bothriochloa pertusa</i>	Indian Bluegrass

Key to the species of *Bothriochloa*

These species are covered in the *Dichanthium* genus.



***Bothriochloa bladhii* subsp. *bladhii* - Forest Bluegrass**

Derivation

bladhii - named after Pehr Johann Bladh who collected in China and South Africa.

Habit

A perennial, tufted grass, the culms are erect or geniculately ascending, 50-150 cm tall.

Inflorescence

The inflorescence is compound; it is commonly purple-coloured. The panicle is 3-14 cm long. It is different than all other of *Bothriochloa* which normally have digitate panicle. Does not have a pitted lower glume.

Habitat

An important and widespread grass of *Eucalyptus* forests and woodlands. It is usually in alluvial soils and is the vicinity of major creeks.



***Bothriochloa decipiens* - Pitted Grass**

Derivation

decipiens - from the Latin word for resembling another species.

Habit

A perennial, tufted grass, culms erect or geniculately ascending, 30-200 cm tall.

Inflorescence

The inflorescence digitate to subdigitate, with ramose branches, the racemes 3-5, 4-7 cm long. The sessile spikelet is awned and with a pit. There are only two species in Townsville which have a pit in the glumes, these are *B. decipiens* and *B. pertusa*.

Habitat

Mostly found on drier soils or hillslopes, a widespread grass in open forest country.



Spikelets



***Bothriochloa ewartiana* - Desert Bluegrass**

Derivation

ewartiana - after the English-born Australian botanist, Albert James Ewart (1872-1937).

Habit

A perennial, tufted grass, culms erect or geniculately ascending, 30-60 cm tall. The leaves are bluish-green when young, becoming reddish.

Inflorescence

The inflorescence digitate to subdigitate, with ramose branches, the racemes. The inflorescence is subdigitate, with ramose branches, the racemes 4-11, and 3.5-7 cm long. Lower glumes not or rarely pitted, without a sub-apical arch of hairs.

Habitat

Generally, found in stream banks, drainage plains and valley slopes. It is found in grasslands and *Eucalyptus* forests.

This species may be confused with *Dichanthium* species.



Spikelets



***Bothriochloa pertusa* - Indian Bluegrass**

Derivation

pertusa - Latin for with a pit, alluding to the deep pit on the abaxial surface of the lower glume.

Habit

A low growing mat forming grass 15-70 cm tall, rooting from lower nodes. The mid-culm nodes are glabrous or bearded. This is a perennial grass that spreads via stolons.

Inflorescence

The inflorescence is subdigitate, the length of the panicle is 2.5-7.5 cm, and racemes are usually 3-8. The lower glumes are pitted (usually needs a hand lens to see it). There are only two species in Townsville which have a pit in the glumes, these are *B. decipiens* and *B. pertusa*.

Habitat

This species is introduced to Australia as a fodder grass. It spreads as a weed everywhere; it is in moderate to low fertility soils.

It is found in many gardens since it grows as a lawn. It grows in many types of soils; it withstands drought and disturbance.



Spikelets



Brachyachne

Native Couches

From the Greek *brachys* (short) and *achne* (scale, chaff), alluding to lemmas shorter than glumes.

Stoloniferous or tufted annuals or perennials. The digitate inflorescence contains 3-6 racemes with the spikelets in 2 close rows on 1 side of the raceme. The glumes are much longer than the single floret.

Subfamily: Chloridoideae; Tribe: Cynodonteae

World = 9, Australia = 5


Townsville species

* *Brachyachne convergens* Common Native Couch



Key to the species of *Brachyachne* and related species

Glumes shorter than floret *Cynodon*
 Glumes longer than floret *Brachyachne convergens*



***Brachyachne convergens* - Common Native Couch**

Derivation

convergens - from the Latin *convergo* (approach), the spikelets two-ranked but appearing to be one-ranked.

Habit

An annual or biennial erect or decumbent grass, the culms 23-60 cm tall. The leaf sheaths are bearded

Inflorescence

The panicle is digitate with 2-5 branches, 2-7 cm long. The glumes are 3-5 mm long.

Habitat

This species grows in a range of habitats including claypans, riverbanks, levees and flood plains and sandplains, often in disturbed areas such as roadsides.



Spikelet



Capillipedium


Scented Tops

From the Latin *capillus* (hair) and *pes* (foot), alluding to the hair-like pedicels.

Tufted perennials with erect, slender culms. The inflorescence is an open panicle with secondary branching obvious. The spikelets are paired; one sessile and awned, one pedicelled and unawned. The sessile spikelet at the end of the branches is accompanied by 2 pedicelled spikelets (terminal triplet). The sessile spikelet is awned

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 22, Australia = 2

Closely related to *Bothriochloa*, with the same pedicel morphology, i.e. with a longitudinal translucent furrow, often purple coloured; in cross section the pedicels are dumbbell shaped. The differences between the genera are shown in the table below.



Bothriochloa bladhii
Inflorescence, usually an arrangement of racemes on a central axis



Racemes with more than 8 spikelet pairs

Capillipedium
Inflorescence more than once-branched



Racemes with 1-8 spikelet pairs

Townsville species

Capillipedium parviflorum Scented Top
Capillipedium spicigerum Scented Top

Key to the species of *Capillipedium*

Racemes 1-2-jointed; lower glume 6-nerved, 2 nerves between keels*Capillipedium parviflorum*
Racemes 3-8-jointed; lower glume 8-9-nerved, 4-5 nerves between keels*Capillipedium spicigerum*



Capillipedium parviflorum



Capillipedium spicigerum

***Capillipedium parviflorum* - Scented Top**

Derivation

parviflorum - from the Latin *parvus* (small) and *flos* (flower); spikelets small or with few florets.

Habit

An erect perennial grass 50-100 cm tall.

Inflorescence

The panicle is open and purple coloured. The sessile spikelet is accompanied with a pedicelled spikelet. These spikelet pairs (sessile and pedicelled spikelets) are grouped 1-2 along the raceme.

Habitat

It is widespread in *Eucalyptus* woodlands.



***Capillipedium spicigerum* - Scented Top**

Derivation

spicigerum - from the Latin *spica* (thorn) and *gero* (carry); inflorescence a spicate panicle.

Habit

An erect perennial grass 50-150 cm tall. The grass is regarded as being of ornamental value.

Inflorescence

The panicle is open and purple coloured. The sessile spikelet is accompanied with a pedicelled spikelet. These spikelet pairs (sessile and pedicelled spikelets) are grouped 3-8 along the raceme.

Habitat

It is widespread in *Eucalyptus* forests and woodlands.



Spikelet



Cenchrus


(including *Pennisetum*)

From the Greek *kenchros* (millet).

Tufted or stoloniferous or rhizomatous annuals or perennials. The inflorescence is a spicate panicle (spike-like with short branches). The spikelets are alike and in groups of one to five and subtended by an involucre of bristles or fused to form a spiny burr. Spikelets fall from plant with bristles attached, usually leaving a bare rachis. The primary bristles are located immediately below the spikelets and are usually noticeably the longest bristles.

Subfamily: Panicoideae; Tribe: Paniceae

World = 133, Australia = 25



Townsville species

Cenchrus brevisetosus

* <i>Cenchrus ciliaris</i>	Buffel Grass
* <i>Cenchrus echinatus</i>	Mossman River Grass, Burr Grass
<i>Cenchrus elymoides</i>	
* <i>Cenchrus pedicellatus</i> subsp. <i>unispiculus</i>	Annual Mission Grass
* <i>Cenchrus pennisetiformis</i>	White Buffel Grass
* <i>Cenchrus setaceus</i>	Fountain Grass
<i>Cenchrus setiger</i>	Birdwood Grass

Key to the species of *Cenchrus* and related genera

1. Spikelets falling at maturity with bristles or spines attached2
- Spikelets falling at maturity without any bristles or spines attached *Setaria*



Setaria



Cenchrus

2. Bristles or spines stiff, joined to some degree at base, forming a burr3
- Bristles fine and thread-like, free to almost the base5
3. Spines connate for more than 1/3 of the distance above the base, forming a more or less globose burr or involucre, which encloses 1 or more spikelets *Cenchrus echinatus*
- Spines connate to 1/3 of the distance above the base, forming a small disc or shallow cup4
4. Spines prolonged beyond the burr none, glabrous or subglabrous, 2-4 mm long *Cenchrus setiger*
- Spines prolonged beyond the burr into distinct, slender, scabrid bristle *Cenchrus elymoides*

- 5. Inner bristles of involucre surrounding spikelets woolly-hairy *Cenchrus pedicellatus* subsp. *unispiculus*
- Inner bristles of involucre surrounding spikelets hairy6



Bristles woolly-hairy



Bristles hairy

- 6. Bristles or spines joined to some degree at the base..... *Cenchrus ciliaris*
- Bristles free or almost free to the base, fine and thread-like *Cenchrus setaceus*



Cenchrus ciliaris



Cenchrus setaceus

***Cenchrus ciliaris* - Buffel Grass**

Derivation

ciliaris - from the Latin *cilium* (eyelid) and *-are* (pertaining to), the glumes or lemmas ciliate on nerves or margins.

Habit

A perennial, tufted grass, the culms are 10-150 cm tall.

Inflorescence

The panicle is solid, a spicate, it is 2-14 cm long. The primary bristles are 10-23 mm, hairy; they are fine and thread-like, free to base. The inner bristles form an uninterrupted ring.

Habitat

This non-native species is widespread, naturalised in dry sites and wet places, in soils from sand to black cracking clay.



Spikelet



***Cenchrus echinatus* - Mossman River Grass, Burr Grass**

Derivation

echinatus - from the Latin *echinatus* meaning spiny, the inflorescence a very condensed panicle and the spikelets or axillary structures are awned, the whole thereby resembling a hedgehog.

Habit

An annual or perennial, tufted grass, the culms are 10-90 cm tall.

Inflorescence

The panicle is solid, a spicate, it is 2-10 cm long. The spines are joined for more than 1/3 of the distance above the base, forming a more or less globose burr.

Habitat

This non-native species grows in disturbed areas.



Spikelet



Cenchrus elymoides

Derivation

elymoides - from the Greek *-oides* (similar to), its resembling *Elymus* with respect to the inflorescence.

Habit

A perennial, tufted grass, the culms are 40-150 cm tall.

Inflorescence

The panicle is solid, a spicate, it is 10.5-16 cm long. The bristles are connate, the distance above the base forming a burr.

Habitat

This species is found in woodlands and coastal grasslands. It occurs on Castle Hill.



Spikelets



***Cenchrus pedicellatus* subsp. *unispiculus* - Annual Mission Grass**

Derivation

pedicellatum - from the Latin *pes* (foot), *-ellus* (diminutive) and *-ata* (possessing), the spikelets shortly stalked.

unispiculus -from the Latin *uni* (one) and *spiculus* (spike), the involucre with one spikelet.

Habit

An annual or perennial, tufted grass, the culms are 30-150 cm tall.

Inflorescence

The panicle is solid, a spicate, it is 15-25 cm long. The primary bristles are 15-25 mm long, woolly-hairy, they are fine and thread-like, free to base.

Habitat

This non-native species grows in woodlands, arid hummock grasslands, and coastal grasslands



Spikelet



***Cenchrus setaceus* - Fountain Grass**

Derivation

setaceus - from the Latin *seta* (bristle) and *-acea* (indicating resemblance), with long bristles in the inflorescence.

Habit

A perennial, tufted grass, the culms are 20-150 cm tall.

Inflorescence

The panicle is solid, a spicate, it is 6-30 cm long. The primary bristles are 26-35 mm, hairy; they are fine and thread-like, free to base.

Habitat

This non-native species has been cultivated as an ornamental; it escapes to become a weed.



Spikelet



***Cenchrus setiger* - Birdwood Grass**

Derivation

setigerus - from the Latin *seta* (bristle) and *gero* (carry), with spikelets subtended by bristles.

Habit

A perennial, tufted grass, the culms are 5-100 cm tall.

Inflorescence

The panicle is solid, a spicate, it is 2-12 cm long. The bristles are connate to 1/3 of the distance above the base, forming a small disc or shallow cup.

Habitat

This non-native species grows in open dry woodlands and grasslands, usually on alkaline soils, sometimes on heavy black clays with impeded drainage.



Spikelet



Chionachne


From Greek *chion* (snow) and *achne* (chaff or scale), in allusion to the pale-coloured glumes of some species.

Annuals or reed-like perennials, rhizomatous or tufted. The inflorescence is subtended by a spathe. The spikelets are unisexual and segregated in different parts of the inflorescence which is a raceme or spike. The female spikelets are at the base and are sessile. The male flowers are at the apex, they are usually paired, one sessile and one pedicelled.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 11, Australia = 2

Townsville species


Chionachne cyathopoda River Grass



Raceme showing male and female spikelets




Female spikelets



Female spikelet



Male spikelets



Male spikelets



***Chionachne cyathopoda* - River Grass**

Derivation

cyathopoda - from the Greek *kyathos* (cup) and *pous* (foot) referring to the shape of the apex of peduncle.

Habit

A tufted perennial grass with reed-like culms, the culms are 2-4 m tall

Inflorescence

The inflorescence is subtended by a spatheole, the racemes are single, 6-12 cm long and bearing a few female spikelets.

Habitat

This species resembles *Megathyrsus maximus*. It grows on creek banks, in sandy or loamy soils.



Male spikelets



Female spikelets

Chloris


Windmill Grasses

From the Greek *chloros* (green), possibly alluding to the greenish flowers and green leaves; alternatively, named for *Chloris* (The Green One), mythological Greek goddess of flowers.


Tufted or stoloniferous perennials or annuals. The inflorescence is digitate or subdigitate, with the branches erect to spreading. The spikelets are solitary, laterally compressed, usually overlapping in 2 rows along 1 side of the slender rachis. Each spikelet consists of 2-4 florets with the basal floret bisexual and the upper florets reduced. The lemmas have 1-3 awns.

Chloris is closely related and easily confused with the genera *Enteropogon* and *Oxychloris*. The genus *Chloris* also has a superficial similarity to the genus *Pseudopogonatherum*.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 131, Australia = 13



Chloris inflata
spikelet




Chloris pumilio
florets

Townsville species

* <i>Chloris gayana</i>	Rhodes Grass
* <i>Chloris inflata</i>	Purpletop Grass
<i>Chloris lobata</i>	
<i>Chloris pectinata</i>	Comb Windmill Grass
<i>Chloris pumilio</i>	
* <i>Chloris virgata</i>	Feathertop Rhodes Grass


Key to the species of *Chloris* and similar species

1. Second and third lemmas very broad, sharply recurved at maturity (wing-like) *Oxychloris scariosa*
Lemmas not very broad, not wing-like 2
2. Lowest lemma dorsally compressed (lying on front or back when placed on a flat surface)
..... *Enteropogon ramosus*
Lowest lemma laterally compressed (lying on the side when placed on a flat surface) *Chloris*



Key to the species of *Chloris*

1. Spikelets blunt at apex2
- Spikelets narrowed to pointed at apex4



2. Upper lemmas inflated*Chloris inflata*
- Upper lemmas not inflated3
3. Spikelet with 2 florets (1 bisexual) *Chloris virgata*
- Spikelet with 3-4 florets (1 or rarely 2 bisexual) *Chloris gayana*
4. Lemma 1-awned5
- Lemma 3-awned6
5. Lowest lemma laterally compressed *Chloris pectinata*
- Lemma dorsally compressed.....*Enteropogon ramosus*
6. Lemma awns subequal, the central slightly longer *Chloris lobata*
- Lemma awns very unequal*Chloris pumilio*

***Chloris gayana* - Rhodes Grass**

Derivation

gayana - in honour of Claude Gay (1800-1873), a French botanist.

Habit

A perennial, erect grass, the culms are 45-120 cm tall.

Inflorescence

The inflorescence is digitate with 6-18 branches, erect or spreading, 5-10 cm long.

Habitat

This non-native species is mostly found on roadsides and other disturbed areas, it prefers heavy black or grey soils but is found on sandy and reddish-brown loamy soils.



Spikelet



***Chloris inflata* - Purpletop Grass**

Derivation

inflata - Latin for swollen, the upper lemmas are swollen.

Habit


An annual or biennial, erect or decumbent grass, the culms are 40-90 cm tall.

Inflorescence

The inflorescence is digitate with 7-16 branches, erect or spreading, 2-9 cm long.

Habitat

This non-native species occurs generally as a weed of disturbed sites, common on roadsides.



Florets



Chloris lobata

Derivation

lobata - from the Latin *lobus* (lobe) and *-ata* (possessing) the lemmas are lobed.

Habit


An annual, erect or prostrate grass, the culms are 15-40 cm tall.

Inflorescence

The inflorescence is digitate with 2-7 branches, ascending or spreading, 1.4-5 cm long.

Habitat

This species grows on a range of soil types but mostly heavy soils in seasonally wet areas such as claypans, river flood flats, creek beds and creek banks.



Florets



***Chloris pectinata* - Comb Windmill Grass**

Derivation

pectinata - from the Latin *pecten* (comb) and *-ata* (possessing), the inflorescence with spikes or racemes bearing a resemblance to a comb.

Habit

An annual, erect or decumbent grass, the culms are 10-60 cm tall.


Inflorescence

The inflorescence is digitate with 3-10 branches, spreading, 3-14 cm long.

Habitat

This species grows in depressions, flood-outs and other watered sites, and extensive cracking clay pans.

Chloris pectinata florets are like *Enteropogon ramosus*, *Chloris pectinata* lowest lemma is laterally compressed.



Florets

Chloris pumilio

Derivation

pumilio - Latin for a dwarf, this grass is small compared with related species.

Habit

An annual, erect or decumbent, the culms are 18-90 cm tall.


Inflorescence

The inflorescence is digitate with 6-10 branches, erect, 2-10 cm long.

Habitat

This species is found mostly in sandy and silty soils, in moist, low-lying areas, also growing on podsolic, reddish brown and grey, brown soils or dark grey loams.

Chloris pumilio can be mistaken for the genus *Pseudopogonatherum*.



Florets

***Chloris virgata* - Feathertop Rhodes Grass**

Derivation

virgata - from the Latin *virga* (broom) and *-ata* (possessing), the inflorescence branches or culms held erect.

Habit

An annual or short-lived perennial grass, the culms are 15-100 cm tall.

Inflorescence

The inflorescence is digitate with 7-19 branches, erect, 3-9 cm long.

Habitat

This non-native species is generally a weed of disturbed ground, especially roadsides. It grows on a variety of soil types such as heavy black soil, grey and brown clays and loams, and red sandy soil.



Florets



Chrysopogon

Golden Beard Grasses


From Greek *chrysos* (golden) and *pogon* (beard), alluding to the golden hairs on the inflorescence.

Annuals or perennials, rhizomatous or stoloniferous or tufted or decumbent. The inflorescence is an open or contracted panicle. The spikelets are borne at the ends of the inflorescence branches. The spikelets in pairs or triplets; one sessile spikelet and 1 or 2 pedicelled spikelets. The sessile spikelet is awned or unawned. Mature spikelets are often purplish coloured.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 67, Australia = 13

Townsville species

* <i>Chrysopogon aciculatus</i>	Mackies Pest
<i>Chrysopogon fallax</i>	Golden Beard Grass
<i>Chrysopogon filipes</i>	Australian Vetiver
* <i>Chrysopogon zizanioides</i>	Vetiver Grass



Key to the species of *Chrysopogon*

- Plants stoloniferous; grows in lawns *Chrysopogon aciculatus*
Plants not stoloniferous, tufted.....2
- Plant 150-300 cm tall; sessile spikelets awnless..... *Chrysopogon zizanioides*
Plant 30-120 cm tall; sessile spikelets awned3
- Racemes with 1 or 2 jointed.....*Chrysopogon fallax*
Racemes 3-many jointed *Chrysopogon filipes*



Inflorescence
Chrysopogon fallax



Racemes



Inflorescence
Chrysopogon filipes



Racemes

***Chrysopogon acicularis* - Mackies Pest**

Derivation

aciculatus - from the Latin *acus* (needle), *-ulus* (diminutive) and *-atus* (possessing), the spikelets with a needle-like callus.

Habit

A perennial, mat forming grass, stolons present, the culms are 20-50 cm tall.

Inflorescence

The panicles are open, ovate, 4-10 cm long, the racemes are 1-2 jointed. The sessile spikelets are awned and are 4-10 mm long.

Habitat

This non-native species grows as a weed in lawns.



Spikelets



***Chrysopogon fallax* - Golden Beard Grass**

Derivation

fallax - Latin for deceptive, it closely resembling another species.

Habit


A perennial, tufted grass, the culms are 30-120 cm tall.

Inflorescence

The panicles are open, 7-20 cm long, the racemes are 1-2 jointed. The sessile spikelets are awned and are 9-14 mm long.

Habitat

This species is widespread on a range of soils, often on floodplains.



Spikelets



Chrysopogon filipes - Australian Vetiver

Derivation

filipes - from the Latin *filum* (thread) and *ped* (foot), the pedicels of racemes slender.

Habit

A perennial, tufted grass, the culms are 60-100 cm tall.

Inflorescence

The panicles are contracted, 15-30 cm long, the racemes are 3-many jointed. The sessile spikelets are awned and are 8-10 mm long.

Habitat

This species grows on banks, floodplains and dry parts of creek or riverbeds.



Spikelets



***Chrysopogon zizanioides* - Vetiver Grass**

Derivation

zizanioides - by the riverside, reflecting the fact that the plant is commonly found along the waterways.

Habit

A perennial, tufted grass, the culms are 150-300 cm tall.

Inflorescence

The panicles are open, lanceolate, 15-40 cm long, the racemes are 3-many jointed. The sessile spikelets are not awned and are 3.5-5 mm long.

Habitat

This non-native species is planted for erosion control, however occasionally it escapes to nearby areas.

Chrysopogon zizanioides is confused with *Melinis* and *Capillipedium*; use the key on the *Melinis* page.



Spikelets




Cleistochloa

From the Greek *kleistos* (enclosed) and *chloe* (grass), referring to some of the spikelets being cleistogamous (flowers which do not open and are self-fertilized in the bud).

Tufted perennial grasses with wiry culms. The inflorescences are of two types, one a terminal raceme, the other is an axillary inflorescence with cleistogamous spikelets. The spikelets in each inflorescence are dissimilar. This genus is endemic to Australia and species grow on low-nutrient sandstone-derived soils.

Subfamily: Panicoideae; Tribe: Paniceae
World = 3, Australia = 3

Townsville species
Cleistochloa subjuncea



Cleistochloa subjuncea

Derivation

subjuncea - from this genus; *sub* (somewhat similar) and *juncea* (rush-like), refers to the habit.

Habit

An erect or slightly spreading perennial grass, the culm is 30-60 cm tall. The leaf sheaths usually have erect white hairs on the back, and when the grass dries off, the leaf sheath loosens from the culm and the leaf blade breaks off at the ligule at maturity.

Inflorescence

The inflorescence is in two forms, one terminal, and the other axillary. The terminal inflorescence is 0.7-3.5 cm long with 2-6 spikelets; the axillary inflorescence is reduced to a single spikelet that is hidden in the leaf sheaths.

Habitat

This species grows on very poor, dry soils.

Old leaf
sheath



Leaf sheath showing
erect white hairs



Terminal spikelets



Axillary spikelet




Cymbopogon

Lemon-scented Grasses

From the Greek words *kumbe* (boat) and *pogon* (beard), referring to many-awned inflorescences and boat-shaped spathes.

Tufted perennials with aromatic (often lemon-scented) shoots and leaves when crushed. The inflorescence is a panicle of short paired racemes each subtended by a reddish spatheole. The spikelets are in pairs, one sessile and one pedicelled. The sessile spikelet is awned. Many of the species are covered with long, soft, white hairs.

The lemon-scented leaves are distinctive and make the identification of this genera easier even when the grass is not flowering. However, there is a grass species in another genera in Townsville with lemon-scented leaves - *Elionurus citreus*. Once all the fluffy spikelets have fallen, this genera can be mistaken for *Themeda*.



Subfamily: Panicoideae; Tribe: Andropogoneae
World = 101, Australia = 12

Townsville species

<i>Cymbopogon ambiguus</i>	Scented Oil Grass, Lemon Grass
<i>Cymbopogon bombycinus</i>	Silky Oil Grass
<i>Cymbopogon nardus</i>	Citronella Grass
<i>Cymbopogon obtectus</i>	Silkyheads
<i>Cymbopogon queenslandicus</i>	
<i>Cymbopogon refractus</i>	Barbed Wire Grass

Key to the species of *Cymbopogon* species

1. Racemes distinctly hairy, hairs 4-7 mm long.....2
Racemes only slightly hairy, hairs less than 3 mm long4
2. Longer callus hairs up to 1.3 mm long *Cymbopogon ambiguus*
Longer callus hairs 3.5-5 mm long3
3. Leaf sheaths to 6 mm wide, rolling back at maturity *Cymbopogon bombycinus*
Leaf sheaths 2-3 mm wide, not rolling back at maturity..... *Cymbopogon obtectus*

Racemes and spikelets distinctly hairy, hairs 4-7 mm long



Cymbopogon ambiguus



Cymbopogon bombycinus



Cymbopogon obtectus

Racemes and spikelets only slightly hairy, hairs less than 3 mm long



Cymbopogon queenslandicus



Cymbopogon refractus



Cymbopogon nardus

Leaf sheaths to 6 mm wide, rolling back at maturity



Leaf sheaths 2-3 mm wide, not rolling back at maturity



4. Plant 150-250 cm.....*Cymbopogon nardus*
Plant 30-150 cm5
5. Pedicelled spikelet with one glume only; sessile spikelet usually awnless (if awned, awn small and imperfect), always distinctly reflexed at maturity*Cymbopogon refractus*
Pedicelled spikelet with 2 glumes; sessile spikelet with an awn 10-12 mm long, usually not reflexed at maturity..... *Cymbopogon queenslandicus*

***Cymbopogon ambiguus* - Scented Oilgrass, Lemon Grass**

Derivation

ambiguus - Latin for uncertain, species that may be readily confused with others or do not necessarily belong to the genus in which they have been placed or are intermediate in characters between other genera.

Habit

A perennial, tufted grass, the culms are 30-150 cm tall.

Inflorescence

The panicle is 15-40 cm long, the spathes 1.5-3.5 cm long. The racemes are erect, distinctly hairy, hairs 4-7 mm long, the longer callus hairs up to 1.3 mm long.

Habitat

This species is very adaptable to different types of soils; it is commonly on rocky outcrops.



Spikelets



***Cymbopogon bombycinus* - Silky Oil Grass**

Derivation

bombycinus - from the Latin *bombyx* (silk) and *-inus* (belonging to), the inflorescence or leaves invested with long silky hairs.

Habit

A perennial, tufted grass, the culms are 30-120 cm tall. The leaf sheaths are rolling back at maturity.

Inflorescence

The panicle is 6-40 cm long, the spathes 2.0-3.5 cm long. The racemes are erect, distinctly hairy, hairs 4-7 mm long, the longer callus hairs 3.5-5 mm long.

Habitat

This species is usually in *Eucalyptus* forest on sandy or stony soils, it grows on hills.



Spikelets



***Cymbopogon nardus* - Citronella Grass**

Derivation

nardus - from the species *Nardus stricta*, another aromatic plant.

Habit

A robust, perennial, tufted grass, the culms are 150-250 cm tall. The leaves are used to extract citronella oil, which used as insect-repelling properties.

Inflorescence

The dense panicle is 15-30 cm long, the spathes 1-1.5 cm long. The racemes are erect, but drooping due the large number of spikelets, which are only slightly hairy. The pedicels are hairy. The sessile spikelet is 4-4.5 mm long awned.

Habitat

This non-native species grows in drains and along roadsides.



Spikelet



***Cymbopogon obtectus* - Silkyheads**

Derivation

obtectus - from the Latin *obtego* (conceal), it is segregated from a closely related species.

Habit

A perennial, tufted grass, the culms are 50-100 cm tall. The leaf sheaths not rolling back at maturity.

Inflorescence

The panicle is 7-25 cm long, the spathes 2.0-3.2 cm long. The racemes are erect and commonly deflexed, and distinctly hairy, hairs 4-7 mm long, the longer callus hairs 3.5-5 mm long.

Habitat

This species is tolerant of a range of environments; it grows on hills.



Spikelets



Cymbopogon queenslandicus

Derivation

queenslandicus - from Queensland.

Habit

A perennial, tufted grass, the culms are 70-150 cm tall.

Inflorescence

The panicle is 25-45 cm long, the spathes 1.8-2.5 cm. The racemes are erect and occasionally deflexed, and are only slightly hairy, hairs less than 3 mm long. The pedicels are hairy.

Habitat

This species occurs in *Eucalyptus* forests, often on stony hillsides.



Spikelets



***Cymbopogon refractus* - Barbed Wire Grass**

Derivation

refractus - Latin for curved back abruptly, it is mostly applied to species whose mature inflorescence branches curve back.

Habit

A perennial, tufted grass, the culms are 30-150 cm tall.

Inflorescence

The panicle is 10-45 cm long, the spathes 1.7-2.5 cm. The racemes are always distinctly reflexed at maturity, and usually glabrous.

Habitat

This species is found in grassland and in the *Eucalyptus* forests and woodlands.

Cymbopogon refractus appears to produce occasional hybrids with other species.



Spikelet



Cynodon

Couchs and Star Grasses

From Greek *kynos* (dog) and *odous* (tooth), alluding to the tooth-like buds of the rhizome.


Stoloniferous and rhizomatous perennials with short, erect flowering stems. The inflorescence is digitate or subdigitate. Spikelets are solitary and on one side of the raceme. The glumes are shorter than the single floret.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 56, Australia = 9

Cynodon dactylon is the common lawn grass (couch) in Townsville. This grass is an extremely variable species, the growth form varies according to local condition. Widely distributed and common in disturbed areas, it can grow to 30 cm tall, when frequently mown it can form a close sward with a prostrate habit.

Townsville species


- | | |
|------------------------------|---------------|
| * <i>Cynodon dactylon</i> | Couch Grass |
| * <i>Cynodon nlemfuensis</i> | Bermuda Grass |
| * <i>Cynodon radiatus</i> | Bermuda Grass |




Spikelet

Key to the species of *Cynodon* and related species

1. Glumes longer than floret *Brachyachne*
- Glumes shorter than floret 2



Glumes longer than floret
Brachyachne



Glumes shorter than floret
Cynodon

2. Rhizomes present; leaf blades 0.5-3 mm long *Cynodon dactylon*
- Rhizomes absent; leaf blade 2-7 mm long 3
3. Culms herbaceous or woody, 20-83 cm high; leaf blades acuminate; anthers orange
..... *Cynodon nlemfuensis*
- Culms wiry, 15-45 cm high; leaf blades acute; anthers yellow or purple *Cynodon radiatus*

***Cynodon dactylon* - Couch Grass**

Derivation

dactylon - from the Greek *daktylos* finger, the inflorescence branches are finger-like.

Habit

A perennial, stoloniferous and rhizomatous grass, the culms are 10-35 cm tall.

Inflorescence

The panicle is digitate with 4-7 branches, 2-5.5 cm long.

Habitat

This non-native species is cultivated extensively as a lawn grass. A weed of roadsides and in pastures.



Spikelet



***Cynodon nlemfuensis* - Bermuda Grass**

Derivation

nlemfuensis - from Nlemfu, Zaire.

Habit

A perennial, stoloniferous grass, the culms are 20-80 cm tall.

Inflorescence

The panicle is digitate or subdigitate with 4-13 branches, 4-10 cm long.

Habitat

This non-native species was introduced as a pasture grass, has become a vigorous weed.



Inflorescence



Cynodon radiatus - Bermuda Grass

Derivation

radiatus - from the Latin *radius* (spoke of a wheel).

Habit

A perennial, stoloniferous grass, the culms are 15-45 cm tall.

Inflorescence


The panicle is digitate with 4-8 branches, 4-9 cm long.

Habitat

This non-native species is a vigorously spreading coloniser of disturbed sites.



Inflorescence



Dactyloctenium

Button Grasses

From the Greek *daktylos* (a finger) and *ktenion* (a little comb), alluding to the digitate inflorescence in which the spikelets have a comb-like arrangement.

Annuals or perennials, rhizomatous or stoloniferous, tufted or decumbent grass and native or naturalised. The inflorescence is digitate, with 2-6 branches which are erect or spreading, the end of the branches project beyond the spikelets. The spikelets are sessile and closely overlapping in two rows on one side of the rachis. The spikelets contain 2 or more bisexual florets.


Sometimes grows in saline habitats or dunes, mostly in dry sandy soils.

Subfamily: Chloridoideae; Tribe: Cynodonteae

World = 17, Australia = 5

Townsville species

- | | |
|-------------------------------------|----------------------|
| * <i>Dactyloctenium aegyptium</i> | Coastal Button Grass |
| <i>Dactyloctenium buchananensis</i> | |
| <i>Dactyloctenium radulans</i> | Native Button Grass |



Key to the species of *Dactyloctenium* and similar species

1. Axis of each raceme ending with a spikelet..... *Eleusine indica*
- Axis of each raceme ending in a point2



Axis of each raceme ending with a spikelet



Axis of each raceme ending in a point

2. Inflorescence branches are 0.4-0.6 cm mm long.....*Dactyloctenium buchananensis*
- Inflorescence branches are at least 1 cm long.....3
3. Inflorescence branches up to 1.5 cm long; spikelets c. 4 mm long.....*Dactyloctenium radulans*
- Inflorescence branches 1-5 cm long, spikelets 2.5-3 cm long *Dactyloctenium aegyptium*



Dactyloctenium buchananensis



Dactyloctenium radulans



Dactyloctenium aegyptium

***Dactyloctenium aegyptium* - Coastal Button Grass**

Derivation

aegyptium - from the Latin *-ium* (characteristic of), from Egypt.

Habit

An annual or short-lived perennial grass, the culms are 40-100 cm tall.

Inflorescence

The inflorescence branches 2-5, touching only near the base, 1.2-6.5 cm long.

Habitat

This non-native species is most common in disturbed areas but also in open *Eucalyptus* forest on a variety of soils, and on coastal dunes.



Inflorescence



Dactyloctenium buchananensis

Derivation

It grows on the edge of Lakes Buchanan and Galilee in central Queensland.

Habit

An ephemeral or annual grass, the culms are 8-40 cm tall.

Inflorescence

The inflorescence branches 8-12, touching for most of length and forming a head-like structure, 0.4-0.6 cm long.

Habitat

This species grows in waters of high salinity, on saline clay; it is reported as an introduction in the Townsville area.



Inflorescence



***Dactyloctenium radulans* - Native Button Grass**

Derivation

radulans - from the Latin *radula* (scraper) and *-ans* (assuming the form of), the leaf-blades are scabrid.

Habit

An ephemeral or annual grass, the culms are 7-33 cm tall.

Inflorescence

The inflorescence branches 3-10, touching for most of length and forming a head-like structure, 0.5-1.5 cm long.

Habitat

This species grows on many soil types, but most often on sand and in seasonally wet areas such as creek banks and flood plains, tolerates saline and swampy conditions.



Inflorescence



Dichanthium


Bluegrasses

From the Greek *dicha* (in two, apart) and *anthos* (flowers), alluding to the two kinds of spikelet pairs in the raceme.

Tufted, decumbent or stoloniferous perennials, often with branched culms. The inflorescence is digitate or subdigitate or a single raceme. The spikelets are in pairs (with terminal triplets), one sessile and one pedicelled. The sessile spikelet is awned, and in one species (*D. fecundum*), the pedicelled spikelet is also sometimes awned.

Dichanthium is closely related to *Bothriochloa*, and the difference between these two genera is described under *Bothriochloa*.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 54, Australia = 11



Townsville species

- * *Dichanthium annulatum*
- * *Dichanthium aristatum*
- Dichanthium fecundum*
- Dichanthium sericeum* subsp. *polystachyum*
- Dichanthium sericeum* subsp. *sericeum*

Sheda Grass
Angleton Grass
Curly Bluegrass
Queensland Bluegrass
Queensland Bluegrass



Key to the species of *Dichanthium*, *Bothriochloa* and *Capillipedium*

1. Inflorescence a panicle of racemes2
 Inflorescence digitate or subdigitate or a single raceme3
2. Inflorescence, usually an arrangement of racemes on a central axis; racemes with more than 8 spikelet pairs *Bothriochloa bladhii* subsp. *bladhii*
 Inflorescence more than once branched; racemes with 1-8 spikelet pairs..... *Capillipedium*

Inflorescence, usually an arrangement of racemes on a central axis
 Racemes with more than 8 spikelet pairs




Bothriochloa bladhii

Inflorescence more than once branched
 Racemes with 1-8 spikelet pairs




Capillipedium


3. Lower glumes pitted4
 Lower glumes not pitted5



Spikelet showing pitted lower glume



4. Culms erect or ascending 30-200 cm tall; nodes glabrous and purple; pedicelled spikelet reduced to .. lower glume *Bothriochloa decipiens*
 Culms erect, ascending or prostrate 15-70 cm tall and with stolons; nodes glabrous or bearded; pedicelled spikelet similar to sessile spikelet, male or sterile *Bothriochloa pertusa*
5. Peduncle below inflorescence covered in downy hairs, sometimes has a single raceme *Dichanthium aristatum*
 Peduncle below inflorescence glabrous.....6



Bothriochloa decipiens



Bothriochloa pertusa



Dichanthium aristatum. Showing downy hairs at the base of the inflorescence

- 6. Racemes bases sessile; lower glume of sessile spikelet with a distinct sub-apical arch of long, fine, simple hairs7
- Raceme bases filiform or sub-sessile; lower glume of sessile spikelet without a sub-apical arch of simple hairs, tubercle-based hairs sometimes present8



Racemes bases sessile



Racemes filiform or sub-sessile

- 7. Lower glume of pedicelled spikelet obovate with subapical arch conspicuous, hairs erect from surface; racemes usually 10 or more*Dichanthium sericeum* subsp. *polystachyum*
- Lower glume of pedicelled spikelet linear to narrowly ovate, hairs not erect from surface; racemes usually 6 or less *Dichanthium sericeum* subsp. *sericeum*
- 8. Pedicels of the companion spikelet with a translucent furrow between thickened margins *Bothriochloa ewartiana*
- Pedicels of the companion spikelet without a translucent mid-line9



Dichanthium sericeum



Bothriochloa ewartiana

9. Spikelets acute; pedicelled spikelets bisexual or male and sometimes awned; stigmas distinctly protruding at maturity *Dichanthium fecundum*
 Spikelets obtuse or truncate; pedicelled spikelets sterile or male; stigmas not distinctly protruding at maturity *Dichanthium annulatum*



Dichanthium fecundum



Dichanthium annulatum



***Dichanthium annulatum* - Sheda Grass**

Derivation

annulatum - from the Latin *annulus* (ring) and *-ata* (possessing).

Habit

A perennial, tufted grass, the culms are decumbent, 25-100 cm tall, the mid-culms nodes are bearded.

Inflorescence

The inflorescence is digitate or subdigitate, with 2 to 9 racemes, 3-7 cm long. The pedicelled spikelet is sterile or male.

Habitat

This non-native grass is found in situations growing in pastureland, roadsides, weedy lawns, sand dunes and open wasteland.



Spikelets



Node



Spikelets pair

***Dichanthium aristatum* - Angleton Grass**

Derivation

aristatum - from the Latin *arista* (bristle) and *-ata* (possessing); the apices of lemmas, paleas or glumes drawn out into a distinct awn.

Habit

A perennial, tufted grass, the culms are decumbent, 20-100 cm tall, the mid-culm nodes are glabrous or pubescent.

Inflorescence

The inflorescence is a digitate or subdigitate, 2-5 racemes, sometimes only one raceme at the end of the season, 3-8 cm long. The pedicelled spikelet is sterile or male. The peduncles are covered in downy hairs.

Habitat

This non-native grass is tolerant of a wide range of soils, it is a vigorous plant with a tendency to dominate and become a weed.



Spikelets



***Dichanthium fecundum* - Curly Bluegrass**

Derivation

fecundum - Latin for fruitful, for producing abundant spikelets.

Habit

A perennial, tufted grass, the culms erect or geniculately ascending, 60-150 cm tall, the mid-culm nodes are bearded.

Inflorescence

The inflorescence is digitate or subdigitate, with 3-7 racemes, sometimes only one raceme, 4-6 cm long. The lower glume can sometimes have a transverse fringe of hairs. The pedicelled spikelet is bisexual or male, sometimes with awns.

Habitat

This species is found in clay loam and alluvial areas and along stream banks, and river flats.



Spikelet



Dichanthium sericeum - Queensland Bluegrass

Derivation

sericeum - from the Latin *sericeus* (silken) which means silky, referring to the hairs on the inflorescence.

Habit

An annual or perennial, tufted grass, culms 10-120 cm tall, the mid-culm nodes are bearded.

Inflorescence

The inflorescence is digitate or subdigitate with 2-10 plus racemes, 1.5-7 cm long. The racemes bases are brief. The lower glume of sessile spikelet has a distinct, sub-apical arch of long, fine, simple hairs. The pedicelled spikelet is sterile.

There are two subspecies found growing in Townsville.

Racemes usually 8 or more *Dichanthium sericeum* subsp. *polystachyum*

Racemes usually 6 or less *Dichanthium sericeum* subsp. *sericeum*

Habitat

Dichanthium sericeum subsp. *polystachyum* - This species is found mainly in open wet grassland on wet soil

Dichanthium sericeum subsp. *sericeum* - This species is found in open grassland on clay soils.



Sessile spikelet
with sub-apical
hairs



subsp. *sericeum*




subsp. *polystachyum*

Digitaria

Finger Grasses


From Latin *digitus* (finger) alluding to the digitate inflorescence and *aria* pertaining to.

Annuals or perennials, rhizomatous or stoloniferous or tufted, decumbent or prostrate. The inflorescence is a raceme, digitate, subdigitate or a once-branched panicle. The spikelets are usually in groups of 2 or 3, the lower glume is absent or small and acute at the apex and the palea (the upper bract of the floret enclosing the flower) of the upper floret almost completely clasps the upper lemma (the outer bract of the floret).




Subfamily: Panicoideae; Tribe: Paniceae

World = 316, Australia = 50



Spikelets



Inflorescences



Townsville species

<i>Digitaria ammobila</i>	Silky Umbrella Grass, Spider Grass
<i>Digitaria bicornis</i>	Hairy Finger Grass
<i>Digitaria breviglumis</i>	
<i>Digitaria brownii</i>	Cotton Panic Grass
* <i>Digitaria ciliaris</i>	Summer Grass, Crab Grass
* <i>Digitaria didactyla</i>	Queensland Bluegrass
<i>Digitaria diffusa</i>	
* <i>Digitaria eriantha</i>	Pangola Grass
<i>Digitaria fumida</i>	
<i>Digitaria gibbosa</i>	
* <i>Digitaria insularis</i>	Sourgrass
<i>Digitaria leucostachya</i>	
<i>Digitaria longiflora</i>	
<i>Digitaria minima</i>	
<i>Digitaria nematostachya</i>	
<i>Digitaria orbata</i>	
<i>Digitaria parviflora</i>	Small Flower Finger Grass
<i>Digitaria ramularis</i>	
* <i>Digitaria violascens</i>	Purple Crabgrass

Key to the species of *Digitaria*

1. Racemes all distinctly stalked; inflorescence 50 cm wide2
- Racemes mostly with spikelets to base3



Racemes all distinctly stalked



Racemes mostly with spikelets to base

2. Spikelets woolly-hairy *Digitaria ammophila*
- Spikelets glabrous or very slightly hairy *Digitaria nematostachya*
3. Spikelets with long hairs exceeding spikelet.....4
- Spikelets glabrous or hairy, if latter then hairs not exceeding spikelet7




Spikelets with long hairs exceeding spikelet




Spikelets glabrous or hairy

4. Annual; inflorescence of 1 raceme *Digitaria gibbosa*
- Perennials; inflorescence of more than 1 raceme5
5. Inflorescence of 8-50 racemes *Digitaria insularis*
- Inflorescence of 1-7 racemes.....6
6. Lower glume $\frac{1}{4}$ - $\frac{1}{2}$ spikelet length, separated to some degree from upper glume..... *Digitaria brownii*
- Lower glume absent or a minute hyaline rim, adjacent to upper glume *Digitaria leucostachya*
7. The upper glume is 10-40% of the length of the spikelet8
- The upper glume is 50-100% of the length of the spikelet 10
8. Inflorescence with 1-4 racemes *Digitaria breviglumis*
- Inflorescence with many racemes, 5-89
9. Lower glume absent*Digitaria orbata*
- Lower glume present *Digitaria fumida*
10. Inflorescence a once-branched open panicle, 2-15 racemes 11
- Inflorescence digitate or subdigitate..... 14



Inflorescence a raceme or once branched



Inflorescence digitate or subdigitate

11. Erect grass 40-150 cm tall 12
Decumbent grass to 10-60 cm tall 13
12. Upper glume 1/2-4/5 spikelet length *Digitaria ramularis*
Upper glume subequal to spikelet length *Digitaria parviflora*
13. Upper glume muricate and non-reflective *Digitaria minima*
Upper glume relatively smooth and reflective *Digitaria diffusa*
14. Spikelets of each spikelet pair different or similar and bristled *Digitaria bicornis*
Spikelets of each pair similar and not bristled 15



Spikelets of each spikelet pair different,
one is hairy, one is glabrous



Spikelets of each pair similar

15. Spikelets clustered or in threes 16
Spikelets arranged in pairs 17



Spikelets usually arranged in threes



Spikelets arranged in pairs

16. Plant usually a stoloniferous perennial; upper lemma pale to yellow at maturity; upper glume 5-7-nerved *Digitaria longiflora*
Plant annual; Upper lemma brown to black at maturity; upper glume 3-nerved *Digitaria violascens*
17. Mat-forming perennials; racemes 2-4 *Digitaria didactyla*
Tufted grass; racemes 2-12 18
18. Annuals *Digitaria ciliaris*
Perennials *Digitaria eriantha*

Inflorescence a once-branched open panicle				
	Racemes	Spikelets	Spikelets	Upper glume length compared to spikelet
<i>Digitaria ammophila</i>	5-15	solitary or pairs	hairy	100%
<i>Digitaria breviglumis</i>	1-4	clustered at each node or in pairs	glabrous	30-40%
<i>Digitaria brownii</i>	1-7	pairs	long hairs	85-100%
<i>Digitaria diffusa</i>	3-10	pairs	glabrous or hairy	100%
<i>Digitaria fumida</i>	(2-)6-10	clustered at each node or in pairs or threes	glabrous	10-25%
<i>Digitaria insularis</i>	8-50	pairs	long hairs	100%
<i>Digitaria minima</i>	2-7	pairs	glabrous	75-85%
<i>Digitaria nematostachya</i>	10-20	pairs	glabrous	50-90%
<i>Digitaria orbata</i>	(2-)6-10	clustered at each node or in pairs	glabrous	10-25%
<i>Digitaria parviflora</i>	4-15	clustered at each node or in pairs	glabrous	100%
<i>Digitaria ramularis</i>	2-10	solitary or clustered at each node or in pairs.	glabrous or pubescent	50-75%

Inflorescence a digitate or subdigitate panicle				
	Racemes	Spikelets	Spikelets	Upper glume length compared to spikelet
<i>Digitaria bicornis</i>	2-4	pairs	glabrous and hairy	66%
<i>Digitaria brownii</i>	1-7	pairs	long hairs	85-100%
<i>Digitaria ciliaris</i>	2-12	pairs	hairy	60-75%
<i>Digitaria didactyla</i>	2-4	pairs	hairy	60-75%
<i>Digitaria eriantha</i>	3-10	pairs	hairy	60-75%
<i>Digitaria leucostachya</i>	1-4	clustered at each node or in pairs	long hairs	100%
<i>Digitaria longiflora</i>	2-4	clustered at each node or in threes	hairy	100%
<i>Digitaria violascens</i>	2-6	clustered at each node or in threes	hairy	80-100%

Inflorescence a raceme				
	Racemes	Spikelets	Spikelets	Upper glume length compared to spikelet
<i>Digitaria breviglumis</i>	1-4	clustered at each node or in pairs	glabrous	30-40%
<i>Digitaria brownii</i>	1-7	pairs	long hairs	85-100%
<i>Digitaria gibbosa</i>	1	clustered at each node or in pairs	long hairs	60-95%
<i>Digitaria leucostachya</i>	1-4	clustered at each node or in pairs	long hairs	100%

***Digitaria ammophila* - Silky Umbrella Grass, Spiker Grass**

Derivation

ammophila - from the Greek *amos* (sand) and *phileo* (love); growing in sandy habitats.

Habit

A perennial, tufted grass, the culms are 15-60 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes 5-15, the lowest whorled, the primary branches are lacking spikelets on the basal portion. The panicle is 10-20 cm long. The spikelets grows in solitary or in pairs, they are densely woolly-hairy, 1.8-2.9 mm long. Upper glume 100% of the length of spikelets.

Habitat

This species grows in a range of habitats, usually grasslands or low open vegetation.



Spikelets



Lower branches of the inflorescence



***Digitaria bicornis* - Hairy Finger Grass**

Derivation

bicornis - two horns, probably the paired spikelets are different

Habit

An annual, tufted grass, the culms are 20-80 cm tall.

Inflorescence

The panicle is digitate, racemes 2-4, 5-10 cm long. The spikelets grows in pairs, 2.9-3.5 mm long. The lower spikelet is glabrous; the upper spikelet has a mix of sericeous (silky) hairs and bristles, both spreading and distinct at maturity. Upper glume is 66% of the length of spikelets.


Habitat

This species grows in a range of habitats, forest, woodlands, shrublands, and grasslands.

This species may be misidentified in the field as *Digitaria ciliaris*.



Spikelets



Digitaria breviglumis

Derivation

breviglumis- from the Latin *brevis* (short) and *gluma* (husk). One or both glumes short with respect to the length of the spikelet.

Habit



A perennial, tufted grass, the culms are 30-60 cm.

Inflorescence

Inflorescence is often as a raceme and also a once-branched panicle with racemose branches. Racemes 1-4, 2-10 cm long. The spikelets are clustered at each node or in pairs, glabrous, 1.3-1.6 mm long. The upper glume 30-40% of the length of spikelets

Habitat

Usually in rocky, stony sites, in crevices, on shallow, often sandy soil.



***Digitaria brownii* - Cotton Panic Grass**

Derivation

brownii - in honour of Robert Brown (1773-1858), a Scottish-born English botanist.

Habit

A perennial, tufted grass, the culms are 30-75 cm tall.

Inflorescence

The panicle is of subdigitate, racemes 1-7, 5-18 cm long. The spikelets grow in pairs, 2.6-3.5 mm long, have long hairs. The upper glume is 85-100% of length of spikelet.

Habitat

This species grows in a range of habitats, forest, woodlands, shrublands, and grasslands.



Spikelet



***Digitaria ciliaris* - Summer Grass, Crab Grass**

Derivation

ciliaris - from the Latin *cilium* (eyelid) and *-are* (pertaining to), the glumes or lemmas ciliate on nerves or margins.

Habit

An annual grass, the culms are 10-100 cm tall.

Inflorescence

The panicle is subdigitate, racemes 2-12, 6-22 cm long. The spikelets grows in pairs, are hairy, The upper glume is 60-75% of the length of spikelets.

Habitat

This non-native species is common weed of cultivation, especially in gardens.



Spikelets



***Digitaria didactyla* - Queensland Bluegrass**

Derivation

didactyla - from the Greek *dis* (twice) and *daktylos* (finger), the inflorescence of two racemes.

Habit


A perennial, mat forming grass, the culms decumbent or prostrate, 10-60 cm, tall.

Inflorescence

The panicle is digitate; racemes 2-4, 3-10 cm long. The spikelets grows in pairs, with fine white hairs along the margins, 2-2.7 mm long. The upper glume is 60-75% of the length of spikelets.

Habitat

This non-native species grow as a lawn grass and now occurs in disturbed sites and is most competitive in moist sandy soils.



Digitaria diffusa

Derivation

diffusa- Latin for widely spaced.

Habit

A perennial, decumbent grass with short rhizomes, the culms decumbent 10-40 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes 3-10, 3-7 cm long, glabrous or hairy. The spikelets grow in pairs, hairy, 1.25-2 mm long. The upper glume is 100% of length of spikelet.

Habitat

This species grows in woodlands and forests.



***Digitaria eriantha* - Pangola Grass**

Derivation

eriantha - from the Greek *erion* (wool) and *anthos* (flower), it has with hairy glumes, lemmas or awns.

Habit

A perennial, tufted grass, the culms are 40-120 cm tall.

Inflorescence

The panicle is digitate or subdigitate, racemes 3-10, 5-20 cm long. The spikelets grow in pairs, are hairy, 2.2-3.5 mm long. The upper glume is 60-75% of the length of spikelets.

Habitat

This non-native species grows relatively well in various soils but grows especially well in moist soils. It is used for hay, grazing or silage.



Spikelet



Digitaria fumida

Derivation

fumida - (unknown).

Habit


A perennial, decumbent grass, the culms are 30-80 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes (2-)6-10 racemes, 3.5-15 cm long. The spikelets are clustered at each node or in pairs or threes, glabrous, 1.3-1.9 mm long. The upper glume is 10-25% of length of spikelet. This species differs from *Digitaria orbata* by having a lower glume and differs from *Digitaria breviglumis* by having lots more of racemes.

Habitat

This species grows in a range of *Eucalyptus* woodlands on mainly sandy soils.



Digitaria gibbosa

Derivation

gibbosa - from the Latin *gibba* (swelling) and *-osa* (indicating abundance), the spikelets swollen asymmetrically.

Habit


An annual, erect grass, the culms are 25-40 cm tall.

Inflorescence

The panicle is a raceme, 4-18 cm long. The spikelets are clustered at each node or pairs, have long hairs, 1.5-2.8 mm long. The lower glume is usually absent, the upper glume is 60-95% of length of spikelet.

Habitat

This species grows in woodlands and coastal grasslands.



Spikelet



***Digitaria insularis* - Sourgrass**

Derivation

insularis - of an island.

Habit

A perennial, tufted grass, the culms are up to 150 cm tall.

Inflorescence


Inflorescence a once-branched panicle with racemose branches. Racemes 8-50, up to 30 cm long, congested. The spikelets grow in pairs, have long hairs, 3.5-5 mm long. The upper glume is 100% of length of spikelet.

Habit

This non-native species is generally in moist conditions, along creeks.



Spikelet



Spikelet



Digitaria leucostachya

Derivation

leucostachya - from the Greek *leukos* (white) and *stachys* (ear of corn), the spikelets are white and sometimes invested with copious white hairs.

Habit

A perennial, densely tufted grass, the culms are 40-100 cm tall.

Inflorescence

The panicle is digitate or subdigitate, racemes 1-4, 12-25 cm long. The spikelets are clustered at each node or in pairs, have long hairs, the lower glume is absent or a minute hyaline rim. The upper glume is 100% of length of spikelet.

Habitat

This species grows in sandy soils in coastal areas.



Spikelets



Digitaria longiflora

Derivation

longiflora - from the Latin *longus* (long) and *flos* (flower), it has long spikelets.

Habit


An annual or perennial, mat forming grass, the culms are 10-60 cm tall.

Inflorescence

The panicle is digitate, racemes 2-4, 1-10 cm long. The spikelets are clustered or in threes, are hairy, 1.2-2.3 mm long. The lower glume is absent or obscure, and the upper glume is 100% of the length of spikelets.

Habitat

This species occurs in diverse habitats.



Spikelets



Digitaria minima

Derivation

minima - Latin for least, the smallest of the known species.

Habit

A perennial, decumbent grass, the culms are 30-60 cm tall.

Inflorescence


Inflorescence a once-branched panicle with racemose branches. Racemes 2-7, 0.7-3.5 cm long. The spikelets grows in pairs, glabrous, 1.2-1.55 mm long. The upper glume is 75-85% of length of spikelet

Habitat

This species grows in woodlands.



Spikelets



Digitaria nematostachya

Derivation

nematostachya - from the Greek *nema* (thread) and *stachys* (ear of corn); the branches of panicle are thread-like.

Habit

A perennial grass, the culms are 35-55 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes 10-20, the lowest whorled, and the primary branches are lacking spikelets on the basal portion. Central inflorescence axis 2.5-11 cm long. The spikelets grow in pairs, glabrous, 1.7-2.1 mm long. The upper glume is 50-90% of length of spikelet.

Habitat

This species grows in woodlands or grasslands.



Spikelets



Lowest branches of the inflorescence

Digitaria orbata

Derivation


orbata - from the Latin *orbo* (deprive), the glumes are reduced or absent.

Habit

A perennial, tufted grass, the culms are 40-120 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes (2-)6-10, 7-16 cm long. The spikelets are clustered at each node or in pairs, glabrous, 1.3-1.9 mm long. The lower glume is absent or obscure, the upper glume is 10-25% of length of spikelet. This species differs from *Digitaria fumida* by not having a lower glume and differs from *Digitaria breviglumis* by having lots more of racemes.



Habitat

This species grows in a range of woodlands on a variety of soils.



***Digitaria parviflora* - Small Flower Finger Grass**

Derivation

parviflora - from the Latin *parvus* (small) and *flos* (flower), the spikelets are small or with few florets.

Habit


A robust, perennial grass, the culms are 50-150 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes 4-15, 6-20 cm long. The spikelets grows solitary or clustered at each node or in pairs, glabrous, 1.6-2.1 mm long. The upper glume is 100% of length of spikelet.

Habitat

This species grows in coastal woodlands; it is common in shaded forests.



Spikelets



Digitaria ramularis

Derivation

ramularis - from the Latin *ramulus* (small branch) and *-are* (pertaining to).
Inflorescence much branched.

Habit

A perennial, erect grass, the culms are 30-90 cm tall.

Inflorescence

Inflorescence a once-branched panicle with racemose branches. Racemes 2-10, 3-13 cm long. The spikelets are solitary or clustered at each node or in pairs, glabrous or pubescent, 1.5-2 mm long. The lower glume is absent or obscure, and the upper glume is 50-75% of the length of spikelets.

Habitat

This non-native species prefers coarse-textured soils in disturbed habitats and woodland margins.



Spikelets



***Digitaria violascens* - Purple Crabgrass**

Derivation

violascens - Latin for 'becoming violet' (significance unknown).

Habit

An annual, mat forming grass, the culms are 20-60 cm tall.

Inflorescence

The panicle is digitate or subdigitate; racemes 2-6, 3-14 cm long. The spikelets are clustered at each node or in threes, hairy, 1.2-2 mm long. The lower glume is absent or obscure, and the upper glume is 80-100% of the length of spikelets.

Habitat

This non-native species prefers coarse-textured soils in disturbed habitats and woodland margins.



Spikelets




Dinebra


From Arabic *denab* (queue), alluding to the long inflorescence, or a corruption of Arabic *danaiba* (a little tail), alluding to acuminate glumes.

Annual or perennial, tufted to decumbent. The inflorescence is a raceme of numerous small spikes which become deflexed (bent or curving downward or backward) at maturity. The spikelets are laterally compressed. The spikelets are solitary and all similar with 1-several florets per spikelet; they are very similar to the genera *Diplachne* and *Eragrostis*. The species *Dinebra neesii* is usually a one-flowered spikelet and can be confused with *Sporobolus* species.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 46, Australia = 11



Dinebra decipiens spikelet



Dinebra neesii spikelet




Dinebra neesii spikelets with one floret

Townsville species


Dinebra decipiens Slender Cane Grass
Dinebra neesii Swamp Grass

Key to the species of *Dinebra* and similar species

1. Inflorescence a once-branched open panicle2
- Inflorescence a panicle with secondary branches5



Inflorescence once branched



Inflorescence with secondary branches

2. Spikelets with usually one florets.....*Dinebra neesii*
- Spikelets with 2-14 florets3
3. Spikelets 5-18 mm long, with 4-14 florets, fertile florets 5-18 mm long*Diplachne fusca*
- Spikelets 1-6 mm long, with 2-9 florets, fertile florets 2.5-6 mm long *Dinebra decipiens*
4. Spikelets with one floret*Sporobolus*
- Spikelets with three or more bisexual florets..... *Eragrostis*

Dinebra decipiens - Slender Cane Grass

Derivation

decipiens - from the Latin *decipio* (deceive), it resembles another species, or in some other way deceptive.

Habit

A perennial, tufted grass, the culms are erect and hollow to wiry, 15-150 cm tall.

Inflorescence

The inflorescence is a once-branched open or dense. The panicle is 5-50 cm long with and the spikelets with 3-9 florets (2-8 fertile florets).

Habitat

This species is found with the margins swamps or creek and riverbanks.



***Dinebra neesii* - Swamp Grass**

Derivation

neesii - in honour of Christian Daniel Nees von Esenbeck (1766-1858), German botanist.

Habit


An annual or biennial grass, tufted, short-lived, the culms are erect and hollow, 60-150 cm tall.

Inflorescence

The inflorescence is a once-branched open or dense panicle. The panicle is 20-40 cm and the spikelets with one (sometimes with two) florets (1 fertile floret).

Habitat

Almost invariably, plants grow on cracking clay plains and floodplains, usually in lakes and similar seasonally flooded depressions. In saline situations, the species is often associated with *Sporobolus virginicus*.



Diplachne

Beetle Grasses


From the Greek *diplos* (double) and *achne* (scale), referring to the two-lobed lemma.

Annuals or perennials, tufted, culms erect, drooping in the upper parts. The inflorescence is a racemose once-branched panicle. The spikelets are solitary and all similar with several florets per spikelet, they are very similar to the genus *Eragrostis*. Lemmas are bifid or minutely notched, very shortly awned or mucronate.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 44 , Australia = 3


Townsville species

- | | |
|--|---------------------|
| <i>Diplachne fusca</i> var. <i>fusca</i> | Brown Beetle Grass |
| * <i>Diplachne fusca</i> var. <i>uninervia</i> | Mexican Sprangletop |




Key to the species of *Diplachne* and similar species

1. Inflorescence a panicle with secondary branches *Eragrostis*
- Inflorescence a once-branched open panicle 2



Inflorescence with
secondary branches



Inflorescence once
branched

2. Spikelets with usually one florets *Dinebra neesii*
- Spikelets with 2-14 florets 3
3. Spikelets 1-6 mm long, with 2-9 florets, fertile florets 2.5-6 mm long *Dinebra decipiens*
- Spikelets 5-18 mm long, with 4-14 florets, fertile florets 5-18 mm long 4
4. Perennial 40-150 cm tall, spikelets 5-13 flowered; lemma of various colours; native common
..... *Diplachne fusca* var. *fusca*
- Annual or biennial to 70 cm tall, spikelets 4-6 flowered; lemma dark green or lead coloured.
non-native not common *Diplachne fusca* var. *uninervia*

***Diplachne fusca* var. *fusca* - Brown Beetle Grass**

Derivation

fusca - Latin for dark or swarthy, the glumes or lemmas are dark brown.

Habit

An erect, tufted perennial or sometimes annual grass, the culms are 25-150 cm tall.

Inflorescence

A panicle of racemes, it is usually fewer than 30 branches. The lemma apex usually acute or acuminate, notched or not, mucronate or not; lemma is usually generally not dark green or lead coloured; the lemma is pubescent or silky about midnerve.

Habitat

This native species is usually in or near water or in depressions, it is salt-tolerant. It flowers usually after rain.



Spikelet



***Diplachne fusca* var. *uninervia* - Mexican Sprangletop**

Derivation

uninervia - from the Latin *unus* (one) and *nervus* (nerve).

Habit

An annual or biennial, sparingly branched grass, the culms 5-70 cm tall.

Inflorescence

A panicle of racemes, often greater than 30 branches. The lemma apex obtuse to truncated, often notched and mucronate; lemma is dark green or lead coloured.

Habitat



This non-native species grows in disturbed areas.



Echinochloa

From Greek *echinos* (hedgehog) and *chloe* (grass), alluding to the echinate (with prickles) inflorescence branches.

Annuals or perennials, tufted or decumbent. Ligule (an outgrowth on the inside of the leaf blade/leaf sheath junction) absent or present. The inflorescence is a once branched or contracted panicle. The spikelets are solitary or clustered, sometimes in distinct rows, awned, mucronate (a sharp, abrupt terminal point) or awnless. The glume and lower lemma usually have stiff, bristly hairs.



Members of this genus are usually found in wet habitats and along roadsides.


Subfamily: Panicoideae; Tribe: Paniceae
World = 60, Australia = 18

Townsville species

- | | |
|----------------------------------|------------------------|
| * <i>Echinochloa colona</i> | Awnless Barnyard Grass |
| * <i>Echinochloa crus-galli</i> | Barnyard Grass |
| * <i>Echinochloa esculenta</i> | Japanese Millet |
| * <i>Echinochloa polystachya</i> | Aleman Grass |

Key to the species of *Echinochloa*

1. Plants perennial; ligule a fringe of long stiff hairs. *Echinochloa polystachya*
- Plants annual; ligule absent 2



2. Spikelets 1.5-3 mm long *Echinochloa colona*
- Spikelets 3-4 mm long 3
3. Racemes simply spaced *Echinochloa crus-galli*
- Racemes closely spaced *Echinochloa esculenta*



Racemes simply spaced



Racemes closely spaced

***Echinochloa colona* - Awnless Barnyard Grass**

Derivation

colona - Latin for colonist, it was planted as a cereal.

Habit

An annual, tufted grass, the culms are 10-100 cm tall. It does not have a ligule.

Inflorescence

The inflorescence is once-branched panicle; the central axis is 1-15 cm. The spikelets are 4-rowed. The spikelets are 15-3 mm long.

Habitat

This non-native species is a weed in waste areas, agricultural lands, along stream banks and around waterholes.



Spikelets



***Echinochloa crus-galli* - Barnyard Grass**

Derivation

crus-galli - from the Latin *crus* (foot) and *gallus* (cock), the panicle resembles a cock's foot.

Habit

An annual, tufted grass, the culms are 25-150 cm tall. It does not have a ligule.

Inflorescence

The inflorescence is a once-branched panicle; the central axis is 2-10 cm. The spikelets are 2-4-rowed. The spikelets are 3-4 mm long.

Habitat

This non-native species is a weed in waste areas, agricultural lands, along stream banks and around waterholes.



Spikelets



***Echinochloa esculenta* - Japanese Millet**

Derivation

esculenta - Latin for edible, the grain is edible.

Habit

An annual, erect grass, the culms are 30-150 cm tall. It does not have a ligule.

Inflorescence

The inflorescence is a once-branched panicle; the central axis is 7-20 cm long. The spikelets are 4-6-rowed. The spikelets are 3-4 mm long.

Habitat

This non-native species is used as a soil stabilizer, often found on roadsides.



Spikelets



***Echinochloa polystachya* - Aleman Grass**

Derivation

polystachya - from the Greek *polys* (many) and *stachys* (ear of corn), the grass has many-branched culms or inflorescences.

Habit

A perennial, decumbent, aquatic or semi-aquatic grass, the culms are 100-300 cm tall. The ligule is a rim of stiff, yellow hairs.

Inflorescence

The inflorescence is a once-branched panicle; the central axis is 15-30 cm long. The spikelets are 4-rowed. The spikelets are 4.5-6 mm long.

Habitat

This non-native species was introduced to Queensland as a ponded pasture due to its palatability and it has now invaded seasonally flooded areas, swamps, lakeshores and rivers due to its ability to grow in deep water and reproduce vegetatively.

The cultivar 'Amity' commonly planted in Queensland differs from the species norm, in having flowering culms 100-200 cm long,



Ectrosia

From the Greek *ectrosis* (miscarriage), referring to the spikelets with only 1 or 2 basal bisexual flowers, with male or empty lemmas above them.

Tufted annuals and perennials. The inflorescence is an open or contracted panicle, sometimes reduced to a few racemes. The spikelets are solitary with 4 or more florets. This genus is readily recognised by the modification of its upper florets into a dispersal device.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 18, Australia = 15

Townsville species

Ectrosia agrostoides

Ectrosia lasioclada

Ectrosia leporina Hare's Foot Grass

Ectrosia nervilemma



Spikelet, the upper sterile florets act as a dispersal device

Key to the species of *Ectrosia*

1. Florets 8-25, all similar, mainly bisexual; awns 0.5-1.5 mm long *Ectrosia lasioclada*
Florets 2-9, lower bisexual or male, upper sterile; awns up to 12 mm long 2
2. Spikelets 2-3mm long; inflorescence open *Ectrosia agrostoides*
Spikelets 3-7mm long; inflorescence ±congested 3
3. Spikelets 0.5-1.5 mm wide *Ectrosia leporina*
Spikelets 2-6 mm wide *Ectrosia nervilemma*



Ectrosia agrostoides

Derivation

agrostoides - agrostology, the study of grasses,

Habit

An annual, tufted grass, the culms are 10-40 cm tall. The leaves are very narrow, 0.2-0.5 mm wide.

Inflorescence

The inflorescence is a spicate panicle, 1-4.5 cm long. The spikelets are 4-5 flowered, with 1 bisexual floret.

Habitat

This species usually grows on shallow sandy soils in open woodland.



Ectrosia lasioclada

Derivation

lasioclada - from the Greek *lasios* (shaggy) and *klados* (branch), the leaf-sheaths are hairy.

Habit

The grass is usually perennial, sometimes annual; the culms are 40-90 cm tall.

Inflorescence

The inflorescence is a compound panicle, 5-20 cm long. The lemmas awns are usually short and do not increase in length.

Habitat

This species is mainly on sandy soils on the edge of clay plans, in savanna woodland or forest.

Ectrosia lasioclada is the most distinctive species; sometimes it is put in the *Eragrostis* genus.



***Ectrosia leporina* - Hare's Foot Grass**

Derivation

leporina - from the Latin *lepus* (hare) and *-ina* (belonging to), which refers to the inflorescence which resembles a hare's foot.

Habit


This grass is a slender annual, the culms are 10 to 90 cm tall.

Inflorescence

The mature inflorescence is green-purple and has a “furry” appearance, hence its common name. The inflorescence is very dense and is 5-15 cm long

Habitat

This species grows in sandy soils and in open forest.



Ectrosia nervilemma

Derivation

nervilemma - the prominent nerves of the lemmas.

Habit

Annual or ephemeral grass, the culms are 12-40 cm tall. The leaves are flat or conduplicate 1-4 mm wide.

Inflorescence

The inflorescence is a compound panicle, 6-12 cm long. The spikelets are 2-8 fertile flowers, 5-11 mm long.

Habitat

This species grows on sandy soils in open woodland.



Spikelet



Eleusine

Eleusis was named after the Greek town where Demeter, the Greek goddess of agriculture was worshipped, and the Eleusinian festivals, concerning the mysteries of the growth of corn, were celebrated.

Tufted annuals or perennials. The inflorescence is digitate or subdigitate. The spikelets are overlapping in two rows on the underside of the rachis; each spikelet contains 3 or more bisexual florets.

Eleusine species are mostly from Africa; the species *Eleusine coracana* (finger millet) is grown in Africa for human food and for making beer.


Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 45, Australia = 3

Townsville species

* *Eleusine indica* Crowsfoot Grass



Spikelets



Key to the species of *Eleusine indica* and similar species

Axis of each raceme ending with a spikelet *Eleusine indica*
Axis of each raceme ending in a point *Dactyloctenium*



Axis of each raceme ending with a spikelet



Axis of each raceme ending in a point

***Eleusine indica* - Crowsfoot Grass**

Derivation

indica - from India.

Habit


An annual or perennial, erect, prostrate or geniculate, the culms 15-90 cm tall.

Inflorescence

The inflorescence is subdigitate or digitate; it is made up of usually 2-6 digitate racemes at the apex, usually with 1 raceme inserted lower on the culm.

Habitat

A non-native species is usually a weed of roadsides, waste ground and cultivated land.



Elionurus

From the Greek *eleuein* (to roll) and *oura* (tail) alluding to the spike-like racemes curling strongly when old.


Tufted perennials, occasionally annuals, leaves sometimes aromatic. The inflorescence is a single raceme or a series of racemes which are subtended by spathe-like leaf sheaths. The spikelets are in pairs, one sessile and one pedicelled.

Subfamily: Panicoideae; Tribe: Andropogoneae

World = 21, Australia = 2

Townsville species

Elionurus citreus Lemon-scented grass



This species could be confused with *Eremochloa bimaculata*, *Lepturus repens* and *Hemarthria uncinata*. Check the key on page 16.



***Elionurus citreus* - Lemon-scented grass**

Derivation

citreus - Latin for pure yellow, spikelets invested in yellow hairs.

Habit


An erect, slender perennial grass, the culms 50-100 cm tall. When the leaves are crushed, they give off a lemon scent. *Cymbopogon* is another grass genus with lemon-scented leaves.

Inflorescence

The inflorescence is a raceme or a series of simple racemes covered with white hairs, the spathes below the racemes are not obvious. When the spikelets mature, the pedicelled spikelet sticks out at right angles to the raceme giving it a distinctive appearance. The lower glume of the sessile spikelet is often asymmetrical; the apex is drawn out and forked at the tip.

Habitat

This species grows in sandy soil along rivers and in coastal areas and sand dunes.



Elytrophorus

From the Greek *elytron* (sheath or husk) and *phero* (to bear), referring to the large outer glume.

Tufted annuals. The inflorescence is a panicle of dense globular clusters borne at intervals along the central axis. The spikelets are strongly laterally compressed with 2-7 bisexual florets. The lemmas are awned.


Subfamily: Danthonioideae; Tribe: Danthonieae
World = 3, Australia = 1

Elytrophorus spicatus is the only Australian species from this genus.

Townsville species

Elytrophorus spicatus

Spike Grass




Key to the species of *Elytrophorus* and similar species


1. Inflorescence an interrupted spicate panicle *Elytrophorus spicatus*
- Inflorescence an open or contracted panicle 2



Interrupted
spicate
panicle



Open panicle



Contracted panicles

2. Inflorescence a contracted panicle; spikelets with one or two basal bisexual flowers, with male or empty lemmas above them, the upper lemmas reduced to awns *Ectrosia*
- Inflorescence an open panicle, spikelets with two bisexual florets only *Eriachne*

***Elytrophorus spicatus* - Spike Grass**

Derivation

spicatus - from the Latin *spica* (thorn) and *-ata* (possessing), refers to the inflorescence which is a spike or spicate panicle.

Habit

A tufted annual grass, the culms are unbranched, arising from the base, 10-40 cm tall.

Inflorescence

The inflorescence is a distinctly interrupted spicate panicle; it is often $\frac{3}{4}$ or more of the entire culm. The panicle is 15-35 cm long and 0.5-0.8 cm wide. The spikelets are numerous in dense clusters; each one spikelet has 5-7 florets.

Habitat

This species occurs on clay soils, they are found in damp soils along creeks and damp hollows and seepages.



Enneapogon

Nineawn Grasses

From the Greek *ennea* (nine) and *pogon* (beard), alluding to the nine, plumose lemma awns.


Tufted perennials or annuals. The inflorescence is a spicate panicle (a spike-like panicle with short branches). The spikelets are solitary. The genus is readily recognised by the nine-awned lemmas and with each spikelet having 2 or more florets. The spreading awns form an attractive circular arrangement.

Species from this genus grow in a variety of habitats, from riverbeds to rocky slopes.

Subfamily: Chloridoideae; Tribe: Pappophoreae
World = 27, Australia = 16

Townsville species

<i>Enneapogon lindleyanus</i>	Nineawn Grass
<i>Enneapogon nigricans</i>	Black heads
<i>Enneapogon polyphyllus</i>	Leafy Nineawn
<i>Enneapogon robustissimus</i>	Nineawn Grass
<i>Enneapogon truncatus</i>	
<i>Enneapogon virens</i>	



Spikelet

Key to the species of *Enneapogon*

1. Body of basal lemma with appressed hairs on its entire surface *Enneapogon virens*
Body of basal lemma totally covered by long, erect or spreading hairs or partly glabrous.....2
2. Body of basal lemma obscured by dorsal hairs which exceed base of awns*Enneapogon polyphyllus*
Body of basal lemma partly exposed as dorsal hairs do not or scarcely reach base of awns3
3. Panicle less than 4.0 cm long*Enneapogon lindleyanus*
Panicle usually more than 4.0 cm4
4. Palea longer than of its lemma by about 1 mm *Enneapogon truncatus*
Palea similar in length to body of its lemma.....
5. Culms pubescent; panicle linear-oblong.....*Enneapogon nigricans*
Culms glabrous; panicle ovate or pyramidal *Enneapogon robustissimus*



***Enneapogon lindleyanus* - Nineawn Grass**

Derivation

lindleyanus - is from John Lindley (1799-1856), an English botanist.

Habit

A perennial or annual, tufted grass, culms 10-65 cm tall.

Inflorescence

The inflorescence is a spicate panicle, 0.5-2.5 cm long, 0.5-1 cm wide (including awns). The spikelet has 3-5 florets (usually 1 bisexual). It is characterised by small, compactly capitate inflorescence.

Habitat

This species usually grows on hills.



Spikelet



***Enneapogon nigricans* - Black-heads**

Derivation

nigricans - from the Latin *nigrico* (become blackish), becoming black with maturity, usually of spikelets.

Habit


A perennial, tufted grass, the culms are 30-70 cm tall.

Inflorescence

The inflorescence is a spicate panicle, 0.8-9.5 cm long, 0.8-2.8 cm wide (including awns). The spikelet has 3-5 florets (1 bisexual). The body of basal lemma partly exposed as dorsal hairs do not or scarcely reach base of awn. The glumes apices are obtuse to acuminate.

Habitat

This species is uncommon in Townsville, it occurs on Magnetic Island. It often grows on red and red-brown earths.



Spikelet



***Enneapogon polyphyllus* - Leafy Nineawn**

Derivation

polyphyllus - from the Greek *polys* (several) and *phyllon* (leaf), the culms many-leaved in comparison with related species.

Habit


A perennial or annual, tufted grass, culms 3-50 cm tall. The leaf blades are involute or convolute.

Inflorescence

The inflorescence is a spicate panicle, linear, mostly 4-9 cm long, 1-2 cm wide (including awns). The spikelet has 4-6 (1 bisexual), the 4th and 5th floret are reduced to lemma and reflexed or horizontal. The body of basal lemma totally covered by long, erect or spreading hairs or partly glabrous. The glumes apices are acute or mucronate.

Habitat

This species grows mainly on sandy or loamy red earth soils.



Spikelet



***Enneapogon robustissimus* - Nineawn Grass**

Derivation

robustissimus - Latin for most robust, the culms are very tall for the genus.

Habit

A perennial, tufted grass, culms 30-110 cm tall.

Inflorescence

The inflorescence is a spicate panicle; it is often pyramidal shape, 1.5-10 cm long, 1.5-2.5 cm wide (including awns). The spikelet has 3-5 florets (1 bisexual)

Habitat

This species grows on sandy soils, near creeks and in gullies on rocky hills.



Spikelet



Enneapogon virens

Derivation

virens - from the Latin *vireo* (be green), leaf-blades or culms unusually bright yellow green.

Habit

A perennial, tufted grass, the culms are 10-60 cm tall.

Inflorescence


The inflorescence is a spicate panicle, 3-10 cm long, 1 cm wide (including awns). The spikelets have 3-4 florets (1 bisexual). The body of basal lemma is hairy all over the back, the hairs not reaching the base of the awns.

Habitat

This species usually grows in open woodland, in well-drained country.



Spikelet



Enteropogon

Windmill Grasses

From the Greek *enteron* (intestine) and *pogon* (a beard), perhaps alluding to the beards on the callus or in the axils of the spikes.


Tufted perennials. The inflorescence is a simple spike, or many spikes arranged digitately and spreading at maturity. The spikelets are solitary, dorsally compressed with 2-3 florets and arranged on one side of the spikes. The lemmas are awned.

Enteropogon is closely related and easily confused with the genera *Chloris*.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 22, Australia = 8

Townsville species

Enteropogon acicularis Curly Windmill Grass
Enteropogon ramosus Curly Windmill Grass



Key to the species of *Enteropogon* and similar species

1. Lowest lemma laterally compressed (lying on the side when placed on a flat surface) *Chloris*
 Lowest lemma dorsally compressed (lying on front or back when placed on a flat surface) 3



Chloris lemma



Enteropogon lemma

2. Leaves arising from the whole culm length; culms branched at most nodes with up to 8 branches;
 spikes 2-8 spreading in a single plane *Enteropogon ramosus*
 Leaves usually arising from the base of the culm; culms usually solitary; spikes 7 or more,
 arranged in several planes *Enteropogon acicularis*



Enteropogon ramosus



Enteropogon acicularis

Enteropogon acicularis - Curly Windmill Grass

Derivation

acicularis - from the Latin, meaning needle-shaped.

Habit

A perennial grass, the many culms are 25-60 cm tall.

Inflorescence

The inflorescence is digitate with 7 or more spikes, 8-17 cm long.

Habitat

This species often grows on heavier-textured red, brown, grey and black soils.



Spikelet



***Enteropogon ramosus* - Curly Windmill Grass**

Derivation

ramosus - from the Latin *ramus* (branch) and *-osa* (abundance), referring to the much-branched inflorescences or culms.

Habit

A perennial, tufted grass, the culms grow to 90 cm tall.

Inflorescence

The inflorescence is digitate with 2-8 spikes, 4-20 cm long.

Habitat

This species grows on sandy loams, red earths and skeletal soils.



Spikelets




Eragrostis

Lovegrasses

There are several possible derivations of the name of which the three following are the most common, all are from the Greek. 1. *eros* (love) and *agrostis* (a grass). 2. *er* (early), because many species are pioneer invaders of bare ground. 3. *eri* (to strengthen a noun), that is a many flowered *Agrostis*.

Annuals or perennials, tufted or decumbent or stoloniferous. The inflorescence is an open or contracted panicle with secondary branching. The spikelets are solitary. The genus is characterised by its usually numerous, identical florets (3 or more), and some species have persistent paleas. The photograph to the right shows the persistent paleas at the base of the spikelet.

Although it is easy to recognise the genus *Eragrostis*, the species are considered difficult to identify. There is a large number of species, and a number of species are polymorphic and intergrade. Superficially similar to *Sporobolus* and *Leptochloa* species.



Spikelet

Species of *Eragrostis* are commonly early invaders of arable land, are often on poor or sandy soils or disturbed ground. There are 25 species in Townsville.

Subfamily: Chloridoideae; Tribe: Cynodonteae

World = 475, Australia = 98




Townsville species

<i>Eragrostis basedowii</i>	Neat Lovegrass
<i>Eragrostis brownii</i>	Brown's Lovegrass
* <i>Eragrostis cilianensis</i>	Stinking Lovegrass
<i>Eragrostis cumingii</i>	Cuming's Lovegrass
* <i>Eragrostis curvula</i>	African Lovegrass
<i>Eragrostis dielsii</i>	Mallee Lovegrass
<i>Eragrostis elongata</i>	Clustered Lovegrass
<i>Eragrostis exigua</i>	Delicate Lovegrass
<i>Eragrostis fallax</i>	
<i>Eragrostis interrupta</i>	
<i>Eragrostis lacunaria</i>	Purple Lovegrass
<i>Eragrostis leptostachya</i>	Paddock Lovegrass
* <i>Eragrostis mexicana</i>	Mexican Lovegrass
* <i>Eragrostis minor</i>	Small Stink Grass
<i>Eragrostis parviflora</i>	Weeping Lovegrass
* <i>Eragrostis pilosa</i>	Soft Lovegrass
<i>Eragrostis pubescens</i>	
<i>Eragrostis schultzii</i>	
<i>Eragrostis sororia</i>	
<i>Eragrostis spartinoides</i>	
<i>Eragrostis stenostachya</i>	
* <i>Eragrostis tenella</i>	Delicate Lovegrass
<i>Eragrostis tenellula</i>	Delicate Lovegrass
* <i>Eragrostis tenuifolia</i>	Elastic Grass
<i>Eragrostis unioides</i>	

Key to the species of *Eragrostis* and similar species

1. Spikelets with one floret *Sporobolus*
Spikelets with three or more florets 2
2. Lemmas with blunted apex *Eragrostis*
Lemmas with pointed apex *Ectrosia lasioclada*



Sporobolus




Ectrosia lasioclada



Eragrostis

Key to species of *Eragrostis*


1. Plants distinctly glandular on culms below nodes or on leaf margins, inflorescence branches, pedicels, glumes or lemmas2
- Plants not or sometimes faintly glandular6




Gland on pedicel



Gland on culms



Gland on inflorescence branches and pedicels



Gland below inflorescence

2. Perennials*Eragrostis leptostachya*
- Annuals3
3. Plants glandular on culms below nodes and/or leaf margins*Eragrostis mexicana*
- Plants glandular on inflorescence branches, pedicels, glumes or lemmas4
4. Spikelets 2-4 mm wide*Eragrostis cilianensis*
- Spikelets less than 2 mm wide5
5. Palea keels as long as the palea and not dividing at apex; glumes keeled *Eragrostis minor*
- Palea keels shorter than the palea and dividing apically into 3 small lobes; glumes rounded on back *Eragrostis pilosa*
6. Spikelets terete or \pm biconvex7
- Spikelets \pm laterally compressed8
7. Lemma with lateral nerves closer to margin than to mid-nerve*Eragrostis lacunaria*
- Lemma with lateral nerves mid-way between margin and mid-nerve or sometimes closer to latter*Eragrostis dielsii*

8. Spikelets to 1 mm wide9
Spikelets wider than 1 mm 11
9. Spikelets 1 mm long, 3-5-flowered *Eragrostis exigua*
Spikelets more than 2 mm long, 4-9-flowered..... 10
10. Inflorescence 2-14 cm long *Eragrostis tenella*
Inflorescence 10-50 cm long..... *Eragrostis tenellula*



Eragrostis tenella



Eragrostis tenellula

11. Lateral nerve of lemma distinct 12
Lateral nerve of lemma indistinct 12



Lateral nerve of lemma distinct



Lateral nerve of lemma indistinct

12. Lemma 2.5-3.5 mm long 13
Lemma less than 2.5 mm long.....3
13. Pedicels less than 0.5 mm long; inhabits coastal dunes..... *Eragrostis interrupta*
Pedicels more than 0.5 mm long; inhabits sclerophyll woodlands and forests *Eragrostis pubescens*



Eragrostis interrupta



Eragrostis pubescens

14. Inflorescence comprising more than half height of plant..... *Eragrostis cumingii*
Inflorescence comprising less than half height of plant..... 15



Eragrostis cumingii

- 15. Spikelet on pedicels 0.1-0.5 mm long 16
- Spikelet on pedicels more than 0.5 mm long 24
- 16. Some spikelets bunched in clusters 17
- Spikelets not bunched in clusters 20



Spikelets grouped together



Spikelets spread through the inflorescence



- 17. Palea prominently ciliate *Eragrostis basedowii*
- Palea glabrous or minutely ciliate 18



- 18. Spikelets 1.5-2 mm wide, 6-14-flowered *Eragrostis elongata*
- Spikelets 1.5-2.2 mm; 9-34 florets *Eragrostis schultzei*
- Spikelets 2-3.3 mm wide, 11-56-flowered 19

19. Florets shed from apex of spikelet downwards, falling entire, each attached to its rachilla segment *Eragrostis sororia*
 Florets ripening and shed from base of spikelet upwards..... *Eragrostis fallax*
20. Florets shed from apex of spikelet downwards, falling entire, each attached to its rachilla segment *Eragrostis brownii*
 Florets ripening and shed from base of spikelet upwards..... 21



Florets shed from apex
of downwards

Florets ripening and shed from
base of spikelets upwards

21. Spikelet on pedicels more than 0.5 mm long *Eragrostis spartinoides*
 Spikelet on pedicels 0.1-0.5 mm long..... 22
22. Spikelets 1-1.5 mm wide *Eragrostis stenostachya*
 Spikelets 1.5-4 mm wide 23
23. Florets closely imbricated; spikelet 2.5-4 m wide *Eragrostis uniolooides*
 Florets not closely imbricated; spikelets 1.5-2.5 wide *Eragrostis spartinoides*



Eragrostis stenostachya



Eragrostis uniolooides



Eragrostis spartinoides

24. Inflorescence \pm weeping *Eragrostis parviflora*
 Inflorescence erect 25
25. Inflorescence branches hairy in their axils *Eragrostis tenuifolia*
 Inflorescence branches not hairy in their axils *Eragrostis curvula*



Eragrostis parviflora



Eragrostis tenuifolia



Eragrostis curvula

Groups of *Eragrostis* species

A	B	C	D	E
Spikelets terete or biconvex	Small spikelets, 1 mm wide, 4-10 florets	Plants distinctly glandular on culms below nodes or on leaf margins, inflorescence branches, pedicels, glumes and/or lemmas	Florets disarticulating from the base of the spikelets	Florets falling entire from spikelets apex downwards
<i>E. dielsii</i> <i>E. lacunaria</i>	<i>E. exigua</i> <i>E. tenella</i> <i>E. tenellula</i>	<i>E. cilianensis</i> <i>E. leptostachya</i> <i>E. mexicana</i> <i>E. minor</i> <i>E. pilosa</i> <i>E. tenella</i>	<i>E. cumingii</i> <i>E. curvula</i> <i>E. fallax</i> <i>E. interrupta</i> <i>E. parviflora</i> <i>E. pilosa</i> <i>E. pubescens</i> <i>E. schultzei</i> <i>E. spartinoide</i> <i>E. stenostachya</i> <i>E. tenuifolia</i> <i>E. unioloides</i>	<i>E. basedowii</i> <i>E. brownii</i> <i>E. elongata</i> <i>E. exigua</i> <i>E. sororia</i> <i>E. tenella</i> <i>E. tenellula</i>

Native *Eragrostis*

Eragrostis basedowii
Eragrostis brownii
Eragrostis cumingii
Eragrostis dielsii
Eragrostis elongata
Eragrostis exigua
Eragrostis fallax
Eragrostis interrupta
Eragrostis lacunaria

Eragrostis leptostachya
Eragrostis parviflora
Eragrostis pubescens
Eragrostis schultzei
Eragrostis sororia
Eragrostis spartinoide
Eragrostis stenostachya
Eragrostis tenellula
Eragrostis unioloides

Weedy *Eragrostis*

Eragrostis cilianensis
Eragrostis curvula
Eragrostis mexicana
Eragrostis minor
Eragrostis pilosa
Eragrostis tenella
Eragrostis tenuifolia

***Eragrostis basedowii* - Neat Lovegrass**

Derivation

basedowii - in honour of Herbert Basedow (1881-1933) a South Australian geologist.

Habit

A tufted annual grass, the culms 5-30 cm tall.

Inflorescence

The panicle are terminal and axillary, 2.5-10 cm long and 1-6 cm wide. The terminal panicles spiciform, sometimes interrupted in lower part. The florets are 10-30, the palea keels with spiny hairs. The rachilla of the spikelet is straight. The spikelets are usually 11-16 mm long.

Habitat

This species occurs in or near watercourses, clay plains, lagoons, saline depressions and sand ridges.



Eragrostis brownii - Brown's Lovegrass

Derivation

brownii - in honour of Robert Brown (1773-1858), a Scots-born English botanist.

Habit

A slender compactly tufted perennial grass, the culms are striate or ribbed, 10-60 cm tall.

Inflorescence

The panicle is open or contracted, 5-30 cm long and 2-20 cm wide. The florets are 8-25, these are falling entire from spikelets apex downwards. The spikelets are 4-12.5 mm long.

Habitat

The species occurs in sandy, red earth, loamy soils; it occurs often on skeletal and stony or gravely hillslopes.

A polymorphic species, highly variable which appears to intergrade with *Eragrostis elongata* and *Eragrostis sororia*. In the past, the name was widely misapplied to several other species e.g. *Eragrostis spartinoide*s *Eragrostis cumingii* and *Eragrostis pubescens*.



Spikelet



***Eragrostis cilianensis* - Stinking Lovegrass**

Derivation

cilianensis - from Cigliano Italy.

Habit

An annual, tufted grass, the culms are erect or ascending 10-90 cm tall, often odorous. The nodes are often purple black, there are pit-like or warty glands on the stems, just below the nodes and on the leaves.

Inflorescence

The panicle is open or dense, 4-30 cm long and 1.5-10 cm wide. The inflorescence are glandular. The spikelets are 5-20 mm long and 2-4 mm wide, the florets are 5-35, these are olive-green.


Habitat

The species is non-native and grows as a weed of lawns, gardens, roadsides and cultivated crops. It occurs in red and brown soils, sandy and clayey loams, heavy clay clays, alluvial and red basaltic soils.

There are similarities to *Eragrostis minor*, the spikelets are wider, 2-4 mm wide.



Spikelet



***Eragrostis cumingii* - Cuming's Lovegrass**

Derivation

cumingii - in honour of Hugh Cuming (1791-1865), English naturalist and traveller.

Habit

An annual, variable in habit, 14-70 cm tall.

Inflorescence

The panicle is open or spiciform, 4-70 cm long and 1.5-11.5 cm wide. It is often the inflorescence comprises half height of plant. The florets are 10-20, and the rachilla of the spikelet is straight. The spikelets are 4.5-35 mm long.

Habitat

The species is widespread in seasonally wet, sometimes saline sites. It occurs in deep often alluvial sands, clays, loams or in shallow soils.

A polymorphic species, they are very variable in size, branching habit and panicle structure. It intergrades with *Eragrostis basedowii*. It is confused with the perennial *Eragrostis spartinoides* and *Eragrostis brownii*.



Spikelet



***Eragrostis curvula* - African Lovegrass**

Derivation

curvula - from the Latin *curvus* (bent) and *-ula* (diminutive) and refers to the curved leaf-blades.

Habit

A densely tufted perennial grass, the culms 30-120 cm tall. The basal leaf sheath are hairy on the back

Inflorescence

The panicle is open or dense, 6-30 cm long and to 20 cm wide. The spikelets are dark olive-grey becoming cream. The florets are 4-13. The lateral nerve of the lemma are indistinct. The spikelets are 4-10 mm long.

Habitat

This non-native species occurs on disturbed sites.

A variable species complex. It is regarded as a serious weed.



Spikelet



The basal leaf sheath are covered in fair hairs

Eragrostis dielsii - Mallee Lovegrass

Derivation

dielsii - in honour of Friedrich Ludwig Emil Diels (1874-1945), a German botanist.

Habit


A tufted annual or short-lived perennial grass, the culms erect or prostrate 3-55 cm tall.

Inflorescence

The panicle is contracted or spiciform, 2-17 cm long and 1-7 cm wide. The spikelets are biconvex or terete, and they are 4.5-50 mm long.

Habitat

This species is found on seasonally flooded clays to sandy and saline soil. In Townsville it is found in the salt pans in Rowes Bay's wetlands.



Spikelet



***Eragrostis elongata* - Clustered Lovegrass**

Derivation

elongata - Latin for elongated. Inflorescence elongated.

Habit

A tufted perennial, culms erect 21-80 cm tall.

Inflorescence

The panicle is a spiciform with spikelets in compact, interrupted clusters, 9-20 cm long and 1-7.5 cm wide. The spikelets are 3-20 mm long and with 6-27 florets.

Habitat

This species grows on a wide range of soil types, often in or near alluvial, well-watered habitats, also on disturbed sites.

Eragrostis elongata can hybridize with *Eragrostis sororia*.



Spikelet



***Eragrostis exigua* - Delicate Lovegrass**

Derivation

exigua - Latin for wanting in size or number, the spikelets or inflorescence branches few.

Habit

An annual, slender, tufted grass, the culms are 15-70 cm tall.

Inflorescence

The panicle is loosely contracted, often occupying 2/3-5/6 of the plant, 18-36 cm long and 3-7 cm wide. The spikelets are 0.8-2 mm long and with 3-7 florets.

Habitat

The species is usually found in seasonally flooded, alluvial habitats with heavy clay (rarely sandy) soils.

It is similar to *Eragrostis tenellula*, but the spikelets are smaller. *Eragrostis exigua* has spikelets 2 mm or shorter and has 3-7 florets.



Eragrostis fallax

Derivation

fallax - Latin for deceptive, closely resembling another species.

Habit

A perennial, densely tufted grass, the culms are 40-90 cm tall.

Inflorescence

The inflorescence are terminal and axillary. The axillary panicle are especially near the base of the plant. The terminal inflorescence is 10-20 cm long and 4-6 cm wide. The spikelets are 2-3.3 mm wide and with 11-56 florets.

Habitat

The species occurs in or nearly seasonally flooded sites (lagoons, swamps, floodplains, streams) in alluvial clay or sandy soils.



Spikelets



Eragrostis interrupta

Derivation

interrupta - Latin for not continuous, spikelets or inflorescence branches clustered at intervals along an axis.

Habit

A perennial, often prostrate, grass, the culms are 50-150 long, often pruinose or glaucous. The leaves are blue green.

Inflorescence

The panicle is contracted or spiciform, usually interrupted, 20-60 cm long and 1-1.5 cm wide. The spikelets are 10-35 mm and with 14-50 florets.

Habitat

The species occur in coastal and subcoastal areas on beach sands and dunes.

The species resemble *Eragrostis pubescens*, but the foliage and panicle are glabrous.



Eragrostis lacunaria - Purple Lovegrass

Derivation

lacunaria - from the Latin *lacuna* (cavity) and *-aria* (pertaining to), surface of grain pitted.

Habit

An annual or short-lived perennial grass, the culms are 12-60 cm tall.

Inflorescence

The panicle is open often comprising about ½ the plant, 4-24 cm long and 2-12 cm wide. The spikelets are biconvex or terete, are often purple-coloured, are 5-22.5 mm long and with 9-43 florets.

Habitat

The species occurs in deep, red or brown, often alluvial loams; often also occurs in skeletal sands on rocky sandstones and granite ridges and hills.



***Eragrostis leptostachya* - Paddock Lovegrass**

Derivation

leptostachya - from the Greek *leptos* (narrow) and *stachys* (ear of corn).

Habit

A perennial, compactly, tufted grass, the culms are 20-90 cm tall. The culms and leaf sheaths are glandular.

Inflorescence

The panicle is open, often comprising about ½ the plant, 9-40 cm long and 3-21 cm wide. The spikelets are 4-14 mm long and with 6-20 florets. The spikelets are often olive-green. The pedicel usually has a glandular band.

Habitat

Occurs in a variety of soils in association with basalt, granite and laterite, in forest and grassland clearings, waste ground and plantations.



Spikelet



***Eragrostis mexicana* - Mexican Lovegrass**

Derivation

mexicana - from Mexico.

Habit

An annual, tufted, erect grass, the culms are 12-120 cm tall. Sometimes have a ring of glands below the brown nodes.

Inflorescence

The panicle is finally loose and open, often occupying 1/2-2/3 of the plant, 10-40 cm long and 2-17 cm wide. The spikelets are 3.5-9.5 mm long and with 5-16 florets.

Habitat

The species is non-native and grows as a weed of habitation.



The ring of glands below the brown node.

Eragrostis minor - Small Stink Grass

Derivation

minor - Latin for smaller, lesser, plants small in comparison with related species.

Habit

An annual, tufted grass, the culms 6-60 cm tall. The leaves often with pit-like or warty glands especially on nerves.

Inflorescence

The panicle is rather dense or open, 4-20 mm long and 1.5-12 cm wide. The spikelets are 3-15 mm long and with 6-20 florets. Often glandular on the divisions and with a single crateriform gland on the pedicel.

Habitat

The non-native species is a weed of railway tracks, habitation and cultivation. This plant is often odorous.

There are similarities to *Eragrostis cilianensis*, the spikelets are narrower, 1.3-2 mm wide.



***Eragrostis parviflora* - Weeping Lovegrass**

Derivation

parviflora- from the Latin *parvus* (small) and *flos* (flower), spikelets small or with few florets.

Habit

An annual or short-lived perennial, the culms 30-90 cm tall.

Inflorescence

The panicle is open and often weeping, 20-60 cm long and 11-30 cm wide. The spikelets are 4-10 mm long and with 5-15 florets. The lateral nerve of lemma are indistinct.

Habitat

The species occur in a wide range of soils, often alluvial, in usually flat to gently sloping country, often in well-watered sites and seasonally flooded flats, sometimes on roadsides, in saline soils and disturbed ground.

This species resembles *Eragrostis pilosa*.



***Eragrostis pilosa* - Soft Lovegrass**

Derivation

pilosa - from the Latin *pilus* (a hair) and *-osa* (abundance), the whole plant or any of its organs invested with long spreading hairs.

Habit

An annual, erect grass, sometimes with glandular striations, tubercles or pustules on culms, leaves and panicles. The culms are 8-50 cm tall.

Inflorescence

The panicle is contracted or open, 5-20 cm long and 1.5-5 cm wide. The axis sometimes with glandular striations, tubercles or pustules clustered or scattered about the basal whorl of branches. The spikelets are 2.5-9.5 mm and with 4-10 florets.

Habitat

The non-native species occurs as a weed of railway tracks, on roadside, in gardens; in sandy alluvial, loams and lateritic soils.

This species resembles *Eragrostis parviflora*, but *Eragrostis pilosa* has the basal panicle branches often whorled.



The glands are showed below the basal whorl of panicle branches.



Eragrostis pubescens

Derivation

pubescens - from the Latin *pubesco* (become hairy), plant whole or in part hairy.

Habit

A perennial, erect or sprawling grass, the culms 100-200 cm tall. The leaves often pilose to hirsute with simple and tubercle-based hairs.

Inflorescence

The panicle is open and interrupted, 46-85 cm long and 5-22 cm wide. The branches often have tubercle-based hairs. The spikelets are 8-50 mm long and with 12-90 florets.

Habitat

The species occurs chiefly in sandy often alluvial soils, on river levees, margins of swamps coastal flats and dunes.

The species resemble *Eragrostis interrupta* and *Eragrostis brownii*.



Eragrostis schultzei

Derivation

schultzei - in honour of Frederick Schultze (period of activity 1869) who collected in northern Australia.

Habit

A perennial, robust, tussock-forming grass, the culms 80-150 cm tall.

Inflorescence

The panicle is spiciform, rather loose to open, and often interrupted, 11-45 cm long and 1-4 cm wide. The spikelets are 5-15 mm long and with 9-35 florets.

Habitat

The species occurs often in low-lying alluvial habitats, in usually sandy or loam soils.



Eragrostis sororia

Derivation

sororia - from the Latin *soror* (sister). Readily confused with related species.

Habit

A perennial, leafy and compact near base, the culms 20-70 cm tall.

Inflorescence

The panicle contracted, usual spiciform often interrupted in the lower part, 5-25 cm long and 1-2.5 cm wide. The spikelets are 6-20 mm long with 13-40 florets. The spikelets are 6-20 mm long with 13-40 florets.

Habitat

The species occurs in flat to undulating country with usually deep sands, clay and sandy loams; it occurs on alluvial soils on river levees and banks and disturbed ground.

A polymorphic species which intergrades into *Eragrostis elongata*, producing plants with a wide range of intermediate characters.



Eragrostis spartinoides

Derivation

spartinoides - superficially similar to *Spartina*, a genus of Poaceae found in Victoria, South Australia and Tasmania.

Habit

A perennial, tufted grass, the culms are 20-85 cm tall.

Inflorescence

The panicle is open, 7-44 cm long and 3-6 cm wide. The spikelets are 5-28 mm and with 8-55 florets. The spikelets are appressed to primary branches.

Habitat

The species occurs in a range of sandy and clayey, often shallow soils.

It is confused with *Eragrostis brownii*, *Eragrostis cumingii* and *Eragrostis schultzei*.



Spikelets



Eragrostis stenostachya

Derivation

stenostachya - from the Greek *stenos* (narrow) and *stachys* (ear of corn), the inflorescence a narrow or spike-like panicle.

Habit

A perennial, erect or stoloniferous grass, often purplish near the base, the culms 20-60 cm tall.

Inflorescence

The panicle is open, spiciform upwards, 5-15 cm long and 2-7.5 cm wide. The panicle and spikelets branches are stiffly spreading or reflexed. The spikelets are 7-21 mm long and with 9-27 florets.

Habitat

The species occurs in saline meadows and on sandy loam soils.



Spikelet



Eragrostis tenella - Delicate Lovegrass

Derivation

tenella - from the Latin *tenellus* (tender or delicate).

Habit

An annual grass, the culms 5-60 cm tall.

Inflorescence

The panicle is open that is sometimes pilose in axils of the axis, 3.5-14 cm long and 0.8-5.5 cm wide. The inflorescence is often sticky because of small glands on the pedicels. The spikelets are 1-2.3 mm long with 4-7 florets.

Habitat

This non-native grass is a weed of gardens, lawns and roadsides; it occurs in sandy soils.



***Eragrostis tenellula* - Delicate Lovegrass**

Derivation

tenellula - from the Latin *tenellus* (tender or delicate) and *-ula* (diminutive).

Habit

An annual, tufted, erect to semi-prostrate grass, the culms 5-50 cm tall.

Inflorescence

The panicle is open, often comprising about $\frac{1}{2}$ to $\frac{3}{4}$ of plant, 6-25 cm long and 1.5-7.5 cm wide. The inflorescence has perfectly arranged, spreading branches. The spikelets 2-5 mm long and with 3-12 florets.

Habitat

The species occur almost always in seasonally flooded sites, often in grassland plains, and in heavy clays or loams.

It is similar to *Eragrostis exigua*, but the spikelets are longer. *Eragrostis tenellula* has spikelets more than 2 mm long and with 3-12 florets.



Eragrostis tenuifolia - Elastic Grass

Derivation

tenuifolia - from the Latin *tenuis* (slender) and *folium* (leaf), the leaf-blades are narrow.

Habit

A compactly leafy, tufted perennial, the culms 15-90 cm tall.

Inflorescence

The panicle is open and erect, 5-28 cm long and 5-15 cm wide. The axils of inflorescence branches and the pedicles with tufts of hairs present at all nodes, and with purple-black, bearded pulvini. The spikelets are olive-green, they are 4-16 mm long with 4-16 florets. The lateral nerve of lemma indistinct.

Habitat

The non-native species occurs as a weed of roadsides, lawns and habitation.

Similar to *Eragrostis pilosa* but the lower nodes are not whorled.



Spikelets



Eragrostis unioloides

Derivation

unioloides - with spikes resembling those of *Uniola*, a genus of Poaceae not found in Australia, but occur in Americas.

Habit

A tufted, annual grass, mostly glabrous, the culms 8-70 cm tall.

Inflorescence

The panicle is open or rather densely branched, 8-20 cm long and 4-10 cm wide. The spikelets are usually yellowish but with reddish-purple a tinge and have a distinct shape. The spikelets are 4-15 cm long and with 9-70 florets.

Habitat

The species occurs in seasonally wet grey sandy silt.



Spikelets



Eremochloa


From Greek *eremos* (solitary) and *chloa* (grass), referring to the single terminal spike.

Tufted perennials. The inflorescence is a spike or raceme. The spikelets are in pairs, one sessile and one pedicelled, the pedicelled spikelet is rudimentary and often reduced to a pedicel.

Subfamily: Panicoideae; Tribe: Andropogoneae.
World = 13, Australia = 3

Townsville species

Eremochloa bimaculata Poverty Grass



This species could be confused with *Lepturus repens*, *Elionurus citreus* and *Hemarthria uncinata*. Check the key on page 16.



***Eremochloa bimaculata* - Poverty Grass**

Derivation

bimaculata - from the Latin *bi* (twice), *macula* (spot or stain) and *-ata* (possessing), i.e. possessing two spots, which appears to be rather baffling?

Habit

An erect, tufted, perennial grass, the culms are 30-80 cm tall. It will have a distinctive rootstock; the basal leaf sheaths are flattened and keeled.

Inflorescence

The inflorescence is a solitary terminal spike. The lower glume of the sessile spikelet has curved spines on the lower margins and wings at the apex. The pedicelled spikelet is reduced to a narrow glume or absent.

Habitat

The species grows in open woodlands and forests.



Spikelets




Eriachne

Wanderrie Grasses

From the Greek *erion* (wool) and *achne* (chaff or scale), possibly referring to the florets being hairy.

Annuals or perennials, rhizomatous or tufted. The inflorescence is an open or contracted panicle. The spikelets are solitary, and the glumes spread at maturity to reveal two bisexual florets which are awned or unawned. The glumes often persist after the florets have fallen and often spread out.

Subfamily: Micrairoideae; Tribe: Eriachneae
World = 62, Australia = 61



Townsville species

Eriachne squarrosa
Eriachne ciliata
Eriachne mucronata
Eriachne obtusa
Eriachne pallescens
Eriachne rara
Eriachne triodioides

Wanderrie Grass
Slender Wanderrie Grass
Mountain Wanderrie Grass
Northern Wanderrie Grass
Wanderrie Grass

Wanderrie Grass

Key to the species of *Eriachne*

1. Lemma unawned or sometimes mucronate (a sharp, abrupt terminal point), mucro to 0.6 mm long2
 Lemma with a distinct (although sometimes short) awn 1.5-30 mm long3
2. Glumes straight when mature; lemma with a mucronate (a sharp, abrupt terminal point) apex*Eriachne mucronata*
 Glumes spread out when mature; lemma with an acute apex*Eriachne obtusa*




Eriachne mucronata



Eriachne obtusa

3. Awns reflexed and curved4
 Awns straight5
4. Inflorescence ±loose*Eriachne rara*
 Inflorescence dense*Eriachne squarrosa*



Eriachne rara



Eriachne squarrosa

5. Perennial, spikelets 7-10.5 mm long (commonly on coastal sand dunes) *Eriachne triodioides*
 Annual, spikelets 2.5-6 mm long6
6. Glumes open wide when mature and light coloured*Eriachne ciliata*
 Glumes not open when mature and dark coloured *Eriachne pallescens*



Eriachne triodioides



Eriachne ciliata



Eriachne pallescens

***Eriachne ciliata* - Slender Wanderrie Grass**

Derivation

ciliata - from the Latin *cilium* (eyelid) and *-ata* (possessing), the plant is hairy overall or in part.

Habit

A slender annual or ephemeral grass, the culms are 10-30 cm tall.

Inflorescence

The panicle is open at maturity, 4-7 cm long and 1.5-4 cm wide. The glumes are open wide when mature and light coloured. The lemma's awn is short, 1.5-3 mm long

Habitat

This species is common of shallow and skeletal sandy or loam soils.



Spikelet



***Eriachne mucronata* - Mountain Wanderrie Grass**

Derivation

mucronata - from the Latin *mucro* (sharp point) and *-ata* (possessing), with glumes or lemmas contracted into a short hard point or a bifid and shortly awned from between the teeth.

Habit

A perennial grass, the culms are 10-80 cm tall, the base is knotty or thickened, pubescent or woolly. This species is highly variable in habit.

Inflorescence

The panicle is usually open 3-8 cm long and 0.5-1 cm wide. The glumes are equal to florets; the lemmas are acute or mucronate. The lemma is 5.8-7 mm long.

Habitat

The species is a widespread and adaptable species, occurs on various soils.

Eriachne mucronata closely resembles *Eriachne obtusa*. This species has lemmas which are mucronate and 11-13 nerves.



Spikelet



***Eriachne obtusa* - Northern Wanderrie Grass**

Derivation

obtusa - Latin for blunt, the glumes or lemmas apically rounded.

Habit

A perennial grass, sometimes glaucous, culms are 30-60 cm tall.

Inflorescence

The panicle is rather contracted, 4-10 cm long and 1.5-3 cm wide. The glumes are open wide when mature and light coloured. The lemmas are not awned, and are 3.8-5 mm long.

Habitat

This species grows in skeletal, shallow and deep soils.

Eriachne obtusa closely resembles *Eriachne mucronata*. This species has lemmas which are muticous and have 5-7 nerves.



Spikelet



***Eriachne pallescens* - Wanderrie Grass**

Derivation

pallescens - from the Latin *pallesco* (become pale), they are losing colour at maturity, especially spikelets.

Habit

A perennial grass, mostly glabrous on culms and foliage; base knotty or slightly thickened, sparsely pubescent. The culms are 25-105 cm tall.

Inflorescence

The panicle is open, sometimes drooping, 6-15 cm long and 3-9 cm wide. The glumes are erect when mature. The lemmas awn is 1.5-8 mm long.

Habitat

This species usually grows in deep sandy soils, often in low-lying sites such as swamps and flood channels. It also grows on coastal dunes and skeletal and shallow often stony soils.



Spikelet



Eriachne rara

Derivation

rara - Latin for far apart, the spikelets far apart in panicle.

Habit

A short-lived perennial grass, the culms are 15-60 cm tall.

Inflorescence

The panicle is open, 3.5-10 cm long and 0.5-5 cm wide. The glumes are erect when mature. The lemmas awn are 13-3 mm long, they are flexuose (bent).

Habitat

This species occurs in deep or shallow sandy, sometimes gritty or stony soils.



Spikelet



Eriachne squarrosa - Wanderrie Grass

Derivation

squarrosa - Latin for spreading at right angles from a common axis.
Inflorescence arms held at right angles to common axis.

Habit

A perennial, tufted grass, the culms are 35-80 cm tall.

Inflorescence


The inflorescence is a spicate panicle, 3-8 cm long, 1.5 cm wide. The glumes are erect when mature. The lemma awns 25-28 mm long and curved.

Habitat

This species grows in often moist, low-laying, sandy flats and depressions under *Melaleuca* species and on flood banks of seasonal streams.



Spikelet



Glumes

***Eriachne triodioides* - Wanderrie Grass**

Derivation

triodioides - resembling *Triodia* species.

Habit

A robust, perennial grass, the culms are 75-120 cm tall, the base not thickened, pubescent.

Inflorescence

The panicle contracted or eventually open, 8-15 cm long, 1.5-6 cm wide (including awns). Lemmas usually awned, the awn is 2-13 mm long.

Habitat

This species is commonly on islands and coastal mainland. It is chiefly of coastal sand dunes and deep, sandy inland sands.



Spikelet



Eriochloa

Spring or Cup Grasses


From Greek *erion* (wool) and *chloe* (grass), referring to the hairy spikelets and pedicels.

Annuals or perennials, stoloniferous or tufted to decumbent. The inflorescence is a once-branched panicle with racemes appressed at first, spreading later. The spikelets are solitary or in pairs. The glumes are unequal, the lower glume usually reduced to a cup-like ring at the base of the spikelet. The genus is distinguishable by this 'cup' which is formed from the lower rachilla (the axis of the spikelet) internode which becomes swollen and fused to the lower glume.

Subfamily: Panicoideae; Tribe: Paniceae
World = 51, Australia = 6

Townsville species

<i>Eriochloa crebra</i>	Tall Cup Grass
<i>Eriochloa procera</i>	Cup Grass
<i>Eriochloa pseudoacrotricha</i>	Early Spring Cup Grass



Key to the species of *Eriochloa*

1. Spikelets with apices drawn out into bristles *Eriochloa pseudoacrotricha*
Spikelets with apices acute to acuminate 2



Apices drawn out into bristles



Apices acute to acuminate

2. Spikelets 3.7-5.5 mm long, lightly overlapping on racemes *Eriochloa crebra*
Spikelets 2.5-4.0 mm long, \pm loose on racemes *Eriochloa procera*

***Eriochloa crebra* - Tall Cup Grass**

Derivation

crebra - from the Latin *creber* (pressed together), the racemes held erect appressed to inflorescence axis.

Habit

A perennial, tufted grass, the culms 35-100 cm.

Inflorescence

A panicle of racemes, the 5-25 racemes are 1.5-5 cm long. The spikelets are 3.7-5.5 mm long, the apices are acute to acuminate.

Habitat

This species usually grows in temporarily wet places such as creek banks and floodplains.



Spikelet



***Eriochloa procera* - Cup Grass**

Derivation

procera - Latin for tall, the culms are tall.

Habit

An annual or perennial, tufted grass, the culms are 20-120 cm tall.

Inflorescence

A panicle of racemes, the 3-10 racemes are 2-10 cm long. The spikelets are 2.5-4 mm long.

Habitat

This species usually grows on disturbed ground in wetter areas.



Spikelet



***Eriochloa pseudoacrotricha* - Early Spring Cup Grass**

Derivation

pseudoacrotricha - from the Greek *pseudos* (false), resembling *Eriochloa racemosa* var. *acrotricha*.

Habit

A perennial, loosely tufted grass, the culms are 20-100 cm tall.

Inflorescence

A panicle of racemes, the 2-10 racemes are 2-10 cm long. The spikelets are 3.6-6 mm long.

Habitat

This species usually associated with temporarily wet places such as creeks and floodplains.



Spikelets



Eulalia

Named to honour Eulalia Delile, a botanical artist.


Tufted perennial grasses. The inflorescence is digitate or subdigitate; the branches are very hairy or silky, often brown or purple. The spikelets are in pairs and similar, one sessile, the other pedicelled, each with one bisexual floret, both awned.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 59, Australia = 6

Townsville species

Eulalia aurea

Silky Browntop



This grass is easily recognised by its fluffy, golden-yellow-brown inflorescences. However, this species could be confused with the closely-related annual grass *Pseudopogonatherum contortum*.

Key to the species of *Eulalia* and related genera

Perennial; awns not intertwining at maturity; sessile spikelets usually more than 3 mm long*Eulalia*
Delicate annual; awns intertwining at maturity; all spikelets very small, 1.5-2.5 mm long
.....*Pseudopogonatherum*



Pseudopogonatherum



Eulalia

***Eulalia aurea* - Silky Browntop**

Derivation

aurea - the Latin for golden-yellow which refers to the spikelets or pedicels or other parts with golden-yellow hairs.

Habit

A perennial, tufted grass with slender, erect stems, the culm is 40-150 cm tall.

Inflorescence

The inflorescence is digitate with 2-3 branches, 3-12 cm long. The racemes are clothed with fluffy, dark brown hairs 1-2 mm long. The spikelets are 3.5-5 mm long,

Habitat

This species is often in ephemeral water courses in drier areas.



Hackelochloa


From Eduard Hackel, and Austrian botanist and *chloa* (grass).

Annual, tufted or solitary. The inflorescence is composed of several racemes. Leaf-blade surface are hirsute, with tubercle-based hairs. The spikelets are in pairs, the sessile spikelet is round and rugose, the pedicelled spikelet is narrow and smooth.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 2, Australia = 1

Townsville species

Hackelochloa granularis Pit Scale Grass



***Hackelochloa granularis* - Pit Scale Grass**

Derivation

granularis - from the Latin *granum* (grain) and *-aris* (pertaining to), it has segments of the inflorescence resemble beads.

Habit

An erect, annual grass, the culms are 10-75 cm tall.

Inflorescence

The panicle is a series of spike-like racemes up to 2.5 cm long, each partly enclosed by spathes. The sessile spikelet is very rounded and with a distinctive, sculptured patterning on the back. The pedicelled spikelet is narrow and is sterile (rarely male).

Habitat

This species usually grows on open sands, coastal salt marshes, clays, moist places and roadsides.



Hemarthria

From Greek *hemi* (half) and *arthron* (jointed) i.e. half-jointed, referring to the raceme internodes articulated but not breaking up.


Perennials, often with stolons and rhizomes. The inflorescence is a single flattened raceme, usually subtended by an inflated leaf sheath. The spikelets in pairs, one sessile and one pedicelled and partially embedded in the rachis, the pedicelled spikelet resembles the sessile spikelet and the pedicel is fused to the internode.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 14, Australia = 2

Townsville species

Hemarthria uncinata

Mat Grass



This species could be confused with *Elionurus citreus*, *Eremochloa bimaculata* and *Lepturus repens*. Check the key on page 16.



***Hemarthria uncinata* - Mat Grass**

Derivation

uncinata - from the Latin *uncinus* (hook) and *-ata* (possessing), the upper glume drawn out into a hook.

Habit

A tufted perennial often rhizomatous or stoloniferous grass, stems prostrate or erect. The culms are 20-100 cm tall and often form a coarse matted sward.

Inflorescence

The inflorescence is a raceme, 6-14 cm long. The lower glume similar to that of the sessile spikelet with a straight or hooked apex; upper asymmetric, narrowly winged.

Habitat

This species grows on a range of soils, usually in damp areas, swamps, near coastal estuaries, sand dunes and stream banks. This species does not grow in Townsville. *Lepturus repens* are often misidentified as this species.



Heteropogon

Spear Grasses


From the Greek *heteros* (different) and *pogon* (beard), referring to the difference between the awnless male and awned female spikelets.

Tufted perennials. The inflorescence is a raceme of paired spikelets. The lower paired spikelets are alike, unawned, and either male or neuter. The upper paired spikelets are dissimilar, one sessile and awned and one pedicelled and unawned. The awns of the sessile spikelet are long and extend above the raceme and when the seeds mature, the awns tangle.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 19, Australia = 2

Townsville species

<i>Heteropogon contortus</i>	Black Spear Grass
<i>Heteropogon triticeus</i>	Giant Spear Grass



Key to the species of *Heteropogon* and similar species

1. Inflorescence spike-like, but composed of 2 erect, closely adpressed racemes*Ischaemum*
Inflorescence is a raceme2



Ischaemum



Sehima nervosum



Heteropogon triticeus



Heteropogon contortus

2. Awns are 20-40 mm long; pedicelled spikelets is the same length as fertile*Sehima nervosum*
Awns are 50-140 mm long; pedicelled spikelet longer than fertile.....3
3. Moderately tall grass (usually less than 1 m tall); racemes less than 6 cm long; fertile spikelets 6-10 mm long*Heteropogon contortus*
Very tall grass (usually greater than 1 m tall); racemes more than 9 cm; fertile spikelets 10-15 mm long *Heteropogon triticeus*

***Heteropogon contortus* - Black Spear Grass**

Derivation

contortus - awns hygroscopic and so twisted when dry.

Habit

A perennial, tufted grass, the culms are 30-100 cm tall.

Inflorescence

The inflorescence is a raceme, 3-6 cm long. The sessile is 6-9 mm long and the pedicelled spikelet is 5-15 mm long. The awns, 50-80 mm long, and the way they become twisted as the seeds mature are a characteristic trait of spear grass.

Habitat

This species is widespread in fire-prone open forest and woodland.



Sessile spikelet



Pedicelled spikelet



Mass of sessile spikelets



This is a raceme of pedicelled spikelets, the seeds have dropped off

***Heteropogon triticeus* - Giant Spear Grass**

Derivation

triticeus - resembling *Triticum* with respect to the inflorescence.

Habit

A perennial, tufted grass, the culms are 100-160 cm tall. The base is fan shaped.

Inflorescence

The inflorescence is a raceme, 9-14 cm long. The sessile is 10-15 mm long and the pedicelled spikelet is 18-25 mm long. The awn is 70-140 mm long.

Habitat

This species grows in *Eucalyptus* forests and woodlands.



This is a raceme of pedicelled spikelets, the seeds have dropped off



Spikelets



Hymenachne


Decumbent, spreading perennial aquatic grasses. The stems produce roots at the nodes. The inflorescence is a spicate panicle (spike-like with short branches). The spikelets are solitary.

Subfamily: Panicoideae; Tribe: Paniceae
World = 21, Australia = 3

Hymenachne invades permanent water bodies and seasonally inundated wetlands.

Townsville species

Hymenachne acutigluma
* *Hymenachne amplexicaulis* Hymenachne



Key to the species of *Hymenachne*

Spikelets 3-4 mm long; leaf-base amplexicaul (having lobes, usually auriculate, that completely surround the stem). *Hymenachne amplexicaulis*
Spikelets 4.5-5.5 mm long; leaf-base not amplexicaul..... *Hymenachne acutigluma*



Hymenachne acutigluma

Derivation

acutigluma - from the Latin *acuo* (sharpen) and *gluma* (husk), the glumes tapering.

Habit

A perennial, aquatic or subaquatic grass. The culms are decumbent, 30-200 cm tall. The stems are spongy, rooting from lower nodes. The leaf-base not amplexicaul.

Inflorescence

The panicle is spiciform, 8-50 cm long and 1–3.5 cm wide. The spikelets are dorsally compressed 4.5-5.5 mm long.

Habitat

This native grass is a swamp grass, more or less independent of rainfall.



Leaf blade bases of *Hymenachne* spp.

Left to right: *H. amplexicaulis*, *H. × calamitosa* and *H. acutigluma*

John Clarkson photograph

Naturally occurring hybridisation between the introduced species *Hymenachne amplexicaulis* and the Australian native species *H. acutigluma* is reported from two widely separated locations in tropical Australia. This has been named *Hymenachne × calamitosa*.

***Hymenachne amplexicaulis* - Hymenachne**

Derivation

amplexicaulis - from the Latin *implexus* (encircling) and *caulis* (stem), the bases of the leaf-blades encircle the stem.

Habit

A perennial, aquatic or subaquatic grass. The culms are decumbent, 200-350 cm tall. The stems are spongy, rooting from lower nodes. The leaf-base is amplexicaul (having lobes, usually auriculate, that completely surround the stem).

Inflorescence

The panicle is spiciform, 10-40 cm long and 1-2 cm wide. The spikelets are dorsally compressed 3-4 mm long.

Habitat

This non-native species was released in Queensland in 1988 for use as 'ponded pasture'. It has since escaped from cultivation and is now considered a Weed of National Significance (WONS). It is regarded as one of the worst weeds in Australia because of its invasiveness, potential for spread, and economic and environmental impacts.



Spikelet



Hyparrhenia

Thatch Grasses

From the Greek words *hypo* (below) and *arrhen* (male), referring to the pair of male spikelets at the base of each raceme.


Tufted perennials or annuals. The inflorescence is composed of a few to many pairs of racemes subtended by spatheoles (small bracts or modified leaves which enclose the inflorescence). The spikelets are in pairs, one sessile and usually awned and the other pedicelled and awnless.

Subfamily: Panicoideae; Tribe: Andropogoneae

World = 78, Australia = 4

Townsville species

* *Hyparrhenia rufa* subsp. *rufa* Thatch Grass



Hyparrhenia rufa subsp. *rufa* grows mainly along roadsides and in disturbed areas. It could be confused with Grader Grass (*Themeda quadrivalvis*), Giant Spear Grass (*Heteropogon triticeus*) and Gamba Grass (*Andropogon gayanus*), especially before flowering.

These species are tall grasses, and their increase in height before flowering is caused by a section of pale yellow culm being pushed upwards until it protrudes well above the green leaf sheath that previously enclosed it. The pale sections of the internodes alternate with the green sheaths giving these grass species a conspicuously banded appearance.

Key to the species of *Hyparrhenia* and similar genera

1. Inflorescence is a raceme *Heteropogon triticeus*
 Inflorescence with branches 2
2. Fertile spikelets within an involucre of four male or barren spikelets *Themeda quadrivalvis*
 Fertile spikelets without an involucre of spikelets 3
3. Spikelets covered with red or brown hairs *Hyparrhenia rufa* subsp. *rufa*
 Spikelets covered with white hairs *Andropogon gayanus*



Spikelets covered with brown or red hairs
Hyparrhenia rufa subsp. *rufa*



Spikelets covered with white hairs
Andropogon gayanus

***Hyparrhenia rufa* - Thatch Grass**

Derivation

rufa - Latin for reddish, the inflorescence is purple to red.

Habit

A perennial, tufted grass, the culms are 30-300 cm tall.

Inflorescence

The panicle is subtended by a spatheole. The rames are paired and are 1.5-4 cm long. The sessile spikelet is 3.5-5.5 mm long. The sessile spikelet is awned.

Habitat

This non-native species is mainly a weed of roadsides and disturbed sites.

There are two subspecies, only one is found in Townsville, *Hyparrhenia rufa* subsp. *rufa*.



Spikelets



Imperata


In honour of Ferrante Imperato (1550–1625), a Neopolitan naturalist/pharmacist.

Perennial grasses with rhizomes; leaves mostly basal. The inflorescence is a fluffy, spicate (spike-like with short inflorescence branches) panicle. The spikelets are solitary or paired, all alike, if paired pedicels unequal.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 18, Australia = 1

Townsville species

Imperata cylindrica Blady Grass



Spikelets



***Imperata cylindrica* - Blady Grass**

Derivation

cylindrica - refers to the shape of the inflorescence.

Habit

A tufted, perennial grass with stiff, erect leaves, the culms is 10-120 cm tall.

Inflorescence

The fluffy, white inflorescence is 3-20 cm long. The very small spikelets are hidden by long, silky, white hairs. The spikelets are 2.2-6 mm long and 1 mm wide.

Habitat

Often grows in damp or weedy places. After burning it rapidly resprouts from the rhizomes.



Ischaemum


From the Greek *ischo* (to restrain) and *haima* (blood), as woolly seeds of this type species were reported as being used to stop bleeding.

Erect or sprawling annuals or perennials. The inflorescence is usually of paired digitate appressed racemes, thus appearing spike-like. At maturity, the racemes split open. The spikelets are paired, one sessile and one pedicelled, and are partially embedded in rachis. The sessile spikelet is awned, the pedicelled spikelet is sometimes awned.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 138, Australia = 15

Townsville species

Ischaemum australe Large Bluegrass
Ischaemum rugosum



Key to the species of *Ischaemum* and similar species

1. Inflorescence is a raceme2
- Inflorescence spike-like, but composed of 2 erect, closely appressed raceme3



Ischaemum



Sehima nervosum



Heteropogon triticeus



Heteropogon contortus

2. Awns are 20-40 mm long; pedicelled spikelet is the same length as fertile *Sehima nervosum*
Awns are 50-140 mm long; pedicelled spikelet longer than fertile..... *Heteropogon*
3. Lower glume of sessile spikelet transversely rugose..... *Ischaemum rugosum*
Lower glume of sessile spikelet not transversely rugose *Ischaemum australe*

***Ischaemum australe* - Large Bluegrass**

Derivation

australe - from the South (Australia).

Habit

A perennial, tufted grass, the culms are 50-150 cm tall.

Inflorescence

The inflorescence is spike-like, but is paired, digitate, appressed racemes, 3-10 cm long. The sessile spikelet is 6-7 mm long and the awn is 8-10 mm long. The lower glume of the sessile spikelet is smooth.

Habitat

This species is usually found in wet situations often confined to poor soils.

There are three varieties

1. Nodes glabrous..... *Ischaemum australe* var. *arundinaceum*
Nodes hairy 2
2. Leaves, rachis internodes and pedicels hairy *Ischaemum australe* var. *villosum*
Leaves, rachis internodes and pedicels glabrous *Ischaemum australe* var. *australe*



Spikelets



Ischaemum rugosum

Derivation

rugosum - from the Latin *ruga* (wrinkle) and *-osa* (abundance), usually with sculptured glumes.

Habit

An annual, decumbent grass, the culms are 10-100 cm tall.

Inflorescence

The inflorescence is spike-like, but is paired, digitate, appressed racemes, 3-14 cm long. The sessile spikelet is 4-6 mm long and the awn is 15-20 mm long. The lower glume of the sessile spikelet is transversely ridged.

Habit

This species grows in wet sites.



Spikelets

There are two varieties

1. Pedicelled spikelet developed *Ischaemum rugosum* var. *rugosum*
Pedicelled spikelet much reduced..... *Ischaemum rugosum* var. *segetum*



Leersia


Named for Johan Daniel Leers, a German apothecary and botanist (1727-1774).

Perennials, rarely annuals usually growing in marshland, stream banks and shallow water. The inflorescence is an open or contracted panicle. The spikelets are solitary and strongly laterally compressed. Glumes are rudimentary, apparently represented by a narrow rim at the pedicel apex. *Leersia* is closely related to rice (*Oryza*).

Subfamily: Ehrhartoideae; Tribe: Oryzeae
World = 27, Australia = 2

Townsville species

Leersia hexandra Swamp Rice Grass



Key to the species of *Leersia* and *Oryza*

Spikelets 1-flowered with a bisexual floret and without sterile florets..... *Leersia*
Spikelets 3-flowered, the lower 2 reduced to sterile lemmas *Oryza*

***Leersia hexandra* - Swamp Rice Grass**

Derivation

hexandra - from the Greek *hexa* (six) and *aner* (man), the florets possess six stamens.

Habit

Aquatic or semi aquatic perennial, rhizomatous grass, the culms are erect or floating on water 0.3-1.5 m tall.

Inflorescence

The inflorescence is a contracted to open panicle, 5-10 cm long and 1-4 cm wide. Each floret has six stamens.

Habitat

This species grows in swamps, on river and creek banks into shallow water, waterholes channels and drains, in sand, clay and peat.



Spikelets




Lepturus

From the Greek *leptos* (slender) and *oura* (tail), referring to the slender inflorescence.

Annuals or perennials, tufted and stoloniferous. Grows on sandy beaches, with some species extending to coastal hinterlands. The inflorescence is a solitary, bilateral spike (almost cylindrical). The spikelets are solitary and partially embedded in the rachis. The lower glume is absent or obscure.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 30, Australia = 5

Townsville species
Lepturus repens



This species could be confused with *Elionurus citreus*, *Eremochloa bimaculata* and *Hemarthria uncinata*. Check the key on page 16.



Lepturus repens

Derivation

repens - from the Latin *repo* (crawl), referring to the well-developed rhizome.

Habit

A tufted and creeping perennial grass, the culms are 10-60 cm tall.

Inflorescence

The inflorescence is a cylindrical spike 3.5-7 cm long and about 3 mm wide. The spikelets are contained within the cavities of the axis and at maturity it is falling apart.

Habitat

A coastal plant growing close to water in sand, especially coral sand, or in shallow soil creeping over rocks, or in *Casuarina* woodland.



Spikelets



Megathyrsus


Guinea Grass or Green Panic

From *mega* (large) and *thyrsus* (a dense flower cluster; much-branched indeterminate inflorescence with pedicellate flowers).

Megathyrsus maximus was previously known as *Panicum maximum* and *Urochloa maxima* and is distinguished by its large open panicle and rugose (wrinkled) fertile floret.

Densely tufted perennials, culms usually erect, 60-250 m tall. The panicle is 12-60 cm long and whorled at the lower nodes. The spikelets are solitary or in pairs.

Subfamily: Panicoideae; Tribe: Paniceae
World = 3, Australia = 3



Megathyrsus maximus is the only Australian species from this genus; *maximus* - Latin for greatest.

It can be confused with the species *Panicum mitchellii*. *Megathyrsus maximus* have a fertile lemma transversely rugulose (with a wrinkled appearance).



Townsville species and varieties

- * *Megathyrsus maximus* var. *coloratus*
- * *Megathyrsus maximus* var. *maximus*
- * *Megathyrsus maximus* var. *maximus* 'Hamil'
- * *Megathyrsus maximus* var. *pubiglumis*

Purple-topped Guinea Grass
Common Guinea Grass
Hamil Grass
Green Panic

***Megathyrsus maximus* - Guinea Grass, Green Panic**

Derivation

maximus - Latin for greatest, the culms are very tall.

Habit

A perennial, tufted grass, the culms are 2.5-3.5 m tall, the cultivar 'Hamil' is 2.5-3.5 m tall.

Inflorescence

The panicle is open, 12-60 cm long. The fertile lemma has a wrinkled appearance.

Habitat

This non-native species has been widely cultivated as a pasture grass, but it is a very common, widespread weed of roadsides and disturbed sites.



Spikelet



Key to *Megathyrsus maximus* varieties

1. Plant 2.5-3.5 m tall *Megathyrsus maximus* var. *maximus* 'Hamil'
- Plant 0.5-2.5 m tall 2
2. Leaf glabrous at sheath and blade..... *Megathyrsus maximus* var. *maximus*
- Leaf hairy at junction of sheaths and blade 3



Leaf glabrous at sheath and blade



Leaf hairy at junction of sheaths and blade

3. Spikelets indistinctly hairy..... *Megathyrsus maximus* var. *pubiglumis*
- Spikelets glabrous *Megathyrsus maximus* var. *coloratus*


Melinis

From the Greek *meline* (a cereal, probably millet).

Stoloniferous or tufted, aromatic annuals or perennials. The inflorescence is an open panicle, red, white or purple coloured. The spikelets are solitary; the lower glume is absent or obscure and the lower lemma has an awn arising from the apical lobes, which is sometimes obscured by hairs on the spikelet.

Subfamily: Panicoideae; Tribe: Paniceae
World = 37, Australia = 2

Based on inflorescence colour and shape, *Melinis* could be confused with *Capillipedium* and *Chrysopogon zizanioides* however the spikelets are very different.



Melinis repens

Melinis minutiflora
Spikelets

Townsville species

- * *Melinis minutiflora* Molasses Grass
- * *Melinis repens* Red Natal Grass

Key to the species of *Melinis* and similar genera

1. Plant 150-300 cm tall *Chrysopogon zizanioides*
Plant 10-140 cm tall 2
2. Spikelets in pairs, one sessile and the other (companion) spikelet pedicelled in threes *Capillipedium*
Spikelets solitary 3
3. Spikelets covered with long, red, silky hairs; leaves not sticky *Melinis repens*
Spikelets more or less hairless; leaves sticky and smelling of molasses *Melinis minutiflora*



Capillipedium spicigerum



Melinis repens



Melinis minutiflora



Chrysopogon zizanioides

***Melinis minutiflora* - Molasses Grass**

Derivation

minutiflora - from the Latin *minutis* (very small) and *flos* (flower), the spikelets or florets are very small.

Habit

A perennial, tufted grass, the culms are 30-120 cm tall. The leaves are minutely to densely cover with glandular hairs exuding drops of viscid oil, with characteristic molasses odour.


Inflorescence

The panicle is compound, dense or open, 6-30 cm long, it is purple in colour. The spikelets are 1.5-2.4 mm long.

Habitat

This non-native species has been cultivated as a pasture grass and for erosion control in warmer and wetter areas. It is now a weed of roadsides, forest margins, open woodlands, pastures, disturbed sites, waste areas.

This species has a small seed with long awn like *Pseudopogonatherum contortum*.



Spikelet



***Melinis repens* - Red Natal Grass**

Derivation

repens - from the Latin *repo* (crawl), the rhizome well developed.

Habit

An annual or perennial, tufted grass, the culms are 20-130 cm tall.

Inflorescence

The panicle is compound, open, 5-20 cm long; it is pink, red or shining white in colour. The spikelets are 3-5 mm long.

Habitat

This non-native species is a very common weed of roadsides, railways, parks, gardens, footpaths, disturbed sites, waste areas, pastures and crops.



Spikelet



Mnesithea

Named after Mnesitheus (4th century BC), a Greek herbalist.

A genus of variable habit ranging from robust perennials to delicate annuals. The inflorescence is a single raceme or panicle and subtended by a spathe. The spikelets are paired, and partially embedded in rachis, which breaks into segments at maturity. The pedicelled spikelet is sometimes different in shape and size from the sessile spikelet.

Subfamily: Panicoideae; Tribe: Andropogoneae
World: = 39, Australia = 4

Townsville species

Mnesithea formosa



Mnesithea rottboellioides Northern Canegrass

Key to the species of *Mnesithea*


Perennial, robust, 100-300 cm tall; the pedicelled spikelet is similar to the sessile spikelet

.....*Mnesithea rottboellioides*

Annual, erect 10-75 cm tall; the pedicelled spikelet is not like the sessile spikelet*Mnesithea formosa*



Mnesithea formosa



Mnesithea rottboellioides

Mnesithea formosa

Derivation

formosa - Latin for handsome, it is attractive in appearance.

Habit

An erect, annual grass, the culms are 10-60 cm tall.


Inflorescence

The panicle is subtended by an inflated leaf-sheath; the racemes are 2-8 cm long. The sessile spikelet is terete and densely white hairy. The pedicelled spikelet is sterile and is reduced to two unequal glumes.


Habitat

This species grows in gravelly silt, alluvium, pink or red clayey sand. Occurs in savanna woodlands.


Mnesithea formosa could be confused with *Schizachyrium fragile*. The spikelets of *Schizachyrium fragile* have awns.



Mnesithea formosa



Schizachyrium fragile



***Mnesithea rottboellioides* - Northern Canegrass**

Derivation

rottboellioides - resembling the genus *Rottboellia* with respect to the inflorescence.

Habit

A perennial, robust grass, the culms are 100-300 cm tall.


Inflorescence

The panicle is terminal and axillary, subtended by a spatheole. The cylindrical racemes, 5-10 cm long, are crowded at one or several nodes. Spikelets on one side of the axis. The pedicelled spikelet is similar to the sessile spikelet.

Habitat

This species grows on hills, in grasslands or open forests.

Mnesithea rottboellioides could be confused with *Ophiuros exaltatus*.



Mnesithea rottboellioides



Ophiuros


From the Greek *ophis* (serpent) and *oura* (tail), alluding to the smooth spikes with the scale-like appressed lower glumes of the spikelets.

Tufted perennials or annuals, culms woody and persistent. The inflorescence is a compound panicle composed of aggregated cylindrical racemes. The spikelets are sunken into the rachis and are borne alternately on opposite sides. The lower glume is sculptured when the spikelet is mature.

Subfamily: Panicoideae; Tribe: Andropogoneae.
World = 17, Australia = 1

Townsville species

Ophiuros exaltatus Cane Grass



This species resembles *Mnesithea rottboellioides*, however *Ophiuros exaltatus* has spikelets that appear to be solitary, and the lower glume is pitted in rows.



***Ophiuros exaltatus* - Cane Grass**

Derivation

exaltatus - from the Latin for raised up, tall.

Habit

A perennial, erect grass, the culms are 150-300 cm tall.

Inflorescence

The inflorescence is a clustered panicle of slender racemes, each subtended by a short spathe.

Habitat

The inflorescence is a clustered panicle of slender racemes, each subtended by a short spathe.



Spikelets




Oplismenus

From the Greek *hoplismenus* (bearing arms), referring to the armed spikelets.

Shade-loving, decumbent annuals or perennials. The inflorescence is a once-branched panicle, with spikelets on one side of a slender axis, or branches are reduced to fascicles (clusters) of spikelets. The lower glume is awned and is often sticky at the tip, and by sticking to passing animals, acts as a fruit dispersal mechanism.

Oplismenus species are commonly found in rainforest or in damp shady places.

Subfamily: Panicoideae; Tribe: Paniceae
World = 47, Australia = 5



Townsville species

Oplismenus aemulus Australian Basket Grass
Oplismenus compositus Running Mountain Grass

Key to the species of *Oplismenus*

1. Most leaves less than 7 times as long as wide2
- Most leaves 10 or more times as long as wide3



Most leaves less than 7 times as long as wide



Most leaves 10 or more times as long as wide

2. Basal racemes more than 4 cm long.....*Oplismenus compositus*
Basal racemes less than 4 cm long *Oplismenus aemulus*
3. Racemes in distinct clusters; leaf blade indumentum dense..... *Oplismenus mollis*
Racemes not in distinct clusters; leaf blade indumentum sparse*Oplismenus imbecillis*



Oplismenus mollis



Oplismenus imbecillis

***Oplismenus aemulus* - Australian Basket Grass**

Derivation

aemulus - Latin for more or less equalling, subtending glumes more or less equal.

Habit

A perennial, prostrate grass, the culms are 5-50 cm long. The leaf-blades 4-18 mm wide

Inflorescence


The inflorescence is a once branched panicle; it has 3-9 racemes, which are 2.5-3.5 cm long.

Habitat

This species is common in shady, damp places in or near rainforest and along streamlines.



Spikelets



***Oplismenus compositus* - Running Mountain Grass**

Derivation

compositus- from the Latin *compono* (bring together).

Habit

A perennial, prostrate grass, the culms are 15-80 cm long. The leaf-blades are 8-27 mm wide.

Inflorescence

The inflorescence is a once branched panicle; it has 4-12 racemes, which are 2.5-11 cm long.

Habitat

This species grows in moist places in rainforests, vine forest, vine thicket and along forest margins.



Spikelet



Oryza

Rice

Latin from the Arabic *uruz* (rice), whence Greek *oruza* is also derived.


Tufted annuals or perennials often growing in open swamps. The inflorescence is an open or contracted panicle. The spikelets are solitary, strongly laterally compressed, usually with a long awn (the cultivated rice *Oryza sativa* is not of awned). The spikelet consists, two rudimentary, scale-like glumes, two basal florets reduced to lemmas and a terminal fertile (bisexual) floret. The fertile floret contains 6 stamens. Most Australian grass genera contain 1-3 stamens, *Leersia hexandra* also has 6 stamens.

Oryza species usually grow in swampy areas or seasonally inundated clay soils. During the dry season the grass dies back and the above-ground parts of the plant are not visible.

Subfamily: Ehrhartoideae; Tribe: Oryzeae
World = 36, Australia = 5


Townsville species

<i>Oryza australiensis</i>	Australian Wild Rice
<i>Oryza meridionalis</i>	Australian Wild Rice




Key to the species of *Leersia* and *Oryza* species

1. Spikelets 1-flowered with a bisexual floret and without sterile florets *Leersia*
Spikelets 3-flowered, the lower 2 reduced to sterile lemmas 2
2. Awn 30-150 mm long; spikelets inserted at an angle to the pedicel; ligule split, apex two lobed *Oryza meridionalis*
Awn 10-60 mm long; spikelets inserted more or less horizontal on their pedicels; ligule not split *Oryza australiensis*



Oryza meridionalis



Oryza australiensis

***Oryza australiensis* - Australian Wild Rice**

Derivation

australiensis - from Australia.

Habit

A perennial, rhizomatous grass, the culms are erect, 0.8-2.5 m tall. The ligule apex is acute, abuse or truncate (shorten or reduce).

Inflorescence

The panicle is open or sometimes contracted, 13-45 cm long. The lemma awn is 10-60 mm long; the spikelet is inserted more or less horizontal on their pedicels.

Habitat

The species grows in or at the edge of water, in swamps, lagoons, creeks, drainage channels or seasonally inundated areas, often in black or grey clays.



***Oryza meridionalis* - Australian Wild Rice**

Derivation

meridionalis - from the Latin *meridies* (meridian) and *-ale* (pertaining to), on the same meridian (line of longitude) as a related species.

Habit

An annual or sometimes perennial grass, the culms are erect to decumbent, 0.3-2 m tall. The ligule apex two lobed.

Inflorescence

The panicle is open or sometimes contracted, 9-30 cm long. The lemma awn is 30-150 mm long; the spikelet is inserted more or less horizontal on their pedicels.

Habitat

The species grows in shallow or sometimes deeper water, in aquatic to drying conditions, on margins of lagoons, waterholes and creeks, include seasonal swamps, often in cracking clay, sometimes in loams and sandy soils.



Spikelet

Oxychloris

Windmill Grasses

From Greek *oxys* (sharp) referring to the pungent callus, and the generic name *Chloris* (in which it was formerly included).

Tufted annuals or short-lived perennials. The inflorescence is digitate with 3-6 spikes. The spikelets are laterally compressed with 4-6 florets and arranged on one side of the rachis. The upper florets are inflated, and each floret is awned, therefore each spikelet has 4-6 awns.

Oxychloris is closely related and easily confused with the genera *Chloris* and *Enteropogon*.


Subfamily: Chloridoideae; Tribe: Cynodonteae

World = 1, Australia = 1

Townsville species


Oxychloris scariosa

Winged Windmill Grass, Winged Chloris



Key to the species of *Oxychloris* and similar species


1. Second and third lemmas very broad, sharply recurved at maturity (wing-like)..... *Oxychloris scariosa*
 Lemmas not very broad, not wing-like.....2
2. Lowest lemma dorsally compressed (lying on front or back when placed on a flat surface).....
*Enteropogon*
 Lowest lemma laterally compressed (lying on the side when placed on a flat surface)*Chloris*



Oxychloris spikelet without glumes



Enteropogon spikelet



Chloris inflata spikelet

***Oxychloris scariosa* - Winged Windmill Grass, Winged Chloris**

Derivation

scariosa - Latin for thin, referring to the texture of the glumes or lemmas.

Habit

An annual or short-lived perennial grass, the culms are 15-50 cm tall.

Inflorescence

The inflorescence is digitate with 3-6 branches, erect to spreading, 1-8-6 cm long. The spikelets are 5.5-9 long with 4-9 florets, the second and third lemmas are very broad and wing-like.

Habitat

This species are usually on clay soils in depressions, often with saline conditions, but also found on sandy and loam soils, especially in disturbed areas such as roadsides.



Spikelet



Inflorescence

Panicum

Latin name for common millet, from Latin word *panis* (bread).

Annuals or perennials, of various habit but commonly tufted. The inflorescence is an open or contracted panicle, with secondary branches. The spikelets are solitary or in pairs, the surfaces of the fertile floret are smooth and often shiny. The lower glume varies from being very short to equal to the spikelet.

Species from this genus are from diverse habitats, including aquatic grasses.

Subfamily: Panicoideae; Tribe: Paniceae

World = 1180, Australia = 40

Members of this genus could be confused with *Urochloa*, *Megathyrsus*, *Whiteochloa* and *Arthrargrostis*.


Townsville species

<i>Panicum decompositum</i> var. <i>decompositum</i>	Australian Millet
<i>Panicum decompositum</i> var. <i>tenuius</i>	Australian Millet
<i>Panicum effusum</i>	Hairy Panic
<i>Panicum laevinode</i>	Pepper Grass
<i>Panicum mindanaense</i>	
<i>Panicum mitchellii</i>	
<i>Panicum paludosum</i>	Swamp Panic
<i>Panicum seminudum</i>	
<i>Panicum simile</i>	Two-coloured Panic
<i>Panicum trichoides</i>	




Key to the species of *Panicum* and related genera

1. Inflorescence once-branched..... *Urochloa*
- Inflorescence with secondary branches.....2




Inflorescence
once branched





Inflorescence with
secondary branches

2. Fertile lemma transversely rugulose (with a wrinkled appearance) *Megathyrsus maximum*
- Fertile lemma smooth and shiny3
3. Spikelets laterally compressed4
- Spikelets dorsally compressed.....5



Spikelets laterally compressed



Spikelets dorsally compressed

4. Plant 45-110 cm tall; grows on sandy, alluvial soils; inflorescence open or contracted, branches persistent *Whiteochloa airoides*
- Plant 17-60 cm; grows on rocky hillsides; inflorescence open; branches deciduous
.....*Arthragrostis deschampsoides*
5. Plants aquatic or growing in swamp.....*Panicum paludosum*
- Plants not aquatic, not growing in swamps,6
6. Spikelets 0.8-1.4 mm long*Panicum trichoides*
- Spikelets 2 mm or more long7
7. Culms 1-2 m tall *Panicum mitchellii*
- Culms less than 1 m tall.....8
8. Leaf blades hairy (use a hand lens).....9
- Leaf blades glabrous 10
9. Perennial; culms vastly hairy..... *Panicum effusum*
- Annual; culms glabrous or slightly hairy; *Panicum seminudum*
10. Inflorescence branches, lower ones usually whorled or concealed by leaf sheath
..... *Panicum decompositum*
- Inflorescence branches, lower ones solitary and spreading 11

11. Lower glume $\frac{1}{8}$ - $\frac{1}{3}$ of spikelet length..... *Panicum laevinode*
Lower glume $\frac{1}{2}$ or more of spikelet length..... 12
12. Lower glume $\frac{1}{2}$ - $\frac{2}{3}$ of spikelet length.....*Panicum simile*
Lower glume \pm equal to spikelet length *Panicum mindanaense*

Pictorial key to the species of *Panicum*

The lower glume length compared with the spikelet length is a good way to identify the species.

Lower glume $\frac{1}{3}$ and less of spikelet length

Found over a wide area of woodlands
A perennial; leaf-blades 25-50 cm long; inflorescence with whorled branches at lower nodes



Panicum decompositum

An annual; leaf-blades 5-20 cm long; inflorescence not with branches whorled



Panicum laevinode

Aquatic or swamp grasses
Rhizomes thick and spongy



Panicum paludosum

Lower glume $\frac{1}{3}$ - $\frac{2}{3}$ of spikelet length

Inflorescence small, 4-20 cm long,
Spikelet small, 1-1.5 mm long
Culms 15-100 cm tall



Panicum trichoides

Inflorescence 8-50 cm long
Spikelet 2-3 mm long
Culms 2-100 cm tall
Culm internodes and base distinctly hairy



Panicum effusum

Inflorescence a fairly narrow panicle
Spikelet 2.25-3.5 mm long
Culms 20-70 cm tall,



Panicum simile

Inflorescence a large open panicle
Spikelet 2.5-3 mm long
Culms 100-200 cm tall



Panicum mitchellii

Lower glume at least $\frac{3}{4}$ of spikelet length

Lower glume shorter than spikelet (0.75-0.95)

Lower glume \pm equal to spikelet length



Panicum seminudum



Panicum mindanaense

***Panicum decompositum* - Australian Millet**

Derivation

decompositum - Latin for much divided, the inflorescence is much branched.

Habit

A perennial, densely tufted grass, the culms are 30-80 cm tall.

Inflorescence

The panicle is open, it is usually whorled at lower nodes, 15-35 cm long. The spikelet is 2.3-3.5 mm long. The lower glume is 25-33% of length of spikelet.

Habitat

The species occur over a wide area of woodlands.



Spikelet

There are two varieties

Spikelets 2.3-2.8 mm long *Panicum decompositum* var. *tenuis*

Spikelets 3-3.5 mm long..... *Panicum decompositum* var. *decompositum*



***Panicum effusum* - Hairy Panic**

Derivation

effusum - Latin for spread out, the inflorescence an open panicle.

Habit

A perennial, tufted grass, the culms are 2-100 cm tall. The stems and leaf sheaths are hairy, they are usually tubercle-based.

Inflorescence


The panicle is open, 8-50 cm long. The spikelet is 2-3 mm long. The lower glume is 50-65 % of length of spikelet.

Habitat

The species is widespread in woodland and on disturbed sites.



Spikelet



***Panicum laevinode* - Pepper Panic**

Derivation

laevinode - Latin for (*laevis*) smooth and (*nodus*) nodes.

Habit

An annual or perennial short-lived, tufted grass, the culms are 15-120 cm tall.

Inflorescence

The panicle is open, 5-30 cm long. The spikelet is 2.3-3.5 mm long. The lower glume is 33-50 % of length of spikelet.

Habitat

The species is spread a widespread on good soils.



Spikelet



Panicum mindanaense

Derivation

mindanaense - from Mindanao Island (Philippines).

Habit

An annual or perennial grass, the culms are 20-80 cm tall.

Inflorescence


The panicle is open, 10-30 cm long. The spikelet is 2-3.5 mm long. The lower glume is \pm equal to spikelet length.

Habitat

This species grows in tropical and subtropical rainforest and grasslands.



Spikelet



Panicum mitchellii

Derivation

mitchellii - in honour of Thomas Livingstone Mitchell (1792-1855), a Scots-born Australian surveyor and explorer.

Habit

A robust, erect, perennial grass, the culms are 100-200 cm tall.

Inflorescence

The panicle is open, 20-40 cm long. The spikelet is 2.5-3 mm long. The lower glume is 50-60% of length of spikelet.

Habitat

This species grows in tropical and subtropical rainforest and in woodlands.



Spikelet



***Panicum paludosum* - Swamp Panic**

Derivation

paludosum - from the Latin *palus* (swamp) and *-osa* (abundance), it is growing in swampy places.

Habit

A perennial, aquatic, spongy grass, the culms are 25-130 cm tall.

Inflorescence

The panicle is open, 10-30 cm long. The spikelet is 3.2-4.1 mm long. The lower glume is 10-25% of length of spikelet.

Habitat

This species grows in still and slowing water.



Spikelet



Panicum seminudum

Derivation

seminudum - from the Latin *semi-* (half) and *nuda* (bare), only part of the plant bearing hairs.

Habit

An erect, annual grass, the culms are 40-90 cm tall. The leaves and culms can be glabrous or hairy.

Inflorescence

The panicle is open, 15-30 cm long. The spikelet is 3.1-3.7 mm long. The lower glume is 75-95% of length of spikelet.

Habitat

This species grows in rainforests and woodlands



Spikelet



***Panicum simile* - Two-coloured Panic**

Derivation

simile - Latin for like, it is readily confused with one or more other species, e.g. *Panicum effusum*.

Habit

A slender, annual grass, the culms are 20-70 cm tall.

Inflorescence

The panicle is open, 4-20 cm long. The spikelet is 2.25-3.5 mm long. The lower glume is 50% of length of spikelet. The spikelets are pale to golden and purple or irregularly blotched with purple.

Habitat

This species grows in poor soil in woodlands.



Spikelet



Panicum trichoides

Derivation

trichoides - from the Greek *thrix* (hair) and *-oides* (resembling), the spikelets are sparsely hirsute.

Habit

An annual grass, the culms are 15-100 cm tall. The leaves are short, 1.5-8 cm long and 0.5-2 cm wide.

Inflorescence

The panicle is open, 4-20 cm long. The spikelet is 1-1.5 mm long. The lower glume is 25-50% of length of spikelet.

Habitat

This species grows well in moist, shaded forested areas at lower elevations.



Spikelet



Paspalidium

From the Greek *eidos* (shape) and *Paspalum* (another grass genus) 'shaped like *Paspalum* or a diminutive of *Paspalum*.


Annuals or perennials (often aquatic), rhizomatous or tufted to decumbent. The inflorescence is usually a once-branched panicle; however, the racemes are appressed to the main axis or a single raceme. The rachis of each raceme extending as a bristle beyond the point of attachment of the last spikelet. The spikelets are solitary or in pairs.

Subfamily: Panicoideae; Tribe: Paniceae
World = 40, Australia = 22

Townsville species


<i>Paspalidium criniforme</i>	
<i>Paspalidium distans</i>	Spreading Panic Grass
<i>Paspalidium flavidum</i>	
<i>Paspalidium gracile</i>	Slender Panic
<i>Paspalidium rarum</i>	Rare Panic
V <i>Paspalidium udum</i>	

Vulnerable Species (V) are threatened plants.




Key to the species of *Paspalidium* and related species

1. Inflorescence terminating in a spikelet *Whiteochloa airoides*
 Inflorescence terminating in a bristle 2



Inflorescence terminating in a bristle




Inflorescence in a spikelet

2. Racemes reduced to 1 or 2 spikelets *Paspalidium rarum*
 Racemes with more than 2 spikelets 3
3. Aquatic or semi-aquatic; culms thick and spongy *Paspalidium udum*
 Terrestrial; culms are slender and not spongy 4




Paspalidium udum culms are spongy

4. Spikelets irregularly arranged on raceme axes 5
 Spikelets regularly arranged on raceme axes 6
5. Spikelets up to 1 mm wide *Paspalidium criniforme*
 Spikelets more than 1 mm wide *Paspalidium gracile*
6. Leaves width 1-6.5 mm; spikelets dorsally compressed, 1.8-2.2 mm long *Paspalidium distans*
 Leaves width 4-8 mm; spikelets laterally compressed or terete, 2.5-3 mm long
 *Paspalidium flavidum*



Spikelet dorsally compressed



Spikelet laterally compressed

T.S. = transverse section

Paspalidium criniforme

Derivation

criniforme - from the Latin *crinum* (hair) and *forma* (appearance).

Habit


A perennial, tufted grass, the culms are 10-60 cm tall. The leaves are 1-3 mm wide.

Inflorescence


The panicle is of racemes 4-6. The inflorescence is 2.5-23 cm long. The spikelet is 2-2.5 mm long.

Habitat

This species grows in forest and woodland.



Spikelet



***Paspalidium distans* - Spreading Panic Grass**

Derivation

distans - from the Latin *disto* (be apart), the spikelets widely separated in inflorescence.

Habit

A perennial, tufted grass, the culms are 30-70 cm tall. The leaves are 1-6.5 mm wide.

Inflorescence

The panicle is of racemes 4-10, distant. The inflorescence is 6-16 cm long. The spikelet is 1.7-2.2 mm long.

Habitat

This species grows in forest, woodland and shrub woodlands.



Spikelet



Paspalidium flavidum

Derivation

flavidum - from the Latin *flavidus* (pale yellow), the spikelets yellow.

Habit

A perennial, tufted grass, the culms are 10-100 cm tall. The leaves are 4-8 mm wide.

Inflorescence

The panicle is of racemes 4-9, distant. The inflorescence is 7-30 cm long. The spikelet is laterally compressed or terete and is 1.9-3 mm long.

Habitat

This species grows in rainforest and woodlands.



Spikelets



***Paspalidium gracile* - Slender Panic**

Derivation

gracile - Latin for slender, the culms or inflorescences slender.

Habit

A perennial, wiry, tufted grass; the culms are 10-90 cm tall. The leaves are 1-4 mm wide.

Inflorescence

The panicle is of racemes 5-10, with a marked constriction. The inflorescence is 5-10 cm long. The spikelet is 2-3 mm long.

Habitat

This species grows in wet sclerophyll forests, woodlands, shrublands and grasslands.



Spikelet



***Paspalidium rarum* - Rare Panic**

Derivation

rarum - Latin for far apart, the spikelets far apart in panicle.

Habit

An annual, tufted grass, the culms are 15-45 cm tall. The leaves are 1-3.5 mm wide.

Inflorescence

The inflorescence of this species is unique; typically, the lateral branches are reduced to a single spikelet. The inflorescence is a raceme, 5-13 cm long. The spikelets are 2.7-3.5 mm long.

Habitat

This species grows in woodlands, shrublands and grasslands.



Spikelet



Paspalidium udum

Derivation

udum - Latin for damp, growing in swampy areas.

Habit

A perennial, semi-aquatic or aquatic grass, with elongated rhizomes. The culms are 40-90 cm tall; they are spongy. The leaves are 5-11 mm wide.

Inflorescence

The panicle of is of racemes 10-16, they are close together. The inflorescence is 12-19 cm long. The spikelet is 2.7-3 mm.

Habitat

This species is considered a vulnerable plant.



Spikelet



Spongy culm



Paspalum

From the Greek word *paspalos* (a kind of millet).



Perennials (usually) or annuals, rhizomatous or stoloniferous or tufted or decumbent. The inflorescence is either digitate or a racemose once-branched panicle. The spikelets are solitary or paired and arranged along one side of rachis (the axis or branch of the inflorescence). The genus is best recognised by its plano-convex spikelets with the upper glume adaxial (the side facing to the rachis), often with a hemispherical or oblong shape. The lower glume is usually absent.

Some species are similar to the genus *Axonopus*; however, this genus has the upper glume abaxial (the side facing away from the rachis).

Subfamily: Panicoideae; Tribe: Paniceae
World = 480 , Australia = 21

Townsville species

* <i>Paspalum conjugatum</i>	Sour Grass, Johnson River
Grass	
* <i>Paspalum dilatatum</i>	Paspalum
<i>Paspalum distichum</i>	Water Couch
* <i>Paspalum notatum</i>	Bahia Grass
<i>Paspalum scrobiculatum</i>	Ditch Millet
<i>Paspalum vaginatum</i>	Saltwater Couch



Key to the species of *Paspalum* and related genera

- Lower floret adjacent to the inflorescence axis*Axonopus*
Lower floret positioned away from the inflorescence axis2
- Inflorescence of 2 racemes at end of a culm, sometimes with a third one below3
Inflorescence of 1 - many racemes along a common elongated axis7
- Spikelets 1.2-1.8 mm long *Paspalum conjugatum*
Spikelets 2.5-4.5 mm long4
- Upper glume finely pubescent*Paspalum distichum*
Upper glume glabrous5
- Plants tufted *Paspalum scrobiculatum*
Plants stoloniferous or rhizomatous6
- Spikelets about twice as long as broad; usually aquatic *Paspalum vaginatum*
Spikelets almost circular *Paspalum notatum*
- Plant usually less than 1 m tall; Spikelet without long silky hairs around margin.....*Paspalum scrobiculatum*
Robust plant 1-2 m tall; Spikelets conspicuously silky hairy around margin, hairs as long as or
exceeding spikelet length*Paspalum dilatatum*

***Paspalum conjugatum* - Sour Grass, Johnson River Grass**

Derivation

conjugatum - from the Latin *conjugo* (yoke together), the inflorescence branches paired.

Habit

A perennial, stoloniferous grass, the culms are 20-100 cm tall.

Inflorescence

The panicle is digitate with 2 branches, 4-17 cm long. The spikelets are 1.2-1.8 mm long.

Habitat

This non-native species grows in wet roadsides or ditches and spreads quickly by stolons.



Spikelets



***Paspalum dilatatum* - Paspalum**

Derivation

dilatatum - from the Latin *differe* (spread abroad) and *-ata* (possessing), the racemes spreading in pseudo-vercils.

Habit

A perennial, tufted grass, the culms are 25-180 cm tall.

Inflorescence

The panicle is once branched of 3-10 racemes, 4-12.5 cm long. The spikelets are 2.7-3.9 mm long.

Habitat

This non-native species is a weed, preferring moist ditches, disturbed areas, and is a common invader of gardens.



Spikelets



***Paspalum distichum* - Water Couch**

Derivation

distichum - from the Greek *distichos* (two-rowed), it has conspicuously two-rowed spikelets or leaves.

Habit

A perennial, stoloniferous grass, the culms are 8-60 cm tall.

Inflorescence

The panicle is digitate with 2 to 3 branches, 1.5-8 cm long. The spikelets are 2.6-4 mm long.

Habitat

This species prefers moist ditches or coastal areas and is associated with fresh or salt water.



Spikelets



***Paspalum notatum* - Bahia Grass**

Derivation

notatum - from the Latin *noto* (mark), the spikelets are multi-coloured.

Habit

A perennial, rhizomatous grass, the culms are 15-75 cm tall.

Inflorescence

The panicle is digitate with 2 to 3 branches, 2.5-13 cm long. The spikelets are 2.5-3.8 mm long.

Habitat

This non-native species can dominate disturbed soils, pastures and roadsides.



Spikelet



***Paspalum scrobiculatum* - Ditch Millet**

Derivation

scrobiculatum - from the Latin *scrobis* (ditch), *-ulus* (diminutive) and *-atus* (possessing), the glumes or lemmas furrowed.

Habit

A perennial tufted grass, the culms are 10-150 cm tall.

Inflorescence

The panicle is digitate with 2 branches or is once branched of 1-20 racemes, 2-15 cm long. The spikelets are 1.4-3 mm long.

Habitat

This species grows in poor soil, damp soils, open cultivated places and wastelands.

This is a widespread, morphologically variable species. This species is notoriously difficult to treat taxonomically as it is possibly an aggregate swarm of apomicts.



Spikelets



***Paspalum vaginatum* - Saltwater Couch**

Derivation

vaginatum - from the Latin *vagina* (sheath) and *-ata* (possessing), the inflorescences concealed or partially concealed in uppermost leaf-sheaths.

Habit

A perennial, stoloniferous and rhizomatous grass, the culms are 8-60 cm tall.

Inflorescence

The panicle is digitate with 2 branches or is once branched of 3-5, 1.5-7.5 cm long. The spikelets are 2.5-4.5 mm long.

Habitat

This species grows in wet areas near saline or brackish water, chiefly coastal habitats. This grass is sometimes called non-native in Australia.



Spikelets



Perotis

From the Greek *peros* (deficient, mutilated) and *ous* (an ear), the lemma is awnless.

Tufted annuals or rarely perennials. The inflorescence is a single raceme or spike. The spikelets are solitary and both glumes are long and awned. The single fertile floret is very small and held within the glumes.


Subfamily: Chloridoideae; Tribe: Cynodonteae

World = 20, Australia = 3

Townsville species

Perotis rara

Comet Grass



***Perotis rara* - Comet Grass**

Derivation

rara - is Latin for far apart and refers to the spikelets being spaced out along the inflorescence.

Habit

An annual grass, the culms are 15-40 cm tall. The leaves are erect.

Inflorescence

The inflorescence is 10-28 cm and about 2.5 cm wide. The spikelets are often pointing downwards at maturity, hence the common name, Comet Grass.

Habitat

This species occurs in a variety of habitats from coastal dunes and vine thickets to open *Eucalyptus* forests, often on sandy soils near streams or waterholes or in other disturbed areas.



Phragmites

Reed Grasses

From the Greek *phragma* (a hedge or screen) and *-ites* (resembling), referring to the way in which the grass grows like a fence along a riverbank.


Tall rhizomatous perennials, often forming dense stands, culms erect, reed-like, 1-5 m tall. Grows in permanently wet places, especially along the banks of slow-running streams and swamps. The inflorescence is a large 10-50 cm, fairly dense feathery panicle. The spikelets are solitary, and comprising of numerous, 3-12 florets; the rachilla (the axis of the grass spikelet) is covered with long silky hairs and the lemma is glabrous. Once the spikelets have fallen, the inflorescence remains on the plant and therefore does not look feathery.

Closely related and similar to *Arundo donax*

Subfamily: Arundinoideae; Tribe: Arundineae
World: = 17, Australia = 2



Townsville species

<i>Phragmites australis</i>	Cane Grass, Bamboo Reed
<i>Phragmites karka</i>	Tropical Reed



Key to the species of *Phragmites* and similar species

1. Lemmas hairy; rachilla not hairy *Arundo donax*
 Lemmas not hairy (glabrous); rachilla (axis of spikelet) hairy 2



2. Leaf blades scabrid beneath (at least in the upper half); the upper glumes are up to 6 mm long *Phragmites karka*
 Leaf blades smooth beneath; the upper glumes longer than 6 mm *Phragmites australis*



Arundo donax



Phragmites karka



Phragmites australis

***Phragmites australis* - Cane Grass, Bamboo Reed**

Derivation

australis - of the south.

Habit

A perennial grass, culms erect, reed-like, 1-3 m tall. The leaf blades are smooth beneath.

Inflorescence

The panicle is erect and dense, with lower branches spikelet-bearing to base, with 200-500 peduncles per sheath; 20-50 cm long, 6-15 cm wide.

Habitat

This grass grows along rivers and streams in temperate zones of the planet; sometimes found in semi-salt water and tidal systems, and along saline springs, commonly along rivers or in floodplains.



Spikelet



***Phragmites karka* - Tropical Reed**

Derivation

karka - origin obscure.

Habit

A perennial grass with culms erect, reed-like, 1.5-4 m tall. The leaf blade surface is scabrid beneath (at least in the upper half).

Inflorescence

The panicle is diffuse and partially drooping, with lower branches for 1-4 cm; 30-50 cm long, 10-20 cm wide.

Habitat

Commonly along rivers or in floodplains.




Pseudopogonatherum

From Greek *pseudo* (false) and *Pogonatherum*, referring to the similarity to the genus *Pogonatherum*.

Tufted, slender annuals, usually to less than 1 m tall, the thin leaves are mostly basal. The inflorescence is digitate or subdigitate and the racemes are erect. The spikelets are in pairs, similar and unevenly pedicelled. The spikelets are very small (1-2.5 mm long) and have long awns (15-30 mm long), they break up at maturity making it difficult to see the pairing.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 6, Australia = 2



Townsville species

Pseudopogonatherum contortum

Key to the species of *Pseudopogonatherum* and similar species

1. Each spikelet consists of 2-4 florets with the basal floret bisexual and the upper florets reduced; fertile florets fall from glumes *Chloris*
Each spikelet consists of 2 florets with the basal floret sterile and the upper floret fertile: fertile floret fall with glumes 2



Chloris



Pseudopogonatherum



Eulalia

2. Perennial; awns not intertwining at maturity; sessile spikelets usually more than 3 mm long *Eulalia*
Delicate annual; awns intertwining at maturity; all spikelets very small, 1.5-2.5 mm long *Pseudopogonatherum*

Pseudopogonatherum contortum

Derivation

contortum - awns are twisted when dry.

Habit

An annual, tufted, slender, erect grass, the culms are 20-100 cm tall.

Inflorescence

The digitate inflorescence consists of 3-20 racemes each 3-7 cm long. The spikelets and awns are dark brown, and the inflorescence branches and spikelets are covered with white hairs. The spikelets are 1.5-2.5 mm long.

Habitat

This species occurs sporadically in open woodland following summer rainfall.

Pseudopogonatherum contortum can be mistaken for *Chloris pumilio* or *Melinis minutiflora*.



Spikelets



Pseudoraphis

Mud Grasses

From the Greek *pseudo* (false) and *raphis* (needle), referring to the bristle-like point protruding beyond the uppermost spikelet.

Perennial aquatic or marsh grasses, culms mostly prostrate and often floating in water. The inflorescence is a racemose or contracted panicle; the slender branches are often reduced to one or two spikelets. The spikelets are solitary, narrow and awned or unawned. The apex of the spikelets is acuminate, and a bristle protrudes beyond the uppermost spikelet. Other genera with a single bristle subtending the spikelet are *Paspalidium* and some species of *Setaria*.

Subfamily: Panicoideae; Tribe: Paniceae
World = 7, Australia = 5

Townsville species

Pseudoraphis spinescens Spiny Mud Grass

Key to the species of *Pseudoraphis spinescens* and related genera


1. Primary branches of panicle alternate on opposite sides of the main axis *Paspalidium*
Primary branches of panicle originate from all sides 2



Primary branches from
one or two sides



Primary branches from all
sides



2. Inflorescence is an open panicle *Pseudoraphis spinescens*
Inflorescence is a constricted panicle *Setaria*



*Pseudoraphis
spinescens*



Setaria

***Pseudoraphis spinescens* - Spiny Mud Grass**

Derivation

spinescens - from the Latin *spinesco* (become thorny), the inflorescence branches terminally pungent.

Habit


A semi-aquatic perennial grass, with the stems floating in shallow water or forming a sward on mud. The culms are 10-50 cm tall.

Inflorescence



The inflorescence is a panicle of racemes; the central inflorescence axis is 3-11 cm long. A bristle is formed beyond the uppermost spikelet.

Habitat

This species grows in shallow water or mud beside rivers and lagoons.



Spikelet at the end of the inflorescence




Sacciolepis

From the Greek *sakkion* (small bag) and *lepis* (scale), alluding to the shape of the upper glume.

Annual or perennials grasses. Recognised by the spiciform (spike-like panicle with short branches) inflorescences and the gibbous (hump-backed), ribbed spikelets. Species from this genus grow in or near water or in wet places.

Subfamily: Panicoideae; Tribe: Paniceae
World = 31, Australia = 2



Townsville species

Sacciolepis indica Indian Cupscale Grass

Sacciolepis indica - Indian Cupscale Grass

Derivation

indica - from India.

Habit

An annual, erect or creeping grass, the culms 10-80 cm tall.

Inflorescence

The panicle is spicate, 1-13 cm long, the spikelets are solitary and 2-3.5 mm long.

Habitat

This species may be found growing among more vigorous grasses which give the plants some support and protection. This grass is found growing in damp areas.



Spikelet



Sarga

Sorghum


The meaning of *Sarga* is not given by the author and so is obscure. Recent studies of the Australian *Sorghum* species, have suggested three distinct lineages, and these species have been divided into three genera, *Sorghum*, *Sarga* and *Vacoparis*.

Tufted annuals or perennials. The inflorescence is an open or contracted panicle. The spikelets are in pairs (with terminal triplets), one sessile and one pedicelled. The mature sessile spikelets are dark reddish brown almost black. The sessile spikelets usually have long awns (1.5-8.5 cm long) and the pedicelled spikelet is well-developed.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 9, Australia = 5

Townsville species

Sarga plumosum



Key to the species of *Sarga*, *Sorghum* and *Vacoparis*

1. Pedicelled spikelet reduced to narrow, linear glumes *Vacoparis laxiflorum*
 Pedicelled spikelet containing empty lemmas or male 2
2. Awn on sessile spikelet 25-80 mm long *Sarga plumosum*
 Awn on sessile spikelet 0-20 mm long *Sorghum*



Vacoparis laxiflorum



Sarga plumosum



Sorghum nitidum
f. *aristatum*



Sorghum x alnum

***Sarga plumosum* - Plume Sorghum**

Derivation

plumosum - from the Latin for feathery, referring to the long hairs giving the pedicels a feathery appearance.

Habit

A perennial, tufted grass, the culms are 100-300 cm tall. The stems have distinctive bearded nodes. Leaves usually have a white midrib and vary in colour from shiny green to blue green.

Inflorescence

The inflorescence is usually a dense panicle 12-45 cm long, with dark red-brown spikelets. The pedicelled spikelets containing empty or male lemmas; the sessile spikelets are 6-18 mm long, the lemma awn is 2.5-8 cm long.

Habitat

This species is found on sands, red earths and heavy loams, it grows in swamps, claypans, watercourses, waterholes and valleys.



Spikelets



Bearded node



Schizachyrium

From the Greek *schizen* (to split) and *achuron* (chaff), alluding to the bilobed lemma of the sessile spikelet. *Schizachyrium* can be pronounced “shize-ah-KIR-ee-um”.

Perennials or annuals, tufted to decumbent. The inflorescence is a single raceme or a panicle with a spathe subtending each raceme. The spathe is sometimes not obvious, and the inflorescence looks like a spike. The spikelets are in pairs, one sessile and one pedicelled, which is much reduced. The sessile spikelet is awned and arises from a bilobed lemma.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 78, Australia = 9

Townsville species

Schizachyrium fragile

Fire Grass, Red Spathe Grass

Schizachyrium occultum

Schizachyrium pseudeulalia


The spikelets are usually clothed in silky white hairs, similar to *Mnesithea formosa*, however *Schizachyrium* spikelets are awned.

Key to the species of *Schizachyrium* and similar species


1. Spikelets not awned *Mnesithea formosa*
Spikelets sessile distinctly awned.....2



Mnesithea formosa spikelets



Schizachyrium fragile spikelets




2. Sessile spikelets ±laterally compressed, 7-13 times as long as wide; leaf-blades 0.6-2 mm wide *Schizachyrium occultum*
Sessile spikelets dorsally compressed 4-8 times as long as wide; leaf-blades 0.8-5.5 mm wide.....3
3. Lower glume of sessile spikelet thin, dry, or flimsy like paper; papery, winged, at least above the middle *Schizachyrium fragile*
Lower glume of sessile spikelet resembling or having the texture of leather), not winged *Schizachyrium pseudeulalia*



Schizachyrium fragile



Schizachyrium pseudeulalia



***Schizachyrium fragile* - Fire Grass, Red Spathe Grass**

Derivation

fragile- Latin for weak, the inflorescences readily disarticulating.

Habit

An annual, tufted grass, the culms are 10-75 cm tall. The leaves are 2-8 cm long and 1-3 mm wide.

Inflorescence

The panicle is nearly completely enclosed in the spathes. The sessile spikelet is dorsally compressed and is 5-8 as long as wide; the lower glume is winged at least above the middle.

Habitat

This species is a fairly common annual on sandy soils from the savanna regions.

Schizachyrium fragile could be confused with *Mnesithea formosa*. The spikelets of *Schizachyrium* have awns



Spikelets



Schizachyrium occultum

Derivation

occultum - from the Latin *occultus* (hidden).

Habit

The annual, slender grass, the culms are 10-30 cm tall. The leaves are 1.5-3.5 cm long and 0.6-2 mm wide.

Inflorescence

The panicle is subtended by a spatheole. The sessile spikelet is laterally compressed and is 9-13 times as long as wide.

Habitat

This species is found in sandy soil in *Eucalyptus* forest. It is usually associated with other *Schizachyrium* species.



Spikelets



Schizachyrium pseudéalalia

Derivation

pseudéalalia - false *Eulalia*, it is often mistaken for *Eulalia*.

Habit


The annual grass, the culms are 30-90 cm tall. The leaves are 1-9 cm long and 0.8-5.5 mm wide.

Inflorescence

The panicle is subtended by a spatheole. The sessile spikelet is dorsally compressed and is 5.5-8 times as long as wide.

Habitat

This species is sandy soil.



Spikelets



Sehima


From the Egyptian vernacular name of the type; named after the Arab sign *Sahim*.

Tufted annuals or perennials. The inflorescence is a single, curved raceme. The spikelets are paired, one sessile and one pedicelled, and are partially embedded in rachis. The sessile spikelet is awned.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 13, Australia = 1

Townsville species

Sehima nervosum White Grass



Key to the species of *Sehima* and similar species

1. Inflorescence is a raceme2
- Inflorescence spike-like, but composed of 2 erect, closely adpressed raceme*Ischaemum*



Ischaemum



Sehima nervosum



Heteropogon triticeus



Heteropogon contortus

2. Awns are 20-40 mm long; pedicelled spikelets is the same length as fertile *Sehima nervosum*
- Awns are 50-140 mm long; pedicelled spikelet longer than fertile..... *Heteropogon*

***Sehima nervosum* - White Grass**

Derivation

nervosum - from the Latin *nervus* (nerve) and *-osa* (abundance) which refers to the conspicuous nerves in the glumes or lemmas.

Habit

A perennial, tufted grass, the culms are 30-100 cm tall.

Inflorescence

The inflorescence is a single raceme, 3-12 cm long. The pedicels and internodes of the inflorescence are densely bearded with white hairs. The sessile spikelet is 6-10 mm long; the awns are 20-40 mm long. This species can be recognised by the lower glume of the pedicelled spikelet it is slightly asymmetrical and strongly nerved.

Habitat

This species is usually found growing in dry areas in poor soils, often associated with *Themeda triandra*.



Spikelet pairs



Pedicelled spikelet showing asymmetric glume




Setaria

Pigeon Grasses

From Latin *seta* (bristle), referring to the bristly inflorescences.

Annuals or perennials, rhizomatous or stoloniferous or tufted or decumbent. The inflorescence is a spicate panicle (spike-like with short branches). The spikelets are solitary, usually with numerous bristles beneath them. The lemma of the upper floret is transversely rugose (wrinkled or with a creased surface).



Setaria species looks similar to *Cenchrus* (including *Pennisetum*), however spikelets fall from the plant without any bristle attached leaving a rachis with the bristles attached.

Subfamily: Panicoideae; Tribe: Paniceae
World = 190, Australia = 28

Townsville species

- * *Setaria pumila* subsp. *subtesselata* Pale Pigeon Grass
- * *Setaria sphacelata* South African Pigeon Grass
- Setaria surgens* Pigeon Grass
- Setaria* sp. Mount Larcom

Key to the species of *Setaria* and related genera

1. Spikelets falling at maturity with bristles or spines attached *Cenchrus*
Spikelets falling at maturity without any bristles or spines attached 2



Cenchrus



Setaria



2. Inflorescence an open panicle *Setaria* sp. Mount Larcom
Inflorescence a tightly contracted false spike 3
3. Inflorescence is 3-50 cm long; perennial with rhizomes *Setaria sphacelata*
Inflorescence is 1-10 cm long; annual, no rhizomes 4
4. Bristles are white cream; lower glume 30-40% of length of spikelet *Setaria surgens*
Bristles are rusty coloured; lower glume 30-70% of length of spikelet
..... *Setaria pumila* subsp. *subtesselata*

***Setaria pumila* subsp. *subtesselata* - Pale Pigeon Grass**

Derivation

pumila - Latin for dwarf or low growing, the habit is typically depauperate.

subtesselata - Latin for somewhat almost a mosaic-like refers the fertile lemma.

Habit

An annual, tufted grass, the culms are 5-130 cm tall.

Inflorescence

The panicle is solid, spiciform, linear, 1-10 cm long. The spikelets are 1.5-2.5 cm, the lower glume 30-70% of length of spikelet.

Habitat

This non-native species is mainly a weed of disturbed sites, roadsides, footpaths and cultivation.



Spikelet



***Setaria sphacelata* - South African Pigeon Grass**

Derivation

sphacelata - Latin for speckled with brown or black, the apices of the fertile lemmas are purple or black.

Habit

A perennial, tufted grass, the culms are 20-300 cm tall.

Inflorescence

The panicle is solid, spiciform, linear, 3-50 cm long. The spikelets are 1.5-3.5 cm long; the lower glume is 25-50% of length of spikelet.

Habitat

This non-native species is a weed of roadsides, waterways, grasslands, open woodlands, parks, disturbed sites and waste areas.



Spikelet



***Setaria surgens* - Pigeon Grass**

Derivation

surgens - from the Latin *surgo* (raise), the culms long, ascending.

Habit

An annual grass, the culms are 20-60 cm tall.

Inflorescence

The panicle is solid, spiciform, linear, 2.5-5 cm long. The spikelets are 2.5-2.9 mm long; the lower glume is 30-40% of length of spikelet.

Habitat

This species grows in woodlands, shrublands and grasslands.



Spikelet



***Setaria* sp. Mount Larcom**

Habit

A perennial grass.

Inflorescence

The panicle is open. Spikelets 3.3-4.0 mm long; upper glume c. 80% of spikelet; bristles > c. 15 mm long.



Spikelet




Sorghum

From *sorgho*, the Italian name for the genus. *Sorghum*, in the broadest sense (*sensu lato*), is a genus is widespread in tropical and subtropical parts of the world. Grain sorghum is an important cereal being a staple food grain in west Africa and widely used for feeding livestock in western countries. It is believed to have been domesticated over 3000 years ago and many varieties have been developed.

Recent studies of the Australian *Sorghum* species, have suggested three distinct lineages, and these species have been divided into three genera, *Sorghum*, *Sarga* and *Vacoparis*.

Sorghum, in the narrowest sense (*sensu stricto*), are robust annuals or perennials. The inflorescence is an open or contracted panicle. The spikelets are in pairs (with terminal triplets), one sessile and one pedicelled. The mature sessile spikelets are usually dark reddish-brown almost black. The sessile spikelets usually have short awns (1-1.5 cm long) or are awnless, and the pedicelled spikelet is well-developed.



Subfamily: Panicoideae; Tribe: Andropogoneae
World = 218, Australia = 7

Townsville species

- * *Sorghum x alnum* Columbia Grass
- * *Sorghum bicolor* Forage Sorghum
- * *Sorghum halepense* Johnson Grass
- Sorghum nitidum*

Key to the species of *Sarga*, *Sorghum* and *Vacoparis*

1. Pedicelled spikelet reduced to narrow, linear glumes *Vacoparis laxiflorum*
Pedicelled spikelet containing empty or male lemmas 2
2. Awn on sessile spikelet 25-80 mm long *Sarga plumosum*
Awn on sessile spikelet 0-20 mm long 3



Vacoparis laxiflorum



Sarga plumosum



Sorghum nitidum



Sorghum x alnum

3. Primary branches of inflorescence not divided, bearing only terminal racemes; culms bearded *Sorghum nitidum*
 Primary branches of inflorescence divided, bearing terminal and lateral racemes; culms not bearded4



Sorghum nitidum inflorescence



Sorghum x alnum inflorescence

4. Racemes not disarticulating at maturity; leaf blades 5-70 mm wide (cultivated plant) *Sorghum bicolor*
 Racemes disarticulating readily at maturity; leaf blades 5-40 mm wide5



Sorghum bicolor inflorescence

5. Plants aggressively rhizomatous, rather robust; sessile spikelet 4-6 x 1.5-2.3 mm; seeds readily shattering at maturity; caryopsis markedly shorter than glumes *Sorghum halepense*
 Plants rhizomatous, but not aggressively; sessile spikelet 5-6.5 x 2.5 mm; seeds not readily shedding; caryopsis only slightly shorter than glumes *Sorghum x alnum*



Sorghum halepense rhizome



Sorghum x alnum rhizome

***Sorghum x alnum* - Columbia Grass**

Derivation

album - Latin for nourishing, a nutritious forage grass.

This species is a hybrid of Sorghum bicolor and Sorghum halepense.

Habit

A perennial grass, the rhizomes are short, the culms are erect, 200-320 cm tall.

Inflorescence

The panicle is open, 20-60 cm long. The pedicelled spikelets are male or neuter; the sessile spikelets are 5-6.5 mm long and 2-5 mm wide. The lemma of the sessile spikelet is awnless or awned. The awn is 8-10 mm long and the column of awn is glabrous.

Habitat

This non-native species is cultivated widely as a forage grass but has the tendency to escape and become weedy.



Spikelets



***Sorghum bicolor* - Forage Sorghum**

Derivation

bicolor - from the Latin *bis* (twice) and *color* (colour), it is two-coloured, usually with respect to spikelets or florets.

Habit

An annual, robust grass, the culms are erect, 100-600 cm tall.

Inflorescence

The panicle is open or contracted, 4-50 cm long. The pedicelled spikelets are male; the sessile spikelets are 3-10 mm long and 1.5-5 mm wide. The lemma of the sessile spikelets is usually awnless.

Habitat

This non-native species is cultivated; it is found in disturbed sites but does not persist.



Spikelets



***Sorghum halepense* - Johnson Grass**

Derivation

Habit

A perennial grass, the rhizomes are elongated; the culms are 50-300 cm tall.

Inflorescence

The panicle is open, 10-55 cm long. The pedicelled spikelets are male or neuter; the sessile spikelets are 4-6 mm long and 1.5-2.3 mm wide. The lemma of the sessile spikelet is awnless or awned. The awns are 10-16 mm long and the column of awn is pubescent.

Habitat

This non-native species is cultivated widely as a forage grass but has the tendency to become an aggressive agricultural weed.



Spikelets



Sorghum nitidum

Derivation

nitidum - from the Latin *niteo* (shine), the spikelets or lemmas are shiny.

Habit

A perennial, short-lived grass, the culms are 100-200 cm tall.

Inflorescence

The panicle is open, 10-30 cm long. The pedicelled spikelet is male or neuter; the sessile spikelets are 3-5.5 mm long and 2-5 mm wide. The lemma of the sessile spikelet is awnless or awned. The awns are 7-17 mm long and the column of awn and the column of awn is glabrous.

Habitat

This species grows on grey and brown loams and sands and in disturbed habitats



Spikelets

There are two forms. The spikelets are awned and awnless forms are recognised as separate taxa:

Spikelets awnless..... *Sorghum nitidum* forma *nitidum*

Spikelets awned *Sorghum nitidum* forma *aristatum*



Spinifex

Beach Spinifex Grasses

Spinifex is from the Latin *spina* (a thorn) and *facere* (to make), alluding to sharp pointed leaves.

Tufted and rhizomatous, dioecious (plants are either male or female) perennials. Female plants (with female or bisexual spikelets) have globose, spiny inflorescences which fall from the plant whole and roll along the beach in the wind. The spikelets of the female plants are awned. Male plants have erect inflorescences consisting of clustered racemes. The spikelets of the male plants are not awned.

The spinifex grasses of inland Australia belong to the genus *Triodia*. Species from the genus *Spinifex* grow on the sand dunes in coastal areas.

Subfamily: Panicoideae; Tribe: Paniceae

World = 7, Australia = 5

Townsville species

Spinifex sericeus Beach Spinifex, Hairy Spinifex



***Spinifex sericeus* - Beach or Hairy Spinifex**

Derivation

sericeus - from the Latin *sericus* (silken) and *-ea* (indicating resemblance), referring to the soft hairs on the leaves and shoots.

Habit

A stout, perennial grass, the culms grow to 30 cm tall with strong creeping rhizomes, with new plants produced at the nodes; leaves are bluish green.

Inflorescence

The inflorescences are of different sex and appearance, on separate plants.

The female bisexual panicle is a large, globose, spiny head of numerous sessile racemes, each of which is reduced to a single spikelet into a long, stout bristle. The spikelets are 12-16 mm long. This panicle becomes detached from the plant at maturity.


The male panicle is a terminal cluster of racemes, subtended by large, partly enclosing, silky-hairy bracts (resembling spathes) The spikelets are 6-10 mm long.

Habitat

This species is common on the sand-dunes of the coast. It is an effective sand-binder in beach reclamation.



Female bisexual inflorescence



Male inflorescence




Sporobolus

Dropseed Grasses & Rat's Tail Grasses

From the Greek *spora* (seed) and *bolos* (throwing), at maturity the seeds are squeezed out of the fruits.

Perennials or annuals, erect to decumbent, usually tufted. The inflorescence is an open or contracted panicle; many species of non-native *Sporobolus* with contracted panicles are becoming serious weeds. The spikelets are solitary. The mature grain becomes sticky when wet.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 270, Australia = 34



Spikelet

Sporobolus it is distinguished from *Eragrostis* by its 1-flowered spikelets and 1-nerved lemmas.

The species *Dinebra neesii* can be confused with *Sporobolus* species.




Townsville species


<i>Sporobolus australasicus</i>	Australian Dropseed
<i>Sporobolus caroli</i>	Fairy Grass
* <i>Sporobolus coromandelianus</i>	Small Dropseed
* <i>Sporobolus fertilis</i>	Giant Parramatta Grass
* <i>Sporobolus jacquemontii</i>	Rat's Tail Grass
<i>Sporobolus lenticularis</i>	
* <i>Sporobolus natalensis</i>	Giant Rat's Tail Grass
<i>Sporobolus sessilis</i>	Tussocky Sporobolus
<i>Sporobolus virginicus</i>	Salt Couch

Key to the species of *Sporobolus* and similar species

1. Inflorescence a one-branched panicle; mature grain is held within the lemma and palea *Dinebra neesii*
 Inflorescence a panicle with secondary branches, open or contracted; mature grain is readily shed from thin lemma and palea 2



Dinebra neesii spikelets



Sporobolus spikelets

2. Lowest node of inflorescence with whorled branches 3
 Lowest node of inflorescence with 1 or 2 branches 6

Lowest node of inflorescence
with whorled branches



Lowest node of inflorescence
with 1 or 2 branches



3. Inflorescence branches only whorled at lowest nodes; the inflorescence is 8-15 cm wide *Sporobolus caroli*
 Most of inflorescence branches whorled; the inflorescence is 2-6 cm wide 4
4. Spikelets uniformly distributed in inflorescence *Sporobolus australasicus*
 Spikelets situated on apical zone of inflorescence branches 5
5. Inflorescence branches not all whorled; leaves 4-5 mm wide; common weed *Sporobolus coromandelianus*
 Inflorescence branches all (except uppermost) whorled; leaves to 4 mm wide; not very common *Sporobolus lenticularis*
6. Inflorescences spiciform to semi-spiciform 7
 Inflorescences spreading 8




Inflorescences spiciform
to semi-spiciform thorough




Inflorescences spreading

7. Inflorescence less than 12 cm long; rhizomatous and stoloniferous coastal sand dune and saline swamp grass..... *Sporobolus virginicus*
 Inflorescence more than 12 cm; tufted grass.....*Sporobolus fertilis*
8. Upper glume $\pm 1/3$ spikelet length (microscopic examination in the laboratory is usually required to confirm identifications); grows 30 to 70 cm tall; common weed in many lawns
 *Sporobolus jacquemontii*
 Upper glume half the spikelet length or more; grasses 40-160 cm tall9



Upper glume $\pm 1/3$
spikelet length



Upper glume half the spikelet length or more



9. Grass 60-150 cm tall; leaf blades 2-4 mm wide*Sporobolus natalensis*
 Grass 10-100 cm tall; leaf blade 5-30 mm wide*Sporobolus sessilis*

***Sporobolus australasicus* - Australian Dropseed**

Derivation

australasicus - from Australia.

Habit


An ephemeral, slender and erect grass, the culms are 20-35 cm tall.

Inflorescence

The inflorescence is an open panicle, 5-13 cm long and 2-6 cm wide. The primary branches are mostly whorled, and the spikelets are distributed equally in the panicle. The seed is globose.

Habitat

This species occurs on a wide range of soils, usually a coloniser in degraded areas



Superficially close to *Sporobolus coromandelianus*, the spikelets arranged differently.

The panicle at the left is *Sporobolus australasicus*. The spikelets are distributed equally in the panicle

The panicle at the right is *Sporobolus coromandelianus*. The spikelets are arranged towards the apices of panicle branches.

***Sporobolus caroli* - Fairy Grass**

Derivation

caroli - in honour of Jean Martin Francois Carolus (1808-1863), a Belgian botanist.

Habit

An ephemeral and erect grass, the culms are 20-75 cm tall.

Inflorescence

The inflorescence is an open panicle, 8-15 cm long and 8-15 cm wide. The primary branches are only whorled at lowest node.

Habitat

This species often occurs on roadsides but also grows in open grasslands and moist situations and over a large range of soil types.



***Sporobolus coromandelianus* - Small Dropseed**

Derivation

coromandelianus - from the Coromandel, that is south-eastern India.

Habit


An ephemeral and geniculate (bent abruptly like a knee joint) grass, the culms are 10-30 cm tall.

Inflorescence

The inflorescence is an open panicle, 5 cm long and 2 cm wide. The primary branches are mostly whorled, and the spikelets are arranged towards the apices of panicle branches.

Habitat

The non-native species is considered as a weed of gardens and roadsides.



Superficially close to *Sporobolus australasicus*, the spikelets arranged differently.

The panicle at the left is *Sporobolus australasicus*. The spikelets are distributed equally in the panicle

The panicle at the right is *Sporobolus coromandelianus*. The spikelets are arranged towards the apices of panicle branches.

***Sporobolus fertilis* - Giant Parramatta Grass**

Derivation

fertilis - Latin for fruitful, this species produce abundant grain.

Habit


A perennial and erect grass, the culms are 80-160 cm tall.

Inflorescence

The inflorescence is a spiciform to semi-spiciform panicle, 25-50 cm long and 0.5-2 cm wide. The upper glume is at least half the spikelet length.

Habitat

This non-native species is a serious weed of pastures and roadsides.



Spikelet



***Sporobolus jacquemontii* - Rat's Tail Grass**

Derivation

jacquemontii - in honour of Victor Jacquemont (1801-1832), a French naturalist and traveller.

Habit

A perennial and erect grass, the culms are 30-75 cm tall.

Inflorescence

The inflorescence is open to contracted and loosely spiciform panicle, 8-25 cm long and 0.5-3 cm wide. The upper glume is less than half the spikelet (microscopic examination in the laboratory is usually required to confirm identifications).

Habitat

This non-native species is a weed in disturbed areas, it is common in lawns.



Spikelet



Sporobolus lenticularis

Derivation

lenticularis - from the Latin *lens* (lentil), *-ulus* (diminutive) and *-aris* (pertaining to), the grain is lenticular.

Habit


An ephemeral and erect grass, the culms are 10-60 cm tall.

Inflorescence

The inflorescence is an open panicle, 3-17 cm long and 2-2.5 cm wide. The primary branches are mostly whorled, and the spikelets are arranged towards the apices of panicle branches. The grain is lenticular.

Habitat

The species is found mainly on sand dune coastal woodland, but also in the highlands on red-brown earths.



Sporobolus natalensis - Giant Rat's Tail Grass

Derivation

natalensis- from Natal, South Africa.

Habit

A perennial and erect grass, the culms are 60-150 cm tall. The leaves are 2-4 mm wide.

Inflorescence

The inflorescence is open to contracted and loosely spiciform panicle, 20-30 cm long and 2-4 cm wide. The upper glume at least half the spikelet length.

Habitat

This non-native species is a weed in disturbed areas.



Spikelet



Sporobolus sessilis - Tussocky Sporobolus

Derivation

sessilis - Latin for sessile, the spikelets are without a stalk.

Habit

A perennial and erect grass, the culms are 40-100 cm tall. The leaves are 5-30 mm wide.

Inflorescence

The inflorescence is open to contracted and loosely spiciform panicle, 14-30 cm long and 0.5-4 cm wide. The upper glume at least half the spikelet length.

Habitat

This species occurs in woodland and native grassland.



***Sporobolus virginicus* - Salt Couch**

Derivation

virginicus - from Virginia, USA.

Habit

A perennial, erect or decumbent, rhizomatous and stoloniferous grass, the culms are to 50 cm tall.

Inflorescence

The inflorescence is a contracted spiciform panicle, 3-15 cm long and 0.5-1.5 cm wide.

Habitat

This species is mainly restricted to maritime habitats with some records also from saline localities areas.



Stenotaphrum

From Greek *stenos* (narrow) and *taphros* (trench), alluding to the cavities in the rachis.

Creeping or prostrate perennials or tufted erect annuals. The inflorescence is a spike or raceme with spikelets embedded in hollows of a broad, flattened axis.

Subfamily: Panicoideae; Tribe: Paniceae

World = 9, Australia = 2

Townsville species

Stenotaphrum secundatum Buffalo Grass

Similar to the common lawn grass Buffalo Grass (*Axonopus compressus*).

Key to the species of *Stenotaphrum* and *Axonopus*

Inflorescence is paired or digitate racemes..... *Axonopus compressus*
Inflorescence is a spike..... *Stenotaphrum secundatum*



Axonopus compressus



Stenotaphrum secundatum



Axonopus compressus



Stenotaphrum secundatum

***Stenotaphrum secundatum* - Buffalo Grass**

Derivation

secundatum - is from the Latin *secundus* (bent to one side) and *-atus* (possessing) and refers to the curved fleshy axis of the inflorescence.

Habit


A mat-forming, prostrate, perennial grass, with smooth, hairless leaves and culms, the culms are 10-40 cm tall.

Inflorescence

The inflorescence is a false, one-sided spike; it is 2-10 cm long, with the spikelets embedded in the axis which is swollen and flat on one side.

Habitat

This species is becoming popular as a lawn grass with many varieties. In Townsville it is only found in cultivation.



Spikelet



Part of inflorescence



Back of inflorescence



Themeda

From the Arabic *thaemed* (transliterated by the author as a depression filled with water). The meaning of the choice of name is not given by the author.


Tufted perennials or annuals. The inflorescence is a panicle of condensed racemes each subtended by spathe (leaf-like bract). The spikelets are in groups of seven which are difficult to see in the field. There are four involucre spikelets at the base, two pedicelled spikelets and a sessile spikelet which is the only one to produce a seed and is the only spikelet that is awned.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 38, Australia = 5

Townsville species

Themeda arguens

* *Themeda quadrivalvis* Grader Grass
Themeda triandra Kangaroo Grass



Members of this genus could be confused with *Cymbopogon* species.



Involucral spikelets
(4 spikelets)



Sessile and pedicelled
spikelets (3 spikelets)



Sessile spikelet
(seed)



Complete spikelets
(7 spikelets)

Key to the species of Themeda species

1. Fertile spikelet with an awn 5-9 cm long and a pungent callus 3-4 mm long *Themeda arguens*
Fertile spikelet with an awn less than 7 cm long and a callus 1-3 mm long 2
2. Perennial; involucre spikelets 6-14 mm long; sessile spikelets 6-11 mm long *Themeda triandra*
Annual; involucre spikelets 4-7 mm long; sessile spikelets 4-6 mm long *Themeda quadrivalvis*



Spikelets, the glumes of the involucre spikelets
are without tubercle-based hairs
Themeda triandra



Spikelets, the glumes of involucre spikelets
with tubercle-based hairs
Themeda quadrivalvis

Themeda arguens

Derivation

arguens - from the Latin *arguo* (content).

Habit

An annual tufted grass, the culms are 20- 120 cm tall.

Inflorescence

The spatheole is 2.5-4 cm long. The involucre spikelets are 6-10 mm long; sessile spikelet is 8-11 mm long; the awn is 50-90 mm long

Habitat

This species grows in disturbed places.



Spikelets, (sessile spikelet on left and the involucre and pedicelled spikelets on the right)



***Themeda quadrivalvis* - Grader Grass**

Derivation

quadrivalvis - from the Latin *quator* (four) and *valva* (leaf of a folding door); the four sessile male spikelets form an involucre below the hermaphrodite spikelet.

Habit

An annual, erect grass, the culms are 30-200 cm tall.

Inflorescence

The spatheole is 1.7-6 cm long. The involucral spikelets are 4-7 mm long; sessile spikelet is 4-6 mm long; the awn is 10-45 mm long.

Habitat

This non-native species is found along roadsides and disturbed areas.

Grader Grass (*Themeda quadrivalvis*) could be confused with Giant Spear Grass (*Heteropogon triticeus*) and *Hyparrhenia rufa*, especially before flowering. These three species are tall grasses, and their increase in height before flowering is caused by a section of pale yellow culm being pushed upwards until it protrudes well above the green leaf sheath that previously enclosed it. The pale sections of the internodes alternate with the green sheaths giving these grass species a conspicuously banded appearance.



Spikelets



***Themeda triandra* - Kangaroo Grass**

Derivation

triandra - from the Greek *treis* (three) and *aner* (man), the sessile spikelet surrounded by three (?) male spikelets.

Habit

A perennial, tufted grass, the culms are 30-130 cm tall. The local grass is not tall, the ones from southern Australia are much taller.

Inflorescence

The spatheole is 1.5-3.5 cm long. The involucral spikelets are 6-14 mm long; sessile spikelet is 6-11 mm long; the awn is 25-70 mm long.

Habitat

This species is widespread, but it recently has become scarce as a result of overgrazing.



Spikelets



Thuarea

Thuarea is named after Aubert du Petit-Thouars (1756-1831), the French botanist and ship's officer.

Creeping, mat-forming perennials which grow on sandy seashores. The inflorescence is a spatheate (with spathe) raceme consisting of 1-2 bisexual spikelets in the lower part and several male spikelets in the upper part.

Subfamily: Panicoideae; Tribe: Paniceae

World = 2, Australia = 1

Townsville species

Thuarea involuta Tropical Beachgrass, Bird Beak Grass



***Thuarea involuta* - Tropical Beachgrass, Bird Beak Grass**

Derivation

involuta from the Latin *involvo* (inroll), referring to the inrolled leaf blades.

Habit


A prostrate, perennial grass, the culms are 12 cm tall. The leaves are velvety-soft.

Inflorescence

The inflorescence is a one-sided raceme with 1 to 2 bisexual flowers at the base and 4 to 6 male flowers at the apex. The short flowering shoots bend down as the seed ripens, and the capsules (formed by the spathe in a water-tight fold) either become buried in the sand or float away in the sea.

Habitat

This species grows on sand dunes, particularly foredunes. It is an efficient sand binder and forms deep roots. It has been planted at Saunders Beach for dune stabilisation.



Ripening seed



Triodia

Porcupine Grasses or Spinifex

Triodia from the Greek *treis* (three) and *odous* (tooth), referring to the 3-toothed or 3-lobed lemmas.

Tufted perennials, mostly hummock-forming, culms wiry. Leaf blades narrow, hard, woody needle-like. The inflorescence is a panicle or single raceme or spike. The spikelets are usually solitary with 4-15 florets. Some species are awned, and some are awnless.

The spinifex grasses of inland Australia belong to the genus *Triodia*. Species from the genus *Spinifex* grow on the sand dunes in coastal areas.


Subfamily: Chloridoideae; Tribe: Triodieae

World = 126, Australia = 96

Townsville species

Triodia stenostachya

Porcupine Grass, Spinifex



***Triodia stenostachya* - Porcupine Grass, Spinifex**

Derivation

stenostachya - from the Greek *stenos* (narrow) and *stachys* (ear of corn), referring to the inflorescence which is a narrow spike or spike-like panicle.

Habit

A tufted, stoloniferous perennial forming dense hummocks, the culms are 30-100 cm tall. The culms and leaves are highly resinous, not aromatic.

Inflorescence

The inflorescence is a contracted panicle, 40 cm long and 3.5-4 cm wide. The spikelets are awnless or shortly awned.

Habitat

Grows on rocky hillslopes and ridges in shallow soils. This species is only found on the top of Castle Hill and on Magnetic Island.



Tripogonella

From Latin *treis* (three) referring to hairs at the base of the three lemma nerves, *pogon* (a beard), and *-ella* (diminutive suffix).

Annual or perennial tufted grass. Inflorescence a single, slender spike, a single raceme or spike. The spikelets are laterally compressed, broadside to the rachis. The genus is quite closely to *Leptochloa* and *Diplachne*.

Subfamily: Chloridoideae; Tribe: Cynodonteae

World = 3, Australia = 1

Townsville species

Tripogonella loliiformis Five Minute Grass, Eight-day Grass



Tripogonella loliiformis - Five Minute Grass

Derivation

loliiformis - resembling *Lolium* in some respect.

Habit


An annual or perennial, tufted grass, culms erect, 3.5-15 cm tall, with dense fibrous butts. This species is termed a 'resurrection plant'. The leaves are mostly basal, 5-7.5 cm long. This grass is hardly noticed because it is very small.

Inflorescence

The inflorescence is an erect raceme, usually 2-9.5 cm long. The spikelets are laterally compressed, 4-12 mm long, 2 mm wide, breaking up at maturity.

Habitat

In a variety of habitats: rocky slopes, plateaux and outcrops of granite and sandstone in skeletal reddish soils with spinifex; on plains in red sand or sandy to clayey loams in open *Acacia* woodlands; depressions and creek lines; floodplains in red to brown clayey soils in open Eucalypt woodland; rocky slopes and outcrops of granite in pockets of sandy loam in eucalypt forest and woodland with a grassy understorey



Morphologically this species is very variable. Throughout the majority of its range, it is characterised by a short dense inflorescence with overlapping spikelets. In the eastern country specimens grade to having a longer, less dense inflorescence. The specimen I collected in Townsville had a short spikelet with about five florets.



Urochloa


Arm Grasses or Signal Grasses

From Greek *oura* (tail) and *chloe* (grass), alluding to the muricate (with short sharp point) lemma of the upper floret. *Brachiaria* (excluding *Brachiaria eruciformis*) species are now included in *Urochloa*, these species do not have a muricate fertile lemma.

Annuals or perennials, rhizomatous or stoloniferous, tufted or decumbent. The inflorescence is a once-branched panicle. The spikelets are solitary, in pairs or clustered. The fertile lemma rugose (wrinkled).

Members of this genus could be confused with *Echinochloa*, *Eriochloa*, *Paspalidium* and *Paspalum* species.

Subfamily: Panicoideae; Tribe: Paniceae
World = 104, Australia = 30





Townsville species

- * *Urochloa distachya*
- Urochloa holosericea* Silkytop Armgrass
- * *Urochloa mosambicensis* Sabi Grass
- * *Urochloa mutica* Para Grass
- Urochloa oligotricha*
- * *Urochloa panicoides* Liverseed Grass
- Urochloa piligera* Hairy Arm Grass
- Urochloa polyphylla*
- Urochloa pubigera* Arm Grass
- * *Urochloa subquadripara* Green Summer Grass

Key to the species of *Urochloa* and related genera

1. Inflorescence terminating in a bristle (inspect carefully since it looks similar to an awn) ... *Paspalidium*
Inflorescence terminating in a spikelet2
2. Spikelet with a bead-like swelling at base (lower glume)*Eriochloa*
Spikelet without a bead-like swelling at base3



3. Inflorescence terminating in a bristle *Echinochloa*
Bead-like swelling at the base of the spikelet4
3. Ligule absent; glumes very unequal; palea tip reflexed*Echinochloa*
Ligule present.....4
4. Lower glume present *Urochloa*
Lower glume absent or very much reduced *Paspalum*

Key to the species of *Urochloa* species


1. Upper lemma obtusely rounded at apex and with distinct mucro (a short sharp point at the end); lower glume abaxial to spikelet bearing axis.....2
- Upper lemma obtuse to acute, usually mucicous (without a point or pointed process); lower glume adaxial to spikelet bearing axis.....4




Upper lemma with a distinct mucro



Upper lemma mucicous



Abaxial of the side of lower glume, facing away the axis



Adaxial of the side of lower glume facing the axis

2. Lower glume c. 1/3 spikelet length*Urochloa panicoides*
- Lower glume at least 1/2 spikelet length3
3. Spikelets tidy; lower glume 3-nerved, rarely sub-5-nerved *Urochloa mosambicensis*
- Spikelets untidy; lower glume 5-7-nerved..... *Urochloa oligotricha*



Spikelets tidy



Spikelets untidy

4. Inflorescence ± open or with spikelets in several untidy rows; lower racemes sometimes branching5
- Inflorescence strictly regular and racemose with spikelets in two regular rows; lower racemes never branching.....6
5. Plant with robust ascending culms 1-4 m tall, with stolons *Urochloa mutica*
- Plant with slender culms 1 m tall, without stolons*Urochloa pubigera*
6. Spikelets up to 3 mm long *Urochloa distachya*
- Spikelets more than 3 mm long8
8. Spikelets hairy9
- Spikelets glabrous 10
9. Spikelets evenly hairy or if with longer apical hairs these are not arranged in a fringe *Urochloa piligera*
- Spikelets with a subapical fringe of silky hairs *Urochloa holosericea*
10. Spikelets reddish; leaf margins serrate*Urochloa polyphylla*
- Spikelets green to yellow *Urochloa subquadripara*

Urochloa distachya

Derivation

distachya - from the Greek *dis* (twice) and *stachys* (an ear of corn), the inflorescences with two branches or with regularly bifurcating branches.

Habit

An annual or perennial, mat-forming grass, the culms are prostrate, 5-20 cm tall.

Inflorescence

The panicle is 1-3.5 cm long; it has 2-3 racemes. The spikelet is glabrous. The lower glume is 33-50% of the length of spikelet.

Habitat

This species is well adapted to sandy loam soil.

Urochloa distachya and *Urochloa subquadriflora* are very much alike, this species has a shorter spikelet, 2.4-3 mm long.



Spikelet



***Urochloa holosericea* - Silkytop Armgrass**

Derivation

holosericea - from the Greek words *holos* (entire) and *sericea* (silky), the plant or spikelets totally invested in dense hairs

Habit

An annual grass, the culms are 15-70 cm tall.

Inflorescence

The panicle is 0.8-3.5 cm long; it has 4-6 racemes. The spikelet is hairy; it has a subapical fringe of silky hairs. The lower glume is 60-70% of the length of spikelet.

Habitat

This species is distributed by woodlands and coastal grasslands.



Spikelet



***Urochloa mosambicensis* - Sabi Grass**

Derivation

mosambicensis - from Mozambique.

Habit

A perennial grass, the culms are 20-150 cm tall.

Inflorescence

The panicle is 2-10 cm long; it has 3-8 racemes. The lower glume is 50-75% of the length of spikelet. The upper lemma has a distinct mucro (a short sharp point at the end).

Habitat

This non-native species is common on roadsides, parks, and disturbed sites.



Spikelet



Upper floret



***Urochloa mutica* - Para Grass**

Derivation

mutica - Latin word for blunt, lacking awns or lemmas and/or glumes truncate.

Habit

A robust, perennial grass, stolons present, the culms are sprawling, 1-4 m long, rooting from lower nodes.

Inflorescence

The panicle is 2-13 cm long; it has 5-20 racemes. The spikelet is glabrous. The lower glume is 25-33% of the length of spikelet.

Habitat

This non-native species grows in wet and submerged habitats.



Spikelet



Urochloa oligotricha

Derivation

oligotricha - from the Greek *oligos* (few) and *thrix* (hair), the plant in whole or part sparsely hairy.

Habit


A perennial grass, the culms are 30-150 cm tall.

Inflorescence

The panicle is 2-12 cm long; it has 6-20 racemes. The lower glume is 66-75% of the length of spikelet. The upper lemma has a distinct mucro (a short sharp point at the end).

Habitat

This non-native species is uncommon. It is found around James Cook University grounds and nearby.



Spikelet



***Urochloa panicoides* - Liverseed Grass**

Derivation

panicoides - resembling *Panicum*, usually with respect to the inflorescence or spikelet.

Habit

An annual grass, the culms are 10-80 cm tall.

Inflorescence

The panicle is 1-7 cm long; it has 2-7 racemes. The lower glume is 30-40% of the length of spikelet. The upper lemma has a distinct mucro (a short sharp point at the end).

Habitat

This non-native species is found on roadsides, parks, and disturbed sites.



Spikelet



Urochloa piligera - Hairy Arm Grass

Derivation


piligera - from the Latin *pilus* (a hair) and *gero* (carry), it is a hairy spikelet.

Habit

An annual grass, the culms are 15-60 cm tall.

Inflorescence

The panicle is 2-6 cm long; it has 3-4 racemes. The spikelet is hairy. The lower glume is 50-70% of the length of spikelet.



Spikelets

Habitat

This species occurs in forests, rainforest, woodlands, shrublands and grasslands.



Urochloa polyphylla

Derivation

polyphylla - from the Greek words *polys* (many) and *phyllon* (leaf), the culms many-leaved in comparison with related species.

Habit

An annual grass, the culms 15-60 cm.

Inflorescence


The panicle is 1.5-4 cm long; it has 3-7 racemes. The spikelet is glabrous. The lower glume is 50-60% of the length of spikelet.

Habitat

This species occurs in rainforest, woodlands and grasslands.



Spikelet



***Urochloa pubigera* - Arm Grass**

Derivation

pubigera - from the Latin word *pubes* (hair that appears at puberty) and *gera* (to bear), the spikelet is slightly hairy.

Habit

An annual grass, the culms 20-60 cm tall.

Inflorescence

The panicle is 3-9 cm long; it has 4-10 racemes. The spikelet looks glabrous but has small hairs. The lower glume is 50-60% of the length of spikelet.

Habitat

This species occurs in rainforest, woodlands and grasslands.



Spikelet



***Urochloa subquadrifera* - Green Summer Grass**

Derivation

subquadrifera - from the Latin *sub* (almost), *quadri* (four) and *fero* (bear), referring to the inflorescence with mostly four racemes.

Habit

An annual or perennial mat-forming grass, the culms are prostrate, 10-150 cm long.


Inflorescence

The panicle is 2-6 cm long; it has 3-5 racemes. The spikelet is glabrous. The lower glume is 33-50% of the length of spikelet.

Habitat

This non-native species occurs widely as a weed. It grows in rainforest, woodlands and grasslands.

Urochloa subquadrifera and *Urochloa distachya* are very much alike, this species has a longer spikelet, 3-3.7 mm long.



Spikelet



Vacoparis


Sorghum

From Latin *vaco* (empty) and *paris* (companion), referring to the greatly reduced pedicelled spikelets. Recent studies of the Australian *Sorghum* species, have suggested three distinct lineages, and these species have been divided into three genera, *Sorghum*, *Sarga* and *Vacoparis*.

Tufted annuals. The inflorescence is an open or contracted panicle. The spikelets are in pairs (with terminal triplets), one sessile and one pedicelled. The mature sessile spikelets are dark reddish brown almost black. The awn of the sessile spikelet 2.5-5.2 cm long, the pedicelled spikelet is reduced to narrow, linear glumes.

Subfamily: Panicoideae; Tribe: Andropogoneae
World = 2, Australia = 2

Townsville species
Vacoparis laxiflorum



Key to the species of *Sarga*, *Sorghum* and *Vacoparis*

1. Pedicelled spikelet reduced to narrow, linear glumes *Vacoparis laxiflorum*
 Pedicelled spikelet containing empty or male lemmas 2
2. Awn on sessile spikelet 25-80 mm long *Sarga plumosum*
 Awn on sessile spikelet 0-20 mm long *Sorghum*



Vacoparis laxiflorum



Sarga plumosum



Sorghum nitidum
f. aristatum



Sorghum x alnum

Vacoparis laxiflorum

Derivation

laxiflorum - from the Latin *laxus* (loose) and *flos* (flower), referring to an open panicle.

Habit

An annual grass, the culms are 100-200 cm tall. The nodes of the culms are glabrous, pubescent or bearded.

Inflorescence

The inflorescence is an open panicle, 5-20 cm long, which hangs to one side. The pedicelled spikelets are only linear glumes; the sessile spikelets are with white or pale brown hairs 5-6.5 mm, the lemma awn is 2.5-1.5 cm long.

Habitat

This species is usually found on heavy clays and alluvial soils along river flats and flooded areas.




Whiteochloa

Whiteochloa is named after the Cyril Tenison White, Queensland Government botanist 1917-1950.

Annual, or short-lived perennials, tufted to decumbent. The inflorescence is an open or contracted panicle. The spikelets are solitary.

Subfamily: Panicoideae; Tribe: Paniceae
World = 6, Australia = 6


Townsville species
Whiteochloa airoides



This species could be confused with *Arthragrostis*, *Panicum*, and *Paspalidium* species.


Key to the species of *Whiteochloa* and related genera

1. Inflorescence terminating in a bristle..... *Paspalidium*
- Inflorescence terminating in a spikelet2




Inflorescence terminating in a bristle


2. Spikelets dorsally compressed..... *Panicum*
- Spikelets laterally compressed3



Spikelets dorsally compressed



Spikelets laterally compressed



3. Plant 45-110 cm tall; grows on sandy, alluvial soils; inflorescence open or contracted, branches persistent *Whiteochloa airoides*
- Plant 17-60 cm; grows on rocky hillsides; inflorescence open; branches deciduous *Arthragrostis deschampsoides*

Whiteochloa airoides

Derivation

airoides - the inflorescences resemble those of genus *Aira*.

Habit

A tufted, perennial, tussock grass, the culms are 45-110 cm tall, with basal leaf sheaths pubescent.

Inflorescence


The inflorescence is an open or contracted panicle 7-25 cm long, 1-10 cm wide, contracted about primary branches. The spikelet is laterally compressed.

Habitat

This species grow in sandy alluvial soil.



Spikelets



Zoysia

Zoysia Grasses

Named for Baron Karl von Zois, 1756-1800, Austrian botanist.

Mat-forming perennials, rhizomatous. The inflorescence is a single raceme with spikelets arranged on all sides. The spikelets have one fertile floret and one glume, the lower glume is absent or obscure.

Zoysia grasses are native to southeast Asia, Australia and New Zealand and are cultivated for lawns and ornamental plants.

Subfamily: Chloridoideae; Tribe: Cynodonteae
World = 16, Australia = 3


There are three principle species which have been used to develop varieties and hybrids:

Zoysia japonica - Japanese Lawn Grass, Korean Lawn Grass

Zoysia matrella - Manila Grass

*Zoysia pacifica*¹ - Korean Lawn Grass, No Mow Grass

The Australian endemic species *Zoysia macrantha* (Prickly Couch Grass), grows in coastal areas between north of Yeppoon, and Port Lincoln, South Australia.



¹ Recent research has identified the improper application of the name *Z. tenuifolia*. *Z. tenuifolia* is endemic to Mauritius, is not represented in *Zoysia* breeding collections, and has not been used in the development of zoysia grasses. The specimens collected in Pacific Rim countries belong to the species *Z. pacifica*.

Key to Zoysia species used for tuftgrass

1. Blades to 0.5 mm diameter; racemes with 3-12 spikelets; peduncles exerted less than 1 cm beyond flag leaf sheath *Zoysia pacifica*
 Blades greater than 0.5 mm width or diameter, stressed or fully hydrated; racemes with 10-50 spikelets; peduncles exerted at least 1 cm beyond flag leaf sheath 2
2. Pedicels 1.6-3.5 mm; spikelets ovate, 1.0-1.4 mm wide; culm internodes 2-10 mm long *Zoysia japonica*
 Pedicels 0.6-1.6 mm long; spikelets lanceolate 0.6-1.0 mm wide; culm internodes, at least a few, over 14 mm long *Zoysia matrella*

***Zoysia japonica* - Japanese Lawngrass, Korean Lawngrass**

Varieties - Empire (SS-500), Palisades, El Toro, UltimateFlora (BA-189), Ozeboy.



***Zoysia matrella* - Manila Grass**

Varieties - Cavalier, Cavier, Shade Tuff (A-1), Layzee (G-4).



***Zoysia pacifica* - Korean Lawngrass, No Mow Grass**



Townsville grasses, Genus and species names

Status

Naturalised (non-native) grasses (*)

Naturalised species are those that are considered to have successfully established populations outside their native range, by reproducing there without cultivation or other human intervention.

Cultivated grasses (#) are only found in gardens.

Vulnerable grasses (V) are threatened plants.

Status	Botanical Name	Common Name
	<i>Alloteropsis cimicina</i>	Annual Cockatoo Grass
	<i>Alloteropsis semialata</i>	Cockatoo Grass
	<i>Ancistrachne uncinulata</i>	Hooky Grass
*	<i>Andropogon gayanus</i>	Gamba Grass
	<i>Aristida acuta</i>	
	<i>Aristida calycina</i>	Dark Wiregrass, Branched Wiregrass
	<i>Aristida gracilipes</i>	
	<i>Aristida holathera</i>	Erect Kerosine Grass
	<i>Aristida jerichoensis</i> var. <i>subspinulifera</i>	Jericho Wiregrass, Feathertop Threawn
	<i>Aristida latifolia</i>	Feathertop Wiregrass
	<i>Aristida pernicioso</i>	
	<i>Aristida queenslandica</i> var. <i>dissimilis</i>	Queensland Wiregrass
	<i>Aristida queenslandica</i> var. <i>queenslandica</i>	Queensland Wiregrass
	<i>Aristida spuria</i>	
	<i>Aristida superpendens</i>	
	<i>Aristida utilis</i>	
	<i>Aristida warburgii</i>	
	<i>Arthrargrostis deschampsoides</i>	
	<i>Arundinella nepalensis</i>	Reed Grass
	<i>Arundinella setosa</i>	Reed Grass
*	<i>Arundo donax</i>	Giant Reed
*	<i>Axonopus compressus</i>	Buffalo Grass, Broadleaf Carpet Grass
	<i>Bothriochloa bladhii</i> subsp. <i>bladhii</i>	Forest Bluegrass
	<i>Bothriochloa decipiens</i>	Pitted Grass
	<i>Bothriochloa ewartiana</i>	Desert Bluegrass
*	<i>Bothriochloa pertusa</i>	Indian Bluegrass
	<i>Brachyachne convergens</i>	Common Native Couch
	<i>Capillipedium parviflorum</i>	Scented Top
	<i>Capillipedium spicigerum</i>	Scented Top
*	<i>Cenchrus ciliaris</i>	Buffel Grass
*	<i>Cenchrus echinatus</i>	Mossman River Grass, Burr Grass
	<i>Cenchrus elymoides</i>	
*	<i>Cenchrus pedicellatus</i> subsp. <i>unispiculus</i>	Annual Mission Grass
*	<i>Cenchrus setaceus</i>	Fountain Grass
*	<i>Cenchrus setiger</i>	Birdwood Grass
	<i>Chionachne cyathopoda</i>	River Grass
*	<i>Chloris gayana</i>	Rhodes Grass
*	<i>Chloris inflata</i>	Purpletop Grass
	<i>Chloris lobata</i>	
	<i>Chloris pectinata</i>	Comb Windmill Grass
	<i>Chloris pumilio</i>	
*	<i>Chloris virgata</i>	Feathertop Rhodes Grass
*	<i>Chrysopogon aciculatus</i>	Mackies Pest
	<i>Chrysopogon fallax</i>	Golden Beard Grass
	<i>Chrysopogon filipes</i>	Australian Vetiver
*	<i>Chrysopogon zizanioides</i>	Vetiver Grass
	<i>Cleistochloa subjuncea</i>	
	<i>Cymbopogon ambiguus</i>	Scented Oilgrass, Lemon Grass

	<i>Cymbopogon bombycinus</i>	Silky Oilgrass
*	<i>Cymbopogon nardus</i>	Citronella grass
	<i>Cymbopogon oblectus</i>	Silkyheads
	<i>Cymbopogon queenslandicus</i>	
	<i>Cymbopogon refractus</i>	Barbed Wire Grass
*	<i>Cynodon dactylon</i>	Couch Grass
*	<i>Cynodon nlemfuensis</i>	Bermuda Grass
*	<i>Cynodon radiatus</i>	Bermuda Grass
*	<i>Dactyloctenium aegyptium</i>	Coastal Button Grass
	<i>Dactyloctenium buehnerianum</i>	
	<i>Dactyloctenium radicans</i>	Native Button Grass
*	<i>Dichanthium annulatum</i>	Sheda Grass
*	<i>Dichanthium aristatum</i>	Angleton Grass
	<i>Dichanthium fecundum</i>	Curly Bluegrass
	<i>Dichanthium sericeum subsp. polystachyum</i>	Queensland Bluegrass
	<i>Dichanthium sericeum subsp. sericeum</i>	Queensland Bluegrass
	<i>Digitaria ammobila</i>	Silky Umbrella Grass, Spider Grass
	<i>Digitaria breviglumis</i>	
	<i>Digitaria bicornis</i>	Hairy Finger Grass
	<i>Digitaria brownii</i>	Cotton Panic Grass
*	<i>Digitaria ciliaris</i>	Summer Grass, Crab Grass
*	<i>Digitaria didactyla</i>	Queensland Bluegrass
	<i>Digitaria diffusa</i>	
*	<i>Digitaria eriantha</i>	Pangola Grass
	<i>Digitaria fumida</i>	
	<i>Digitaria gibbosa</i>	
*	<i>Digitaria insularis</i>	Sourgrass
	<i>Digitaria leucostachya</i>	
	<i>Digitaria longiflora</i>	
	<i>Digitaria minima</i>	
	<i>Digitaria nematostachya</i>	
	<i>Digitaria orbata</i>	
	<i>Digitaria parviflora</i>	Small Flower Finger Grass
	<i>Digitaria ramularis</i>	
*	<i>Digitaria violascens</i>	Purple Crabgrass
	<i>Dinebra decipiens</i>	Slender Cane Grass
	<i>Dinebra neesii</i>	Swamp Grass
	<i>Diplachne fusca</i> var. <i>fusca</i>	Brown Beetle Grass
*	<i>Diplachne fusca</i> var. <i>uninervia</i>	Mexican Sprangletop
*	<i>Echinochloa colona</i>	Awnless Barnyard Grass
*	<i>Echinochloa crus-galli</i>	Barnyard Grass
*	<i>Echinochloa esculenta</i>	Japanese Millet
*	<i>Echinochloa polystachya</i>	Aleman Grass
	<i>Ectrosia agrostoides</i>	
	<i>Ectrosia lasioclada</i>	
	<i>Ectrosia leporina</i>	Hare's Foot Grass
	<i>Ectrosia nervilemma</i>	
*	<i>Eleusine indica</i>	Crowsfoot Grass
	<i>Elionurus citreus</i>	Lemon-scented Grass
	<i>Elytrophorus spicatus</i>	Spike Grass
	<i>Enneapogon lindleyanus</i>	Nineawn Grass
	<i>Enneapogon nigricans</i>	Black heads
	<i>Enneapogon polyphyllus</i>	Leafy Nineawn
	<i>Enneapogon robustissimus</i>	Nineawn Grass
	<i>Enneapogon virens</i>	
	<i>Enteropogon acicularis</i>	Curly Windmill Grass
	<i>Enteropogon ramosus</i>	Curly Windmill Grass
	<i>Eragrostis basedowii</i>	Neat Lovegrass
	<i>Eragrostis brownii</i>	Brown's Lovegrass
*	<i>Eragrostis cilianensis</i>	Stinking Lovegrass
	<i>Eragrostis cumingii</i>	Cuming's Lovegrass

*	<i>Eragrostis curvula</i>	African Lovegrass
	<i>Eragrostis dielsii</i>	Mallee Lovegrass
	<i>Eragrostis elongata</i>	Clustered Lovegrass
	<i>Eragrostis exigua</i>	
	<i>Eragrostis fallax</i>	
	<i>Eragrostis interrupta</i>	
	<i>Eragrostis lacunaria</i>	Purple Lovegrass
	<i>Eragrostis leptostachya</i>	Paddock Lovegrass
*	<i>Eragrostis mexicana</i>	Mexican Lovegrass
*	<i>Eragrostis minor</i>	Small Stink Grass
	<i>Eragrostis parviflora</i>	Weeping Lovegrass
*	<i>Eragrostis pilosa</i>	Soft Lovegrass
	<i>Eragrostis pubescens</i>	
	<i>Eragrostis schultzei</i>	
	<i>Eragrostis sororia</i>	
	<i>Eragrostis spartinoides</i>	
	<i>Eragrostis stenostachya</i>	
*	<i>Eragrostis tenella</i>	Delicate Lovegrass
	<i>Eragrostis tenellula</i>	Delicate Lovegrass
*	<i>Eragrostis tenuifolia</i>	Elastic Grass
	<i>Eragrostis unioides</i>	
	<i>Eremochloa bimaculata</i>	Poverty Grass
	<i>Eriachne ciliata</i>	Slender Wanderrie Grass
	<i>Eriachne mucronata</i>	Mountain Wanderrie Grass
	<i>Eriachne obtusa</i>	Northern Wanderrie Grass
	<i>Eriachne pallescens</i>	Wanderrie Grass
	<i>Eriachne rara</i>	
	<i>Eriachne squarrosa</i>	Wanderrie Grass
	<i>Eriachne triodioides</i>	Wanderrie Grass
	<i>Eriochloa crebra</i>	Tall Cupgrass
	<i>Eriochloa procera</i>	Cup Grass
	<i>Eriochloa pseudoacrotricha</i>	Early Spring Cupgrass
	<i>Eulalia aurea</i>	Silky Browntop
	<i>Hackelochloa granularis</i>	Pit Scale Grass
	<i>Hemarthria uncinata</i>	Mat Grass
	<i>Heteropogon contortus</i>	Black Spear Grass
	<i>Heteropogon triticeus</i>	Giant Spear Grass
	<i>Hymenachne acutigluma</i>	
*	<i>Hymenachne amplexicaulis</i>	Hymenachne
*	<i>Hyparrhenia rufa</i> subsp. <i>rufa</i>	Thatch Grass
	<i>Imperata cylindrica</i>	Blady Grass
	<i>Ischaemum australe</i> var. <i>arundinaceum</i>	Large Bluegrass
	<i>Ischaemum australe</i> var. <i>australe</i>	Large Bluegrass
	<i>Ischaemum australe</i> var. <i>villosum</i>	Large Bluegrass
	<i>Ischaemum rugosum</i> var. <i>rugosum</i>	
	<i>Ischaemum rugosum</i> var. <i>segetum</i>	
	<i>Leersia hexandra</i>	Swamp Rice Grass
	<i>Lepturus repens</i>	
*	<i>Megathyrsus maximus</i> var. <i>coloratus</i>	Purple-topped Guinea Grass
*	<i>Megathyrsus maximus</i> var. <i>maximus</i>	Common Guinea Grass
*	<i>Megathyrsus maximus</i> var. <i>maximus</i> 'Hamil'	Hamil Grass
*	<i>Megathyrsus maximus</i> var. <i>pubiglumis</i>	Guinea Grass
*	<i>Melinis minutiflora</i>	Molasses Grass
*	<i>Melinis repens</i>	Red Natal Grass
	<i>Mnesithea formosa</i>	
	<i>Mnesithea rottboelliioides</i>	Northern Cane Grass
	<i>Ophiuros exaltatus</i>	Cane Grass
	<i>Oplismenus aemulus</i>	Australian Basket Grass
	<i>Oplismenus compositus</i>	Running Mountain Grass
	<i>Oryza australiensis</i>	Australian Wild Rice
	<i>Oryza meridionalis</i>	Australian Wild Rice

	<i>Oxychloris scariosa</i>	Winged Windmill Grass
	<i>Panicum decompositum</i> var. <i>decompositum</i>	Australian Millet
	<i>Panicum decompositum</i> var. <i>tenuius</i>	Australian Millet
	<i>Panicum effusum</i>	Hairy Panic
	<i>Panicum laevinode</i>	Pepper Grass
	<i>Panicum mindanaense</i>	
	<i>Panicum mitchellii</i>	
	<i>Panicum paludosum</i>	Swamp Panic
	<i>Panicum seminudum</i> var. <i>cairnsianum</i>	
	<i>Panicum simile</i>	Two-coloured Panic
	<i>Panicum trichoides</i>	
	<i>Paspalidium criniforme</i>	
	<i>Paspalidium distans</i>	Spreading Panic Grass
	<i>Paspalidium flavidum</i>	
	<i>Paspalidium gracile</i>	Slender Panic
	<i>Paspalidium rarum</i>	Rare Panic
V	<i>Paspalidium udum</i>	
*	<i>Paspalum conjugatum</i>	Sour Grass, Johnson River Grass
*	<i>Paspalum dilatatum</i>	Paspalum
	<i>Paspalum distichum</i>	Water Couch
*	<i>Paspalum notatum</i>	Bahia Grass
	<i>Paspalum scrobiculatum</i>	Ditch Millet
	<i>Paspalum vaginatum</i>	Saltwater Couch
	<i>Perotis rara</i>	Comet Grass
	<i>Phragmites australis</i>	Cane Grass, Bamboo Reed
	<i>Phragmites karka</i>	Tropical Reed
	<i>Pseudopogonatherum contortum</i>	
	<i>Pseudoraphis spinescens</i>	Spiny Mud Grass
	<i>Sacciolepis indica</i>	Indian Cupscale Grass
	<i>Sarga plumosum</i>	Plume Sorghum
	<i>Schizachyrium fragile</i>	Fire Grass, Red Spathe Grass
	<i>Schizachyrium occultum</i>	
	<i>Schizachyrium pseudeulalia</i>	
	<i>Sehima nervosum</i>	White Grass
	<i>Setaria</i> sp. Mount Larcom	
*	<i>Setaria pumila</i> subsp. <i>subtesselata</i>	Pale Pigeon Grass
*	<i>Setaria sphacelata</i>	South African Pigeon Grass
	<i>Setaria surgens</i>	Pigeon Grass
*	<i>Sorghum bicolor</i>	Forage Sorghum
*	<i>Sorghum halepense</i>	Johnson Grass
	<i>Sorghum nitidum</i>	
*	<i>Sorghum x alnum</i>	Columbia Grass
	<i>Spinifex sericeus</i>	Beach Spinifex, Hairy Spinifex
	<i>Sporobolus australasicus</i>	Australian Dropseed
	<i>Sporobolus caroli</i>	Fairy Grass
*	<i>Sporobolus coromandelianus</i>	Small Dropseed
*	<i>Sporobolus fertilis</i>	Giant Parramatta Grass
*	<i>Sporobolus jacquemontii</i>	Rat's Tail Grass
	<i>Sporobolus lenticularis</i>	
*	<i>Sporobolus natalensis</i>	Giant Rat's Tail Grass
	<i>Sporobolus sessilis</i>	Tussocky Sporobolus
	<i>Sporobolus virginicus</i>	Salt Couch
#	<i>Stenotaphrum secundatum</i>	Buffalo Grass
	<i>Themeda arguens</i>	
*	<i>Themeda quadrivalvis</i>	Grader Grass
	<i>Themeda triandra</i>	Kangaroo Grass
	<i>Thuarea involuta</i>	Tropical Beachgrass, Bird Beak Grass
	<i>Triodia stenostachya</i>	Porcupine Grass, Spinifex
	<i>Tripogonella loliiformis</i>	Five Minute Grass, Eight-day Grass
*	<i>Urochloa distachya</i>	
	<i>Urochloa holosericea</i> subsp. <i>holosericea</i>	Silkytop Armgrass

*	<i>Urochloa mosambicensis</i>	Sabi Grass
*	<i>Urochloa mutica</i>	Para Grass
	<i>Urochloa oligotricha</i>	
*	<i>Urochloa panicoides</i>	Liverseed Grass
	<i>Urochloa piligera</i>	Hairy Arm Grass
	<i>Urochloa polyphylla</i>	
	<i>Urochloa pubigera</i>	Arm Grass
*	<i>Urochloa subquadrifera</i>	Green Summer Grass
	<i>Vacoparis laxiflorum</i>	
	<i>Whiteochloa airoides</i>	Creeping Panic
#	<i>Zoysia japonica</i>	Japanese Lawn Grass, Korean Lawn Grass
#	<i>Zoysia matrella</i>	Manila Grass
#	<i>Zoysia pacifica</i>	No Mow Grass, Korean Lawn Grass

Townville grasses, common names

Status	Common Name	Botanical Name
*	African Lovegrass	<i>Eragrostis curvula</i>
*	Aleman Grass	<i>Echinochloa polystachya</i>
*	Angleton Grass	<i>Dichanthium aristatum</i>
	Annual Cockatoo Grass	<i>Alloteropsis cimicina</i>
*	Annual Mission Grass	<i>Cenchrus pedicellatus subsp. unispiculus</i>
	Arm Grass	<i>Urochloa pubigera</i>
	Australian Dropseed	<i>Sporobolus australasicus</i>
	Australian Millet	<i>Panicum decompositum var. decompositum</i>
	Australian Millet	<i>Panicum decompositum var. tenuius</i>
	Australian Vetiver	<i>Chrysopogon filipes</i>
	Australian Wild Rice	<i>Oryza australiensis</i>
	Australian Wild Rice	<i>Oryza meridionalis</i>
*	Awnless Barnyard Grass	<i>Echinochloa colona</i>
*	Bahia Grass	<i>Paspalum notatum</i>
	Bamboo Reed	<i>Phragmites australis</i>
	Barbed Wire Grass	<i>Cymbopogon refractus</i>
*	Barnyard Grass	<i>Echinochloa crus-galli</i>
	Beach Spinifex	<i>Spinifex sericeus</i>
*	Bermuda Grass	<i>Cynodon nlemfuensis</i>
*	Bermuda Grass	<i>Cynodon radiatus</i>
	Bird Beak Grass	<i>Thuarea involuta</i>
*	Birdwood Grass	<i>Cenchrus setiger</i>
	Black Spear Grass	<i>Heteropogon contortus</i>
	Blady Grass	<i>Imperata cylindrica</i>
	Branched Wire Grass	<i>Aristida calycina</i>
*	Broadleaf Carpet Grass	<i>Axonopus compressus</i>
	Brown Beetle Grass	<i>Diplachne fusca var. fusca</i>
	Brown's Lovegrass	<i>Eragrostis brownii</i>
*	Buffalo Grass	<i>Axonopus compressus</i>
#	Buffalo Grass	<i>Stenotaphrum secundatum</i>
*	Buffel Grass	<i>Cenchrus ciliaris</i>
*	Burr Grass	<i>Cenchrus echinatus</i>
	Cane Grass	<i>Phragmites australis</i>
	Cane Grass	<i>Ophiuros exaltatus</i>
	Clustered Lovegrass	<i>Eragrostis elongata</i>
*	Coastal Button Grass	<i>Dactyloctenium aegyptium</i>
	Cockatoo Grass	<i>Alloteropsis semialata</i>
*	Columbia Grass	<i>Sorghum x alnum</i>
	Comb Windmill Grass	<i>Chloris pectinata</i>
	Comet Grass	<i>Perotis rara</i>

	Common Native Couch	<i>Brachyachne convergens</i>
*	Common Guinea Grass	<i>Megathyrsus maximus</i> var. <i>maximus</i>
	Cotton Panic Grass	<i>Digitaria brownii</i>
*	Couch Grass	<i>Cynodon dactylon</i>
*	Crab Grass	<i>Digitaria ciliaris</i>
	Creeping Panic	<i>Whiteochloa airoides</i>
*	Crowsfoot Grass	<i>Eleusine indica</i>
	Cuming's Lovegrass	<i>Eragrostis cumingii</i>
	Cup Grass	<i>Eriochloa procera</i>
	Curly Bluegrass	<i>Dichanthium fecundum</i>
	Curly Windmill Grass	<i>Enteropogon acicularis</i>
	Curly Windmill Grass	<i>Enteropogon ramosus</i>
*	Citronella grass	<i>Cymbopogon nardus</i>
	Dark Wiregrass	<i>Aristida calycina</i>
*	Delicate Lovegrass	<i>Eragrostis tenella</i>
	Delicate Lovegrass	<i>Eragrostis tenellula</i>
	Desert Bluegrass	<i>Bothriochloa ewartiana</i>
	Ditch Millet	<i>Paspalum scrobiculatum</i>
	Early Spring Cupgrass	<i>Eriochloa pseudoacrotricha</i>
	Eight-day Grass	<i>Tripogonella loliiformis</i>
*	Elastic Grass	<i>Eragrostis tenuifolia</i>
	Erect Kerosine Grass	<i>Aristida holathera</i>
	Fairy Grass	<i>Sporobolus caroli</i>
*	Feathertop Rhodes Grass	<i>Chloris virgata</i>
	Feathertop Wiregrass	<i>Aristida latifolia</i>
	Feathertop Threeawn	<i>Aristida jerichoensis</i> var. <i>subspinulifera</i>
	Fire Grass	<i>Schizachyrium fragile</i>
	Five Minute Grass	<i>Tripogonella loliiformis</i>
*	Forage Sorghum	<i>Sorghum bicolor</i>
	Forest Bluegrass	<i>Bothriochloa bladhii</i> subsp. <i>bladhii</i>
*	Fountain Grass	<i>Cenchrus setaceus</i>
*	Gamba Grass	<i>Andropogon gayanus</i>
*	Giant Parramatta Grass	<i>Sporobolus fertilis</i>
*	Giant Rat's Tail Grass	<i>Sporobolus natalensis</i>
*	Giant Reed	<i>Arundo donax</i>
	Giant Spear Grass	<i>Heteropogon triticeus</i>
	Golden Beard Grass	<i>Chrysopogon fallax</i>
*	Grader Grass	<i>Themeda quadrivalvis</i>
*	Green Panic	<i>Megathyrsus maximus</i> var. <i>pubiglumis</i>
*	Green Summer Grass	<i>Urochloa subquadripara</i>
*	Guinea Grass	<i>Megathyrsus maximus</i> var. <i>maximus</i>
	Hairy Arm Grass	<i>Urochloa piligera</i>
	Hairy Finger Grass	<i>Digitaria bicornis</i>
	Hairy Panic	<i>Panicum effusum</i>
	Hairy Spinifex	<i>Spinifex sericeus</i>
*	Hamil Grass	<i>Megathyrsus maximus</i> var. <i>maximus</i> 'Hamil'
	Hare's Foot Grass	<i>Ectrosia leporina</i>
	Hooky Grass	<i>Ancistrachne uncinulata</i>
*	Hymenachne	<i>Hymenachne amplexicaulis</i>
*	Indian Bluegrass	<i>Bothriochloa pertusa</i>
	Indian Cupscale Grass	<i>Sacciolepis indica</i>
#	Japanese Lawn Grass	<i>Zoysia japonica</i>
*	Japanese Millet	<i>Echinochloa esculenta</i>
	Jericho Wiregrass	<i>Aristida jerichoensis</i> var. <i>subspinulifera</i>
*	Johnson Grass	<i>Sorghum halepense</i>
*	Johnson River Grass	<i>Paspalum conjugatum</i>
	Kangaroo Grass	<i>Themeda triandra</i>
#	Korean Lawn Grass	<i>Zoysia japonica</i>
#	Korean Lawn Grass	<i>Zoysia pacifica</i>
	Large Bluegrass	<i>Ischaemum australe</i> var. <i>arundinaceum</i>
	Large Bluegrass	<i>Ischaemum australe</i> var. <i>australe</i>

	Large Bluegrass	<i>Ischaemum australe</i> var. <i>villosum</i>
	Leafy Nineawn	<i>Enneapogon polyphyllus</i>
	Lemon Grass	<i>Cymbopogon ambiguus</i>
	Lemon-scented Grass	<i>Elionurus citreus</i>
*	Liverseed Grass	<i>Urochloa panicoides</i>
*	Mackies Pest	<i>Chrysopogon aciculatus</i>
	Mallee Lovegrass	<i>Eragrostis dielsii</i>
#	Manila Grass	<i>Zoysia matrella</i>
	Mat Grass	<i>Hemarthria uncinata</i>
*	Mexican Lovegrass	<i>Eragrostis mexicana</i>
	Millet Panic	<i>Panicum mindanaense</i>
*	Molasses Grass	<i>Melinis minutiflora</i>
*	Mossman River Grass	<i>Cenchrus echinatus</i>
	Mountain Wanderrie Grass	<i>Eriachne mucronata</i>
	Native Button Grass	<i>Dactyloctenium radulans</i>
	Neat Lovegrass	<i>Eragrostis basedowii</i>
	Niggerheads	<i>Enneapogon nigricans</i>
#	No Mow Grass	<i>Zoysia pacifica</i>
	Northern Cane Grass	<i>Mnesithea rottboellioides</i>
	Northern Wanderrie Grass	<i>Eriachne obtusa</i>
	Paddock Lovegrass	<i>Eragrostis leptostachya</i>
*	Pale Pigeon Grass	<i>Setaria pumila</i> subsp. <i>subtesselata</i>
*	Pangola Grass	<i>Digitaria eriantha</i>
*	Para Grass	<i>Urochloa mutica</i>
	Pepper Grass	<i>Panicum laevinode</i>
	Pigeon Grass	<i>Setaria surgens</i>
	Pit Scale Grass	<i>Hackelochloa granularis</i>
	Pitted Grass	<i>Bothriochloa decipiens</i>
	Plume Sorghum	<i>Sarga plumosum</i>
	Poverty Grass	<i>Eremochloa bimaculata</i>
*	Purple Crabgrass	<i>Digitaria violascens</i>
	Purple Lovegrass	<i>Eragrostis lacunaria</i>
*	Purple-topped Guinea Grass	<i>Megathyrsus maximus</i> var. <i>coloratus</i>
*	Purpletop Grass	<i>Chloris inflata</i>
	Queensland Bluegrass	<i>Dichanthium sericeum</i> subsp. <i>polystachyum</i>
	Queensland Bluegrass	<i>Dichanthium sericeum</i> subsp. <i>sericeum</i>
*	Queensland Bluegrass	<i>Digitaria didactyla</i>
	Queensland Wiregrass	<i>Aristida queenslandica</i> var. <i>dissimilis</i>
	Queensland Wiregrass	<i>Aristida queenslandica</i> var. <i>queenslandica</i>
	Porcupine Grass	<i>Triodia stenostachya</i>
	Rare Panic	<i>Paspalidium rarum</i>
*	Rat's Tail Grass	<i>Sporobolus jacquemontii</i>
*	Red Natal Grass	<i>Melinis repens</i>
	Red Spathe Grass	<i>Schizachyrium fragile</i>
	Reed Grass	<i>Arundinella nepalensis</i>
	Reed Grass	<i>Arundinella setosa</i>
*	Rhodes Grass	<i>Chloris gayana</i>
	River Grass	<i>Chionachne cyathopoda</i>
*	Sabi Grass	<i>Urochloa mosambicensis</i>
	Salt Couch	<i>Sporobolus virginicus</i>
	Saltwater Couch	<i>Paspalum vaginatum</i>
	Scented Oil Grass	<i>Cymbopogon ambiguus</i>
	Scented Top	<i>Capillipedium parviflorum</i>
	Scented Top	<i>Capillipedium spicigerum</i>
*	Sheda Grass	<i>Dichanthium annulatum</i>
	Silky Browntop	<i>Eulalia aurea</i>
	Silky Oil Grass	<i>Cymbopogon bombycinus</i>
	Silky Umbrella Grass	<i>Digitaria ammophila</i>
	Silkyheads	<i>Cymbopogon obtectus</i>
	Silkytop Armgrass	<i>Urochloa holosericea</i> subsp. <i>holosericea</i>
	Slender Cane grass	<i>Dinebra decipiens</i>

	Slender Panic	<i>Paspalidium gracile</i>
	Slender Wanderrie Grass	<i>Eriachne ciliata</i>
*	Small Dropseed	<i>Sporobolus coromandelianus</i>
*	Small Stink grass	<i>Eragrostis minor</i>
	Small Flower Finger Grass	<i>Digitaria parviflora</i>
*	Soft Lovegrass	<i>Eragrostis pilosa</i>
*	Sourgrass	<i>Digitaria insularis</i>
*	Sour Grass	<i>Paspalum conjugatum</i>
*	South African Pigeon Grass	<i>Setaria sphacelata</i>
	Spider Grass	<i>Digitaria ammophila</i>
	Spike Grass	<i>Elytrophorus spicatus</i>
	Spinifex	<i>Triodia stenostachya</i>
	Spiny Mud Grass	<i>Pseudoraphis spinescens</i>
*	Stinking Lovegrass	<i>Eragrostis cilianensis</i>
*	Summer Grass	<i>Digitaria ciliaris</i>
	Swamp Grass	<i>Dinebra neesii</i>
	Swamp Panic	<i>Panicum paludosum</i>
	Swamp Rice Grass	<i>Leersia hexandra</i>
	Tall Cupgrass	<i>Eriochloa crebra</i>
*	Thatch Grass	<i>Hyparrhenia rufa subsp. rufa</i>
	Tropical Beachgrass	<i>Thuarea involuta</i>
	Tropical Reed	<i>Phragmites karka</i>
	Tussocky Sporobolus	<i>Sporobolus sessilis</i>
	Twirly Windmill Grass	<i>Enteropogon ramosus</i>
	Two-coloured Panic	<i>Panicum simile</i>
*	Vetiver Grass	<i>Chrysopogon zizanioides</i>
	Wanderrie Grass	<i>Eriachne pallescens</i>
	Wanderrie Grass	<i>Eriachne squarrosa</i>
	Wanderrie Grass	<i>Eriachne triodioides</i>
	Water Couch	<i>Paspalum distichum</i>
	Weeping Lovegrass	<i>Eragrostis parviflora</i>
	White Grass	<i>Sehima nervosum</i>
	Winged Windmill Grass	<i>Oxychloris scariosa</i>
#	Zoysia Grass	<i>Zoysia</i>

References

- AVH (Australian Virtual Herbarium) <https://avh.ala.org.au/> (Specimens scanned from various Herbaria).
- Blake ST (1940), Studies on Queensland grasses. *Proceedings of the Royal Society of Queensland* 51: 169-176.
- Blake ST (1974), Revision of the genera *Cymbopogon* and *Schizachyrium* (Gramineae) in Australia. *Contributions From The Queensland Herbarium* 17:1-70.
- Clayton WD, Renvoize SA (1986), *Genera Graminum: Grasses of the World*. Royal Botanical Gardens, Kew.
- Gardner CA (1952), *Flora of Western Australia Vol. 1, Gramineae Part 1*. Government Printer, Perth.
- Gilliland HB, Holttum RE, Bor NL, Burkill HM (1971), *A Revised Flora of Malaya Volume III Grasses of Malaya*. Government Printing Office, Singapore.
- Henrard JT (1933), *A Monograph of the genus Aristida*. Mededeelingen van Rijks Herbarium, Leiden.
- Jacobs SWL, Whalley RDB, Wheeler DJB (2008), *Grasses of New South Wales* (Fourth edition). University of New England, Armidale.
- Jessop J, Dashorst GRM, James FM (2006), *Grasses of South Australia*. Wakefield Press, Kent Town.
- Lazarides M (1970), *The Grasses of Central Australia*. Australian National University Press, Canberra.
- Lazarides M (1978), The genus *Whiteochloa* C.E.Hubbard (Poaceae, Paniceae). *Brunonia* 1:69-93.
- Lazarides M (1992), A Revision of Australian Chlorideae (Gramineae). *Australian Journal of Botany Supplementary Series* 2(5):1-51.
- Lazarides M (1984), New taxa of tropical Australia grasses (Poaceae). *Nuytsia* 5: 273-303.
- Lazarides M (1997), A Revision of *Eragrostis* (Eragrostideae, Eleisininae, Poaceae) in Australia. *Australian Systematic Botany* 10: 77-187.
- Mallett K (Ed.) (2005), *Flora of Australia Volume 44B, Poaceae 3*. ABRIS/CSIRO Publishing, Melbourne.
- Mallett K, Orchard AE (Eds.) (2002), *Flora of Australia Volume 43, Poaceae 1: Introduction and Atlas*. ABRIS/CSIRO Publishing, Melbourne.
- Peterson PM, Romaschenko K, Snow N, Johnson G (2012), A molecular phylogeny and classification of *Leptochloa* (Poaceae: Chloridoideae: Chlorideae) *sensu lato* and related genera. *Annals of Botany* 109: 1317-1329.
- PlantNET New South Wales Flora online.
<https://plantnet.rbgsyd.nsw.gov.au/cgi-bin/NSWfl.pl?page=nswfl&lvl=fm&name=Poaceae>
- Sharp D, Simon BK (2002), *AusGrass: grasses of Australia*. CD-ROM, Version 1.0. Australian Biological Resources Study, Canberra, and Environment Protection Agency, Brisbane.
- Simon BK (1992), A Revision of the Genus *Aristida* (Poaceae) in Australia. *Australian Systematic Botany* 5: 129-226.
- Simon BK, Jacobs SWL (1999), Revision of the genus *Sporobolus* (Poaceae, Chloridoideae) in Australia. *Australian Systematic Botany* 12: 375-448.
- Simon BK (2015) AusGrass2 <https://ausgrass2.myspecies.info/>

- Snow N, Peterson PM (2012), Nomenclatural notes on *Dinebra*, *Diplachne*, *Disakisperma* and *Leptochloa* (Poaceae: Chloridoideae). *Phytoneuron* 71: 1-2
- Snow N, Simon BK (1999), Taxonomic status and Australian distribution of the weedy neotropical grass *Leptochloa fusca* subsp. *uninervia*, with updated key to Australian *Leptochloa* (Poaceae, Chloridoideae). *Austrobaileya* 5: 299-305.
- Soenarko S (1977), The genus *Cymbopogon* Sprengel (Gramineae). *Reinwardtia* 9: 225-375.
- Spangler RE (2003), Taxonomy of *Sarga*, *Sorghum* and *Vacoparis* (Poaceae: Andropogoneae). *Australian Systematic Botany* 16: 279-299.
- Tothill JC, Hacker JB (1983), *The grasses of southern Queensland*. University of Queensland Press, St Lucia.
- WFO The World Flora Online. <https://www.worldfloraonline.org/>
- Wheeler DJB, Rye BL, Koch BL, Wilson AJG (1992), *Flora of the Kimberley Region*. Western Australian Herbarium, Perth.
- Wilson A (Ed.) (2009), *Flora of Australia Volume 44A, Poaceae 2*. ABRS/CSIRO Australia, Melbourne.