

Research



**Cite this article:** Farnan H, Yeeles P, Lach L. 2023 Sublethal doses of insecticide reduce thermal tolerance of a stingless bee and are not avoided in a resource choice test. *R. Soc. Open Sci.* **10**: 230949. <https://doi.org/10.1098/rsos.230949>

Received: 5 July 2023  
Accepted: 24 October 2023

**Subject Category:**  
Ecology, conservation and global change biology

**Subject Areas:**  
ecology

**Keywords:**  
Apidae, heat stress, imidacloprid, fipronil, *Tetragonula hockingsi*, critical thermal maximum

**Author for correspondence:**  
Holly Farnan  
e-mail: [holly.farnan@my.jcu.edu.au](mailto:holly.farnan@my.jcu.edu.au)

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6922106>.

# Sublethal doses of insecticide reduce thermal tolerance of a stingless bee and are not avoided in a resource choice test

Holly Farnan, Peter Yeeles and Lori Lach

College of Science and Engineering, James Cook University, PO Box 6811, Cairns, Queensland 4870, Australia

HF, 0000-0001-8890-4992; PY, 0000-0003-4719-4511; LL, 0000-0001-5137-5185

Insecticides and climate change are among the multiple stressors that bees face, but little is known about their synergistic effects, especially for non-*Apis* bee species. In laboratory experiments, we tested whether the stingless bee *Tetragonula hockingsi* avoids insecticide in sucrose solutions and how *T. hockingsi* responds to insecticide and heat stress combined. We found that *T. hockingsi* neither preferred nor avoided sucrose solutions with either low ( $2.5 \times 10^{-4}$  ng  $\mu\text{l}^{-1}$  imidacloprid or  $1.0 \times 10^{-4}$  ng  $\mu\text{l}^{-1}$  fipronil) or high ( $2.5 \times 10^{-3}$  ng  $\mu\text{l}^{-1}$  imidacloprid or  $1.0 \times 10^{-3}$  ng  $\mu\text{l}^{-1}$  fipronil) insecticide concentrations when offered alongside sucrose without insecticide. In our combined stress experiment, the smallest dose of imidacloprid ( $7.5 \times 10^{-4}$  ng) did not significantly affect thermal tolerance ( $CT_{\text{max}}$ ). However,  $CT_{\text{max}}$  significantly reduced by  $0.8^\circ\text{C}$  ( $\pm 0.16$  SE) and by  $0.5^\circ\text{C}$  ( $\pm 0.16$  SE) when bees were fed as little as  $7.5 \times 10^{-3}$  ng of imidacloprid or  $3.0 \times 10^{-4}$  ng of fipronil, respectively, and as much as  $1.5^\circ\text{C}$  ( $\pm 0.16$  SE) and  $1.2^\circ\text{C}$  ( $\pm 0.16$  SE) when bees were fed  $7.5 \times 10^{-2}$  ng of imidacloprid or  $3.0 \times 10^{-2}$  ng of fipronil, respectively. Predictions of temperature increase, and increased insecticide use in the tropics suggest that *T. hockingsi* will be at increased risk of the effects of both stressors in the future.

## 1. Introduction

Bees are critical components of natural and agricultural ecosystems [1], and concern is growing about declines in their populations [2–4]. These declines are likely driven by myriad stressors including habitat loss, pathogens and parasites, competition from introduced species, poor nutrition and insecticide exposure [1,2]. The effects of these stressors will likely be exacerbated by global climate change [3,5–7].

Insecticides have been blamed for bee deaths across the globe [8–10]. Bees are exposed to insecticides in the nectar and pollen of treated plants [11] and exposure is expected to become more prolific with increased insecticide use under agricultural intensification [12,13]. Imidacloprid is a neonicotinoid, which is the most widely used class of insecticides worldwide. Of the insecticide classes, neonicotinoids are the most often implicated in bee declines [3,4]. Neonicotinoids are neurotoxins derived from the natural compound nicotine and cause overstimulation, paralysis and death [14]. Sublethal effects, such as difficulty in learning and decreased foraging and homing ability, have also been observed in *Apis mellifera* and *Bombus terrestris* [15–17]. Fipronil is another widely used insecticide with lethal and sublethal effects on insect pollinators [18]. Fipronil is part of the phenylpyrazole chemical family and is an entirely synthetic insecticide that inhibits  $\gamma$ -aminobutyric acid (GABA) receptors in insects, leading to excess neural activity causing insects to experience muscle and nerve hyperexcitability, paralysis, and eventually death [19,20]. Fipronil also has sublethal effects on bees, for example reduced motor activity in *A. mellifera* [21], reduced climbing speed in the stingless bee *Melipona scutellaris* [22], and increased cell cytotoxicity of the mushroom bodies associated with memory in the stingless bee *Scaptotrigona postica* [23].

Lethal and sublethal effects of insecticides on bees could be reduced if bees avoided foraging on nectar and pollen contaminated with insecticides [24,25]; however, evidence that bees avoid consuming insecticide-contaminated resources is limited and conflicting. Individual *B. terrestris* workers given sucrose treated with the neonicotinoid imidacloprid ( $10 \mu\text{g kg}^{-1}$  and  $100 \mu\text{g kg}^{-1}$ ) subsequently reduced their consumption of resources containing imidacloprid over a four-day period [26]. Individual *Bombus impatiens* workers exposed to both field relevant doses of imidacloprid in sucrose and control sucrose did not prefer sucrose containing imidacloprid ( $0.25 \mu\text{g kg}^{-1}$ ,  $1 \mu\text{g kg}^{-1}$ ,  $5 \mu\text{g kg}^{-1}$  and  $10 \mu\text{g kg}^{-1}$ ) across a series of time points [27]. Avoidance of neonicotinoids has also been demonstrated by pollinating beetles and flies offered a choice between a sublethal concentration of imidacloprid ( $1.0 \mu\text{g l}^{-1}$ ,  $0.1 \mu\text{g l}^{-1}$  and  $0.01 \mu\text{g l}^{-1}$ ) and a control in pan traps [28] and by gravid mosquitoes that failed to lay eggs in water with either imidacloprid or chlorpyrifos [29]. In contrast, *A. mellifera* and *B. terrestris* given a choice between a sucrose solution and a sucrose solution with a sublethal dose of either of the neonicotinoids imidacloprid, thiamethoxam or clothianidin (1 nM, 10 nM, 100 nM and 1000 nM) consumed more sucrose solution with either imidacloprid or thiamethoxam than sucrose alone, possibly because of the pharmacological action of these compounds on nicotinic acetylcholine receptors in the bees' brains [30]. *Bombus terrestris audax* increased visits to insecticide-laced sucrose feeders, indicating a preference for thiamethoxam at  $2 \mu\text{g kg}^{-1}$  and  $11 \mu\text{g kg}^{-1}$  over untreated sucrose [31]. Studies that test whether other groups of social bees, such as stingless bees (Meliponini), prefer or avoid neonicotinoids are limited. The stingless bee *Nanotrigona perilampoides* consumed more sucrose containing imidacloprid (LC<sub>20</sub>—a lethal concentration that kills 20% of test subjects) and insecticide free sucrose than sucrose containing imidacloprid (LC<sub>10</sub>—a lethal concentration that kills 10% of the test subjects) [32]. The authors attribute their results to bees exposed to LC<sub>10</sub> attempting to consume as little sugar as possible as a behavioural defence mechanism to avoid imidacloprid intoxication, whereas bees exposed to LC<sub>20</sub> may have experienced physiological stress, and sucrose consumption may have been necessary for them to meet energy requirements for metabolic pathways and detoxifying capabilities [32]. In contrast, the stingless bee *Tetragonula laeviceps* chose honey over both honey containing the insecticides alpha-cypermethrin and spinetoram and the insecticides alone in a Y-tube olfactometer dual choice aroma assay [33]. Given the equivocal results and paucity of studies, further research into insecticide preference/avoidance in stingless bees in particular is warranted.

Heat stress associated with climate change-driven extreme heat events is another major stressor that creates challenges for bees, including impacts on foraging activity, pollination services, task-related physiology, immunocompetence, reproductive capacity, growth and development of bees [34–37]. For example, heat stress damages the fertility of queen bees and the digestive tracts of worker bees [38,39] and can trigger malformations of the proboscis, stinger, wings and legs of *Apis mellifera carnica* [40]. Further, bee communities and their composition are shifting with climate change [41,42]. These shifts may be arising due to mismatches between environmental temperatures and organisms' physiological tolerances [43,44].

Insecticide toxicity and heat stress may have interactive effects, but to date they have not been well studied in many insects including bees. Within the small number of studies investigating these interactive effects, the results are equivocal. Heat stress and several insecticides, including imidacloprid and fipronil, act on the nervous system [45–52]. There is evidence from *A. mellifera* that the combined effect of insecticide exposure and heat stress could result in higher heat tolerance or synergize and cause higher mortality. For example, acute oral exposure to the neonicotinoids imidacloprid and acetamiprid increased thermal tolerance of *A. mellifera* by as much as  $4.3^\circ\text{C}$  [53]. Conversely, *A. mellifera* fed three concentrations (0 ppb, 5 ppb and 20 ppb) of imidacloprid and maintained at temperatures ( $26^\circ\text{C}$  (below



**Figure 1.** *Tetragonula hockingsi* collecting pollen from flower. Photo credit: Campbell Simpson, 2021.

optimal) and 38°C (above optimal)) were more susceptible to imidacloprid, with significantly higher mortality compared to the control (32°C) and showed altered gene regulation [54]. Another study demonstrated that *A. mellifera* colonies treated with imidacloprid (20 ppb for 14 days) and high temperature (41°C for 6 h) exhibited altered metabolic pathways [55]. One of the principal mechanisms used by insects to escape adverse effects of both natural and synthetic toxins, such as nicotine and the neonicotinoids, is metabolic resistance [51]. However, *A. mellifera* have fewer numbers of detoxifying genes than other insects, leaving them more sensitive to some insecticides [56]. Further, it is metabolically and energetically costly for bees to detoxify toxins [51]; this could lead to other impacts on the bees' health. With predicted increases in temperatures and insecticide use globally [57–59], research that further elucidates the effect of these combined stressors is essential.


While a growing body of work is revealing the effects of stressors on *A. mellifera*, we know relatively little about how the thousands of other bee species respond to environmental stressors. Stingless bees are the most diverse group of eusocial bees with over 500 described species distributed throughout the tropical and subtropical regions of Africa, Asia, Oceania, and the Americas [60,61]. They provide pollination services to a wide diversity of tropical plants and crops [60,62]. Several stingless bee species have increased sensitivity to insecticide toxicity compared to *A. mellifera* due to differences in body size, physiology, behaviour, and metabolism [23,63,64], and laboratory tests have shown that neonicotinoids are among the most toxic compounds to stingless bees [65–67]. Most research focus has been on the genera *Melipona*, *Scaptotrigona* and *Nannotrigona*, with effects on the 55 other melipone genera little studied [63]. Furthermore, as tropical species, stingless bees are likely to tolerate a narrower range of temperatures and may be living close to their upper thermal limits compared to temperate species [68]. This may result in stingless bees being more susceptible to the effects of climate change related heat stress.

We assessed how the stingless bee *Tetragonula hockingsi* responds to insecticide laced sucrose (mimicking floral nectar) and then investigated how *T. hockingsi* responds to the combined stress of insecticide exposure and heat stress. We asked whether *T. hockingsi* avoided, preferred, or remained indifferent to both high and low sublethal concentrations of either the neonicotinoid imidacloprid or fipronil when given the choice between sucrose solutions with and without one of these insecticides. We then investigated the effect of three different sublethal doses of these insecticides on thermal tolerance of *T. hockingsi* (figure 1).

## 2. Material and methods

### 2.1. Study organism

*T. hockingsi* are distributed widely across the Australian tropics and provide pollination services for many native plants and crops such as mango, avocado, and macadamia [60]. *T. hockingsi* is one of the two



**Figure 2.** Diagram of experimental arena (240 ml, 119 mm diameter  $\times$  38 mm height) fitted with 0.2 ml Eppendorf vials with different shades indicating treatment types.

species of stingless bees commonly used for commercial crop pollination in Australia [62]. They possess common stingless bee traits such as being active year-round and nesting in various cavities including hollow trees, buildings and rock crevices [60,69]. On average, there are approximately 10 000 workers in each nest [70]. Nests are insulated by an involucre (an enveloping membrane to protect and insulate the brood composed of propolis, cerumen and resins) [69] and *T. hockingsi* can modify nest temperatures by active ventilation at the entrances. This involves workers fanning their wings while facing outward towards the entrance to draw cool air into the nest [69].

## 2.2. Field collection and laboratory conditions

For the choice tests, we collected adult *T. hockingsi* from one hive, located at the Cairns Botanic Gardens (16° 53' 57.882" S, 145° 44' 49.998" E) and two hives at Kewarra Beach (16° 78' 43.58" S, 145° 68' 85" E) in northern Queensland, Australia in June and July of 2021. Bees were collected fresh on the day of each assay between 08.00 and 08.30 in which ambient temperatures were between 21°C and 24°C. Hives were selected based on confirmed identification of *T. hockingsi* via genetic analysis (primers described in [71]) to amplify a 299 bp fragment of the mitochondrial gene cytochrome oxidase I (mt-COI) in *T. hockingsi* (sequences: *Barhock\_R*: AAGCCGAATCCTGGAAGAA and *T\_hock\_COI\_spec\_F5*: GAATTCATCTATTCTTGGA) by Dr Ros Gloag, The University of Sydney. We collected exiting foragers by placing our experimental arenas (240 ml, 119 mm diameter  $\times$  38 mm height polypropylene containers) over the hive entrance. We fitted arenas with a mesh window (approx. 60 mm  $\times$  20 mm) that could be opened and closed to add or remove foragers as necessary to achieve 12 bees per arena. Piloting revealed 12 bees per arena was sufficiently high so that bees would not huddle together and low enough to enable all bees to have access to the solutions during the experiment. Arenas were fitted with eight 0.2 ml Eppendorf vials, each with two 1 mm holes for feeding, based on the design used by Kessler *et al.* [30]. Vials were evenly spaced around the wall of the arenas and inserted horizontally (figure 2). Empty feeding tubes were fitted during field collection to prevent bees from escaping from holes. The use of a single arena for field collection and the choice tests reduced handling of bees. Bees in each arena were initially provided with a piece of cotton wool soaked in 2 ml of 25% (w/v) sucrose solution immediately after capture. Piloting revealed that this concentration was sufficiently attractive to *T. hockingsi* and is within the range of sugar concentration in nectar [72]. Bees were then transported inside arenas covered with a loose dark cloth to a temperature- and humidity-controlled chamber (WiseCube model TEMI850) set at 27°C and 50% relative humidity whereby they remained for an acclimatization period of 3 h. After this period, sucrose-soaked cotton



was removed, and experimental treatments were applied between 11.00 and 11.30 (depending on when they were collected). After 24 h the experiments were ended, and data were collected.

For the thermal tolerance experiment, we collected adult *T. hockingsi* throughout July and August of 2021 from the same Cairns Botanic Gardens hive as for the choice tests. Using the same method and container design as detailed above, we captured approximately 40 exiting foragers on each collection day for eight days from which we arbitrarily selected 24 for that day's experimental replicates (temperature ramp bees  $n = 12$ , temperature control bees  $n = 12$ ). After the eight days, temperature ramped bees  $n = 85$  (12 bees  $\times$  6 insecticide treatments + 13 control bees (1  $\times$  extra bee ran on day 6)) and temperature control bees  $n = 84$  (12 bees  $\times$  6 insecticide treatments + 12 control bees). We captured more bees than required to allow for replacement in case of death due to handling effects; however, in all cases extra bees were not needed as there were no mortalities prior to experiments. Collections occurred 24 h prior to each experiment, between 08.00 and 08.30. Following capture, bees were maintained under the same conditions as detailed above for an acclimation period of 24 h. We removed the sucrose-soaked cotton between 08.00 and 08.30 the next day to fast the bees for 1 h prior to running the experiment.

### 2.3. Selection of insecticides, concentrations and exposure route

For both the choice tests and determination of thermal tolerance, we separately tested two insecticides: the neonicotinoid imidacloprid (Pestanal, Sigma-Aldrich) and fipronil (Termidor Residual 100 g fipronil/L, BASF Australia Ltd), a synthetic neurotoxin. We chose these insecticides because they are widely used, have been implicated in mass mortalities of bees, can be systemic, and bees are easily exposed to them in the nectar and pollen of treated crops [73,74]. For the choice tests, we tested low ( $2.5 \times 10^{-4}$  ng  $\mu\text{l}^{-1}$  and  $1.0 \times 10^{-4}$  ng  $\mu\text{l}^{-1}$ ) and high ( $2.5 \times 10^{-3}$  ng  $\mu\text{l}^{-1}$  and  $1.0 \times 10^{-3}$  ng  $\mu\text{l}^{-1}$ ) concentrations of imidacloprid and fipronil, respectively. For the thermal tolerance experiment, we tested the same low and high concentrations of each insecticide and added a third very high ( $2.5 \times 10^{-2}$  ng  $\mu\text{l}^{-1}$  and  $1.0 \times 10^{-2}$  ng  $\mu\text{l}^{-1}$ ) concentration for imidacloprid and fipronil, respectively (hereafter referred to as low, high and very high for each insecticide). When selecting insecticide concentrations, we evaluated three approaches. Firstly, we considered basing concentrations from amounts of residues found in nectar and pollen; however, there were no data available for Australia. We then considered basing concentrations on known toxicities for the species; however, at the time of the experiments there were no published toxicities for any *Tetragonula* species for either insecticide. Finally, we decided to use studies of other bee species and scaled concentrations to body size [23,33,75] (electronic supplementary material, appendix 1). We chose oral exposure over contact exposure because oral exposure is the most likely route of exposure for bees [76]. Bees commonly consume insecticide residues that have been applied as seed treatments, soil treatments, and foliar sprays. Insecticide residues may be expressed in plant nectar and pollen, guttation fluid, and honeydew [77].

We prepared fresh 200 ml of 25% (w/v) sucrose solution daily of which we set aside 50 ml as our control sucrose solution. We then added either 1278 ng of imidacloprid or 500 ng of Termidor (fipronil) to separate beakers containing 50 ml of sucrose solution, to obtain the solutions with the highest concentrations we tested. We then diluted these solutions to obtain the lower concentrations. We vortexed each treatment for 20 s (WiseMix model VM-10 vortex) and then immediately pipetted 0.2 ml of the solution into its corresponding vial and gently tapped each vial to ensure that no air bubbles were present. Each arena contained two vials of each treatment to allow for more opportunity for bees to feed. All vials were arranged in an alternating pattern so that no two of the same treatments were next to each other and the same arrangement of vials was set for each arena. Two vials of water were added to each arena in case bees required water for hydration or thermoregulation [30]. Vials containing water were not included in data analysis. We calculated the amount of solution consumed from each tube as the difference in the mass of each vial after 24 h using a microbalance (A&D model HR-200, accuracy  $\pm 0.01$  mg) minus the average evaporation control for the respective treatment [30]. For each insecticide, we repeated the experiment with 30 cohorts (10 from each hive) of 12 bees each. Each trial was conducted on a fresh cohort of bees with the same arrangement of vials. After each experiment, we held all bees for an additional 72 h to confirm that our concentrations were truly sublethal. All 720 bees survived this period.

### 2.4. Determination of heat tolerance

To assess thermal tolerance, we used the measure of critical thermal maxima ( $CT_{\text{max}}$ , a measure of the highest temperature at which an organism can maintain neuromuscular function [78]). Following

acclimation, we transferred temperature ramp bees and control bees to individual 2 ml Eppendorf vials with three 1 mm air holes. We piloted different sized vials and found that this vial size minimized stress and allowed bees to avoid incidental contact with the treatment solution. We determined the mass of each individual bee by pre-weighing vials using a microbalance (A&D model HR-200, readability(mg): 0.1) and then reweighing once the bee was inside.

After a 1 h fasting period. Each bee was given 3  $\mu\text{l}$  of either 25% (w/v) sucrose solution or a low, high, or very high concentration of imidacloprid ( $2.5 \times 10^{-4} \text{ ng } \mu\text{l}^{-1}$ ,  $2.5 \times 10^{-3} \text{ ng } \mu\text{l}^{-1}$  and  $2.5 \times 10^{-2} \text{ ng } \mu\text{l}^{-1}$ , respectively) or fipronil ( $1.0 \times 10^{-4} \text{ ng } \mu\text{l}^{-1}$ ,  $1.0 \times 10^{-3} \text{ ng } \mu\text{l}^{-1}$  and  $1.0 \times 10^{-2} \text{ ng } \mu\text{l}^{-1}$ , respectively) in 25% (w/v) sucrose. The 3  $\mu\text{l}$  volume given at these concentrations yielded doses of  $7.5 \times 10^{-4} \text{ ng}$ ,  $7.5 \times 10^{-3} \text{ ng}$  and  $7.5 \times 10^{-2} \text{ ng}$  of imidacloprid and  $3.0 \times 10^{-4} \text{ ng}$ ,  $3.0 \times 10^{-3} \text{ ng}$  and  $3.0 \times 10^{-2} \text{ ng}$  of fipronil. Bees were assigned to treatments randomly, and the observer (H.F. for all trials) was kept blind to treatment throughout the experiment. We prepared all solutions fresh daily and agitated them for 20 s using a vortex (WiseMix model VM-10) prior to administering to the bees. Each treatment was applied to four bees each day (=28 bees total across the seven solutions) with half of the bees temperature ramped and half controlling for the ramping (kept at 26°C in a temperature-controlled cabinet). Five bees did not completely consume the solution after 6 h and were excluded from the experiment. An additional five bees were run on another day to achieve 12 bees for each treatment.

We determined the  $CT_{\text{max}}$  of *T. hockingsi* by ramping a temperature increase from an ambient temperature at a set rate [79]. Each day, we transferred the bees to be temperature ramped to individual 5 ml vials sealed with parafilm to prevent water leakage while submerged. Vials were placed into randomized positions in a vial rack positioned in a water bath made by attaching a temperature-controlled immersion heater (Westinghouse WHSV01K) to a 9 l rectangular plastic tub, following a similar design used by Nacko *et al.* [80] to measure the  $CT_{\text{max}}$  of *T. hockingsi* [80] (electronic supplementary material, appendix 2). The temperature-controlled immersion heater included a jet that kept water well mixed, and we confirmed this by using two thermocouple probes at opposite sides of the bath (at depths in line with the highest and lowest vials) to check that temperature was homogeneous throughout the water bath. We stabilized the water bath temperature for 15 min at 26°C prior to the experiment. The water temperature was calibrated against a HOBO temperature logger (model MX2202, accuracy  $\pm 0.5^\circ\text{C}$ ) that was placed in the middle of the tub. We ramped bees from an acclimation temperature of  $26.2 (\pm 0.2 \text{ SD})^\circ\text{C}$  by increments of  $0.5^\circ\text{C}$  every 2 min until the  $CT_{\text{max}}$  was reached for all individuals. We identified  $CT_{\text{max}}$  as the temperature at which an individual became unresponsive to a stimulus [81,82], which was in this case a single flick of the vial (vial was removed from water bath to administer stimulus and replaced if bee responded [81,83,84]). During the temperature ramp, bees exhibited typical indicators of heat stress, initially including wing fluttering, or extending and holding still a single or both wings [82]. As bees approached their  $CT_{\text{max}}$ , they were unable to right themselves and lost muscular coordination and began spasming. Spasms were often fast and whole-bodied initially, before becoming slower uncontrolled movements of the limbs. Bees eventually became still with head and abdomen adducted, however continued to respond to stimulus through the movement of antennae and limbs. We checked for responses at every  $2^\circ\text{C}$  increase until  $38^\circ\text{C}$ , if necessary (i.e. we only provided stimulus if individuals were motionless), then at  $0.5^\circ\text{C}$  increments beyond this.

We held the temperature ramp control bees ( $n = 14$  for each day) at the constant temperature of  $26^\circ\text{C}$  in a temperature-controlled cabinet, to assess whether mortality would occur due to insecticide treatment only, over the same period that the temperature ramp experiment took place. These bees had received the same insecticide and sucrose control doses as the bees that were temperature ramped. To ensure that temperature control bees experienced the same handling effects as their temperature-ramped counterparts, they were also transferred into individual 5 ml vials sealed with parafilm and stratified in a vial rack prior to placement in the temperature-controlled cabinet.

## 2.5. Statistical analysis

We conducted statistical analysis in R v. 1.4.1103 [85]. For the choice tests, we used a generalized linear model (GLM) [85] with a quasibinomial error distribution for each insecticide to compare the amount of each sucrose solution consumed (the combined value of two vials of the same treatment for each arena) expressed as a proportion of the total amount consumed from the six vials of solutions (i.e. excluding water) in each arena (hereafter referred to as relative proportion). Treatment (i.e. sucrose, water, low concentration of insecticide and high concentration of insecticide), hive, and the interaction between treatment and hive were fixed effects. We tested each model against a null model using a type 3

**Table 1.** Summary of statistical analyses for both experiments showing ANOVA output of generalized linear models for imidacloprid and fipronil choice tests and linear mixed models for comparison of  $CT_{max}$ . Values of  $p$  less than 0.05 are in bold.

	insecticide		Chisq	d.f.	$p$ value
choice tests	imidacloprid	treatment	2.4423	2	0.29
		hive	8.7794	2	<b>0.0124</b>
		treatment:hive	15.7919	4	<b>0.0033</b>
	fipronil	treatment	7.2931	2	<b>0.0261</b>
		hive	1.8829	2	0.39
		treatment:hive	7.2948	4	0.12
comparison of $CT_{max}$	imidacloprid	(intercept)	10493.3187	1	<b>&lt;0.001</b>
		treatment	127.8725	3	<b>&lt;0.001</b>
		bee mass	0.1977	1	0.66
	fipronil	(intercept)	13663.1184	1	<b>&lt;0.001</b>
		treatment	69.7993	3	<b>&lt;0.001</b>
		bee mass	0.6383	1	0.42

ANOVA from the ‘car’ R package [86]. We conducted *post hoc* Tukey HSD tests (emmeans v. 1.5.2-1) [87] to determine whether there were significant differences across treatments and hives.

We used a linear mixed model with a Gaussian error distribution for each insecticide to test whether insecticide exposure influenced  $CT_{max}$  for *T. hockingsi* workers, with  $CT_{max}$  as the response variable and insecticide treatment and bee mass as fixed effects (LMM; lme4 v. 1.1-25) [88]. We initially included an interaction between insecticide dose and bee mass and dropped it from the model when they did not improve model fit by more than 2  $\Delta$ AIC [89]. To account for variation among days, we included ‘day of assay’ as a random effect. We conducted *post hoc* Tukey HSD tests (emmeans v. 1.5.2-1) [87] to determine whether there were significant differences in  $CT_{max}$  across sucrose-control and insecticide treatments. Analysis was conducted on temperature-ramped bees only. No analysis was needed for the temperature control bees as they had a 100% survival rate.


We tested all models to ensure they met assumptions of homogeneity of variance, and independence and normality of residuals by plotting and visually inspecting our data using the ‘qqnorm’ functions in R [85] against the error distributions.

### 3. Results

#### 3.1. Choice tests

For the imidacloprid choice tests, the relative proportion consumed did not vary significantly among treatments, but there was a significant variation by hive and a significant interaction between treatment and hive (table 1), reflecting that hives responded to treatments differently. *Post hoc* comparisons among solutions within hives revealed that the relative proportion of each treatment consumed did not differ significantly among treatments for bees from the Botanic Gardens hive or Kewarra Beach hive 1. However, bees from Kewarra Beach hive 2 consumed a significantly higher relative proportion of insecticide-free sucrose than sucrose containing a low concentration of imidacloprid (Tukey HSD;  $p=0.001$ ) and sucrose containing a high concentration of imidacloprid (Tukey HSD;  $p=0.0007$ ) (figure 3a).

For the fipronil choice tests, the relative proportion consumed varied significantly among treatments, but not by hive (table 1). There was no significant hive by treatment interaction (table 1). *Post hoc* comparisons among treatments within hives revealed that *T. hockingsi* from the Botanic Gardens hive consumed a significantly higher relative proportion of sucrose with a low concentration of fipronil than sucrose with a high concentration of fipronil (Tukey HSD;  $p=0.0315$ ; figure 3b). The relative proportion of insecticide-free sucrose consumed did not differ significantly from solutions with either low or high fipronil concentrations for any of the hives (figure 3b).




**Figure 3.** Mean ( $\pm 95\%$  CI) amount of each treatment consumed as a proportion of the total amount of sucrose-based solutions consumed in each arena, across hives for (a) imidacloprid concentrations (= IMD, low =  $2.5 \times 10^{-4}$  ng  $\mu\text{l}^{-1}$ , high =  $2.5 \times 10^{-3}$  ng  $\mu\text{l}^{-1}$ ) and (b) fipronil concentrations (= FIP, low =  $1.0 \times 10^{-4}$  ng  $\mu\text{l}^{-1}$ , high =  $1.0 \times 10^{-3}$  ng  $\mu\text{l}^{-1}$ ). Error bars without a letter in common above denote significant differences among treatments (Tukey HSD;  $p < 0.005$ ).

The findings from our choice experiments suggest that *T. hockingsi* do not avoid sublethal concentrations of imidacloprid or fipronil, possibly due to an inability to detect such substances or due to indifference, either of which suggest that *T. hockingsi* will not actively avoid exposure to these insecticides in nectar.

### 3.2. Comparison of thermal tolerance

Consumption of imidacloprid and fipronil each significantly decreased the thermal tolerance of *T. hockingsi* individuals (figure 4, table 1). For imidacloprid, *post hoc* pairwise comparisons revealed a significantly lower mean  $CT_{\text{max}}$  in bees that received the high and very high doses compared to bees that received sucrose only (Tukey HSD;  $p < 0.0001$  for both comparisons) and bees that received the low dose (Tukey HSD;  $p = 0.0002$ ,  $p < 0.0001$ , respectively) (figure 4a).  $CT_{\text{max}}$  was also lower in bees





**Figure 4.** Mean ( $\pm 95\%$  CI)  $CT_{\max}$  for *Tetragonula hockingsi* workers (from Cairns Botanic Gardens hive) given (a) sucrose solution and low ( $7.5 \times 10^{-4}$  ng), high ( $7.5 \times 10^{-3}$  ng) and very high ( $7.5 \times 10^{-2}$  ng) doses of imidacloprid in sucrose (IMD,  $n = 12$  for all treatment groups), and (b) sucrose control solution and low ( $3.0 \times 10^{-4}$  ng), high ( $3.0 \times 10^{-3}$  ng) and very high ( $3.0 \times 10^{-2}$  ng) doses of fipronil (FIP,  $n = 12$  for all treatment groups). Points indicate individual bees. Error bars in the same panel without a letter in common above them significantly differ ( $p < 0.05$ ).

that received a very high dose compared to bees that received a high dose (Tukey HSD;  $p = 0.0004$ ) (figure 4a).  $CT_{\max}$  did not differ between bees that received sucrose only and bees that received a low dose of imidacloprid (figure 4a).

For fipronil, *post hoc* pairwise comparisons revealed a significantly lower mean  $CT_{\max}$  between the sucrose control bees and those that received low (Tukey HSD;  $p = 0.0022$ ), high (Tukey HSD;  $p = 0.0003$ ) and very high doses (Tukey HSD;  $p < 0.0001$ ) (figure 4b). *Post hoc* comparisons also revealed significantly lower mean  $CT_{\max}$  values between bees that received low and very high doses (Tukey HSD;  $p = 0.0004$ ) and bees that received high and very high doses (Tukey HSD;  $p = 0.0008$ ) (figure 4b).  $CT_{\max}$  did not differ between bees that received low and high doses of fipronil (figure 4b).

Mean bee mass was  $8.4 (\pm 10$  SD) mg. Mass did not have a significant effect on  $CT_{\max}$  of *T. hockingsi* individuals exposed to either insecticide (table 1).

All 84 bees that were exposed to the seven insecticide treatments but not the temperature ramp survived for the duration of the temperature ramping experiment conducted the same day (approx. 120 min).

## 4. Discussion

Insecticide exposure and climate change are widely acknowledged as key threats to bees, yet there has been little work investigating their interactive and potentially synergistic effects, especially for non-*Apis* bee species. Furthermore, little is known about whether bees can avoid these stressors. Our experiments revealed no consistent avoidance of either imidacloprid- or fipronil-laced sucrose solutions by *T. hockingsi*, as bees from only one out of three hives avoided imidacloprid and none avoided fipronil. We also found that the  $CT_{\max}$  (hereafter termed thermal tolerance) of individuals was significantly reduced by  $0.8^{\circ}\text{C}$  ( $\pm 0.16$  SE) and by  $0.5^{\circ}\text{C}$  ( $\pm 0.16$  SE) when bees were fed as little as  $7.5 \times 10^{-3}$  ng of imidacloprid or  $3.0 \times 10^{-4}$  ng of fipronil, respectively, and as much as  $1.5^{\circ}\text{C}$  ( $\pm 0.16$  SE) and  $1.2^{\circ}\text{C}$  ( $\pm 0.16$  SE) when bees were fed  $7.5 \times 10^{-2}$  ng of imidacloprid or  $3.0 \times 10^{-2}$  ng of fipronil, respectively. Our demonstration of diminished tolerance of heat stress at field relevant insecticide doses combined with predictions of increases in mean and extreme temperatures throughout the range of *T. hockingsi* [57,90] suggest that they will be at increased risk of the effects of both stressors in the future.

Stingless bee responses to insecticides broadly vary, with evidence of avoidance or repellence [32,33,91,92], exclusion of foragers exposed to insecticides [93], not avoiding individuals from their colony treated with insecticide [94] and not rejecting resources containing insecticides [95]. Demonstrations of stingless bee responses to imidacloprid are limited, and we are only aware of two other studies that tested a response of a stingless bee to imidacloprid. The first study demonstrated

that *Nannotrigona aff. testaceicornis* antennated rather than avoided individuals from their colony that were treated topically with a sublethal dose of imidacloprid ( $3.5 \times 10^{-1}$  ng per bee) [94]. The second study showed that *N. perilampoides* consumed more sucrose containing imidacloprid (LC<sub>20</sub>—a lethal concentration that kills 20% of test subjects) and insecticide free sucrose than sucrose containing imidacloprid (LC<sub>10</sub>—a lethal concentration that kills 10% of the test subjects) [32].

While we should not expect that all stingless bee species will respond to insecticides in the same way, the experimental designs in some of these studies provided more opportunity for the bees to detect insecticide via olfaction without gustation. Our study was designed more to test for responses to insecticide in nectar, and therefore a design that included a gustatory response was appropriate. Our findings raise concern given the high potential for stingless bees to forage upon flowering crops treated with insecticides in the field [62,76].

Our findings differ from previous work on other insects that found neonicotinoids to have an attractant [30,31], repellent [28], or antifeedant effect [26]. These differences in findings may be due to a concentration dependent effect, as neonicotinoids may act as an attractant in some concentrations but elicit a neutral response at others [30]. Differences in findings may also be due to the responses measured, experimental design, and insect physiology between stingless bees and other species. Differential responses to insecticides may also depend on the amount of previous exposure bees have experienced. For example, strong repellent effects of low concentrations of neonicotinoids in flies and beetles may be due to the widespread use of neonicotinoids in agricultural settings over previous years, leading to a strong selection for their avoidance [28]. Our results for *T. hockingsi* did not reveal strong repellent effects and align with the work of Muth *et al.* [27], where *B. impatiens* did not show a preference towards consuming either neonicotinoid-containing solutions or sucrose solutions.

Consumption of 3  $\mu$ l of the same concentrations and insecticides that *T. hockingsi* failed to consistently avoid in our nectar choice experiments resulted in significantly lower thermal tolerance. The thermal tolerance of imidacloprid exposed bees declined by 0.8–1.5°C, and the thermal tolerance of fipronil exposed bees declined by 0.5–1.2°C. The survival of the temperature ramp control bees indicates that mortality in the temperature ramp experiment was due to the combination of insecticide exposure and thermal stress, and not just insecticide exposure alone. Our findings contrast with a recent study in which sublethal doses of imidacloprid increased thermal tolerance in *A. mellifera* [53]. The thermal tolerance of *A. mellifera* that received doses ranging from 0.18 ng to 3.6 ng of imidacloprid were on average 2.6°C to 4.3°C greater than the control group. The authors theorized that their results may be due to sublethal doses of imidacloprid activating a stress response in the bees, which in turn may facilitate greater heat resistance. The differences in results between our study and Gonzalez *et al.* [53] may be due to differences in body size and physiology between *T. hockingsi* and *A. mellifera*. For example, when comparing the sensitivity of bee species to 158 different pesticides, bees with greater mean weight (*B. terrestris* and *A. mellifera*) had lower sensitivity than smaller bees (*Nomia melanderi* and *Megachile rotundata*) [96]. Moreover, insect body mass often positively correlates with thermal tolerance [97–101], although not in all cases [102,103]. As *T. hockingsi* have a body mass approximately 1/15 that of *A. mellifera* [75], it is possible that *T. hockingsi* are more susceptible to the effects of insecticides and heat stress simply due to their smaller size [96]. Furthermore, responses to insecticide exposure and heat stress could be dictated at the gene level. For example, non-optimal ambient temperatures aggravate imidacloprid toxicity and affect *A. mellifera* gene regulation [54]. Further documentation of stingless bee biology will help us to elucidate their responses to the combined stress of insecticide exposure and heat stress in the future.

While the findings from both our choice assays and our thermal tolerance experiments provide important proxies for what may occur in the field, the responses of bees under field conditions may vary. For example, our choice assay only allowed for a 24 h exposure period to insecticides, whereas in the field, bees may develop preference for or avoidance of insecticide in nectar over longer time periods. CT<sub>max</sub> is suggested to be a useful predictor of species' responses to climate warming in regions with relatively warm baseline temperatures where many species are already close to their upper thermal limits [104]. Given the tropical distribution of *T. hockingsi*, it is reasonable to assume that CT<sub>max</sub> may be a reliable predictor of thermal tolerance for the species. However, there are some limitations of applying CT<sub>max</sub> values directly to ambient temperatures. The temperature ramping protocols expose bees to a more rapid rise in temperature than is likely to occur under natural conditions [105]. Moreover, bees in a temperature ramping experiment cannot move to cooler microclimates or benefit from hive thermoregulation strategies such as fanning [106]. Another limitation of extrapolating ambient survival from CT<sub>max</sub> values is that differences in thermal tolerance might occur among populations or among castes of social insects like bees [107]. Nevertheless, CT<sub>max</sub>

values provide an indicative measurement for what organisms may experience in the field under extreme thermal stress events. These measurements may help us to predict how the species will respond to changes in climate when also challenged by insecticide exposure.

Tropical rainforest temperatures have been increasing rapidly since the 1970s [57] and are predicted to continue, with global climate models projecting average temperature increases of 1–4°C at lower latitudes [90]. These projections suggest that global warming of less than 1°C will cause tropical regions to experience extreme conditions (i.e. temperatures exceeding 2 standard deviations from the mean) sooner than other regions of the globe [90]. Tropical areas in which *T. hockingsi* are distributed are already experiencing extreme heat waves. For example, the city of Cairns, which is well within the geographic range of this species, experienced an extreme heatwave in 2018 and recorded a high temperature of 42.6°C [108]. Furthermore, tropical species are expected to have low tolerance to increased temperatures given the relatively narrow range of temperatures they typically experience [109]. The tropical distribution of stingless bees puts them further at risk given the trends of increasing insecticide use in the tropics in particular, and the potential for combined effects of insecticide exposure and heat stress [10,96,110]. The results from this study provide evidence of how *T. hockingsi* may respond to the combined stress of changes in climate and insecticide exposure.

**Ethics.** This work did not require ethical approval from a human subject or animal welfare committee.

**Data accessibility.** Data are available at Research Data JCU (doi:10.25903/v799-7v58): <https://research.jcu.edu.au/data/published/173e64c0342611ed907d2d60f024bc99>.

Electronic supplementary material is available online [111].

**Declaration of AI use.** We have not used AI-assisted technologies in creating this article.

**Authors' Contributions.** H.F.: conceptualization, data curation, formal analysis, funding acquisition, methodology, writing—original draft, writing—review and editing; P.Y.: conceptualization, supervision, writing—review and editing; L.L.: conceptualization, supervision, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

**Conflict of interest declaration.** We declare we have no competing interests.

**Funding.** This work was supported by the Skyrail Rainforest Foundation Research Fund and the James Cook University Honours SSA Fund.

**Acknowledgements.** We thank Emma Carmichael for assistance with pilots and Lisl Mohr, Indigo Spence, Grace Kruger-Ilingworth and Sophi Ushay for assistance conducting assays and Steve Williams and Cairns Botanic Gardens for providing bees. We are grateful to Ros Gloag for genetically confirming species identification and James Cook University laboratory technicians for sourcing equipment and assisting with serial dilutions.

## References

- Goulson D, Nicholls E, Botías C, Rotheray EL. 2015 Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* **347**, 1255957. (doi:10.1126/science.1255957)
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353. (doi:10.1016/j.tree.2010.01.007)
- Goulson D. 2013 An overview of the environmental risks posed by neonicotinoid insecticides. *J. Appl. Ecol.* **50**, 977–987. (doi:10.1111/1365-2664.12111)
- Pisa LW *et al.* 2015 Effects of neonicotinoids and fipronil on non-target invertebrates. *Environ. Sci. Pollut. Res. Int.* **22**, 68–102. (doi:10.1007/s11356-014-3471-x)
- Gill RJ, Ramos-Rodriguez O, Raine NE. 2012 Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature (London)* **491**, 105–108. (doi:10.1038/nature11585)
- Cameron SA, Sadd BM. 2020 Global trends in bumble bee health. *Annu. Rev. Entomol.* **65**, 209–232. (doi:10.1146/annurev-ento-011118-111847)
- Kennedy CM *et al.* 2013 A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecol. Lett.* **16**, 584–599. (doi:10.1111/ele.12082)
- Doublet V, Labarussias M, De Miranda JR, Moritz RFA, Paxton RJ. 2015 Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle. *Environ. Microbiol.* **17**, 969–983. (doi:10.1111/1462-2920.12426)
- Hrynko I, Kaczyński P, Łozowicka B. 2021 A global study of pesticides in bees: QuEChERS as a sample preparation methodology for their analysis—critical review and perspective. *Sci. Total Environ.* **792**, 148385. (doi:10.1016/j.scitotenv.2021.148385)
- Sanchez-Bayo F, Goka K. 2014 Pesticide residues and bees: a risk assessment. *PLoS ONE* **9**, e94482. (doi:10.1371/journal.pone.0094482)
- Simon-Delso N *et al.* 2015 Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ. Sci. Pollut. Res. Int.* **22**, 5–34. (doi:10.1007/s11356-014-3470-y)
- National Research Council. 2000 *The future role of pesticides in US agriculture*. Washington, DC: National Academy Press.
- Waltz M. 2020 'How it is grown doesn't matter, as long as it's on the table': pesticide use, uncertainty and future aspirations. *Anthropol. Today* **36**, 25–28. (doi:10.1111/1467-8322.12621)
- Tomizawa M, Millar NS, Casida JE. 2005 Pharmacological profiles of recombinant and native insect nicotinic acetylcholine receptors. *Insect Biochem. Mol. Biol.* **35**, 1347–1355. (doi:10.1016/j.ibmb.2005.08.006)
- Yang EC, Chuang YC, Chen YL, Chang LH. 2008 Abnormal foraging behavior induced by sublethal dosage of imidacloprid in the honey bee (Hymenoptera: Apidae). *J. Econ. Entomol.* **101**, 1743–1748. (doi:10.1603/0022-0493-101.6.1743)
- Mommaerts V, Reynders S, Boulet J, Besard L, Sterk G, Smagghe G. 2010 Risk assessment for side-effects of neonicotinoids against bumblebees with and without impairing foraging behavior. *Ecotoxicology (London)* **19**, 207–215. (doi:10.1007/s10646-009-0406-2)

17. Cresswell JE, Thompson HM. 2012 Comment on 'A common pesticide decreases foraging success and survival in honey bees'. *Science* **337**, 1453. (doi:10.1126/science.1224618)
18. El Hassani AK, Dacher M, Gauthier M, Armengaud C. 2005 Effects of sublethal doses of fipronil on the behavior of the honeybee (*Apis mellifera*). *Pharmacol. Biochem. Behav.* **82**, 30–39. (doi:10.1016/j.pbb.2005.07.008)
19. Cole LM, Nicholson RA, Casida JE. 1993 Action of phenylpyrazole insecticides at the GABA-gated chloride channel. *Pestic. Biochem. Physiol.* **46**, 47–54. (doi:10.1006/pest.1993.1035)
20. Gunasekara AS, Truong T, Goh KS, Spurlock F, Tjeerdema RS. 2007 Environmental fate and toxicology of fipronil. *J. Pesticide Sci.* **32**, 189–199. (doi:10.1584/jpestics.R07-02)
21. Zaluski R, Kadri SM, Alonso DP, Martins Ribolla PE, De Oliveira Osi R. 2015 Fipronil promotes motor and behavioral changes in honey bees (*Apis mellifera*) and affects the development of colonies exposed to sublethal doses. *Environ. Toxicol. Chem.* **34**, 1062–1069. (doi:10.1002/etc.2889)
22. De Moraes CR *et al.* 2018 Ecotoxicological effects of the insecticide fipronil in Brazilian native stingless bees *Melipona scutellaris* (Apidae: Meliponini). *Chemosphere* **206**, 632–642. (doi:10.1016/j.chemosphere.2018.04.153)
23. Jacob CRO, Soares HM, Carvalho SM, Nocelli RF, Malaspina O. 2013 Acute toxicity of fipronil to the stingless bee *Scaptotrigona postica* Latreille. *Bull. Environ. Contam. Toxicol.* **90**, 69–72. (doi:10.1007/s00128-012-0892-4)
24. Department for Environment, Food and Rural Affairs. 2013 *An assessment of key evidence about neonicotinoids and bees*. London, UK: DEFRA.
25. Godfray HCJ, Blacquiere T, Field LM, Hails RS, Potts SG, Raine NE, Vanbergen AJ, McLean AR. 2015 A restatement of recent advances in the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proc. R. Soc. B* **282**, 20151821. (doi:10.1098/rspb.2015.1821)
26. Thompson HM, Wilkins S, Harkin S, Milner S, Walters KFA. 2015 Neonicotinoids and bumblebees (*Bombus terrestris*): effects on nectar consumption in individual workers. *Pest Manag. Sci.* **71**, 946–950. (doi:10.1002/ps.3868)
27. Muth F, Gaxiola R, Leonard A. 2020 No evidence for neonicotinoid preferences in the bumblebee *Bombus impatiens*. *R. Soc. Open Sci.* **7**, 191883. (doi:10.1098/rsos.191883)
28. Easton AH, Goulson D. 2013 The neonicotinoid insecticide imidacloprid repels pollinating flies and beetles at field-realistic concentrations. *PLoS ONE* **8**, e54819. (doi:10.1371/journal.pone.0054819)
29. Al Naggar Y, Geasy JP, Kholy SE. 2019 Sublethal effects of chronic exposure to chlorpyrifos or imidacloprid insecticides or their binary mixtures on *Culex pipiens* mosquitoes. *Physiol. Entomol.* **44**, 123–132. (doi:10.1111/phen.12278)
30. Kessler S, Tiedeken EJ, Simcock KL, Derveau S, Mitchell J, Softley S, Radcliffe A, Stout JC, Wright GA. 2015 Bees prefer foods containing neonicotinoid pesticides. *Nature (London)* **521**, 74–76. (doi:10.1038/nature14414)
31. Arce AN, Ramos Rodrigues A, Yu J, Colgan TJ, Wurm Y, Gill RJ. 2018 Foraging bumblebees acquire a preference for neonicotinoid-treated food with prolonged exposure. *Proc. R. Soc. B* **285**, 20180655. (doi:10.1098/rspb.2018.0655)
32. Al Naggar Y *et al.* 2022 The insecticide imidacloprid decreases *Nannotrigona* stingless bee survival and food consumption and modulates the expression of detoxification and immune-related genes. *Insects (Basel, Switzerland)* **13**, 972. (doi:10.3390/insects13110972)
33. Mubin N, Nurulama L. 2022 Attractiveness and toxicity of two insecticides to *Tetragonula laeviceps* (Apidae: Meliponinae). *IOP Conf. Ser. Earth Environ. Sci.* **974**, 12015. (doi:10.1088/1755-1315/974/1/012015)
34. Zhao H, Li G, Guo D, Li H, Liu Q, Xu B, Guo X. 2021 Response mechanisms to heat stress in bees. *Apidologie* **52**, 388–399. (doi:10.1007/s13592-020-00830-w)
35. Alqarni AS, Ali H, Iqbal J, Oways AA, Smith BH. 2019 Expression of heat shock proteins in adult honey bee (*Apis mellifera* L.) workers under hot-arid subtropical ecosystems. *Saudi J. Biol. Sci.* **26**, 1372–1376. (doi:10.1016/j.sjbs.2019.08.017)
36. Bordier C *et al.* 2017 Colony adaptive response to simulated heat waves and consequences at the individual level in honeybees (*Apis mellifera*). *Sci. Rep.* **7**, 3760. (doi:10.1038/s41598-017-03944-x)
37. Greenop A, Mica-Hawkyard N, Walkington S, Wilby A, Cook SM, Pywell RF, Woodcock BA. 2020 Equivocal evidence for colony level stress effects on bumble bee pollination services. *Insects* **11**, 191. (doi:10.3390/insects11030191)
38. McAfee A, Chapman A, Higo H, Underwood R, Milone J, Foster LJ, Guarna MM, Tarp DR, Pettis JS. 2020 Vulnerability of honey bee queens to heat-induced loss of fertility. *Nat. Sustain.* **3**, 367–376. (doi:10.1038/s41893-020-0493-x)
39. Bach DM, Holzmann MA, Wague F, Miranda JLL, Lopatkin AJ, Mansfield JH, Snow JW. 2021 Thermal stress induces tissue damage and a broad shift in regenerative signaling pathways in the honey bee digestive tract. *J. Exp. Biol.* **224**, jeb242262. (doi:10.1242/jeb.242262)
40. Groh C, Tautz J, Rössler W. 2004 Synaptic organization in the adult honey bee brain is influenced by brood-temperature control during pupal development. *Proc. Natl Acad. Sci. USA* **101**, 4268–4273. (doi:10.1073/pnas.0400773101)
41. Kerr JT *et al.* 2015 Climate change impacts on bumblebees converge across continents. *Science* **349**, 177–180. (doi:10.1126/science.aaa7031)
42. Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R. 2013 Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proc. Natl Acad. Sci. USA* **110**, 4656–4660. (doi:10.1073/pnas.1218503110)
43. Walther G-R. 2010 Community and ecosystem responses to recent climate change. *Phil. Trans. R. Soc. B* **365**, 2019–2024. (doi:10.1098/rstb.2010.0021)
44. Helmuth B, Kingsolver JG, Carrington E. 2005 Biophysics, physiological ecology, and climate change: does mechanism matter? *Annu. Rev. Physiol.* **67**, 177–201. (doi:10.1146/annurev.physiol.67.040403.105027)
45. Jørgensen LB, Robertson RM, Overgaard J. 2020 Neural dysfunction correlates with heat coma and CT<sub>max</sub> in *Drosophila* but does not set the boundaries for heat stress survival. *J. Exp. Biol.* **223**, jeb218750. (doi:10.1242/jeb.218750)
46. Thany SH (ed.). 2010 *Insect nicotinic acetylcholine receptors*, 1st edn. New York, NY: Springer.
47. Fairbrother A, Purdy J, Anderson T, Fell R. 2014 Risks of neonicotinoid insecticides to honeybees. *Environ. Toxicol. Chem.* **33**, 719–731. (doi:10.1002/etc.2527)
48. Privitt JJ, Van Nest BN, Fahrback SE. 2023 Altered synaptic organization in the mushroom bodies of honey bees exposed as foragers to the pesticide fipronil. *Front. Bee Sci.* **1**, 1219991. (doi:10.3389/frbee.2023.1219991)
49. Li X, Ma W, Jiang Y. 2023 Expression patterns of heat shock protein genes and antioxidant genes in *Apis cerana cerana* (Hymenoptera: Apidae) under heat stress. *J. Entomol. Sci.* **58**, 95–103. (doi:10.18474/JES22-27)
50. Li G, Zhang S, Wang H, Liang L, Liu Z, Wang Y, Xu B, Zhao H. 2022 Differential expression characterisation of the heat shock proteins DnaJB6, DnaJshv, DnaJB13, and DnaJB14 in *Apis cerana cerana* under various stress conditions. *Front. Ecol. Evol.* **10**, 873791. (doi:10.3389/fevo.2022.873791)
51. Rand EE, Smit S, Beukes M, Apostolides Z, Pirk CWW, Nicolson SW. 2015 Detoxification mechanisms of honey bees (*Apis mellifera*) resulting in tolerance of dietary nicotine. *Sci. Rep.* **5**, 11779. (doi:10.1038/srep11779)
52. Rajak P, Roy S. 2018 Heat shock proteins and pesticide stress. In *Regulation of heat shock protein responses* (eds A Asea, P Kaur), pp. 27–40. Cham, Switzerland: Springer International Publishing. (doi:10.1007/978-3-319-74715-6\_2)
53. Gonzalez VH *et al.* 2022 Acute exposure to sublethal doses of neonicotinoid insecticides increases heat tolerance in honey bees. *PLoS ONE* **17**, e0240950. (doi:10.1371/journal.pone.0240950)
54. Alburaki M, Madella S, Cook SC. 2023 Non-optimal ambient temperatures aggravate insecticide toxicity and affect honey bees *Apis mellifera* L. gene regulation. *Sci. Rep.* **13**, 3931. (doi:10.1038/s41598-023-30264-0)
55. Kim S, Susie C, Si HL. 2022 Synergistic effects of imidacloprid and high temperature on honey bee colonies. *Apidologie* **53**, 67. (doi:10.1007/s13592-022-00980-z)
56. Gong Y, Diao Q. 2017 Current knowledge of detoxification mechanisms of xenobiotic in honey bees. *Ecotoxicology* **26**, 1–12. (doi:10.1007/s10646-016-1742-7)
57. Malhi Y, Wright J. 2004 Spatial patterns and recent trends in the climate of tropical rainforest regions. *Phil. Trans. R. Soc. Lond. B* **359**, 311–329. (doi:10.1098/rstb.2003.1433)
58. Lewis SE, Silburn DM, Kookana RS, Shaw M. 2016 Pesticide behavior, fate, and effects in the tropics: an overview of the current state of



- knowledge. *J. Agric. Food Chem.* **64**, 3917–3924. (doi:10.1021/acs.jafc.6b01320)
59. Sanchez-Bayo F, Hyne RV. 2011 Comparison of environmental risks of pesticides between tropical and nontropical regions. *Integr. Environ. Assess. Manag.* **7**, 577–586. (doi:10.1002/ieam.189)
60. Heard TA. 1999 The role of stingless bees in crop pollination. *Annu. Rev. Entomol.* **44**, 183–206. (doi:10.1146/annurev.ento.44.1.183)
61. Gruter C. 2020 Stingless bees: their behaviour, ecology and evolution. In *Fascinating life sciences*, 1st edn. Cham, Switzerland: Springer.
62. Vit P, Pedro SRM, Roubik D. 2013 *Pot-honey: a legacy of stingless bees*, 1st edn. New York, NY: Springer New York.
63. Tomé HVV, Martins GF, Lima MAP, Campos LAO, Guedes RNC. 2012 Imidacloprid-induced impairment of mushroom bodies and behavior of the native stingless bee *Melipona quadrifasciata anthidioides*. *PLoS ONE* **7**, e38406. (doi:10.1371/journal.pone.0038406)
64. Jacob CRO, Malaquias JB, Zanardi OZ, Silva CAS, Jacob JFO, Yamamoto PT. 2019 Oral acute toxicity and impact of neonicotinoids on *Apis mellifera* L. and *Scaptotrigona postica* Latreille (Hymenoptera: Apidae). *Ecotoxicology (London)* **28**, 744–753. (doi:10.1007/s10646-019-02070-w)
65. Rafael Valdovinos-Nunez G, Quezada-Euán JJ, Ancona-Xiu P, Moo-Valle H, Carmona A, Ruiz Sanchez E. 2009 Comparative toxicity of pesticides to stingless bees (Hymenoptera: Apidae: Meliponini). *J. Econ. Entomol.* **102**, 1737–1742. (doi:10.1603/029.102.0502)
66. Lima MAP, Martins GF, Oliveira EE, Guedes RNC. 2016 Agrochemical-induced stress in stingless bees: peculiarities, underlying basis, and challenges. *J. Comp. Physiol. A* **202**, 733–747. (doi:10.1007/s00359-016-1110-3)
67. Rosa AD, Teixeira JS, Vollet-Neto A, Queiroz EP, Blochstein B, Pires CS, Imperatriz-Fonseca VL. 2016 Consumption of the neonicotinoid thiamethoxam during the larval stage affects the survival and development of the stingless bee, *Scaptotrigona aff. depilis*. *Apidologie* **47**, 729–738. (doi:10.1007/s13592-015-0424-4)
68. Deutsch CA, Tewksbury JJ, Huey RB, Sheldon KS, Ghalambor CK, Haak DC, Martin PR. 2008 Impacts of climate warming on terrestrial ectotherms across latitude. *Proc. Natl Acad. Sci. USA* **105**, 6668–6672. (doi:10.1073/pnas.0709472105)
69. Heard T. 2016 *The Australian native bee book: keeping stingless bee hives for pets, pollination and sugarbag honey*. West End, Australia: Sugarbag Bees.
70. Brito RM, Schaerf TM, Myerscough MR, Heard TA, Oldroyd BP. 2012 Brood comb construction by the stingless bees *Tetragonula hockingsi* and *Tetragonula carbonaria*. *Swarm Intelligence* **6**, 151–176. (doi:10.1007/s11721-012-0068-1)
71. Paul G, Bartels L, Bueno FG, Law G, Heard T, Chapman N, Buchmann G, Lim J, Gloag R. 2023 Shifting range in a stingless bee leads to pre-mating reproductive interference between species. *Conserv. Genet.* **24**, 449–459. (doi:10.1007/s10592-023-01512-7)
72. Nicolson S. 2022 Sweet solutions: nectar chemistry and quality. *Phil. Trans. R. Soc. B* **377**, 20210163. (doi:10.1098/rstb.2021.0163)
73. Dibartolomeis M, Kegley S, Mineau P, Radford R, Klein K. 2019 An assessment of acute insecticide toxicity loading (AITL) of chemical pesticides used on agricultural land in the United States. *PLoS ONE* **14**, e0220029. (doi:10.1371/journal.pone.0220029)
74. Australian Pesticides and Veterinary Medicines Authority. 2017 *Roadmap for insect pollinator risk assessment in Australia*. Kingston, Australia: APVMA.
75. Purkiss T, Lach L. 2019 Pathogen spillover from *Apis mellifera* to a stingless bee. *Proc. R. Soc. B* **286**, 20191071. (doi:10.1098/rspb.2019.1071)
76. Bonmatin J-M *et al.* 2015 Environmental fate and exposure; neonicotinoids and fipronil. *Environ. Sci. Pollut. Res. Int.* **22**, 35–67. (doi:10.1007/s11356-014-3332-7)
77. Bernal J, Garrido-Bailón E, Del Nozal MJ, González-Porto AV, Martín-Hernández R, Diego JC, Jiménez JJ, Bernal JL, Higes M. 2010 Overview of pesticide residues in stored pollen and their potential effect on bee colony (*Apis mellifera*) losses in Spain. *J. Econ. Entomol.* **103**, 1964–1971. (doi:10.1603/EC10235)
78. Lutterschmidt WI, Hutchison VH. 1997 The critical thermal maximum: history and critique. *Can. J. Zool.* **75**, 1561–1574. (doi:10.1139/z97-783)
79. Terblanche JS, Deere JA, Clusella-Trullas S, Janion C, Chown SL. 2007 Critical thermal limits depend on methodological context. *Proc. R. Soc. B* **274**, 2935–2943. (doi:10.1098/rspb.2007.0985)
80. Nacko S, Hall MA, Gloag R, Lynch KE, Spooner-Hart RN, Cook JM, Riegler M. 2023 Heat stress survival and thermal tolerance of Australian stingless bees. *J. Therm. Biol.* **117**, 103671. (doi:10.1016/j.jtherbio.2023.103671)
81. O'Donnell S, Bulova S, Caponeria V, Oxman K, Giladi I. 2020 Species differ in worker body size effects on critical thermal limits in seed-harvesting desert ants (*Messor ebeninus* and *M. arenarius*). *Insectes Soc.* **67**, 473–479. (doi:10.1007/s00040-020-00782-5)
82. Oyen KJ, Dillon ME. 2018 Critical thermal limits of bumblebees (*Bombus impatiens*) are marked by stereotypical behaviors and are unchanged by acclimation, age or feeding status. *J. Exp. Biol.* **221**, jeb.165589. (doi:10.1242/jeb.165589)
83. Woon JS, Atkinson D, Adu-Bredu S, Eggleton P, Parr CL. 2022 Termites have wider thermal limits to cope with environmental conditions in savannas. *J. Anim. Ecol.* **91**, 766–779. (doi:10.1111/1365-2656.13673)
84. Leahy L, Scheffers BR, Williams SE, Andersen AN. 2022 Arboreality drives heat tolerance while elevation drives cold tolerance in tropical rainforest ants. *Ecology (Durham)* **103**, e03549. (doi:10.1002/ecy.3549)
85. R Core Team. 2021 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
86. Fox J, Sanford W. 2018 *An R companion to applied regression*, 3rd edn. Thousand Oaks, CA: SAGE Publications.
87. Lenth R *et al.* 2023 R package 'emmeans' v. 1.5.2-1.
88. Bates D *et al.* 2020 R package 'lme4' v. 1.1-25.
89. Burnham KP, Anderson DR. 2004 Multimodel inference: understanding AIC and BIC in model selection. *Sociol. Methods Res.* **33**, 261–304. (doi:10.1177/0049124104268644)
90. Beaumont LJ, Pitman A, Perkins S, Zimmermann NE, Yoccoz NG, Thuiller W. 2011 Impacts of climate change on the world's most exceptional ecoregions. *Proc. Natl Acad. Sci. USA* **108**, 2306–2311. (doi:10.1073/pnas.1007217108)
91. Gómez-Escobar E, Liedo P, Montoya P, Vandame R, Sánchez D. 2014 Behavioral response of two species of stingless bees and the honey bee (Hymenoptera: Apidae) to GF-120. *J. Econ. Entomol.* **107**, 1447–1449. (doi:10.1603/EC13490)
92. Bernardes RC, Tomé HVV, Barbosa WF, Guedes RNC, Lima MAP. 2017 Azadirachtin-induced antifeeding in Neotropical stingless bees. *Apidologie* **48**, 275–285. (doi:10.1007/s13592-016-0473-3)
93. Almeida FCR, Magalhães DM, Favaris AP, Rodríguez J, Azevedo KE, Bento JM, Alves DA. 2022 Side effects of a fungus-based biopesticide on stingless bee guarding behaviour. *Chemosphere* **287**, 132147. (doi:10.1016/j.chemosphere.2021.132147)
94. Matos WB, Santos ACC, Lima APS, Santana EDR, Silva JE, Blank AF, Araújo APA, Bacci L. 2021 Potential source of ecofriendly insecticides: essential oil induces avoidance and cause lower impairment on the activity of a stingless bee than organosynthetic insecticides, in laboratory. *Ecotoxicol. Environ. Saf.* **209**, 111764. (doi:10.1016/j.ecoenv.2020.111764)
95. Sánchez D, Solórzano EDJ, Liedo P, Vandame R. 2012 Effect of the natural pesticide spinosad (GF-120 formulation) on the foraging behavior of *Plebeia moureana* (Hymenoptera: Apidae). *J. Econ. Entomol.* **105**, 1234–1237. (doi:10.1603/EC12047)
96. Devillers J, Decourtye A, Budzinski H, Pham-Delegue MH, Cluzeau S, Maurin G. 2003 Comparative toxicity and hazards of pesticides to Apis and non-Apis bees. A chemometrical study. *SAR QSAR Environ. Res.* **14**, 389–403. (doi:10.1080/10629360310001623980)
97. Käfer H *et al.* 2020 Temperature tolerance and thermal environment of European seed bugs. *Insects (Basel, Switzerland)* **11**, 197. (doi:10.3390/insects11030197)
98. Gallego B, Verdú JR, Carrascal LM, Lobo JM. 2016 A protocol for analysing thermal stress in insects using infrared thermography. *J. Therm. Biol.* **56**, 113–121. (doi:10.1016/j.jtherbio.2015.12.006)
99. Nyamukondiwa C, Chidawanyika F, Machekano H, Mutamiswa R, Sands B, Mgidiswa N, Wall R. 2018 Climate variability differentially impacts thermal fitness traits in three coprophagous beetle species. *PLoS ONE* **13**, e0198610. (doi:10.1371/journal.pone.0198610)
100. Baudier KM, Mudd AE, Erickson SC, O'Donnell S. 2015 Microhabitat and body size effects on heat tolerance: implications for responses to climate change (army ants: Formicidae, Ecitoninae).



*J. Anim. Ecol.* **84**, 1322–1330. (doi:10.1111/1365-2656.12388)

101. Ribeiro PL, Camacho A, Navas CA. 2012 Considerations for assessing maximum critical temperatures in small ectothermic animals: insights from leaf-cutting ants. *PLoS ONE* **7**, e32083. (doi:10.1371/journal.pone.0032083)
102. Verble-Pearson RM, Gifford ME, Yanoviak SP. 2015 Variation in thermal tolerance of North American ants. *J. Therm. Biol.* **48**, 65–68. (doi:10.1016/j.jtherbio.2014.12.006)
103. Hemmings Z, Andrew NR. 2017 Effects of microclimate and species identity on body temperature and thermal tolerance of ants (Hymenoptera: Formicidae). *Austral Entomol.* **56**, 104–114. (doi:10.1111/aen.12215)
104. Diamond SE, Nichols LM, McCoy N, Hirsch C, Pelini SL, Sanders NJ, Ellison AM, Gotelli NU, Dunn RR. 2012 A physiological trait-based approach to predicting the responses of species to experimental climate warming. *Ecology (Durham)* **93**, 2305–2312. (doi:10.1890/11-2296.1)
105. Terblanche JS, Hoffmann AA, Mitchell KA, Rako L, Le Roux PC, Chown SL. 2011 Ecologically relevant measures of tolerance to potentially lethal temperatures. *J. Exp. Biol.* **214**, 3713–3725. (doi:10.1242/jeb.061283)
106. Engels W, Rosenkranz P, Engels E. 1995 Thermoregulation in the nest of the Neotropical stingless bee *Scaptotrigona postica* and a hypothesis on the evolution of temperature homeostasis in highly eusocial bees. *Stud. Neotrop. Fauna Environ.* **30**, 193–205. (doi:10.1080/01650529509360958)
107. Maebe K, De Baets A, Vandamme P, Vereecken NJ, Michez D, Smaghe G. 2021 Impact of intraspecific variation on measurements of thermal tolerance in bumble bees. *J. Therm. Biol.* **99**, 103002. (doi:10.1016/j.jtherbio.2021.103002)
108. Australian Bureau of Meteorology. 2021 Monthly climate statistics for 'CAIRNS AERO'. Extremes and records.
109. Menzel F, Feldmeyer B. 2021 How does climate change affect social insects? *Curr. Opin. Insect Sci.* **46**, 10–15. (doi:10.1016/j.cois.2021.01.005)
110. Scopel E *et al.* 2012 Conservation agriculture cropping systems in temperate and tropical conditions, performances and impacts. A review. *Agron. Sustain. Dev.* **33**, 113–130. (doi:10.1007/s13593-012-0106-9)
111. Farnan H, Yeeles P, Lach L. 2023 Sublethal doses of insecticide reduce thermal tolerance of a stingless bee and are not avoided in a resource choice test. *Figshare*. (doi:10.6084/m9.figshare.c.6922106)