

Geophysical Research Letters[®]



RESEARCH LETTER

10.1029/2024GL112903

Key Points:

- This study is the first to report dissolved carbon concentrations, stable isotopes ratio & radiocarbon content from *Acacia* peat plantations
- Dissolved carbon concentrations in this study were among the highest reported, making *Acacia* peat plantations a carbon hotspot
- Oxidation ecosystem respiration, rather than methanogenesis, was the main CO₂ source

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

P. Taillardat,
pierre.taillardat@ntu.edu.sg

Citation:

Taillardat, P., Moore, J., Sasmito, S., Evans, C. D., Alfina, T., Lok, S., et al. (2025). Methane and carbon dioxide production and emission pathways in the belowground and draining water bodies of a tropical peatland plantation forest. *Geophysical Research Letters*, 52, e2024GL112903. <https://doi.org/10.1029/2024GL112903>

Received 17 OCT 2024

Accepted 18 JAN 2025

Author Contributions:

Conceptualization: Pierre Taillardat, Jared Moore, Sigit Sasmito, Chris D. Evans, Aditya Bandla, Chandra S. Deshmukh, Sofyan Kurnianto, Sanjay Swarup, Massimo Lupascu, David Taylor

Data curation: Pierre Taillardat

Formal analysis: Pierre Taillardat









Funding acquisition: Sanjay Swarup, Massimo Lupascu, David Taylor

Investigation: Pierre Taillardat, Jared Moore, Sigit Sasmito, Tiara Alfina, Sophie Lok, Muhandianto Cahya, Rama Kant Dubey

© 2025. The Author(s).

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](https://creativecommons.org/licenses/by/4.0/), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Methane and Carbon Dioxide Production and Emission Pathways in the Belowground and Draining Water Bodies of a Tropical Peatland Plantation Forest

Pierre Taillardat^{1,2} , Jared Moore¹, Sigit Sasmito^{1,3}, Chris D. Evans⁴, Tiara Alfina⁵ , Sophie Lok¹, Aditya Bandla¹ , Muhandianto Cahya⁵, Chandra S. Deshmukh⁶ , Rama Kant Dubey¹, Sofyan Kurnianto⁶, Sanjay Swarup¹, Suria Tarigan⁵ , Muh Taufik⁷ , Massimo Lupascu^{1,8} , and David Taylor⁸ 

¹NUS Environment Research Institute, National University of Singapore, Singapore, Singapore, ²Asian School of the Environment, Nanyang Technological University, Singapore, Singapore, ³Centre for Tropical Water and Aquatic Ecosystem Research, James Cook University, Townsville, QLD, Australia, ⁴UK Centre for Ecology and Hydrology, Bangor, UK, ⁵Department of Soil Science and Land Resource, IPB University, Bogor, Indonesia, ⁶Asia Pacific Resources International Ltd., Pekanbaru, Indonesia, ⁷Department of Geophysics and Meteorology, IPB University, Bogor, Indonesia, ⁸Department of Geography, National University of Singapore, Singapore, Singapore

Abstract Peatlands are crucial yet vulnerable carbon stores. Here, we investigated carbon biogeochemical processes in tropical peatlands converted to plantations. We measured carbon dioxide (CO₂) and methane (CH₄) concentrations, stable isotope ratios and radiocarbon content in an experimental *Acacia crassicarpa* plantation in Sumatra, Indonesia. We found exceptionally high levels of dissolved organic carbon (DOC), CO₂, and CH₄ in porewater and drainage networks, indicating that *Acacia* plantations are carbon hotspots due to their high productivity and exposed carbon-dense substrates. Stable isotope models revealed that while CO₂ and CH₄ are produced belowground, CH₄ contribution was lower than in natural undrained peatlands. Radiocarbon analysis suggested that remobilized carbon contributed to the carbon pool, with a median age of ~470 years before present. These findings constrain the links between land-use, water table levels, and carbon dynamics, with implications for carbon management in plantation peatlands.

Plain Language Summary Peatlands keep a lot of carbon in their deep, wet soils, but their conversion to other land-uses leads to important quantities of carbon emitted to the atmosphere, contributing to global warming. In Southeast Asia, large areas of tropical peatlands have been converted to plantations. Yet, data on carbon production and emissions in these systems remain limited. This study measured the concentration, composition and age of carbon in the water and the soil in an experimental *Acacia* plantation site. We found high concentrations of, on average, century-old carbon in the water, similar to reports from other managed and degraded peatlands. However, in comparison to natural peatlands, we found that there was less methane (a strong greenhouse gas) proportional to carbon dioxide (the most common greenhouse gas) being produced in the studied plantations. Our findings show that it is important to carefully monitor and manage the carbon exchange in peatland plantations, to understand their environmental impact and discuss their role in the carbon cycle and the climate.

1. Introduction

Wetlands cover about 2% of Earth's land yet store a large amount of organic carbon (Blakemore, 2018; Fluet-Chouinard et al., 2023; Poulter et al., 2021) due to vegetation productivity surpassing slow organic matter decomposition in anoxic, waterlogged soils (Temming et al., 2022). Peatlands, the predominant vegetated wetland type, are rapidly declining due to land-use changes (LUC; Fluet-Chouinard et al., 2023). Between 2001 and 2022, approximately 41% of Southeast Asia's peatland forests were impacted by land-use change (Sasmito et al., 2025). Conversion to tree plantations for fiberwood, rubber and palm oil was among the most common land-use changes (Sasmito et al., 2025). *Acacia* plantations, like other managed tree plantations on peatlands, are drained to maintain water table levels (WTLs) below root depth to support plant productivity, impacting carbon dioxide (CO₂) and methane (CH₄) production and release (Figure 1). WTL is managed using artificial drainage networks, including first-order channels (ditches), second-order channels (canals), and dams. Ditches are shallow and slow-moving, while canals are larger and deeper.

Methodology: Pierre Taillardat, Sigit Sasmito

Project administration: Pierre Taillardat, Jared Moore, Sigit Sasmito, Tiara Alfina, Muhardianto Cahya, Chandra S. Deshmukh, Sofyan Kurnianto, Sanjay Swarup, Massimo Lupascu, David Taylor

Resources: Chandra S. Deshmukh, Sofyan Kurnianto, Sanjay Swarup

Supervision: Sanjay Swarup, Massimo Lupascu, David Taylor

Visualization: Pierre Taillardat

Writing – original draft: Pierre Taillardat

Writing – review & editing:

Pierre Taillardat, Jared Moore, Sigit Sasmito, Chris D. Evans, Aditya Bandla, Chandra S. Deshmukh, Sofyan Kurnianto, Suria Tarigan, Muh Taufik, Massimo Lupascu

WTL affects the redox environment, influencing peat organic matter (OM) production, decomposition and preservation (Figure 1). A shallower WTL creates anaerobic conditions near the surface, promoting dissolved organic carbon (DOC) production (Kellerman et al., 2014; Lynch et al., 2019), and OM degradation through acetoclastic methanogenesis (AM) and hydrogenotrophic methanogenesis (HM), resulting in CH₄ emissions typical of undisturbed peatlands (Holmes et al., 2015; Neubauer & Magonigal, 2022). Lowering WTL through drainage exposes upper peat layers to aerobic conditions, increasing CO₂ production through respiration and reducing CH₄ via methane oxidation (Figure 1). Conversely, raising WTLs (e.g., Hooijer et al., 2024; Parish et al., 2019) can decrease CO₂ emissions but increase CH₄ emissions (Evans et al., 2021). While WTL impacts on C-GHG emissions are well-documented (Carlson et al., 2015; Evans et al., 2021; Günther et al., 2020; Huang et al., 2021; Zou et al., 2022), subsurface biogeochemical processes in response to WTL shifts in tropical peatland plantations remain poorly understood.

Aquatic carbon dynamics in tropical peatlands remain uncertain due to a predominant focus on terrestrial GHG emissions (e.g., Deshmukh et al., 2020, 2021, 2023; Jauhiainen et al., 2008; Jauhiainen et al., 2012; Lupascu et al., 2020) and the exchange of peat-derived carbon and C-GHG within drainage networks (e.g., Martin et al., 2018; Moore et al., 2013; Nichols & Martin, 2021; Wit et al., 2018). The conceptual model in Figure 1 outlines pathways and δ¹³C isotopic fractionations related to DOC, CO₂, and CH₄. Carbon input from peat porewaters to drainage networks may include particulate organic carbon, DOC, dissolved CO₂ and CH₄. In surface water, DOC can be mineralized via photochemical oxidation or bacterial metabolism, converting into CO₂, which may either evade the water surface or flow downstream (Somers et al., 2023). Similarly, CH₄ can be oxidized to CO₂ within the water column or released directly into the atmosphere. Because of its low solubility, CH₄ quickly evades or is released through ebullition from deeper water levels, while plants on vegetated surfaces may transport CH₄ emissions via their stems (Akhtar et al., 2021). Limited studies on carbon dynamics in ditches and canals draining managed tropical peatlands indicate high rates of organic carbon cycling and CH₄ production (Bowen et al., 2024; Deshmukh et al., 2020; Jauhiainen & Silvennoinen, 2012; Somers et al., 2023). Larger-scale assessments suggest modified peatland drainage networks could be significant CH₄ emissions hotspots globally (Peacock et al., 2021).

Understanding the factors controlling CH₄ and CO₂ production and transport pathways is crucial for evaluating how land-use change affects C-GHG emissions from peatlands. This study aimed to examine carbon and C-GHG production and transport pathways in the belowground and draining water bodies of an industrial short-rotation fiberwood (*Acacia crassicarpa*) plantation on peat, focusing on (a) the distribution, composition and age of dissolved carbon in an *Acacia* plantation; and (b) the influence of WTL on below-ground C-GHG production and emission pathways.

2. Study Site and Experimental Design

The study was conducted within a 647,000 ha ombrotrophic peatland converted into an industrial *Acacia* plantation in the Kampar Peninsula, Riau Province, Sumatra, Indonesia (Figure 2). The peatland, over 5,100 years old (Hapsari et al., 2022), experienced 2,707 mm of cumulative rainfall in 2022, 53% wetter than the 2014–2021 average (Deshmukh et al., 2021). September to November were the wettest months (>300 mm month⁻¹). January to February and May to August were the driest (<200 mm month⁻¹).

Within the experimental plantation, samples were collected in a 34-ha WTL manipulation experimental plot with a 0.52 m elevation gradient, causing southward water flow (Figure 2). This created a deeper WTL at the higher slope (Water Table Trial Deeper, WTT_Deeper) and a shallower WTL at the lower slope (WTT_Shallower). Despite efforts to maintain WTL near constant year-round by adjusting an outlet dam, variability occurred: mean WTLs were -0.23 ± 0.17 m for WTT_Shallower (negative values indicate water levels below the surface) and -0.67 ± 0.19 m for WTT_Deep (Figure S1 in Supporting Information S1). While the experimental plot had specific WTLs and management, factors such as plant density, fertilization regime, silvicultural practices, and *Acacia* species were consistent with the rest of the plantation. The plot featured a network of narrow, shallow and partly shaded primary drainage channels (ditches) with slow-moving to stagnant water conveying water toward wider, navigable secondary channels (canals) mostly exposed to sunlight. During the driest period, ditches and canals are not surface-connected but remain linked through porewater transfers year-round.




Figure 1. Conceptual model of biogeochemical processes in peat porewater and surface water. “Resp.” = OM respiration, the main degradation pathway in the peat oxic layer. “AM” = acetoclastic methanogenesis, an anoxic pathway producing CO₂ and CH₄. “HM” = hydrogenotrophic methanogenesis, another anoxic pathway, part of CO₂ reduction, producing CO₂ and CH₄ (Corbett et al., 2013). Isotope fractionation factors (α) and expected isotopic compositions ($\delta^{13}\text{C}$) are from Holmes et al. (2015).




Figure 2. (a) Study area in the region and (b) in the Kampar Peninsula; (c) experimental site (0.516545°N; 102.107758°E) satellite imagery with the sampling stations along the North-to-South elevation gradient. (d) Conceptual model of the study site cross-section. Note that the water flow goes from North to South (right to left in the model).

3. Methods

3.1. Data Collection and Analysis

Porewater and surface water samples were collected in February, May, August, and November 2022. Triplicated porewater samples were taken from wells at WTT_Deeper and WTT_Shallower, while surface water samples were collected from ditches and canals banks (Figure 2).

For DOC concentration, water samples were filtered (45 μM cellulose acetate; Whatman, USA) and stored refrigerated in 15 mL PP tubes until analysis using a vario TOC cube analyzer (Elementar, Germany). Acidification was initially deemed unnecessary due to the natural acidity of the water ($\text{pH} < 3.25$). However, this resulted in an approximate 10% overestimation of DOC concentration as compared to the same samples acidified to $\text{pH} < 2$ from residual DIC (Figure S2 in Supporting Information S1). Therefore, a -10% correction was applied to non-acidified samples. Additional filtered water samples were collected in 250 mL Nalgene HDPE bottles and 40 mL glass vials for DO^{14}C analysis at the André E. Lalonde AMS Laboratory and $\delta^{13}\text{C}$ -DOC at the Ján Veizer Stable Isotope Laboratory (University of Ottawa, Canada).

Dissolved CO_2 , CH_4 , and their stable isotope composition ($\delta^{13}\text{C}\text{-CO}_2$ and $\delta^{13}\text{C}\text{-CH}_4$) were sampled using the headspace technique (e.g., Taillardat et al., 2022) and analyzed by cavity ring-down spectroscopy (G2201-i, Picarro Inc, USA). The pCO_2 and pCH_4 (ppm and μatm) were converted to concentrations (μM and mgC L^{-1}) based on water temperature and gas solubility coefficients (Goldenfum, 2011). The headspace technique was also used for dissolved $^{14}\text{C}\text{-CO}_2$ following Garnett et al. (2016) and Murseli et al. (2019). Radiocarbon analyses were performed on an Ionplus AG MICADAS (Mini Carbon Dating System).

Statistical differences in dissolved carbon concentrations, $\delta^{13}\text{C}$, and F^{14}C values across peat porewater sites (WTT_Shallower and WTT_Deeper) and surface waters were examined using non-parametric Kruskal-Wallis and Dunn-Bonferroni post-hoc tests. Statistical analyses and graphs were created in R version 4.4.0 (R Core Team, 2024).

3.2. Stable Isotope Mass Balance Models

Carbon stable isotope ratio ($\delta^{13}\text{C}$) and variations in coexisting dissolved carbon forms offer process-level insights (e.g., Evans et al., 2014; Holmes et al., 2015; Moore et al., 2013; Throckmorton et al., 2015). The difference between $\delta^{13}\text{C}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-CO}_2$ provides the isotope separation ϵ_c of $\text{CO}_2\text{-CH}_4$ that can identify the dominant methanogenic pathway between HM and AM (Figure 1; Whiticar, 1999). The apparent fractionation factors between CO_2 and CH_4 (α) were calculated using Equation 1:

$$\alpha = \frac{\delta^{13}\text{C} - \text{CO}_2 + 1000}{\delta^{13}\text{C} - \text{CH}_4 + 1000} \quad (1)$$

3.2.1. Estimating CO_2 Production From Methanogenesis and Respiration

$\delta^{13}\text{C}\text{-CO}_2$ resulting from methanogenesis ($\delta^{13}\text{C}\text{-CO}_2\text{-meth}$) can be calculated when $\delta^{13}\text{C}\text{-CH}_4$ and $\delta^{13}\text{C}\text{-DOC}$ (or $\delta^{13}\text{C}\text{-OM}$) are known (Equation 2) under two assumptions. First, AM produces equal amounts of CH_4 and CO_2 . In contrast, HM produces CH_4 while consuming equimolar CO_2 but is part of the CO_2 reduction process. This CO_2 reduction originates from non-fractionating OM respiration and releases twice the amount of CO_2 over the amount of produced CH_4 . As a result, when the chemical reactions of both methanogenesis pathways are fully accounted for, they both produce equimolar amounts of CH_4 and CO_2 . However, these pathways do not yield similar $\delta^{13}\text{C}\text{-CH}_4$ values (Figure 1). Second, measured $\delta^{13}\text{C}\text{-CH}_4$ in peat porewaters approximate the CH_4 initially produced from methanogenesis (Corbett et al., 2013; Throckmorton et al., 2015).

$$\delta^{13}\text{C}\text{-CO}_2\text{-meth} = \frac{(\delta^{13}\text{C} - \text{DOC}) - (0.5) \cdot (\delta^{13}\text{C} - \text{CH}_4)}{0.5} \quad (2)$$

Non-methanogenic CO_2 production has the same $\delta^{13}\text{C}$ value as the OM substrate or, in this case, subsurface porewater $\delta^{13}\text{C}\text{-DOC}$ (Throckmorton et al., 2015). After determining $\delta^{13}\text{C} - \text{CO}_2\text{-meth}$ (Equation 3), the fraction

of CO₂ from either isotopic fractionating methanogenesis ($f\text{CO}_{2\text{-meth}}$) or non-fractionating OM respiration and fermentation ($f\text{CO}_{2\text{-OM decay}}$) is determined using Equations 3 and 4:

$$(\delta^{13}\text{C}_{\text{-CO}_2}) = (\delta^{13}\text{C}_{\text{-DOC}}) \cdot (f\text{CO}_{2\text{-OM decay}}) + (\delta^{13}\text{C}_{\text{-CO}_2\text{-meth}}) \cdot (f\text{CO}_{2\text{-meth}}) \quad (3)$$

and

$$f\text{CO}_{2\text{-OM decay}} + f\text{CO}_{2\text{-meth}} = 1 \quad (4)$$

Solving these allows the estimation of $f\text{CO}_{2\text{-meth}}$ and $f\text{CO}_{2\text{-OM decay}}$ (Corbett et al., 2013; Throckmorton et al., 2015). Finally, the quantity of CO₂ produced via methanogenesis ($\text{CO}_{2\text{-meth}}$) can be determined using Equation 5:

$$f\text{CO}_{2\text{-meth}} \cdot \text{CO}_{2\text{-porewater}} = \text{CO}_{2\text{-meth}} \quad (5)$$

where $\text{CO}_{2\text{-porewater}}$ is the measured CO₂ concentration in porewater.

3.2.2. Estimating CH₄ Loss via Ebullition and Plant-Mediated Transport

Since methanogenesis yields equimolar amounts of CH₄ and CO₂, comparing the amount of CO₂ produced via methanogenesis ($\text{CO}_{2\text{-meth}} = \text{CH}_{4\text{-meth}}$) with measured CH₄ concentration ($\text{CH}_{4\text{-measured}}$) allows estimation of CH₄ lost through ebullition or oxidation ($\text{CH}_{4\text{-trans}}$) using Equation 6 (Corbett et al., 2013; Throckmorton et al., 2015).

$$\text{CH}_{4\text{-trans}} = \text{CO}_{2\text{-meth}} - \text{CH}_{4\text{-measured}} \quad (6)$$

3.3. Systematic Literature Review

A systematic review synthesized DOC, dissolved CO₂ and CH₄ concentrations from tropical peat porewater, ditches and canals. A literature search using the Scopus database on 17 April 2024 yielded used here from 50 independent studies and 397 individual study sites were used. Most studies did not provide comprehensive carbon values for all sampling types along the water continuum. A detailed description of the literature review methodology is provided in the Supporting Information S1.

4. Results and Discussion

4.1. Dissolved Carbon Concentrations

The dominant form of dissolved carbon in peat porewater was DOC (~76%; median = 127.1 mgC L⁻¹; range: 104.6 to 144.0 mgC L⁻¹), followed by dissolved CO₂ (~24%; median = 39.6 mgC L⁻¹; range: 20.0 to 62.6 mgC L⁻¹) while dissolved CH₄ made up less than 1%, with a median of 1.4 mgC L⁻¹ (range: 0.1 to 8.3 mgC L⁻¹). There were no significant differences in dissolved carbon concentration between WTT_Shallow compared to WTT_Deep (Figures 3a–3c). In surface water (ditches and canals), DOC comprised over 92% of total dissolved carbon, with dissolved CO₂ at 8%, and CH₄ was negligible. Total dissolved carbon (DOC + CO₂ + CH₄) was higher in ditches than in canals, with statistical significance for DOC and CO₂, but not CH₄ (Figures 3a–3c). Dissolved CO₂ and CH₄ levels were 4 and 92 times lower, respectively, in surface water than in peat porewater, while DOC remained within the same order of magnitude (Figures 3a–3c). No clear seasonal trend emerged between the different forms (Figure S3 in Supporting Information S1), likely due to consistent warm, humid conditions throughout the year.

DOC concentrations at our study site were in the highest range reported from tropical peatland settings (Figure 4a). While average DOC concentrations in peatland canals across the tropics are ~46 mgC L⁻¹, we measured DOC concentrations consistently >100 mgC L⁻¹ (Figure 3a). Previously, the highest values (>65 mgC L⁻¹) were from an undisturbed peat swamp forest (Moore et al., 2013), a logged one (Waldron et al., 2019), an oil palm plantation (Waldron et al., 2019), and a Sago palm plantation (Miyamoto et al., 2009). Two explanations were considered. First, plantation trees are highly productive, likely releasing substantial DOC through root




Figure 3. Dissolved carbon concentrations (a–c, i) $\delta^{13}\text{C}$ (d–f) and F^{14}C values (g, h) in peat porewater (WTT_Shallow, WTT_Deep), ditch and canal. Boxplot medians (thick lines), quartile (box limits), whiskers (1.5 times the interquartile range (IQR)), and data points (dots) are shown. Different lowercase red letters above whiskers indicate significant differences ($p < 0.05$; e.g., *a* and *a* are statistically the same; *a* and *b* are statistically different) between groups. Note the exponential y-axis in panel i.

exudation and leaching from fresh litter (Deshmukh et al., 2020, 2023). Second, the ditch conditions in plantations may prevent DOC degradation. About 30% of the DOC in peatland canals may be degraded through photooxidation (Bowen et al., 2024). Since the studied ditches are slow-moving, shallow and partly shaded (Figure 2d), DOC may remain undegraded until reaching wider, deeper canals downstream where incoming radiation is higher (Figure 3a). Our study is only the fourth to report total carbon concentrations from ditches in tropical peatlands (Cook et al., 2018; Manning et al., 2019; Moore et al., 2013), although the importance of ditches on carbon cycling and C-GHG emissions is globally acknowledged (Peacock et al., 2021). Therefore, the lack of data from




Figure 4. Dissolved carbon concentrations (a–c), $\delta^{13}\text{C}$ (d–f), $F^{14}\text{C}$ values (g, h) and CO_2/CH_4 ratio (i) in peat porewater, ditches and canals from our systematic review. Boxplots show medians (thick lines), quartiles (box limits), whiskers (1.5x IQR), and individual data points from the literature (gray dots) and our study (black dots). Different lowercase red letters above whiskers indicate significant differences ($p < 0.05$) between groups. Note the exponential y-axis on panel (i).

ditches in managed tropical peatlands may be a missing link in understanding peat-derived carbon processes along the aquatic continuum (Evans & Taillardat, 2024).

Few studies have reported dissolved CO_2 and CH_4 concentrations in tropical peatlands (Figures 4b and 4c). Our systematic review found no studies on dissolved CO_2 and only two on dissolved CH_4 in canals draining peatland, with CH_4 concentrations ranging from 0.5 to 248 μgCL^{-1} (Somers et al., 2023; Waldron et al., 2019). Comparisons with global-scale data sets covering wetlands and water bodies more broadly suggest that dissolved CO_2 and CH_4 at our study site are among the highest reported; Liu et al. (2022) reported global dissolved CO_2 values in streams and rivers ranging from 34 to 27,205 μatm , while tropical values ranged from 141 to 22,899 μatm , well




Figure 5. (a) Results from the stable isotope mixing models reporting the contribution of methanogenesis and ecosystem respiration to dissolved CO₂ production. (b) Proportion of CH₄ loss pathways, including diffusion or oxidation versus plant-mediated and ebullition pathways. Bars represent median values, with error bars indicating the first (Q1) and third quartiles (Q3). (c) Crossplot of $\delta^{13}\text{C}$ values in porewaters from WTT_Shallower and WTT_Deepier, ditch and canal. Dot size reflects CH₄ concentration. The oxidation line is from Knorr et al. (2009). The hydrogenotrophic and acetoclastic areas are from Negandhi et al. (2019).

below our measurements (10,984–107,018 μatm or 4.20–40.71 mgC L^{-1}). Similarly, our CH₄ levels are among the highest in the Global River Methane Database (GRiMeDB; Stanley et al. (2023)), even though tropical systems are underrepresented in this database. Although peatland drainage minimizes terrestrial CH₄ emissions (Evans et al., 2021; Günther et al., 2020; Huang et al., 2021; Zou et al., 2022), our results confirm that drainage contributes to additional C-GHG emissions via the aquatic pathway (Deshmukh et al., 2020; Peacock et al., 2021). Further measurements of dissolved CO₂ and CH₄ concentrations and fluxes from peatland drainage networks are necessary to incorporate them into global carbon assessments (e.g., Rocher-ros et al., 2023; Rosentreter et al., 2021).

Radiocarbon analysis showed F¹⁴C values between 0.90 and 0.97, suggesting carbon predominantly fixed before 1955 CE (Figures 3g and 3h). DOC contained significantly more ¹⁴C-enriched samples than CO₂, but the indicative mean ages for both were ~470 years before present. Our results align with previous peatland studies, where F¹⁴C-CO₂ is often lower than F¹⁴C-DOC (Figures 4g and 4h) presumably due to methanogenesis producing a fraction of the total CO₂ from deeper, older anaerobic layers compared to respiration in the aerobic upper layer (Dean et al., 2023). Another explanation could be the different physical properties between the DOC and CO₂. DOC can only leave the peat via the drainage system, whereas younger CO₂ produced near the peat surface may escape to the atmosphere. There was no substantial difference in mean F¹⁴C value between WTT_Shallower and WTT_Deepier (Figures 3g and 3h). These results support the rationale that deeper WTLs in degraded and managed peatlands expose older OM to decomposition or dissolution (Cook et al., 2018; Evans et al., 2014; Moore et al., 2013), despite the high *Acacia* productivity in our case.

4.2. Water Level Influences Below-Ground Gas Production Magnitude and Emission Pathways

The $\delta^{13}\text{C}$ values measured at our site (Figures 4d–4f) are consistent with existing data for tropical peatlands. The porewater $\delta^{13}\text{C}\text{-CO}_2$ ($\sim -23\text{‰}$) is similar to values from forested areas with C_3 -derived OM degradation (Campeau et al., 2017a, 2017b; Hutchins et al., 2020) and a tropical wetland in Panama (Holmes et al., 2015). Using an isotope mass balance model (Equations 3 and 4), our results indicate that ecosystem respiration, a non-fractionating pathway, was the main source of CO_2 at both sites (Figure 5a). The predominance of respiration ($>80\%$) contrasts with undisturbed boreal peatlands, where most dissolved CO_2 in porewater is from methanogenesis, indicated by a median porewater $\delta^{13}\text{C}\text{-CO}_2$ of -3.8‰ (Taillardat et al., 2022) or $\delta^{13}\text{C}\text{-DIC}$ of -4.6‰ (Campeau et al., 2018). The importance of ecosystem respiration can be explained by increasing oxic layer depth from drainage and the year-round warm, humid conditions leading to respiration rates greater than those at higher latitudes.

The significantly lower $\delta^{13}\text{C}\text{-CO}_2$ and CO_2/CH_4 ratio at WTT_Shallower compared to WTT_Deeper (Figures 3e and 3i), support the hypothesis that WTL influences belowground gas production. According to our mass balance model, methanogenesis was proportionally more important at WTT_Shallower, even though $>80\%$ of CO_2 was from oxic respiration (Figure 5a). Increased hypoxia suitable for methanogenesis and reduced methanotrophy are expected with a shallower WTL (Dean et al., 2018). Access to fresher, more labile OM may further stimulate degradation in upper peat layers, explaining higher CO_2 and CH_4 concentrations at WTT_Shallower.

The very negative $\delta^{13}\text{C}\text{-CH}_4$ ($< -70\text{‰}$) at both WTTs suggests that methanogenesis was predominantly through the HM pathway (Figure 5c), which is consistent with undisturbed tropical peatlands (Holmes et al., 2015). Our analysis of CH_4 loss pathways (Equation 6) suggests that most dissolved CH_4 in peat porewater was lost via ebullition and plant-mediated transport rather than being available for diffusion, oxidation, or lateral export to ditches, with little variation between WTTs (Figure 5b). This aligns with findings from higher latitudes (Corbett et al., 2013; Throckmorton et al., 2015). The under-documented importance of ebullition and plant-mediated emissions could potentially explain global CH_4 budget discrepancies (Saunio et al., 2020) and requires further investigations (Bodmer et al., 2023).

5. Conclusion

In reporting among the highest published value for DOC, dissolved CO_2 , and CH_4 concentrations, our study highlights that *Acacia* plantations on tropical peat are significant carbon production, transformation and emission hotspots. This is due to a combination of high productivity and carbon-dense substrates exposed to remobilization, combined factors that are also found on other fast-growing tree plantations on peat. Our findings also suggest that within the experimental forest plantation, WTL influenced carbon-GHG production, with higher dissolved CH_4 concentrations relative to CO_2 in areas where the WTL was shallower, where surrounding organic matter was younger and potentially more labile. Findings from this study are essential to understanding carbon cycling in *Acacia* plantations on peat and developing water management practices that can reduce emissions while maintaining commercial productivity.

Data Availability Statement

Data and code to reproduce the analysis, results and figures are available in the Zenodo repository (Taillardat, 2025; Taillardat et al., 2025).

References

- Akhtar, H., Lupascu, M., Sukri, R. S., Smith, T. E. L., Cobb, A. R., & Swarup, S. (2021). Significant sedge-mediated methane emissions from degraded tropical peatlands. *Environmental Research Letters*, 16(1), 014002. <https://doi.org/10.1088/1748-9326/abc7dc>
- Blakemore, R. J. (2018). Non-flat earth recalibrated for terrain and topsoil. *Soil Systems*, 2(4), 1–39. <https://doi.org/10.3390/soilsystems2040064>
- Bodmer, P., Vroom, R. J. E., Stepina, T., del Giorgio, P. A., & Kosten, S. (2023). Methane dynamics in vegetated habitats in inland waters: Quantification, regulation, and global significance. *Frontiers in Water*, 5, 1332968. <https://doi.org/10.3389/frwa.2023.1332968>
- Bowen, J. L., Wahyudjo, P. J., Anshari, G. Z., Aluwihare, L. I., & Hoyt, A. M. (2024). Canal networks regulate aquatic carbon export from degraded tropical peatlands. *Nature Geoscience*, 17(3), 213–218. <https://doi.org/10.1038/s41561-024-01383-8>
- Campeau, A., Bishop, K., Nilsson, M. B., Klemmedtsson, L., Laudon, H., Leith, F. I., et al. (2018). Stable carbon isotopes reveal soil-stream DIC linkages in contrasting headwater catchments. *Journal of Geophysical Research: Biogeosciences*, 123(1), 149–167. <https://doi.org/10.1002/2017JG004083>
- Campeau, A., Bishop, K. H., Billett, M. F., Garnett, M. H., Laudon, H., Leach, J. A., et al. (2017a). Aquatic export of young dissolved and gaseous carbon from a pristine boreal fen: Implications for peat carbon stock stability. *Global Change Biology*, 23(12), 5523–5536. <https://doi.org/10.1111/gcb.13815>

Acknowledgments

We thank Asia Pacific Resources International Limited for study site access and supporting the research, and Nurul Pertiwi for her assistance with Figure 2c. This research was supported by the Temasek Foundation as part of the Integrated Tropical Peatlands Research Programme at NUS Environmental Research Institute (A-0001182-00-00).

- Campeau, A., Wallin, M. B., Giesler, R., Löfgren, S., Mörh, C. M., Schiff, S., et al. (2017b). Multiple sources and sinks of dissolved inorganic carbon across Swedish streams, refocusing the lens of stable C isotopes. *Scientific Reports*, 7(1), 1–14. <https://doi.org/10.1038/s41598-017-09049-9>
- Carlson, K. M., Goodman, L. K., & May-Tobin, C. C. (2015). Modeling relationships between water table depth and peat soil carbon loss in Southeast Asian plantations. *Environmental Research Letters*, 10(7), 074006. <https://doi.org/10.1088/1748-9326/10/7/074006>
- Cook, S., Whelan, M. J., Evans, C. D., Gauci, V., Peacock, M., Garnett, M. H., et al. (2018). Fluvial organic carbon fluxes from oil palm plantations on tropical peatland. *Biogeosciences*, 15(24), 7435–7450. <https://doi.org/10.5194/bg-15-7435-2018>
- Corbett, J. E., Tfaily, M. M., Burdige, D. J., Cooper, W. T., Glaser, P. H., & Chanton, J. P. (2013). Partitioning pathways of CO₂ production in peatlands with stable carbon isotopes. *Biogeochemistry*, 114(1–3), 327–340. <https://doi.org/10.1007/s10533-012-9813-1>
- Dean, J. F., Billett, M. F., Turner, T. E., Garnett, M. H., Andersen, R., McKenzie, R. M., et al. (2023). Peatland pools are tightly coupled to the contemporary carbon cycle. *Global Change Biology*, 30(10), 1–16. <https://doi.org/10.1111/gcb.16999>
- Dean, J. F., Middelburg, J. J., Röckmann, T., Aerts, R., Blauw, L. G., Egger, M., et al. (2018). Methane feedbacks to the global climate system in a warmer world. *Reviews of Geophysics*, 56(1), 207–250. <https://doi.org/10.1002/2017RG000559>
- Deshmukh, C. S., Julius, D., Desai, A. R., Asyhari, A., Page, S. E., Nardi, N., et al. (2021). Conservation slows down emission increase from a tropical peatland in Indonesia. *Nature Geoscience*, 14(7), 484–490. <https://doi.org/10.1038/s41561-021-00785-2>
- Deshmukh, C. S., Julius, D., Evans, C. D., Nardi, Susanto, A. P., Page, S. E., et al. (2020). Impact of forest plantation on methane emissions from tropical peatland. *Global Change Biology*, 26(4), 2477–2495. <https://doi.org/10.1111/gcb.15019>
- Deshmukh, C. S., Susanto, A. P., Nardi, N., Nurholis, N., Kurnianto, S., Suardiwerianto, Y., et al. (2023). Net greenhouse gas balance of fibre wood plantation on peat in Indonesia. *Nature*, 616(4), 740–746. <https://doi.org/10.1038/s41586-023-05860-9>
- Evans, C. D., Page, S. E., Jones, T., Moore, S., Gauci, V., Laiho, R., et al. (2014). Contrasting vulnerability of drained tropical and high-latitude peatlands to fluvial loss of stored carbon. *Global Biogeochemical Cycles*, 28(11), 1215–1234. <https://doi.org/10.1002/2013GB004782>
- Evans, C. D., Peacock, M., Baird, A. J., Artz, R. R. E., Burden, A., Callaghan, N., et al. (2021). Overriding water table control on managed peatland greenhouse gas emissions. *Nature*, 593(7860), 548–552. <https://doi.org/10.1038/s41586-021-03523-1>
- Evans, C. D., & Taillardat, P. (2024). Light on dark water. *Nature Geoscience*, 17(3), 175–176. <https://doi.org/10.1038/s41561-024-01376-7>
- Fluet-Chouinard, E., Stocker, B. D., Zhang, Z., Malhotra, A., Melton, J. R., Poulter, B., et al. (2023). Extensive global wetland loss over the past three centuries. *Nature*, 614(7947), 281–286. <https://doi.org/10.1038/s41586-022-05572-6>
- Garnett, M. H., Gulliver, P., & Billett, M. F. (2016). A rapid method to collect methane from peatland streams for radiocarbon analysis. *Ecology*, 9(1), 113–121. <https://doi.org/10.1002/eco.1617>
- Goldenfum, J. A. (2011). GHG measurement guidelines for freshwater reservoirs.
- Günther, A., Barthelmes, A., Huth, V., Joosten, H., Jurasinski, G., Koebsch, F., & Couwenberg, J. (2020). Prompt rewetting of drained peatlands reduces climate warming despite methane emissions. *Nature Communications*, 11(1), 1–5. <https://doi.org/10.1038/s41467-020-15499-z>
- Hapsari, K. A., Jennerjahn, T., Hari, S., Eko, N., & Behling, H. (2022). Sea level rise and climate change acting as interactive stressors on development and dynamics of tropical peatlands in coastal Sumatra and South Borneo since the Last Glacial Maximum. *Global Change Biology*, 28(1), 3459–3479. <https://doi.org/10.1111/gcb.16131>
- Holmes, M. E., Chanton, J. P., Tfaily, M. M., & Orgam, A. (2015). CO₂ and CH₄ isotope compositions and production pathways in a tropical peatland. *Global Biogeochemical Cycles*, 29(1), 1–18. <https://doi.org/10.1111/1462-2920.13280>
- Hooijer, A., Vernimmen, R., Mulyadi, D., Triantomo, V., Lampela, M., Agusti, R., et al. (2024). Benefits of tropical peatland rewetting for subsidence reduction and forest regrowth: Results from a large - Scale restoration trial. *Scientific Reports*, 14(5), 1–12. <https://doi.org/10.1038/s41598-024-60462-3>
- Huang, Y., Ciais, P., Luo, Y., Zhu, D., Wang, Y., Qiu, C., et al. (2021). Tradeoff of CO₂ and CH₄ emissions from global peatlands under water-table drawdown. *Nature Climate Change*, 11(7), 618–622. <https://doi.org/10.1038/s41558-021-01059-w>
- Hutchins, R. H. S., Tank, S. E., Olefeldt, D., Quinton, W. L., Spence, C., Dion, N., et al. (2020). Fluvial CO₂ and CH₄ patterns across wildfire-disturbed ecozones of subarctic Canada: Current status and implications for future change. *Global Change Biology*, 26(4), 2304–2319. <https://doi.org/10.1111/gcb.14960>
- Jauhiainen, J., Hooijer, A., & Page, S. E. (2012). Carbon dioxide emissions from an Acacia plantation on peatland in Sumatra, Indonesia. *Biogeosciences*, 9(2), 617–630. <https://doi.org/10.5194/bg-9-617-2012>
- Jauhiainen, J., Limin, S., Silvennoinen, H., & Vasander, H. (2008). Carbon dioxide and methane fluxes in drained tropical peat before and after hydrological restoration. *Ecology*, 89(12), 3503–3514. <https://doi.org/10.1890/07-2038.1>
- Jauhiainen, J., & Silvennoinen, H. (2012). Diffusion GHG fluxes at tropical peatland drainage canal water surfaces. *Suoseura — Finnish Peatland Society*, 63(3–4), 93–105.
- Kellerman, A. M., Dittmar, T., Kothawala, D. N., & Tranvik, L. J. (2014). Chemodiversity of dissolved organic matter in lakes driven by climate and hydrology. *Nature Communications*, 5(5), 1–8. <https://doi.org/10.1038/ncomms4804>
- Knorr, K. H., Lischeid, G., & Blodau, C. (2009). Dynamics of redox processes in a minerotrophic fen exposed to a water table manipulation. *Geoderma*, 153(3–4), 379–392. <https://doi.org/10.1016/j.geoderma.2009.08.023>
- Liu, S., Kuhn, C., Amatulli, G., Aho, K., Butman, D. E., Allen, G. H., et al. (2022). The importance of hydrology in routing terrestrial carbon to the atmosphere via global streams and rivers. *Proceedings of the National Academy of Sciences of the United States of America*, 119(11), 1–9. <https://doi.org/10.1073/pnas.2106322119>
- Lupascu, M., Akhtar, H., Smith, T. E. L., & Sukri, R. S. (2020). Post-fire carbon dynamics in the tropical peat swamp forests of Brunei reveal long-term elevated CH₄ flux. *Global Change Biology*, 26(9), 5125–5145. <https://doi.org/10.1111/gcb.15195>
- Lynch, L. M., Suttin, N. A., Feghel, T. S., Boot, C. M., Covino, T. P., & Wallenstein, M. D. (2019). River channel connectivity shifts metabolite composition and dissolved organic matter chemistry. *Nature Communications*, 10(1), 459. <https://doi.org/10.1038/s41467-019-08406-8>
- Manning, F. C., Kho, L. K., Hill, T. C., Cornulier, T., & Teh, Y. A. (2019). Carbon emissions from oil palm plantations on peat soil. *Frontiers in Forests and Global Change*, 2(8), 37. <https://doi.org/10.3389/ffgc.2019.00037>
- Martin, P., Cherukuru, N., Tan, A. S. Y., Sanwlan, N., Mujahid, A., & Müller, M. (2018). Distribution and cycling of terrigenous dissolved organic carbon in peatland-draining rivers and coastal waters of Sarawak, Borneo. *Biogeosciences*, 15(22), 6847–6865. <https://doi.org/10.5194/bg-15-6847-2018>
- Miyamoto, E., Matsuda, S., Ando, H., Kakuda, K. I., Jong, F. S., & Watanabe, A. (2009). Effect of sago palm (*Metroxylon sagu* Rottb.) cultivation on the chemical properties of soil and water in tropical peat soil ecosystem. *Nutrient Cycling in Agroecosystems*, 85(2), 157–167. <https://doi.org/10.1007/s10705-009-9255-x>
- Moore, S., Evans, C. D., Page, S. E., Garnett, M. H., Jones, T. G., Freeman, C., et al. (2013). Deep instability of deforested tropical peatlands revealed by fluvial organic carbon fluxes. *Nature*, 493(7434), 660–663. <https://doi.org/10.1038/nature11818>

- Murseli, S., Middlestead, P., St-Jean, G., Zhao, X., Jean, C., Crann, C. A., et al. (2019). The preparation of water (DIC, DOC) and gas (CO₂, CH₄) samples for radiocarbon analysis at AEL-AMS, Ottawa, Canada. *Radiocarbon*, *61*(5), 1563–1571. <https://doi.org/10.1017/rdc.2019.14>
- Negandhi, K., Edwards, G., Kelleway, J. J., Howard, D., Safari, D., & Saintilan, N. (2019). Blue carbon potential of coastal wetland restoration varies with inundation and rainfall. *Scientific Reports*, *9*(1), 1–9. <https://doi.org/10.1038/s41598-019-40763-8>
- Neubauer, S. C., & Megonigal, J. P. (2022). Biogeochemistry of wetland carbon preservation and flux. In K. Krauss, Z. Zhu, & C. L. Stagg (Eds.), *Wetland carbon and environmental management* (pp. 33–71). <https://doi.org/10.1002/9781119639305.ch3>
- Nichols, R. S., & Martin, P. (2021). Low biodegradability of dissolved organic matter from Southeast Asian peat-draining rivers. *Journal of Geophysical Research: Biogeosciences*, *126*(6), e2020JG006182. <https://doi.org/10.1029/2020JG006182>
- Parish, F., Lew, S. Y., Faizuddin, M., & Giesen, W. (2019). *RSP0 manual on best management practices (BMPs) for management and rehabilitation of peatlands*. Kuala Lumpur.
- Peacock, M., Audet, J., Bastviken, D., Futter, M. N., Gauci, V., Grinham, A., et al. (2021). Global importance of methane emissions from drainage ditches and canals. *Environmental Research Letters*, *16*(4), 044010. <https://doi.org/10.1088/1748-9326/abeb36>
- Poulter, B., Fluet-Chouinard, E., Hugelius, G., Koven, C., Fatoyinbo, L., Page, S. E., et al. (2021). A review of global wetland carbon stocks and management challenges. *Geophysical Monograph Series*, 1–20. <https://doi.org/10.1002/9781119639305.ch1>
- R Core Team. (2024). A language and environment for statistical computing. Retrieved from <https://www.r-project.org/>
- Rocher-roc, G., Stanley, E. H., Loken, L. C., Casson, N. J., Raymond, P. A., Liu, S., et al. (2023). Global methane emissions from rivers and streams. *Nature*, *621*(7979), 530–535. <https://doi.org/10.1038/s41586-023-06344-6>
- Rosentreter, J. A., Borges, A. V., Deemer, B. R., Holgerson, M. A., Liu, S., Song, C., et al. (2021). Half of global methane emissions come from highly variable aquatic ecosystem sources. *Nature Geoscience*, *14*(4), 225–230. <https://doi.org/10.1038/s41561-021-00715-2>
- Sasmito, S. D., Taillardat, P., Adinugroho, W. C., Krisnawati, H., Novita, N., Fatoyinbo, T., et al. (2025). Half of land use carbon emissions in Southeast Asia can be mitigated through peat swamp forest and mangrove conservation and restoration. *Nature Communications*, *16*, 745. <https://doi.org/10.1038/s41467-025-55892-0>
- Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., et al. (2020). The global methane budget 2000–2017. *Earth System Science Data*, *12*(3), 1561–1623. <https://doi.org/10.5194/essd-12-1561-2020>
- Somers, L. D., Hoyt, A., Cobb, A. R., Insin, S., Suhup, M. A. A., bin, H., et al. (2023). Processes controlling methane emissions from a tropical peatland drainage canal. *Journal of Geophysical Research: Biogeosciences*, *128*(3), 1–20. <https://doi.org/10.1029/2022jg007194>
- Stanley, E. H., Loken, L. C., Casson, N. J., Oliver, S. K., Sponseller, R. A., Wallin, M. B., et al. (2023). GRiMeDB: The Global River Methane database of concentrations and fluxes. *Earth System Science Data*, *15*(7), 2879–2926. <https://doi.org/10.5194/essd-15-2879-2023>
- Taillardat, P. (2025). Methane and carbon dioxide production and emission pathways in the belowground and draining water bodies of a tropical peatland plantation forest (v1.0.1) [Software]. *Zenodo*. <https://doi.org/10.5281/zenodo.14649234>
- Taillardat, P., Bodmer, P., Deblois, C. P., Ponçot, A., Prijac, A., Riahi, K., et al. (2022). Carbon dioxide and methane dynamics in a peatland headwater stream: Origins, processes and implications. *Journal of Geophysical Research: Biogeosciences*, *127*(7), 1–27. <https://doi.org/10.1029/2022JG006855>
- Taillardat, P., Moore, J., Sasmito, S. D., Evans, C. D., Alfina, T., Lok, S., et al. (2025). Methane and carbon dioxide production and emission pathways in the belowground and draining water bodies of a tropical peatland plantation forest [Dataset]. *Zenodo*. <https://doi.org/10.5281/zenodo.14626004>
- Temmink, R. J. M., Lamers, L. P. M., Angelini, C., Bouma, T. J., Fritz, C., van de Koppel, J., et al. (2022). Recovering wetland biogeomorphic feedbacks to restore the world's biotic carbon hotspots. *Science*, *376*(6593), eabn1479. <https://doi.org/10.1126/science.abn1479>
- Throckmorton, H. M., Heikoop, J. M., Newman, B. D., Altmann, G. L., Conrad, M. S., Muss, J. D., et al. (2015). Pathways and transformations of dissolved methane and dissolved inorganic carbon in Arctic tundra watersheds: Evidence from analysis of stable isotopes. *Global Biogeochemical Cycles*, *29*(11), 1893–1910. <https://doi.org/10.1002/2014GB005044>. Received
- Waldron, S., Vihermaa, L., Evers, S., Garnett, M. H., Newton, J., & Henderson, A. C. G. (2019). C mobilisation in disturbed tropical peat swamps: Old DOC can fuel the fluvial efflux of old carbon dioxide, but site recovery can occur. *Scientific Reports*, *9*(1), 1–12. <https://doi.org/10.1038/s41598-019-46534-9>
- Whiticar, M. J. (1999). Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane. *Chemical Geology*, *161*(1), 291–314. [https://doi.org/10.1016/S0009-2541\(99\)00092-3](https://doi.org/10.1016/S0009-2541(99)00092-3)
- Wit, F., Rixen, T., Baum, A., Pranowo, W. S., & Hutahaean, A. A. (2018). The Invisible Carbon Footprint as a hidden impact of peatland degradation inducing marine carbonate dissolution in Sumatra, Indonesia. *Scientific Reports*, *8*(1), 1–11. <https://doi.org/10.1038/s41598-018-35769-7>
- Zou, J., Ziegler, A. D., Chen, D., Mcnicol, G., Ciaia, P., Jiang, X., et al. (2022). Rewetting global wetlands effectively reduces major greenhouse gas emissions. *Nature Geoscience*, *15*(8), 627–632. <https://doi.org/10.1038/s41561-022-00989-0>