












RESEARCH PAPER

Hypoxia and ischemic stroke modify cerebrovascular tone by upregulating endothelial BK(Ca) channels—Lessons from rat, pig, mouse, and human

Christian Staehr^{1,2,3}  | Victoria Hinkley^{3,4}  | Vladimir V. Matchkov¹  |
 Rajkumar Rajanathan¹  | Line Mathilde B. Hansen¹  | Yvonne Eiby⁴  |
 Nathan Luque^{5,6}  | Ian Wright^{4,7}  | Stella T. Bjorkman⁴  | Stephanie M. Miller⁴  |
 Rohan S. Grimley^{8,9,10}  | Andrew Dettrick⁸  | Kirat Chand⁴  | Hong L. Nguyen¹¹  |
 Nicole M. Jones¹¹  | Tim V. Murphy⁶  | Shaun L. Sandow^{3,4} 

¹Department of Biomedicine, Health, Aarhus University, Aarhus, Denmark

²Department of Anaesthesiology and Intensive Care, Aarhus University Hospital, Aarhus, Denmark

³Biomedical Science, School of Health, University of the Sunshine Coast, Maroochydore, Queensland, Australia

⁴UQ Centre for Clinical Research, School of Clinical Medicine, Faculty of Medicine, University of Queensland, St Lucia, Queensland, Australia

⁵Blacktown Clinical School, Faculty of Medicine, Western Sydney University, Blacktown, New South Wales, Australia

⁶Department of Physiology, School of Biomedical Sciences, Faculty of Medicine, University of New South Wales, Sydney, New South Wales, Australia

⁷School of Clinical Medicine, Faculty of Medicine, James Cook University, Townsville, Queensland, Australia

⁸Sunshine Coast University Hospital, Birtinya, Queensland, Australia

⁹School of Medicine, Griffith University, Birtinya, Queensland, Australia

¹⁰Stroke and Ageing Research, Department of Medicine, School of Clinical Sciences at Monash Health, Monash University, Clayton, Victoria, Australia

¹¹Department of Pharmacology, School of Biomedical Sciences, Faculty of Medicine, University of New South Wales, Sydney, New South Wales, Australia

Correspondence

Christian Staehr, Department of Anaesthesiology and Intensive Care, Aarhus University Hospital, Palle Juul-Jensens Boulevard 99, Aarhus N 8200, Denmark.

Email: chst@clin.au.dk

Funding information

Gerhard Brøndstedts Rejselegat; University of the Sunshine Coast SPARK; BRAIN Foundation; Knud Højgaard's Fond; Riisfort Fonden; Danmarks Frie Forskningsfond; Lundbeck Foundation, Grant/Award

Abstract

Aim: In animal models and human cerebral arteries, the changes in endothelial cell (EC)-large conductance calcium-activated potassium channel (BK_{Ca}) distribution, expression, and function were determined in hypoxia and ischemic stroke. The hypothesis that hypoxia and ischemic stroke induce EC-BK_{Ca} in cerebral arteries was examined.

Methods: Immunohistochemistry analyzed BK_{Ca} expression in EC and smooth muscle (SM) of the middle-cerebral artery (MCA) from rat, piglet, and mouse, and pial arteriole of human. Pressure myography with pharmacological intervention

Christian Staehr and Victoria Hinkley joint first authors.

Tim V Murphy and Shaun L Sandow joint last authors.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Acta Physiologica* published by John Wiley & Sons Ltd on behalf of Scandinavian Physiological Society.

Number: R344-2020-952 and R412-2022-449; Danish Cardiovascular Academy, Grant/Award Number: PDC5-2024004-DCA; Dagmar Marshalls Fond; Augustinus Fonden, Grant/Award Number: #21-2471; Helga og Peter Kornings Fond, Grant/Award Number: 472123-006 #55; Independent Research Fund Denmark, Grant/Award Number: 8020-00084B; Novo Nordisk Foundation, Grant/Award Number: NNF20SA0067242

characterized EC-BK_{Ca} and TRPV4 function in rat MCA. Electron microscopy determined caveolae density and vessel properties in rat and mouse MCA.

Results: In rat, pig, and human cerebral vessels, EC-BK_{Ca} was absent in normoxia; present after *chronic* (rat) and *acute* hypoxia (pig), post-ischemic stroke in human vessels, and after endothelin-1-induced stroke in rats. Mouse MCA EC-BK_{Ca} expression increased after *acute* hypoxia. In rat MCA post-hypoxia and stroke, EC and SMC caveolae density increased, with reduced medial thickness, and unchanged diameter. Caveolae and BK_{Ca} did not colocalize. In rat MCA, iberiotoxin (IbTx) potentiated pressure-induced tone in hypoxia/stroke, but not in normoxia. In normoxia, overall MCA tone was unaffected by endothelial removal, but was increased in hypoxia/stroke, where there was no additive effect of endothelial removal and IbTx on tone. Functional TRPV4 was expressed in EC of rat MCA post-stroke.

Conclusions: In post-hypoxia/stroke, but not in normoxia, EC-BK_{Ca} contribute to the regulation of MCA tone. Identifying unique up- and downstream signaling mechanisms associated with EC-BK_{Ca} is a potential therapeutic target to control blood flow post-hypoxia/stroke.

KEYWORDS

BK_{Ca}, blood flow, endothelium, hypoxia, ion channel, stroke

1 | INTRODUCTION

Arterial endothelial cells (EC) regulate vascular tone and blood flow, with small and intermediate conductance calcium-activated potassium channel (S/IK_{Ca}) expression and function being key to their dilator and blood flow activity in health and disease.¹⁻³ In a similar, but distinct manner, intact artery EC-large (B)K_{Ca} expression and function can range from absent to a prominent role, depending on the artery, species, and state; including in disease.⁴ In vascular ECs, the BK_{Ca}α pore-forming and β1-regulatory/auxiliary subunits are often (but not always) reported at a transcriptional mRNA level, and depending on the vascular bed, are not always translated to protein.⁴ This differential expression suggests a potential for rapid EC-BK_{Ca} upregulation in specific beds and states, noting that EC isolation and culture induce BK_{Ca} expression.⁴

While EC-BK_{Ca} are reported in a limited number of intact vascular beds, species, strains, and states, including in disease (Table 1), they are generally absent in control (e.g., normoxia) but can be present/upregulated by hypoxia (48 h, as *chronic*⁵⁻⁸), as it is shown in skeletal muscle arteries of adult Sprague–Dawley (SD) rats.⁵⁻⁸ This upregulation is associated with a hyperpolarized smooth muscle (SM) membrane potential, impaired myogenic tone, and reduced contraction,⁵⁻⁸ and thus increased dilation. In hypoxia, elevated EC-BK_{Ca} may be an adaptive response to disease-related vascular dysfunction. Increased vascular

EC-BK_{Ca} may thus be pertinent in states of reduced tissue oxygenation such as the occlusion of ischemic stroke, myocardial infarction, and pulmonary embolism. In the cerebral circulation, reduced oxygen has significant effects on brain function and anatomy, including core and penumbra development, typical of ischemic stroke.⁹

The distribution and function of specific components of spatially localized EC-signaling microdomains, including aspects of S/I/BK_{Ca} and related calcium store and accessory proteins, can differ between species, beds, and states, including in disease. Such domains occur at non-caveolae and/or caveolae-associated membrane; the latter primarily being omega-shaped invaginations ~80 nm at their widest point.^{10,11} In ECs of intact gracilis muscle artery of chronic hypoxic SD rats, the transient receptor potential vanilloid type 4 (TRPV4) calcium channel can form a functional microdomain with BK_{Ca} and caveolin-1, the caveolae-linked eNOS scaffolding protein.¹² Caveolae can be integral in regulating receptor, channel, and effector function and distribution,^{10,11} including those associated with normal and altered vessel activity in hypoxia and stroke. In artery ECs and SMCs, caveolae may in part facilitate the differential compartmentalization of S/I/BK_{Ca} and TRPV4 channels, with this spatial localization facilitating key aspects of vascular dilator and constrictor signaling mechanism/s.² Multiple factors can modulate BK_{Ca} expression and activity, and these include access to calcium, sensitivity to messengers including IP₃ and

TABLE 1 Large conductance calcium-activated potassium channels (BK_{Ca}) occur in *intact* vessel endothelium of limited species, beds, and states.

| Species/strain/sex (m/f)/age | Vessel type/form | Method/s | Refs. |
|----------------------------------|--------------------------------------------|---------------------------|-------|
| Human | | | |
| n.s. | Internal mammary artery | IHC, myography, WB | [37] |
| n.s. | Internal thoracic artery | IHC, myography, q-PCR, WB | [38] |
| m, 4; f, 4; 67 years | Mesenteric artery, 3rd order, colon cancer | rt-PCR | [18] |
| m, 2, f, 5; 57 years | Absent in non-cancer patients | | |
| m, 22, f, 7; 66 years | Pulmonary artery | Myography | [39] |
| n.s. | Saphenous vein | IHC, q-PCR, myography, WB | [38] |
| Mouse | | | |
| C57BL6 and *KO, male, 8–12 weeks | Coronary arteriole | IHC | [40] |
| Pig | | | |
| n.s. | Renal (conduit) artery | Myography | [41] |
| Rat | | | |
| Wistar, male, n.s. | Left anterior descending coronary artery | Myography | [42] |
| SHR, male, 7 mth | Mesenteric artery, superior, and | Myography, WB | [43] |
| WKY, male, 7 mth | Absent <i>in</i> WKY, superior | Myography, WB | [43] |
| Wistar, n.s. | Mesenteric artery, 1st order | Myography | [44] |
| SD, male, n.s. | Mesenteric artery, 3rd order | Myography | [45] |
| SD, n.s. | Mesenteric artery, 4-5th order | Myography | [46] |

Abbreviations: *KO, knock out, TRP ankyrin and vanilloid type 1; IHC, immunohistochemistry; m/f, male/female; mRNA, via rt-/qPCR; n.s., not stated; SD, Sprague–Dawley; SHR, spontaneously hypertensive rat; WB, Western blot; WKY, Wistar Kyoto.

GTP-binding proteins, and reactive oxygen species,¹³ as well as oxygen tension, shear stress, growth factors, and hormones such as insulin.¹⁴

The significance of EC-BK_{Ca} in the middle cerebral artery (MCA) of mouse, pig, and rat models, and human pial arterioles, was determined with a focus on hypoxia and ischemic stroke. The hypothesis that hypoxia-ischemic stroke induces EC-BK_{Ca} expression in intact MCA and pial arterioles, and that caveolae compartmentalize BK_{Ca}, was examined. To test these proposals, a rat model of *chronic* hypoxia and ischemic stroke was assessed for MCA function, and for BK_{Ca} and TRPV4 expression. Cerebrovascular BK_{Ca} expression was also examined in acute hypoxia in piglet and mouse models, and to provide translational insight, in ischemic stroke patients. The data

have implications for understanding how hypoxia affects cerebrovascular BK_{Ca} activity in EC and SMC function and blood flow.

2 | RESULTS

2.1 | *Chronic* (5 days) hypoxia and ischemic stroke induce BK_{Ca} expression and upregulate TRPV4 in the endothelium of rat distal MCA

Semi-quantitative assessment of fluorescence density demonstrates differential BK_{Ca}α and β1 appearance in ECs and SMCs of rat MCA (Figure 1). In ECs of normoxic




FIGURE 1 Immunohistochemical (IHC) distribution and expression of large conductance calcium-activated potassium ($BK_{Ca\alpha}$ and β_1 subunits) and transient receptor potential vanilloid type 4 channel (TRPV4) in a distal middle-cerebral artery from rats under normoxia, *chronic* (5 days) hypoxia and ischemic stroke conditions. Endothelial cell (EC)- $BK_{Ca\alpha/\beta_1}$ expression is absent in normoxia and upregulated in a diffuse and punctate manner in the cell membrane in chronic hypoxia and ischemic stroke (A, B, respectively). Diffuse and punctate $BK_{Ca\alpha/\beta_1}$ occur on the smooth muscle cell (SMC) membrane of normoxia, hypoxia, and ischemic stroke (A, B, respectively, upper insets; green). Diffuse, intermittent punctate TRPV4 occurs in the ECs and SMCs of normoxia, hypoxia, and ischemic stroke (C; green), with density increased in hypoxia and stroke (C). Cell orientation relative to the longitudinal vessel axis indicates EC/SMC layer patency (A–C, and insets), with EC long axis, left to right. IHC insets (lower central panels in A, B, and far-right semi-quantitative panels in A–C) show staining with secondary antibody only, and with 10-fold peptide excess blocking primary label (A–C, right IHC panels, lower insets). Double arrows, main and inset panels indicate the same vessel region, but different focal planes. Averaged fluorescence density values for tissue stained with secondary antibody only and with corresponding primary antibodies are shown in the bar graphs in the right panel. $n = 6–8$ each for normoxia, and 6 each for hypoxia and stroke, each from different animals. $p < 0.05$, cf. secondary only*, and normoxia[†] cf. hypoxia or stroke. Bar, 25 μm .

MCA, both $BK_{Ca\alpha}$ and β_1 were absent, and their expression was upregulated after hypoxia and ischemic stroke (Figure 1A,B; $p < 0.05$). Conversely, no change in $BK_{Ca\alpha}$ and β_1 expression was observed in the SMCs after hypoxia or ischemic stroke, compared to normoxic MCA (Figure 1A,B).

Semi-quantitative fluorescence density shows increased EC-TRPV4 expression after hypoxia and ischemic stroke, compared to normoxia (Figure 1C). SMC-TRPV4 did not differ between normoxic and hypoxic conditions; although a decrease in stroke compared to matched control occurred (Figure 1C).

2.2 | Ischemic stroke causes differential changes in caveolae density in endothelial cells of distal rat MCA

Conventional TEM determined gross characteristics of MCAs. The diameters of distal MCA were comparable between control and ischemic stroke rats (Table S3). However, after ischemic stroke, distal MCA SMC layers were reduced, associated with a thinner arterial wall compared to control.

The density and colocalization of the prevalent caveolae scaffolding coat protein (caveolin-1) and BK_{Ca} in ECs

and SMCs of the distal rat MCA were determined using TEM and immunoEM (Figure S1; Table S1). The density of luminal and abluminal caveolae in ECs and SMCs was increased in the distal MCA after ischemic stroke compared to control ($p < 0.05$; Table S3). ImmunoEM revealed $BK_{Ca}\alpha$ expression in ECs of the distal MCA after ischemic stroke, in contrast to normoxic conditions where no endothelial $BK_{Ca}\alpha$ was seen (Figure S1).

2.3 | Ischemic stroke induces BK_{Ca} -dependent changes in the myogenic tone of the rat distal MCA

The myogenic tone of rat distal MCAs with intact endothelium at different transmural pressures demonstrated no difference between rats undergoing sham surgery and those with ischemic stroke at any given transmural pressure, for example, at 40, 80, and 120 mmHg (Figure 2A–C). This tone was stable and reproducible during pressure

steps over at least 1 h of experimental duration (not shown) suggesting no time-dependent changes in tone during pharmacological interventions. Incubation (10–20 min) with the BK_{Ca} inhibitor, iberiotoxin (IbTx; 0.1 μ M) at 80 mmHg slightly constricted sham distal MCA (from $65.6 \pm 1.8\%$ to $58.9 \pm 2.2\%$ of maximal diameter, $p = 0.0516$, $n = 6$) and this effect was potentiated for arteries after ischemic stroke (from $62.1 \pm 7.1\%$ to $45.4 \pm 4.7\%$ of maximal diameter, $p = 0.0019$, $n = 4$). Thus, although no difference in MCA tone between stroke and sham was seen prior to IbTx incubation, in the presence of IbTx, the tone of distal MCAs from ischemic stroke rats was higher than in the sham group ($p = 0.0227$, 2-way ANOVA; Figure 2A–C).

Air bubble-induced EC removal reduced the tone of distal MCA from rats in the sham group at 80, but not 40 or 120 mmHg, compared to vessels from the same group with intact endothelium (Figure 2D). Conversely, EC denudation of distal MCAs from ischemic stroke rats increased tone at all three transmural pressures. Studies using bradykinin and 5-hydroxytryptamine confirmed endothelial




FIGURE 2 Pressurized distal middle cerebral artery (MCA) from control-normoxia and hypoxia-stroke rats developed myogenic tone. Representative traces for vascular diameter assessments of MCAs from sham-operated rats (A) and after ischemic stroke (B). The same MCA diameter was assessed under control conditions, in the presence of IbTx, and as fully relaxed in calcium-free solution at 40 mmHg, 80 mmHg, and 120 mmHg, as indicated. The vascular tone (C and D) was calculated as a percent of the active diameter of the fully relaxed diameter of the same artery. There was no difference in the amount of tone between sham and stroke MCA. However, while iberiotoxin (IbTx; BK_{Ca} inhibitor; 0.1 μ M) had no effect on tone in the sham group (control, $n = 6$; IbTx, $n = 4$), it contracted MCA after stroke at 80 and 120 mmHg (C; $n = 4$, each for control and IbTx). Endothelial removal (air bolus) reduced tone in the sham group at 80 mmHg ($n = 6$, control; $n = 4$, endothelium removed) and increased tone at all pressures in stroke rats ($n = 4$; D). After endothelial removal, IbTx continued to have no effect on the tone of MCA from the sham group, but now had no effect on the MCA from stroke rats too ($n = 4$, for all) and on the endothelium-denuded stroke group, in the absence and presence of IbTx (E). *significance of intervention (IbTx or -endothelium) on myogenic tone cf. control conditions, as $p < 0.05$, via two-way ANOVA followed by Tukey's multiple comparisons test.

ablation and subsequent functional integrity of SMCs, respectively (Figure S2).

Following ischemic stroke, distal MCAs with intact endothelium also exhibited greater tone in the presence of IbTx at 120 mmHg compared to the baseline conditions (Figure 2E). This IbTx effect was not observed in distal MCAs in the sham group. Notably, the presence of IbTx did not change vascular tone at different transmural pressures in endothelium-denuded distal MCAs of both sham and ischemic stroke rats (Figure 2E), as there was no additive effect.

2.4 | Functional interaction of TRPV4 and BK_{Ca} in rat distal MCA

The functional contribution of TRPV4 in rat distal MCA and its interaction with BK_{Ca} were assessed (Figure 3; Table S4). Relaxation to the TRPV4 agonist GSK1016790A (0.1–1 mM) exhibited no difference between distal MCAs from sham and ischemic stroke rats (Figure 3A; Table S4). Pre-incubation with the TRPV4 antagonist, HC-067047, did not affect the vascular tone of MCAs from sham and ischemic stroke rats (from $65.7 \pm 1.9\%$ to $64.9 \pm 3.5\%$ and from $62.5 \pm 2.8\%$ to $57.7 \pm 8.6\%$, respectively, $n=4$) but abolished the GSK1016790A-induced vasodilation in both groups (Figure 3A). The GSK1016790A-induced MCA dilation in the sham group was unaffected by IbTx or endothelium denudation (Figure 3B; Table S4). However, after stroke, the GSK1016790A-induced vasodilation was reduced after endothelium denudation (Figure 3C;

Table S4). Moreover, vasodilation was nearly abolished by IbTx in the endothelial-denuded arteries of ischemic stroke rats, suggesting a functional interplay between BK_{Ca} and TRPV4 channels in this state. However, the effect of functional antagonisms, that is, the suppression of GSK1016790A-induced vasodilation because of increased myogenic tone, cannot be excluded.

2.5 | Acute (30–50 min) hypoxia induces BK_{Ca}α expression in the endothelium of piglet distal-central MCA

Semi-quantitative fluorescence data show the absence of BK_{Ca}α in ECs and SMCs of piglet MCAs under normoxic conditions (Figure 4). Following acute (30–50 min) hypoxia, BK_{Ca}α expression was present in ECs and SMCs of the piglet central-distal MCA compared to the normoxic control (Figure 4; $p < 0.05$; for vessel segment location, see Figure S3). Notably, for co-incubated tissue, the SMCs of the adult sow MCA showed BK_{Ca}α expression (Figure 4) as a positive control for the relatively low SMC expression in piglet MCA.

2.6 | Acute (120 min) hypoxia induces BK_{Ca}α expression in the endothelium of mouse distal MCA

Semi-quantitative fluorescence data show BK_{Ca}α in ECs and SMCs of mouse distal MCA under normoxic




FIGURE 3 Distal middle-cerebral artery (MCA) endothelial TRPV4 activity is increased after ischemic stroke. Effect of the TRPV4 activator GSK1016790A on the diameter of rat-isolated, pressurized MCA from sham and stroke rats ($n=9$, each) in the absence and presence of the TRPV4 antagonist HC067047 (0.3 μ M; A; $n=4$, each). Effects on MCA from sham (B; $n=6$) and stroke rats (C; $n=6$) following removal of the endothelium (B and C, $n=6$ and 5, respectively) or in the presence of the BK_{Ca} blocker iberitoxin (IbTx, 0.1 μ M; B,C, $n=6$, each), or removal of the endothelium and IbTx combined (C; $n=4$). Data expressed as % of maximum diameter at 80 mmHg. * $p < 0.05$, two-way ANOVA with Tukey's test. For (A), * indicates $p < 0.05$, as significance of HC067047 effect on response to GSK1016790A cf. MCA from sham and stroke rats, respectively, under control conditions. For (C), * indicates $p < 0.05$, as significance of the effect of endothelium removal (–endothelium) or of a combination of IbTx and endothelium denudation (–endothelium + IbTx) cf. responses of MCA from stroke rats under control conditions. Data compared with two-way ANOVA followed by Tukey's multiple comparisons test. For pEC50 and Emax data, see Table S4.




FIGURE 5 Immunohistochemical distribution and expression of $BK_{Ca}\alpha$ in mouse distal middle-cerebral artery in normoxia-control and *acute* (120 min) hypoxia conditions. Whole-mount vessel endothelium (EC) and smooth muscle cells (SMCs) express diffuse $BK_{Ca}\alpha$ under normoxia (A) and hypoxia (B) conditions (green). Propidium iodide (PI; red) nuclear and cell orientation relative to the longitudinal vessel axis indicates EC/SM layer patency (A, B, insets), with EC long axis, left to right. Single (0.25 μ m) 'slices' show brighter fluorescence at the SMC abluminal surface in hypoxia cf. normoxia (dashed line boxes, cf. A,B); and also at the abluminal over the luminal SMC surface (A, B, cf. upper arrows and lower arrowheads). SMC abluminal and luminal surface fluorescence intensity is \sim 2.7 and 1.6-fold increased in hypoxia cf. normoxia, respectively ($n = 4$; see also Figure S4; Table S5). Boxed regions of SM-EC border/vessel edge (A, B, lower left panels) are enlarged (A, B, upper panels; see also Figure S3). In SMCs, a higher level localization of $BK_{Ca}\alpha$ to cell borders occurs (A, B, lower right panels, e.g., arrows with asterisk), noting that these images are from the SMC-adventitial border region; and hence, some pial-related cell labelling[‡]. Staining with secondary antibody only (A, lower right panel, upper inset; C) and with 10-fold peptide excess blocking primary label (B, lower right panel, upper inset). Double arrows, main and inset panels (A,B) indicate the same vessel region and focal plane, but different label. Averaged fluorescence intensity (C) as in representative images. $n = 7$ and 4, for normoxia and hypoxia, each from a different animal. $p < 0.05$, as significant cf. secondary only*, and normoxia[†] cf. hypoxia EC, respectively. Bar, 25 μ m.

to low (mouse) in the normoxic group. The disparity in MCA EC expression across species in both normoxia and hypoxia may be due to differences in basic signaling pathways, including in the stimuli and local functional MCA environment. Indeed, the mechanism/s for *de novo* EC- BK_{Ca} expression and function may also arise from differences in the model disease state, such as the length of the acute (30–50 and 120 min for pig and mouse, respectively) and chronic (5 days, rat) ischemia protocols, and the idiosyncrasies therein.

In rat post-stroke MCA, the appearance of EC- BK_{Ca} contributes to the regulation of myogenic tone, while in a similar manner, in *in vivo* mouse MCA, dilation increases with hypoxia, 24 h after ischemic stroke.¹⁵ Another study showed that increased endothelial-derived hyperpolarization and nitric oxide pathways underlie amplified vasodilation of isolated rat MCA following asphyxical cardiac arrest.¹⁶ Thus, under hypoxia-related pathological conditions, including ischemic stroke and cardiac arrest, potentiation of cerebrovascular dilation via increased EC- BK_{Ca} expression is consistent with the mechanisms facilitating improved blood flow, at least in specific cerebral arteries and brain regions.

While BK_{Ca} mRNA and protein are predominantly absent in intact artery endothelium,^{4,17} robust data show that select *mouse*, *pig*, and *rat* models and human arteries have EC- BK_{Ca} -mediated function that can reflect BK_{Ca} expression¹⁸ (see also Table 1). In the same species, the present data show that EC- BK_{Ca} can be upregulated in

the cerebral vasculature by hypoxia. These data are consistent with the absence of rat skeletal muscle artery EC- BK_{Ca} in normoxia and its appearance following hypoxia in gracilis and femoral vessels.^{5–8} However, the specific mechanism/s underlying the upregulation of EC- BK_{Ca} are unknown. Indeed, hypoxia may affect several vascular mediators such as reactive oxygen species and protein kinase C-dependent modulation of BK_{Ca} function,¹⁷ altered growth factors,¹⁴ acid–base disturbances,¹⁹ the (transcription) hypoxia-inducible factor (HIF) and related oxygen tension changes,²⁰ as well as changed mechanosensory activity associated with altered blood flow.²¹ While there is significant overlap between these options, it is unknown whether SMC- BK_{Ca} expression and function may or may not reflect that in EC- BK_{Ca} . If this is the case (at least in some states), as for example, in pulmonary artery SMCs, a low oxygen tension environment and related changes in HIF may enhance BK_{Ca} -mediated expression and dilation.²⁰

Although the overall expression and distribution of SMC- BK_{Ca} in MCA differ between rat, piglet, and mouse under normoxic conditions, SMC- $BK_{Ca}\alpha$ and β_1 are unchanged by chronic hypoxia and ischemic stroke in rat or, for $BK_{Ca}\alpha$, by acute hypoxia in mouse. In mouse MCA, abluminal SMC- $BK_{Ca}\alpha$ expression is higher than luminal in both normoxia and hypoxia, as is expression in hypoxia over normoxia when determined separately at the luminal and abluminal sites. Similar to rat MCA EC- BK_{Ca} , this differential SMC- BK_{Ca} expression and distribution imply

a potential for related distinct SMC-BK_{Ca} functional contribution; in this case at the abluminal versus luminal sites; although what activity this may confer is unknown. In contrast to the rat and mouse MCA, overall SMC-BK_{Ca}α expression is increased in the piglet after acute hypoxia; an issue that may relate to the immature developmental state of the piglets compared to the adult rat and mouse.²² Accordingly, in the MCA of adult (sow) pig, SMC-BK_{Ca}α was expressed, and is absent in EC.

While BK_{Ca}α protein is absent in the endothelium of porcine and rabbit aorta, its corresponding mRNA is present.^{23,24} At face value, this suggests an ability to facilitate rapid initiation of BK_{Ca} protein translation in ECs exposed to specific stimuli, such as occurs in disease, including hypoxia. In the present study, data from chronic compared to acute hypoxia in rats versus piglets and mice, respectively, suggest a potential quantitative association between the level of hypoxia and elevated MCA EC-BK_{Ca}α expression (compare EC-MCA BK_{Ca} expression in Figures 1, 4 and 5 for rat, pig(let) and mouse, respectively).

In the present study, the expression and function of both TRPV4 and BK_{Ca} were upregulated in rat MCA after ischemic stroke, although EC-TRPV4 did not directly activate EC-BK_{Ca}. However, the combination of endothelium denudation and BK_{Ca} inhibition had a strong synergistic effect in inhibiting TRPV4-induced MCA dilation after stroke. This synergistic effect has to be further validated; potential functional antagonism of an elevated vascular tone for TRPV4-induced vasodilation cannot be completely excluded. Thus, the TRPV4-BK_{Ca} functional axis may contribute to the increased dilatory action of EC-BK_{Ca} via a secondary mechanism associated with TRPV4 mediating EC-Ca²⁺ influx and/or by generating the Ca²⁺ sparks that activate BK_{Ca}, as occurs in SMCs.²⁵ Interestingly, similarly to rat MCA SMC-BK_{Ca} and SMC-TRPV4 was unchanged by hypoxia, but in contrast, SMC-TRPV4 was reduced after stroke compared to normoxia, suggesting a disparity in the regulation of EC and SMC-TRPV4. The basal role of TRPV4 in mediating MCA EC Ca²⁺ is consistent with that of TRPC3 in facilitating EC-Ca²⁺ influx in rat mesenteric artery²⁶; although their activation of endogenous ligands, such as epoxyeicosatrienoic acids and α₁-adrenergic receptors via noradrenaline,²⁷ remains to be determined.

The importance of caveolae, caveolins, and cavins (the key caveolae scaffolding and structural proteins), as membrane microdomain component responses to hypoxia, altered blood flow, and pressure is well recognized.^{10,28,29} Key factors include caveolae distribution, density, and composition; and transcriptional control of the key components. Several studies suggest that caveolae are a site, and in some cases imply the only site, for S/I/BK_{Ca} location in the ECs and/or SMC membrane in the arterial wall.^{30,31} However, while there may be some association of

caveolae and BK_{Ca}, this is not universal, and immunoEM data in the present study suggest that BK_{Ca} are predominantly present at non-caveolae sites in both EC and SMCs of MCA, as also inferred from the lack of correlation between caveolae and BK_{Ca} density, expression, and distribution. Hence, the present data suggest that EC caveolae density increases after chronic, but not acute, hypoxia, while BK_{Ca} expression increases under both conditions. Notably, channels such as BK_{Ca} and TRPV4 have been suggested to be confined to external regions of caveolae-membrane, interacting with caveolin only when caveolae disassemble (Parton RG, pers. comm). Consistent with this, in *cultured* commercial passage 2–5 bovine aortic EC, the BK_{Ca} membrane current activation occurs exclusively in non-caveolar membrane fractions.³² Of note, BK_{Ca} and caveolin-1 have been suggested to colocalize in the SMCs of cremaster and the ECs of chronic hypoxic gracilis skeletal muscle arteries.^{5,31} However, the confocal and immunoEM resolution used was insufficient to be conclusive on whether such ion channels and other signaling proteins occur *within* or *around* caveolae, or both, with further high-resolution work required to clarify this.

4 | MATERIALS AND METHODS

Detailed methods are available as Supporting Information.

Notably, to ensure that the use of a single experimental model did not bias the experimental findings and facilitate observation of potential similarities and differences in the signaling pathways therein to broaden data relevance, the study was performed in multiple species and experimental models.

5 | LIMITATIONS

Targeting EC-BK_{Ca} in MCA is a potential mechanism to facilitate cerebral blood flow control in specific disease states. However, as SMC-BK_{Ca} expression and function are ubiquitous in adult arteries,^{17,25} differentiating EC- and SMC-BK_{Ca} targets when both are expressed is problematic, and thus control of EC-BK_{Ca} in hypoxia/stroke is likely limited to modulating signaling pathways up- or downstream of its primary BK_{Ca} site of activity. Thus, subsequent studies will clarify the presence of distinct EC versus SMC signaling mechanisms as treatment targets, which include specific and distinct EC and SMC calcium modulation pathways, such as TRP and/or IP₃R subtype activity, where EC-, SMC-, and vascular-bed-specific expression can occur.^{33,34}

An additional limitation relates to the ideal need to further examine the MCA in the acute hypoxia mouse

and pig models in the chronic state to further clarify the progression of EC-BK_{Ca} expression and function, per the present rat data. Indeed, the investigation of hypoxic effects on BK_{Ca}-EC expression and function in pulmonary and cardiac vessels, like that of the MCA, is also an area of interest and limitation, whereby hypoxia-related injury may have critical pathophysiological significance for embolism and infarction, respectively.

A technical limitation of the present data relates to submembranous caveolae numbers, in that these may be overestimated due to the potential for membrane folding making enclosed caveolae-like dimensions (appearing as ~80 nm diameter spheres³⁵); thus, suggesting caution in interpreting that data set. Additionally, it would be ideal to record EC membrane potential in intact arteries, particularly in relation to determining the detailed aspects of artery EC dilator signaling in hypoxia and normoxia. However, unfortunately, the technical requirements for this are beyond the scope of the present study.

6 | CONCLUSIONS

In the present study, both hypoxia and ischemic stroke induce EC-BK_{Ca} expression in the MCA, contributing to the regulation of cerebrovascular myogenic tone to facilitate blood flow as an adaptive mechanism in disease. Modulation of EC-controlled perfusion is an important focus for the development of treatments to improve outcomes after ischemic stroke and other cerebral hypoxia-related disorders.³⁶

7 | THERAPEUTIC IMPLICATIONS

Modulating EC-BK_{Ca} is a potential therapeutic target to control blood flow in hypoxic and ischemic stroke states; albeit, doing so directly is limited due to ubiquitous SMC-BK_{Ca} expression. Hence, further focus on determining EC and SMC-specific up- and downstream pathways related to BK_{Ca} control is a priority to develop therapy to control dilation and thereby mediate blood flow.

AUTHOR CONTRIBUTIONS

Christian Staehr: (*equal); conceptualization; formal analysis; funding acquisition; investigation; methodology; project administration; resources; supervision; validation; visualization; writing—original draft preparation; writing—review and editing. Victoria Hinkley: (*equal); investigation; methodology; project administration; validation; visualization; writing—original draft preparation; writing—review and editing. Vladimir V.

Matchkov: conceptualization; funding acquisition; investigation; methodology; project administration; resources; supervision; validation; writing—original draft preparation; writing—review and editing. Rajkumar Rajanathan: conceptualization; methodology; resources; writing—original draft preparation; writing—review and editing. Line Mathilde B. Hansen: investigation; methodology; resources; writing—original draft preparation; writing—review and editing. Yvonne Eiby: conceptualization; funding acquisition; methodology; resources; supervision; writing—review and editing. Nathan Luque: investigation; methodology; visualization; writing—original draft preparation; Ian Wright: methodology; resources; supervision; writing—review and editing. Stella T. Bjorkman: funding acquisition; methodology; resources; writing—review and editing. Stephanie M. Miller: funding acquisition; methodology; resources; writing—review and editing. Rohan S. Grimley: funding acquisition; writing—review and editing. Andrew Dettrick: funding acquisition; writing—review and editing. Kirat Chand: investigation; methodology; writing—original draft preparation; writing—review and editing. Hong L. Nguyen: conceptualization; investigation; methodology; project administration; writing—original draft preparation; writing—review and editing. Nicole M. Jones: conceptualization; investigation; methodology; project administration; resources; supervision; writing—original draft preparation; writing—review and editing. Tim V. Murphy: (lead; #equal); conceptualization (lead; #equal); formal analysis; funding acquisition; investigation; methodology; project administration; resources; supervision; validation; visualization; writing—original draft preparation; writing—review and editing. Shaun L. Sandow: (lead; #equal); conceptualization; formal analysis; funding acquisition; investigation; methodology; project administration; resources; supervision; validation; visualization; writing—original draft preparation; writing—review and editing.

ACKNOWLEDGMENTS

We thank Rob Parton (University of Queensland; UQ) for enlightening input on caveolae, Matthias Floetenmeyer (UQ) for his efficiency in keeping the TEM running, Maria Garcia (Merck, USA) for the BK β_1 antibody, and Heather McCann (Sydney Brain Bank) for efforts providing human samples.

FUNDING INFORMATION

Work was supported by the Brain Foundation (Australia), the University of the Sunshine Coast SPARK fund to S.L.S.; Lundbeck Foundation R344-2020-952, R412-2022-449, and Independent Research Fund Denmark 8020-00084B to V.V.M.; and Riisfort, Knud Højgaard's, Helga & Peter Korning's (472123-006 #55), Augustinus (#21-2471), and

Dagmar Marshall Foundations, Danish Cardiovascular Academy (PDC5-2024004-DCA; funded by the Novo Nordisk Foundation, grant number NNF20SA0067242, and the Danish Heart Foundation), and Gerhard Brønsted's travel grant to C.S.

CONFLICT OF INTEREST STATEMENT

None.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

PATIENT CONSENT STATEMENT

Human experiments were approved by institutional review boards and participants gave informed consent.

ORCID


Christian Staehr  <https://orcid.org/0000-0001-5431-4409>

Victoria Hinkley  <https://orcid.org/0000-0003-3117-2676>

Vladimir V. Matchkov  <https://orcid.org/0000-0002-3303-1095>

Rajkumar Rajanathan  <https://orcid.org/0000-0002-5002-5244>

Rajkumar Rajanathan  <https://orcid.org/0000-0002-5002-5244>

Line Mathilde B. Hansen  <https://orcid.org/0000-0003-2116-578X>

Line Mathilde B. Hansen  <https://orcid.org/0000-0003-2116-578X>

Yvonne Eiby  <https://orcid.org/0000-0003-0079-7796>

Nathan Luque  <https://orcid.org/0000-0003-3392-566X>

Ian Wright  <https://orcid.org/0000-0001-5337-0134>

Stella T. Bjorkman  <https://orcid.org/0000-0002-6545-3962>


Stella T. Bjorkman  <https://orcid.org/0000-0002-6545-3962>

Stephanie M. Miller  <https://orcid.org/0000-0002-5565-2001>

Stephanie M. Miller  <https://orcid.org/0000-0002-5565-2001>

Rohan S. Grimley  <https://orcid.org/0000-0002-7006-6908>

Rohan S. Grimley  <https://orcid.org/0000-0002-7006-6908>

Andrew Detrick  <https://orcid.org/0000-0003-2403-6027>

Andrew Detrick  <https://orcid.org/0000-0003-2403-6027>

Kirat Chand  <https://orcid.org/0000-0001-7739-2098>

Kirat Chand  <https://orcid.org/0000-0001-7739-2098>

Hong L. Nguyen  <https://orcid.org/0000-0003-1046-4733>

Hong L. Nguyen  <https://orcid.org/0000-0003-1046-4733>


Nicole M. Jones  <https://orcid.org/0000-0001-6738-6022>

Nicole M. Jones  <https://orcid.org/0000-0001-6738-6022>

Tim V. Murphy  <https://orcid.org/0000-0002-0110-9548>

Tim V. Murphy  <https://orcid.org/0000-0002-0110-9548>

Shaun L. Sandow  <https://orcid.org/0000-0002-0753-2742>

Shaun L. Sandow  <https://orcid.org/0000-0002-0753-2742>

REFERENCES

1. Vanhoutte PM, Shimokawa H, Feletou M, Tang EH. Endothelial dysfunction and vascular disease - a 30th anniversary update. *Acta Physiol (Oxford)*. 2017;219:22-96. doi:10.1111/apha.12646
2. Murphy TV, Sandow SL. Agonist-evoked endothelial Ca²⁺ signaling microdomains. *Curr Opin Pharmacol*. 2019;45:8-15.
3. Garland CJ, Dora KA. EDH: endothelium-dependent hyperpolarization and microvascular signalling. *Acta Physiol (Oxford)*. 2017;219:152-161. doi:10.1111/apha.12649
4. Sandow SL, Grayson TH. Limits of isolation and culture: intact vascular endothelium and BKCa. *Am J Physiol Heart Circ Physiol*. 2009;297:H1-H7. doi:10.1152/ajpheart.00042.2009
5. Riddle MA, Hughes JM, Walker BR. Role of caveolin-1 in endothelial BKCa channel regulation of vasoreactivity. *Am J Physiol Cell Physiol*. 2011;301:C1404-C1414. doi:10.1152/ajpcell.00013.2011
6. Riddle MA, Walker BR. Regulation of endothelial BK channels by heme oxygenase-derived carbon monoxide and caveolin-1. *Am J Physiol Cell Physiol*. 2012;303:C92-C101. doi:10.1152/ajpcell.00356.2011
7. Hughes JM, Riddle MA, Paffett ML, Gonzalez Bosc LV, Walker BR. Novel role of endothelial BKCa channels in altered vasoreactivity following hypoxia. *Am J Physiol*. 2010;299:H1439-H1450. doi:10.1152/ajpheart.00124.2010
8. Hughes JM, Walker BR. Role of endothelial large conductance Ca²⁺-activated potassium channels in blunted myogenic responsiveness of small skeletal muscle arteries after 48 hour hypoxic exposure. *FASEB J*. 2006;20:738.733. A1164.
9. Markus R, Reutens DC, Kazui S, et al. Hypoxic tissue in ischaemic stroke: persistence and clinical consequences of spontaneous survival. *Brain*. 2004;127:1427-1436. doi:10.1093/brain/awh162
10. Parton RG, Collins BM. The structure of caveolin finally takes shape. *Sci Adv*. 2022;8:eabq6985. doi:10.1126/sciadv.abq6985
11. Patel HH, Murray F, Insel PA. Caveolae as organizers of pharmacologically relevant signal transduction molecules. *Annu Rev Pharmacol Toxicol*. 2008;48:359-391. doi:10.1146/annurev.pharmtox.48.121506.124841
12. Naik JS, Walker BR. Endothelial-dependent dilation following chronic hypoxia involves TRPV4-mediated activation of endothelial BK channels. *Pflugers Arch*. 2018;470:633-648. doi:10.1007/s00424-018-2112-5
13. Nilius B, Droogmans G. Ion channels and their functional role in vascular endothelium. *Physiol Rev*. 2001;81:1415-1459.
14. Wiecha J, Reineker K, Reitmayer M, et al. Modulation of Ca²⁺-activated K⁺ channels in human vascular cells by insulin and basic fibroblast growth factor. *Growth Hormon IGF Res*. 1998;8:175-181.
15. Staehr C, Giblin JT, Gutiérrez-Jiménez E, et al. Neurovascular uncoupling is linked to microcirculatory dysfunction in regions outside the ischemic core following stroke. *J Am Heart Assoc*. 2023;12:e029527. doi:10.1161/JAHA.123.029527
16. Hansen FB, Esteves GV, Mogensen S, et al. Increased cerebral endothelium-dependent vasodilation in rats in the postcardiac arrest period. *J Appl Physiol*. 2021;131:1311-1327. doi:10.1152/jappphysiol.00373.2021
17. Krishnamoorthy-Natarajan G, Koide M. BK channels in the vascular system. *Int Rev Neurobiol*. 2016;216:401-438.
18. Köhler R, Degenhardt C, Kühn M, Runkel N, Paul M, Hoyer J. Expression and function of endothelial Ca²⁺-activated K⁺ channels in human mesenteric artery. *Circ Res*. 2000;87:496-503. doi:10.1161/01.RES.87.6.496
19. Wen D, Cornelius RJ, Sansom SC. Interacting influence of diuretics and diet on BK channel-regulated K homeostasis. *Curr Opin Pharmacol*. 2014;15:28-32. doi:10.1016/j.coph.2013.11.001
20. Resnik E, Herron J, Fu R, Ivy DD, Cornfield DN. Oxygen tension modulates the expression of pulmonary vascular BKCa channel α - and β -subunits. *Am J Physiol Lung Cell Mol Physiol*. 2006;290:L761-L768. doi:10.1152/ajplung.00283.2005

21. Barvitenko N, Ashrafuzzaman M, Lawen A, et al. Endothelial cell plasma membrane biomechanics mediates effects of pro-inflammatory factors on endothelial mechanosensors: vicious circle formation in atherogenic inflammation. *Membranes*. 2022;12(2):205. doi:[10.3390/membranes12020205](https://doi.org/10.3390/membranes12020205)
22. Shvetsova AA, Gaynullina DK, Tarasova OS, Schubert R. Remodeling of arterial tone regulation in postnatal development: focus on smooth muscle cell potassium channels. *Int J Mol Sci*. 2021;22(11):5413. doi:[10.3390/ijms22115413](https://doi.org/10.3390/ijms22115413)
23. Papassotiropoulos J, Köhler R, Prenen J, et al. Endothelial K⁺ channel lacks the Ca²⁺ sensitivity-regulating β subunit. *FASEB J*. 2000;14:885-894. doi:[10.1096/fasebj.14.7.885](https://doi.org/10.1096/fasebj.14.7.885)
24. Rusko J, Tanzi F, van Breemen C, Adams DJ. Calcium-activated potassium channels in native endothelial cells from rabbit aorta: conductance, Ca²⁺ sensitivity and block. *J Physiol*. 1992;455:601-621.
25. Hill-Eubanks DC, Werner ME, Heppner TJ, Nelson MT. Calcium signaling in smooth muscle. *Cold Spring Harb Perspect Biol*. 2011;3(9):a004549. doi:[10.1101/cshperspect.a004549](https://doi.org/10.1101/cshperspect.a004549)
26. Senadheera S, Kim Y, Grayson TH, et al. Transient receptor potential canonical type 3 channels facilitate endothelium-derived hyperpolarization-mediated resistance artery vasodilator activity. *Cardiovasc Res*. 2012;95(4):439-447. doi:[10.1093/cvr/cvs208](https://doi.org/10.1093/cvr/cvs208)
27. White JP, Cibelli M, Urban L, Nilius B, McGeown JG, Nagy I. TRPV4: molecular conductor of a diverse orchestra. *Physiol Rev*. 2016;96:911-973. doi:[10.1152/physrev.00016.2015](https://doi.org/10.1152/physrev.00016.2015)
28. Grayson TH, Ohms SJ, Brackenbury TD, et al. Vascular microarray profiling in two models of hypertension identifies caveolin-1, Rgs2 and Rgs5 as antihypertensive targets. *BMC Genomics*. 2007;8:404. doi:[10.1186/1471-2164-8-404](https://doi.org/10.1186/1471-2164-8-404)
29. Mathew R. Critical role of caveolin-1 loss/dysfunction in pulmonary hypertension. *Med Sci*. 2021;9:58.
30. Kohler R, Degenhardt C, Kuhn M, Runkel N, Paul M, Hoyer J. Expression and function of endothelial Ca(2+)-activated K(+) channels in human mesenteric artery: a single-cell reverse transcriptase-polymerase chain reaction and electrophysiological study in situ. *Circ Res*. 2000;87:496-503.
31. Howitt L, Grayson TH, Morris MJ, Sandow SL, Murphy TV. Dietary obesity increases NO and inhibits BKCa-mediated, endothelium-dependent dilation in rat cremaster muscle artery: association with caveolins and caveolae. *Am J Phys*. 2012;302:H2426-H2476. doi:[10.1152/ajpheart.00965.2011](https://doi.org/10.1152/ajpheart.00965.2011)
32. Wang XL, Ye D, Peterson TE, et al. Caveolae targeting and regulation of large conductance Ca²⁺-activated K⁺ channels in vascular endothelial cells. *J Biol Chem*. 2005;280:11656-11664.
33. Grayson TH, Haddock RE, Murray TP, Wojcikiewicz RJH, Hill CE. Inositol 1,4,5-triphosphate receptor subtypes are differentially distributed between smooth muscle and endothelial cell layers of rat arteries. *Cell Calcium*. 2004;36:447-458.
34. Walker RL, Hume JR, Horowitz B. Differential expression and alternative splicing of TRP channel genes in smooth muscles. *Am J Physiol Cell Physiol*. 2001;280:C1184-C1192. doi:[10.1152/ajpcell.2001.280.5.C1184](https://doi.org/10.1152/ajpcell.2001.280.5.C1184)
35. Bundgaard M, Hagman P, Crone C. The three-dimensional organization of plasmalemmal vesicular profiles in the endothelium of rat heart capillaries. *Microvasc Res*. 1983;25:358-368. doi:[10.1016/0026-2862\(83\)90025-0](https://doi.org/10.1016/0026-2862(83)90025-0)
36. Andjelkovic AV, Xiang J, Stamatovic SM, et al. Endothelial targets in stroke: translating animal models to human. *Arterioscler Thromb Vasc Biol*. 2019;39:2240-2247. doi:[10.1161/ATVBAHA.119.312816](https://doi.org/10.1161/ATVBAHA.119.312816)
37. Sun WT, Xue HM, Hou HT, et al. Homocysteine alters vasoreactivity of human internal mammary artery by affecting the KCa channel family. *Ann Transl Med*. 2021;9:625-638.
38. Sun WT, Hou HT, Chen HX, et al. Calcium-activated potassium channel family in coronary artery bypass grafts. *J Thorac Cardiovasc Surg*. 2019;161:399-409. doi:[10.1016/j.jtcvs.2019.11.016](https://doi.org/10.1016/j.jtcvs.2019.11.016)
39. Karpinska O, Baranowska-Kuczko M, Malinowska B, et al. Mechanisms of l-alpha-lysophosphatidylinositol-induced relaxation in human pulmonary arteries. *Life Sci*. 2018;192:38-45. doi:[10.1016/j.lfs.2017.11.020](https://doi.org/10.1016/j.lfs.2017.11.020)
40. Sinharoy P, Bratz IN, Sinha S, Showalter LE, Andrei SR, Damron DS. TRPA1 and TRPV1 contribute to propofol-mediated antagonism of U46619-induced constriction in murine coronary arteries. *PLoS One*. 2017;12(6):e0180106. doi:[10.1371/journal.pone.0180106](https://doi.org/10.1371/journal.pone.0180106)
41. Brakemeier S, Eichler I, Knorr A, Fassheber T, Kohler R, Hoyer J. Modulation of Ca²⁺-activated K⁺ channel in renal artery endothelium in situ by nitric oxide and reactive oxygen species. *Kidney Int*. 2003;64:199-207. doi:[10.1046/j.1523-1755.2003.00051.x](https://doi.org/10.1046/j.1523-1755.2003.00051.x)
42. Bang L, Boesgaard S, Nielsen-Kudsk JE, Vejlsstrup NG, Aldershvile J. Nitroglycerin-mediated vasorelaxation is modulated by endothelial calcium-activated potassium channels. *Cardiovasc Res*. 1999;43:772-778. doi:[10.1016/s0008-6363\(99\)00116-9](https://doi.org/10.1016/s0008-6363(99)00116-9)
43. Ando M, Matsumoto T, Kobayashi S, Iguchi M, Taguchi K, Kobayashi T. Differential participation of calcium-activated potassium channel in endothelium-dependent hyperpolarization-type relaxation in superior mesenteric arteries of spontaneously hypertensive rats. *Can J Physiol Pharmacol*. 2018;96:839-844. doi:[10.1139/cjpp-2017-0557](https://doi.org/10.1139/cjpp-2017-0557)
44. Al Suleimani YM, Hiley CR. The GPR55 agonist lysophosphatidylinositol relaxes rat mesenteric resistance artery and induces Ca(2+) release in rat mesenteric artery endothelial cells. *Br J Pharmacol*. 2015;172:3043-3057. doi:[10.1111/bph.13107](https://doi.org/10.1111/bph.13107)
45. Begg M, Mo FM, Offertaler L, et al. G protein-coupled endothelial receptor for atypical cannabinoid ligands modulates a Ca²⁺-dependent K⁺ current. *J Biol Chem*. 2003;278:46188-46194. doi:[10.1074/jbc.M307258200](https://doi.org/10.1074/jbc.M307258200)
46. Naik JS, Osmond JM, Walker BR, Kanagy NL. Hydrogen sulfide-induced vasodilation mediated by endothelial TRPV4 channels. *Am J Physiol Heart Circ Physiol*. 2016;311:H1437-H1444. doi:[10.1152/ajpheart.00465.2016](https://doi.org/10.1152/ajpheart.00465.2016)

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Staehr C, Hinkley V, Matchkov VV, et al. Hypoxia and ischemic stroke modify cerebrovascular tone by upregulating endothelial BK(Ca) channels—Lessons from rat, pig, mouse, and human. *Acta Physiol*. 2025;241:e70030. doi:[10.1111/apha.70030](https://doi.org/10.1111/apha.70030)