

Clinical Study

Development of an internationally agreed national minimum dataset for low back pain: a modified Delphi study

Bayden J. McKenzie, PhD^{a,*}, Giovanni E. Ferreira, PhD^{b,c}, Romi Haas, PhD^a,
Alexandra Gorelik, MSc^a, Chris G. Maher, DMedSc^{b,c},
Rachelle Buchbinder, PhD^a, on behalf of the International Low Back Pain
Minimum Dataset Delphi Panel[#]

^a Musculoskeletal Health and Wiser Health Care Units, School of Public Health and Preventive Medicine, Monash University, Level 4, 553 St Kilda Rd, Melbourne VIC 3004, Australia

^b Institute for Musculoskeletal Health, Sydney Local Health District, Level 10 North, King George V Building, Royal Prince Alfred Hospital, 83/117 Missenden Rd, Camperdown NSW 2050, Australia

^c Sydney School of Public Health, The University of Sydney, Sydney NSW 2006, Australia

Received 9 March 2025; revised 9 September 2025; accepted 7 October 2025

ABSTRACT

BACKGROUND: Low back pain is the leading cause of disability worldwide and the burden is expected to rise due to increases in ageing and population growth. The 2018 *Lancet* Low Back Pain Series proposed urgent actions to reverse the rising trend of disability due to low back pain including the need for a set of data indicators to routinely monitor progress in achieving these actions worldwide.

PURPOSE: To reach international consensus on a national minimum dataset of low back pain indicators that could be used across all countries to monitor progress in improving care and reducing disability from low back pain.

STUDY DESIGN: Modified Delphi study.

SAMPLE: A total of 19 participants attended a preliminary workshop at the 2023 International Back and Neck Pain Forum, The Netherlands. Subsequently, 305 and 339 participants (researchers, clinicians, policy makers, educators, and consumers) completed Delphi surveys Rounds 1 and 2, respectively.

OUTCOME MEASURES: A 9-point Likert scale rated importance and feasibility of low back pain indicators (1 = not important or feasible; 9 = extremely important or feasible, 0 unsure). In Round 2, indicators that achieved consensus on importance and feasibility were ranked “most important” (top-ranked) to “least important.”

FDA device/drug status: Not applicable.

Author disclosures: **BJM:** Grants: Wiser Healthcare - ECR Seeding grant for conference travel (B); Monash University - Study away travel grant (conference) (B); Australia and New Zealand LBP Research Network - travel grant (conference) (A); Fellowship Support: PhD Scholarship - Chiropractic Australia Research Foundation (E); Australia and New Zealand LBP Research Network (D). **GEF:** Research Support (Investigator Salary, Staff/Materials): National Health and Medical Research Council and University of Sydney (G, Paid directly to institution/employer); Grants: National Health and Medical Research Council (I, Paid directly to institution/employer). **RH:** Speaking and/or Teaching Arrangements: Monash University (B); Grants: Arthritis Australia (D, Paid directly to institution/employer). **AG:** Nothing to disclose. **CGM:** Grants: Various Australian government agencies eg NHMRC, MRFF (I, Paid directly to institution/employer); Fellowship Support: NHMRC (I, Paid directly to institution/

employer). **RB:** Royalties: Plantar Fasciitis (B); Grants: Australian National Health and Medical Research Council (NHMRC), Medical Research Future Fund (MRFF), Australian government, HCF Foundation, Arthritis Australia (I, Paid directly to institution/employer); Fellowship Support: Australian National Health and Medical Research Council (NHMRC) (I, Paid directly to institution/employer).

*Corresponding author. Musculoskeletal Health and Wiser Health Care Units, School of Public Health and Preventive Medicine, Monash University, Level 4, 553 St Kilda Rd, Melbourne VIC 3004, Australia.

E-mail address: bayden.mckenzie@monash.edu (B.J. McKenzie).

[#] Names and affiliations of the International Low Back Pain Minimum Dataset Delphi Panel members (n=188) are reported at the end of this article.

METHODS: Workshop participants were divided into four groups and asked to independently consider the importance and feasibility of 105 indicators identified from previous reviews divided into six themes. Participants could remove indicators considered unimportant or not feasible. This required unanimous agreement from independent workshop groups. Participants could also suggest improvements to the wording of indicators, the best unit of measure and additional missing indicators.

RESULTS: Thirty-eight indicators were recommended by workshop participants for inclusion in the Delphi study. Survey responses over two rounds reached consensus for 21 indicators (11 burden, 10 care) ranked from most to least important after reaching consensus on importance and feasibility in Round 1. Number of work days lost and number of opioid prescriptions for low back pain were the highest ranked indicators for burden and care, respectively. Importance rankings were similar across subgroups comparing high-income and low- and middle-income countries, and consumers and nonconsumers.

CONCLUSION: We reached international consensus that 21 indicators could be used to monitor progress in improving care and outcomes for people with low back pain globally. Future work is needed to confirm the acceptability and feasibility of these indicators across countries, and, if implemented, to determine their value over time. © 2025 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>)

Keywords:

Back pain; Benchmarking; Burden of disease; Chronic disease indicators; Dataset; Delphi technique; Health care; Low back pain; Population surveillance; Quality indicators

Introduction

Low back pain is the leading cause of disability worldwide and the burden is expected to rise in coming decades due to increases in ageing and population growth [1]. This will result in even greater healthcare costs, productivity losses and early retirements [2]. The 2018 *Lancet* Low Back Pain Series proposed urgent actions to address this problem, including the need for a set of data indicators to routinely monitor progress in implementing the actions and resulting improvements in back pain burden across the world [3–5].

A national minimum dataset refers to a minimum number of agreed consistent data indicators that could be mandated for routine collection within a country [6]. While collection of health-related data is often vast and complex, national minimum datasets promote streamlined data collection across different health services for benchmarking pertinent data, monitoring trends over time, and guiding health service improvements at a population level [7]. These datasets are routinely used by many national healthcare services [8–10], and have been developed across a range of health fields including nursing [11,12], mental health [13], bio-surveillance [14] and trauma [15].

Clinical care standards for low back pain have been developed in Australia, Canada and the UK, each proposing indicators for quality improvement [16–18]. However, their limited scope highlights the need for an internationally agreed minimum dataset that could serve as a global surveillance tool. Current standards focus on care indicators for low back pain but do not incorporate measures of low back pain burden. The Australian and Canadian clinical care standards monitor acute low back pain care but lack guidance for monitoring care for chronic low back pain

[16,17]. Previous standards are also tailored to specific settings. For example, the Australian standard covers primary care and emergency departments, while the Canadian standard is directed toward primary and community-based care [16,17]. Furthermore, their development for specific national health systems limits their applicability globally.

A national minimum dataset of low back pain indicators could be used by all countries to measure progress in reducing low back pain burden and improving its care. It could also be used to benchmark and make comparisons between countries. Indicators included in the minimum dataset would need to be fit for purpose and feasible to routinely collect in as many countries as possible [19].

As a first step in informing a minimum dataset of indicators, we previously identified 280 indicators that have been used to measure appropriateness of health care for low back pain globally [20]. To inform feasibility, we also performed a scoping review of all low back pain burden and care indicators routinely collected in Australia [21]. We identified 90 burden and 18 care indicators, of which about half are publicly available. Informed by these data, the aim of this study was to reach international consensus on a national minimum dataset that could be used across all countries to monitor progress in improving care and reducing disability from low back pain across all countries.

Methods

Informed by the two previous reviews [20,21], and a preliminary 90-minute workshop at the 18th International Forum for Back and Neck Pain Research in Primary Care in 2023 in Groningen, The Netherlands, we performed a modified international Delphi study. The Delphi method uses a structured approach to achieve consensus among a

group of invited participants (Delphi panel) who are asked to complete a survey over multiple rounds [22]. We used a modified Delphi process to gain consensus as it does not require participants to physically meet, and it provides an equal voice to all participants [23].

Preliminary workshop

There were 19 workshop participants comprised of back and neck pain researchers and clinician-researchers. The aim of the workshop was to obtain participants' views about the importance and feasibility in their country of residence of the low back pain burden and care indicators identified from our prior reviews. After reducing the number of indicators to remove duplication and overlapping indicators, we grouped the final 101 into six themes: intervention (n=31), burden (n=23), healthcare utilization (n=23), work (n=18), self-reported outcomes (n=6) and added an environmental impact indicator [24] (see Appendix A). Three broad indicators for pharmaceutical (n=2) and nonpharmacological (n=1) interventions were also included as there were insufficient indicators relating to these types of low back pain treatments.

To aid discussion participants were divided into four groups. Each group independently considered the themes in different order to ensure that at least two groups considered each theme. Participants were provided with pens and printed lists of indicators arranged by theme. They were invited to suggest improvements to the wording of listed indicators, the best unit of measure for each indicator, and any additional indicators missing from the list. All suggestions by participants were written directly onto the pages of listed indicators. Participants were asked to draw a line through indicators that were considered unimportant or not feasible. Indicators were excluded from the first Delphi survey round when there was unanimous agreement between independent workshop groups that an indicator was unimportant and not feasible to collect. All lists of indicators were collected by investigators at the end of the workshop for analysis.

Modified Delphi

We used a nonprobabilistic purposive and snowball sampling approach to invite participation. We aimed to include at least 160 participants with clinical and research experience in the low back pain field and lived experience with low back pain from both high- and low- and middle-income countries. This target sample size exceeds the recommended sample size for gaining consensus among Delphi panel members with consideration for attrition [25], and ensuring the reliability (replicability) of results [26].

We invited all authors of the 2018 *Lancet* Low Back Pain Series [3–5], the working group of the Australian Low Back Pain Clinical Care Standard [16], and authors of high-quality low back pain clinical practice guidelines identified from a previous study [27]. We also invited registrants

of the 2023 International Forum for Back and Neck Pain Research in Primary Care, members of the Australian and New Zealand Musculoskeletal (ANZMUSC) Clinical Trials Network [28], the Consortium for Low Back Pain in the Low- and Middle-Income Countries [29], the UK National Spine Network [30] and the USA Back Pain Consortium Research Program [31]. The invitees represented a range of professionals from backgrounds including general practice, rheumatology, physiotherapy, occupational therapy, orthopedic and neurosurgery, chiropractic, osteopathy, rehabilitation and pain.

Consumers with a lived experience of low back pain were also invited to participate using internal advertising via special interest groups of the Australian and New Zealand Low Back Pain Research Network (ANZBACK) [32], ANZMUSC [28], Arthritis Australia [33], Arthritis New Zealand [34] and Musculoskeletal Health Australia [35]. Advertisements seeking participants were also posted on social media platform “X” via Cochrane Musculoskeletal [36]. Australian healthcare policy makers identified from government websites in our previous scoping review [21] and networks of coauthors were invited to participate via email.

Invitations to the first Delphi round were sent via email from October 19, 2023 to November 10, 2023. The invitation included an online survey link and Explanatory Statement. A follow-up email was sent to all invitees who did not respond to the initial invite after two and four weeks. Invitations to the second round were sent on 14 December 2023 using the same process. Participants of the first Delphi round were invited to participate in the second round and asked to distribute an online survey link to their networks. Consumer participants were invited to participate in both survey rounds using internal advertising from consumer advocacy groups and social media advertisements. Each online survey link was closed five weeks after the first email was circulated.

Data collection

The survey was conducted using Qualtrics (Qualtrics, Provo, UT) web-based software. For each survey round we asked participants to provide their name, email address and demographic information (age, gender [woman, man, non-binary/gender diverse, gender not listed, prefer not to say], country of residence) and background (researcher, clinician, policy maker, educator [option added for Round 2], consumer and other [specified using a free text box]). For background, participants could indicate all that apply.

Survey Round 1

In Round 1, participants were presented with the low back pain indicators shortlisted from the workshop, presented in random order. For each indicator they were asked to rate 1) its importance for inclusion in a national minimum dataset to monitor the burden and care of people with

low back pain on a 9-point Likert scale where 1 = not at all important to 9 = extremely important; and 2) how feasible it would be to collect this indicator in their country of residence on a 9-point Likert scale where 1 = not feasible to 9 = feasible. If they were unsure of feasibility, they were asked to indicate this with a 0.

A free text box was provided at the end of the survey asking participants to propose additional indicators that were missing in a free text box. Participants were required to respond to all survey items (excluding the free text box).

A priori we considered an indicator important or feasible if it was given a score of 7 to 9, a score of 4 to 6 was deemed unclear, and a score of 1 to 3 was deemed unimportant or not feasible. We determined, *a priori*, that consensus for importance would be reached if at least 60% of respondents rated an indicator as important (≥ 7 on 1–9 Likert). Similar thresholds have been widely used in Delphi studies to determine consensus on survey items, although definitions of consensus vary [37–41]. For feasibility, we set a lower *a priori* threshold and deemed consensus would be reached if at least 30% of respondents rated an indicator as feasible. This cut-off which is lower than standard practice was purposefully chosen to identify indicators that may be feasible in high-income countries and potentially aspirational for middle- and low-income countries [42].

Survey Round 2

Participants in Round 2 were presented with indicators that had reached consensus for importance in Round 1, including the percentage of participants that rated each indicator important and feasible (≥ 7 on 1–9 Likert). They were presented as two lists (burden or care indicators) and in random order to prevent order effects bias. We used a broad definition of burden and care indicators as defined in our recent review [21]. Burden indicators could include self-reported burden, (eg, self-reported measures from population surveys), health services burden (eg, number of people receiving hospital care for low back pain), disease burden (eg, disability-adjusted life-years), financial burden (eg, direct costs for treating low back pain), work disability burden (eg, time off work), medication burden (eg, medications currently being taken by people with low back pain) and time burden (eg, wait times on a waiting list to receive care). Care indicators related to the delivery of health care within relevant settings or facilities (eg, the number and proportion of people who received a specific type of care or health service for low back pain, including any care for work-related injuries). Consistent with our Australian indicators review [21], for indicators that could be classified as both burden and care indicators we classified them as “burden” when the indicator could be used to measure or assess the impact of low back pain on health care services, workplaces or across populations (eg, number of people admitted to hospital with a primary discharge diagnosis of low back pain), and “care” when the indicator could be used to measure specific care people with low back pain

received (eg, number of people who receive lumbar spine decompression surgery).

For each group, participants were asked to rank the indicators from “most important” (top-ranked) to “least important.” We also planned to present any new indicators identified in Survey 1 for rating importance and feasibility as per Survey 1 provided they had not already been excluded at the workshop. If they reached consensus for importance and feasibility, we planned to perform a third round to add the new indicators into the rankings.

We did not revisit rating the feasibility of survey items in Round 2 unless there were new indicators proposed by participants of Round 1. Given the diversity of countries and health system contexts represented, we were concerned that showing aggregated Round 1 feasibility scores could bias responses in Round 2. Specifically, participants might align their ratings with the group mean rather than maintaining their own perspective based on their country’s context.

Data analysis

We present participant demographic data (age group, gender, country of residence and World Bank classification of countries according to income) and background descriptively.

For Survey 1, we present the number (percent) of respondents who rated the low back pain indicators “important,” “unclear” or “unimportant,” and “feasible,” “unclear,” “not feasible,” or “I do not know.” Free text responses were reviewed and inductively coded.

We analyzed survey responses for Round 2 using a weighted rank order scale, where indicators ranked by participants as being “most important” were assigned a higher value than the next ranked indicators in the list (eg, an indicator ranked 1 (most important) was assigned a weight of 10, an indicator ranked 2 (second most important) was weighted 9 etc.). We calculated the total weighted scores of each indicator and present the weighted mean scores (sum of total weighted scores from all participants divided by the number of participant responses) ranked in order of importance within the burden and care lists of indicators. Due to different numbers of indicators within the two theme lists, weighted mean scores were standardized to allow direct comparisons between the groups.

We performed two subgroup analyses: ranked importance of indicators of participants from high-income versus low- and middle-income countries and ranked importance of indicators of participants who identified as being consumers alone versus those who did not. Round 1 feasibility ratings for the two subgroups were also compared. All analyses were performed using Microsoft Excel and Stata 17 (StataCorp LLP, College Station, TX, USA).

Ethics

Ethics approval was obtained from Monash University Human Research Ethics Committee (MUHREC) (Project

ID: 39106). An Explanatory Statement outlining the details of the research project was provided to all potential participants, clearly stating that participating in the workshop or responding to survey rounds was considered consent to participate in the study and having de-identified responses included in the analyses (see Appendix B). Consent was implied by participation in the online Delphi survey(s) and participants could withdraw from the study at any time.

Authorship was offered to all participants who completed both rounds of the Delphi survey and workshop attendees who completed at least one round. All authors were also required to review and approve the final manuscript. All collected data were anonymized and stored on a password protected secure server only accessible by the

investigator team. Data will be retained for five years from the date of publication.

Results

Workshop participants considered 67 indicators were unimportant or not feasible for a national minimum dataset (Fig. 1). After making wording modifications for uniformity, the remaining 35 indicators as well as three additional indicators suggested by workshop participants were included in the first Delphi round (see Appendix C). It was agreed that “number per 100,000 population” was the best unit of measurement to express a scaled proportion of people or events for relevant indicators but for the purpose of the surveys, we used number of people or events.




Fig. 1. Flowchart of the modified Delphi study.

Survey Round 1

Of 680 people invited to participate in the modified Delphi study, 305 participants from 31 countries completed the first survey round and 339 (including an additional 35 consumer participants) participated in the second round (Table 1). Australia (n=157, 51%) was the most represented country, followed by The Netherlands (n=19, 6%), New Zealand (n=18, 6%), United Kingdom (n=15, 5%) and the United States (n=15, 5%). Most participants were from high-income countries (n=269, 88%). Two hundred and thirty-eight (78%) participants were researchers, 190 (62%) were clinicians, 15 (5%) were educators, 11 (4%) were policy makers and 60 (20%) had lived experience of low back pain. 176 (58%) reported dual roles (eg, researcher and clinician), while 21 (7%) indicated they were consumers only.

The results of Survey Round 1 are graphically reported as Figs. 2 and 3 and tabulated in detail as Appendices D and E, respectively. Twenty-three (61%) of the 38 indicators included in Survey Round 1, reached consensus for both importance and feasibility. We elected to remove two indicators: one was deemed unimportant and excluded by participants during the preliminary workshop but was erroneously included in the survey (number of people with low back pain who report restriction in activities of daily living [ADLs]), and the other indicator (number of emergency department attendances for a recorded primary discharge diagnosis of low back pain) was considered to be a similar measure to another indicator included in Round 1 (number of hospital admissions for a primary discharge diagnosis of low back pain). There was no consensus on importance of 14 indicators and one indicator reached consensus for importance but not feasibility. There were an additional 103 indicators suggested by survey participants as free text, but all were excluded as they had been considered and discarded at the preliminary workshop.

The 13 burden and 10 care indicators that reached consensus for both importance and feasibility are shown in Table 2. Eight burden indicators related to work days lost or workers' compensation claims for low back pain, two related to hospital admissions or emergency department attendances, and the remaining indicators related to restriction in ADLs (n=1), people who report feeling confident self-managing their low back pain (n=1), and hospital outpatient clinic attendances (n=1). Care indicators related to imaging of the lumbar spine (n=3), pharmaceutical prescriptions (n=3), specialist referrals (n=2) and surgical treatment (n=2), hospital admission (n=1), hospital outpatient care (n=1) and healthcare costs of approved workers' compensation claims (n=1). Feasibility ratings of burden indicators that reached consensus varied from 30% to 73%, which relate to the number of people who report taking time off work due to low back pain and the number of hospital admissions with a primary discharge diagnosis of low back pain, respectively. The care indicators rated most

Table 1
Characteristics of Delphi participants

Characteristics	Round 1 (n=305) N (%)	Round 2 (n=339) N (%)
Gender		
Woman	137 (44.9)	171 (50.4)
Man	167 (54.8)	165 (48.7)
Nonbinary or gender not listed	1 (0.3)	3 (0.8)
Background (multiple responses were permitted)		
Researcher	238 (78.0)	236 (69.6)
Clinician	190 (62.3)	190 (56.1)
Policy maker	11 (3.6)	14 (4.1)
Educator	15 (4.9)	119 (35.1)
Consumer	60 (19.7)	90 (26.6)
(person with lived experience of low back pain)		
Consumer only	21 (6.9)	56 (15.9)
Other	6 (2.0)*	5 (1.5)*
Country of residence with 5 or more participants		
Australia	157 (51.5)	187 (55.2)
The Netherlands	19 (6.2)	24 (7.1)
New Zealand	18 (5.9)	21 (6.2)
United Kingdom	15 (4.9)	17 (5.0)
United States of America	15 (4.9)	16 (4.7)
Brazil	13 (4.3)	12 (3.5)
Denmark	8 (2.6)	8 (2.4)
Norway	8 (2.6)	7 (2.1)
Nepal	8 (2.6)	6 (1.8)
Canada	6 (2.0)	5 (1.5)
Switzerland	5 (1.6)	5 (1.5)
Belgium	5 (1.6)	4 (1.2)
Other†	28 (9)	27 (8)
World Bank classification of economies of participants' country of residence		
High-income countries	269 (88.2)	304 (89.7)
Low- and middle-income countries	36 (11.8)	35 (10.3)
Age group (y)		
18–29	14 (4.6)	14 (4.1)
30–39	76 (24.9)	79 (23.3)
40–49	88 (28.9)	88 (26.0)
50–59	69 (22.6)	80 (23.6)
60+	57 (18.7)	77 (22.7)
Not reported	1 (0.3)	1 (0.3)

* Other in Round 1: consumer director (n=1), hospital advisory committee member (n=1), manager (n=2), patient experience or usability professional (n=1) and practice consultant (n=1). Other in Round 2: consumer director (n=1), clinical advisor (n=1), clinical manager (n=1), dental assistant (n=1) and regulatory body member (n=1).

† Other countries: France (Round 1 n=3; Round 2 n=2), South Africa (R1 n=2; R2 n=4), India (R1 n=2; R2 n=3), Italy (n=2 both rounds), Finland (R1 n=2; R2 n=1), Sweden (R1 n=2; R2 n=1), Bangladesh (n=1 both rounds), Colombia (n=1 both rounds), Indonesia (n=1 both rounds), Ireland (n=1 both rounds), Israel (n=1 both rounds), Lebanon (n=1 both rounds), Mexico (n=1 both rounds), Nigeria (n=1 both rounds), Pakistan (n=1 both rounds), Portugal (n=1 both rounds), Tanzania (n=1 both rounds), Ethiopia (R1 n=1), Saudi Arabia (R1 n=1), Bahrain (R2 n=1), Burundi (R2 n=1), country not specified (R1 n=2; R2 n=1).

feasible to collect related to the number of lumbar fusion surgeries (78%) and lumbar decompression and fusion surgeries (77%). The number of referrals to medical specialists for low back pain was rated by participants as the least feasible care indicator to collect (47%).




Fig. 2. Percentage of respondents that indicated items were important, unclear and or not important in Survey Round 1, n=305. Vertical line represents the *a priori* consensus threshold for importance. LBP, low back pain; ADLs, activities of daily living; MRIs, magnetic resonance imaging scans; CT, computed tomography.




Fig. 3. Percentage of respondents that indicated items were feasible, unclear or not feasible or don't know in their countries Survey Round 1, n=305. Vertical line represents the *a priori* consensus threshold for feasibility. LBP, low back pain; MRIs, magnetic resonance imaging scans; CT, computed tomography; ADLs, activities of daily living.

Table 2

Indicators that met consensus criteria for importance* and feasibility† in Round 1

Indicator	Number (%) that rated indicators important	Number (%) that rated indicators feasible
Burden indicators (N=13)		
Number of work days lost because of low back pain (LBP)	273 (90)	109 (36)
Number of people who are permanently unable to work due to LBP	272 (89)	142 (47)
Number of people with low back pain who report restriction in activities of daily living	269 (88)	117 (38)
Number of people who report taking time off work due to LBP	251 (82)	93 (30)
Number of hospital admissions for a primary discharge diagnosis of LBP‡	219 (72)	224 (73)
Number of people who report feeling confident about self-managing LBP	212 (70)	102 (33)
Total healthcare cost of approved workers' compensation claims for LBP‡	205 (67)	122 (40)
Number of emergency department attendances for a primary discharge diagnosis of LBP	203 (67)	209 (69)
Number of hospital outpatient clinic attendances for people with LBP‡	199 (65)	179 (59)
Duration of compensated time off work for approved workers' compensation claims for LBP	199 (65)	152 (50)
Workers' compensation claimants for LBP who return to their preinjury status at 12 mo	193 (63)	125 (41)
Total direct cost of approved workers' compensation claims for LBP	189 (62)	143 (47)
Workers' compensation claimants for LBP who return to their preinjury status at 3 mo	185 (61)	130 (43)
Care indicators (N=10)		
Number of opioid prescriptions for LBP	251 (82)	166 (54)
Number of referrals to surgical specialists for LBP	221 (72)	150 (49)
Number of referrals to medical specialists for LBP	215 (70)	143 (47)
Number of lumbar spine MRIs	210 (69)	218 (71)
Number of lumbar spine X-rays	200 (66)	209 (69)
Number of antidepressant prescriptions for LBP	193 (63)	147 (48)
Number of lumbar decompression and fusion surgeries	191 (63)	236 (77)
Number of lumbar spine CT scans	191 (63)	218 (71)
Number of muscle relaxant prescriptions for LBP	185 (61)	151 (50)
Number of lumbar fusion surgeries	183 (60)	238 (78)

* At least 60% of respondents rated indicator as important (≥ 7 on 1–9 Likert).† At least 30% of respondents rated an indicator as feasible (≥ 7 on 1–9 Likert).

‡ Indicators could also be considered care indicators.

Survey Round 2

Of 745 people invited to complete survey Round 2, 339 from 31 countries completed survey Round 2. They had similar characteristics to participants in Round 1 (see Table 1). Twenty-one of the 23 indicators that reached consensus in Round 1 (11 burden, 10 care indicators) were presented for importance ranking in Round 2, and two were discarded for reasons outlined in Survey Round 1.

Table 3 presents the overall importance rank and weighted mean importance scores of the low back pain burden indicators from the Round 2 survey, as well as the importance rank, weighted mean importance scores and number (%) participants that considered indicators were feasible in their country stratified by country income level and participant category (not consumers or consumers). Number of work days lost because of low back pain was the highest ranked burden indicator (weighted mean rank score 7.66) followed by number of people who are permanently unable to work due to low back pain (weighted mean rank score 7.18), number of people who report taking time off work due to low back pain (weighted mean rank score 6.32), number of hospital admissions for a primary discharge diagnosis of low back pain (weighted mean rank score 6.16) and number of hospital outpatient clinic attendances for people with low back pain (weighted mean ranked score 5.7).

Importance rankings were similar across subgroups. Participants from low- and middle-income countries ranked the same five indicators in their top 5 as participants from high-income countries as did both consumers and nonconsumers, although there were slight variations in the order. There were also differences in feasibility rankings between high- and low- and middle-income country participants. For example, only 19% of participants from low- and middle-income countries considered the number of work days lost because of low back pain to be a feasible indicator to collect in their countries compared with 38% of participants from high-income countries.

Table 4 presents the results of Round 2 for the low back pain care indicators. The top indicator was number of opioid prescriptions for low back pain (weighted mean score 7.19) followed by number of medical specialist referrals (weighted mean score 6.38), number of lumbar spine MRIs (weighted mean score 6.11) and X-rays (weighted mean score 5.93) and number of surgical referrals (weighted mean score 5.77). Participants from low- and middle-income countries ranked the same 5 care indicators in their top 5 as participants from high-income countries but the order varied. Medical specialist referrals were the most important care indicator nominated by low- and middle-income country participants followed by lumbar spine X-rays and surgical referrals. Consumer and nonconsumer participants had different importance rankings. For

Table 3

Overall importance rank and weighted mean importance scores of low back pain burden indicators in Round 2 and importance rank, weighted mean importance scores and number (%) participants that considered indicator was feasible in their country stratified by country income level and participant category (not consumers or consumers)

Burden indicators	All participants (n=339)		Participants from HICs (n=304)			Participants from LMICs (n=35)			Not consumers (n=283)			Consumers (n=56)		
	Importance rank	Weighted mean score [†]	Importance rank	Weighted mean score [†]	N (%) that rated indicator feasible* (n=269)	Importance rank	Weighted mean score	N (%) that rated indicator feasible* (n=36)	Importance rank	Weighted mean score [†]	N (%) that rated indicator feasible* (n=284)	Importance rank	Weighted mean score [†]	N (%) that rated indicator feasible* (n=21)
Number of work days lost because of low back pain	1	7.66	1	7.7	102 (38)	1	7.37	7 (19)	1	7.69	104 (37)	1	7.54	5 (24)
Number of people who are permanently unable to work due to low back pain	2	7.18	2	7.25	129 (48)	3	6.57	13 (36)	2	7.25	134 (47)	4	6.86	8 (38)
Number of people who report taking time off work due to low back pain	3	6.32	3	6.36	85 (32)	5	5.87	8 (22)	4	6.13	86 (30)	2	7.25	7 (33)
Number of hospital admissions for a primary discharge diagnosis of low back pain	4	6.16	4	6.18	205 (76)	4	6	19 (53)	3	6.14	212 (75)	5	6.32	12 (57)
Number of hospital outpatient clinic attendances for people with low back pain (eg, consultation with a specialist, physiotherapist etc.)	5	5.7	5	5.58	157 (58)	2	6.67	22 (61)	5	5.46	168 (59)	3	6.88	11 (52)
Total healthcare cost of approved workers' compensation claims for low back pain (injury)	6	5.11	6	5.12	114 (42)	7	5.06	8 (22)	6	5.28	111 (39)	7	4.24	11 (52)
Number of people who report feeling confident about self-managing their low back pain	7	4.87	7	4.88	92 (34)	8	4.85	10 (28)	8	4.75	96 (34)	6	5.54	6 (28)
Total direct cost of approved workers' compensation claims for low back pain (injury)	8	4.69	8	4.64	136 (51)	6	5.19	7 (19)	7	4.9	131 (46)	11	3.64	12 (57)
Duration of compensated time off work for approved workers' compensation claims for low back pain (injury)	9	4.33	9	4.26	143 (53)	9	4.83	9 (25)	9	4.45	142 (50)	10	3.7	10 (48)
Number of workers' compensation claimants for low back pain (injury) who return to their preinjury status at 3 mo	10	4.09	10	4.09	125 (46)	10	4.05	5 (14)	10	4.08	120 (42)	8	4.11	10 (48)
Number of workers' compensation claimants for low back pain (injury) who return to their preinjury status at 12 mo	11	3.88	11	3.93	121 (45)	11	3.51	4 (11)	11	3.87	116 (41)	9	3.93	9 (43)

HICs, high-income countries; LMICs, low- and middle-income countries.

* Feasible if scored ≥ 7 on 1–9 Likert scale in Round 1 (n=305).

[†] Weighted mean scores for the 11 burden indicators were standardized to a 10-item equivalent weighting to allow comparison with the weighted mean scores of the 10 care indicators.

Table 4
Overall importance rank and weighted mean importance scores of low back pain care indicators in Round 2 and importance rank, weighted mean importance scores and number (%) participants that considered indicator was feasible in their country stratified by country income level and participant category (not consumers or consumers)

Care indicators	All participants (n=339)			Participants from HICs (n=304)			Participants from LMICs (n=35)			Not consumers (n=283)			Consumers (n=56)		
	Importance rank	Weighted mean score		Importance rank	Weighted mean score	N (%) of participants rated as feasible to collect*	Importance rank	Weighted mean score	N (%) of participants rated as feasible to collect*	Importance rank	Weighted mean score	N (%) of participants rated as feasible to collect*	Importance rank	Weighted mean score	N (%) of participants rated as feasible to collect*
Number of opioid prescriptions for low back pain	1	7.19	1	1	7.33	150 (56)	5	5.94	16 (44)	1	7.52	154 (54)	6	5.5	12 (57)
Number of referrals to medical specialists for low back pain	2	6.38	2	2	6.19	133 (49)	1	8.09	10 (28)	2	6.23	132 (46)	1	7.16	11 (52)
Number of lumbar spine MRIs	3	6.11	3	3	6.12	194 (72)	4	6.06	24 (67)	3	6.07	204 (72)	3	6.29	14 (67)
Number of lumbar spine X-rays	4	5.93	4	4	5.88	185 (69)	2	6.34	21 (67)	4	5.84	197 (69)	5	6.18	12 (57)
Number of referrals to surgical specialists for low back pain	5	5.77	5	5	5.71	139 (52)	3	6.29	11 (31)	5	5.84	139 (49)	7	5.41	11 (52)
Number of lumbar spine CT scans	6	5.27	6	6	5.37	195 (72)	8	4.49	23 (64)	6	5.08	204 (72)	4	6.27	14 (67)
Number of lumbar decompression and fusion surgeries	7	4.71	7	7	4.77	215 (80)	9	4.14	21 (58)	7	4.91	221 (78)	9	3.68	15 (71)
Number of muscle relaxant prescriptions for low back pain	8	4.7	8	8	4.57	139 (52)	6	5.8	12 (33)	10	4.38	140 (49)	2	6.3	11 (52)
Number of antidepressant prescriptions for low back pain	9	4.5	9	10	4.49	135 (50)	7	4.6	12 (33)	9	4.41	138 (49)	8	4.96	9 (43)
Number of lumbar fusion surgeries	10	4.4	10	8	4.59	217 (81)	10	3.26	21 (58)	8	4.69	223 (79)	10	3.25	15 (71)

HICs, high-income countries; LMICs, low- and middle-income countries.

* Feasible if scored ≥ 7 on 1–9 Likert scale in Round 1 (n=305).

consumers, medical specialist referral was the most important care indicator followed by number of muscle relaxant prescriptions, lumbar spine MRIs, CTs and X-rays. Number of opioid prescriptions was the top ranked indicator for both high-income country participants and nonconsumers, it was ranked of less importance by both low- and middle-income country participants and consumers (fifth and sixth, respectively).

Discussion

We identified 21 indicators that may be suitable for a national minimum dataset to monitor progress in reducing low back pain burden and improving its care. Overall, the top ranked burden indicators were number of work days lost or number of people unable to work due to low back pain, and the number of hospital admissions and hospital outpatient clinic attendances. The top ranked care indicators were number of opioid prescriptions, medical and surgical specialist referrals and lumbar spine imaging. The burden indicator results were similar across subgroups although the rank order and feasibility scores varied. There was greater variability in the importance ranking and scores between high-income and low- and middle-income country participants and between consumers and nonconsumers for the care indicators. Feasibility results also varied across subgroups.

While we aimed to develop a national minimum dataset of indicators, not all the 21 ranked indicators that reached consensus may be feasible or necessary in less resourced countries. For example, it may not be necessary to collect all three lumbar spine imaging indicators (X-ray, CT and MRI) but countries could choose the imaging modality of most relevance to their setting. Additionally, countries without a workers' compensation insurance scheme may not be able to include work-related indicators but could include other proxy indicators sourced from other data.

The different importance rankings for opioid prescriptions for low back pain given by high- and low- and middle-income country participants may reflect lower rates of opioid prescriptions in low- and middle-income countries [43]. Similarly, its lower importance ranking among consumers compared with other participants may have several explanations. Consumers may be less aware of the risk-benefit profile of opioids for low back pain and some consumers have also pushed back against greater opioid regulation [44,45].

Clinical care standards for low back pain published in Australia, Canada, and the United Kingdom have proposed indicators to support the improvement of health care through monitoring various aspects of care over time [16–18]. Of the 21 shortlisted indicators ranked on importance in our Delphi study, 7 (33%) were recommended for routine collection by at least one country. These include percentage of people who receive spinal imaging for low back pain [16–18], percentage of people who are given opioids for low

back pain [16–18], percentage of people referred to a medical specialist or provider for low back pain [17], percentage of people who report feeling confident about self-managing their low back pain [17] and number of work days lost for people with acute low back pain [17]. Our review of low back pain indicators in Australia identified five of these indicators (number of people referred for X-rays/CT/MRI, number of referrals to medical specialists and number of work days lost because of low back pain) that are routinely collected at national and state or territory levels [21]. While the Australian Commission on Safety and Quality in Health Care (ACSQHC) has proposed 12 quality indicators for determining whether the quality of low back pain care improves over time, there is a need for accessible indicators to be revisited regularly.

The Canadian clinical care standard recommends using national data sources for 2 indicators shortlisted in this Delphi study: (1) the percentage of people who receive diagnostic imaging for low back pain [46,47] and (2) the percentage of people who are given opioids for low back pain [46–48]. The majority (94%, $n=29$) of indicators from the Canadian and UK clinical care standards suggest that indicators be collected from “local sources” (eg, service protocols and patient notes), however, it is unclear whether existing low back pain indicators are routinely collected in these jurisdictions [17,18].

The feasibility of collecting several of our shortlisted indicators was demonstrated in a recent study from Belgium [49]. Using a registry of general practice data, the authors established statistical benchmarks of four care indicators, including five indicators ranked in Round 2: 3 imaging indicators, 1 opioid prescriptions indicator and 1 prescription of medical leave indicator [49].

We have developed a short list of low back pain indicators that countries worldwide could choose to include in a national minimum dataset to monitor progress in reducing the burden of low back pain and improving its care over time. Previous consensus efforts have focused primarily on patient outcomes for use in primary care and clinical research rather than for benchmarking at a population level. For example, the US National Institutes of Health (NIH) Task Force on Research Standards developed a minimum dataset of self-reported chronic low back pain indicators to promote standardization across grant applications, clinical studies and epidemiological studies [50]. Similarly, Verbung and colleagues used a modified RAND-UCLA Delphi approach to develop a set of low back pain outcome measures for monitoring aspects of primary care such as shared-decision making and internal quality improvement [51], and Clement and colleagues conducted a six-round Delphi process to recommend a standardized set of outcome measures for supporting quality improvement in degenerative low back conditions [52]. In contrast, our indicators are intended to support consistent monitoring and benchmarking at a population level.

Strengths and limitations

Our study has several strengths. Our international Delphi panel included low back pain researchers, clinicians, policy makers, educators and people with lived experience of low back pain from 33 countries, including 14 low- and middle-income countries. While there is little guidance for what expertise qualifies people to participate in Delphi studies [23], our large sample size included people with knowledge and interest in low back pain across a range of settings. Our combined purposive and snowballing approach led to over 300 participants in each survey round, with consistent response rates of approximately 45% in Rounds 1 and 2 and people with lived experience of low back pain were well represented. The modified-Delphi approach that we used, including a preliminary workshop to refine an extensive list of indicators, was consistent with studies in other health fields and could be used to develop burden and care indicators for other health conditions [53,54].

Our study has some limitations. First, most indicators included in Round 1 were derived from studies in high-income countries and may not be applicable to other settings. Second, participants rated the feasibility of indicator collection in their own country, which may not accurately reflect their true feasibility. Third, we removed two indicators that met consensus thresholds in Round 1 and this may not have been appropriate. One (number of people with low back pain who report restriction in ADLs) was excluded during the preliminary workshop as participants considered it unimportant for use in a national minimum dataset, but it was erroneously included in the first survey round. The other (number of emergency department attendances for a recorded primary discharge diagnosis of low back pain) was initially viewed as being similar to the number of hospital admissions but on reflection, captures a distinct aspect of hospital system burden. We recommend that both indicators be reconsidered in future work. Additionally, other pertinent indicators for monitoring low back pain burden and care may also have been excluded during the preliminary workshop phase of this study.

Another limitation of this study is that only a small proportion of participants in the Delphi survey rounds were from low- and middle-income countries (12% and 10% of participants in Rounds 1 and 2, respectively), which may affect the applicability of the 21 shortlisted indicators in these countries. This imbalance likely reflects the larger pool of researchers, researcher-clinicians, clinicians, policy makers and consumers engaged in back pain research in high-income countries. To address this, we invited all potential participants we could identify from low- and middle-income countries, including members of the Consortium for Low Back Pain in the Low- and Middle-Income Countries [29], and used snowball sampling to increase participation further. Assessing the relevance of the shortlisted indicators in low- and middle-income countries will form

part of the next phase of work where the indicators will be refined and piloted internationally.

Implications and future work

A national minimum dataset of low back pain indicators is urgently needed to monitor care and outcomes, and this surveillance tool could have the potential for international adoption. Routine collection of some or all 21 indicators could serve as an effective benchmarking tool and for monitoring changes over time. Future work is required to understand the acceptability and feasibility of collecting a national minimum dataset of indicators that can be used by all countries. Leveraging the indicators shortlisted from this study, the next step will be to refine the dataset by defining a core set of essential indicators that all countries should collect, while allowing flexibility for additional country-specific indicators. This will require collaborative discussion between researchers, data custodians, health care providers, policy makers and consumers within countries that are actively seeking to reduce the burden of low back pain and improve its care as a national priority. These discussions should focus on determining which indicators are feasible for routine collection, how data can be standardized across different health systems and how best to integrate these indicators into national health monitoring frameworks. Additionally, the expansion of future work to include real-world feasibility studies like the study in Belgium that evaluated low back pain indicators in general practice settings [49], would also be valuable for determining indicators that can be practically implemented on a larger scale.

Conclusion

We developed a minimum dataset of 21 low back pain indicators that could be used to monitor progress in improving care and outcomes for people with low back pain globally. Future work is needed to confirm the acceptability and feasibility of these indicators across countries, and, if implemented, to determine their value over time.

International low back pain minimum dataset Delphi panel

Ilana N. Ackerman¹, Bronwen Jane Ackermann², Andrea Martina Aegerter^{3,4}, Carlo Ammendolia^{5,6}, Toni Andary^{7,8}, Iben Axen⁹, Habibu Salisu Badamasi¹⁰, Florian Bailly¹¹, Claire Barrett^{12,13}, Paul F. Beattie¹⁴, Danielle Barbosa Benfica¹⁵, Danielle Berkovic¹, Belinda R. Beck^{16,17}, Noa Ben Ami¹⁸, Amber M. Beynon¹⁹, Mark D. Bishop²⁰, Lars Christian Haugli Bråten²¹, Andrew M. Briggs¹⁹, Cristy Brooks^{22,23,24}, Benjamin T. Brown²⁵, Paul Buntine^{26,27}, Jose A. Calvache²⁸, Gary Cantwell²⁹, Joel Carmichael^{30,31}, Katy Carnachan, Aidan G. Cashin^{32,33}, Romain Champagne³⁴, Wei-Ju Chang^{33,32}, Bo-Jhen Chen³⁵, Guillaume Christe³⁶, Natalie C. Clohesy, Mary-Lynne Cochrane,

Michel W. Coppieters^{16,37}, Leticia Amaral Correa²⁵, Nathalia Costa³⁸, Amanda J. Cross³⁹, Ben Darlow⁴⁰, Anita Davis, Katie de Luca⁴¹, Marcos de Noronha⁴², Nils-Bo de Vos Andersen⁴³, Kim Delbaere^{44,33}, Michael Di Donato⁴⁵, Chris Dickson⁴⁶, Clermont E. Dionne⁴⁷, Aron Simon Downie²⁵, Kieran E. Fallon⁴⁸, Matthew Fernandez⁴¹, Carol Ann Flavell⁴⁹, Simon D. French²⁵, Julie M. Fritz⁵⁰, Fernanda Bono Fukushima⁵¹, G. Shankar Ganesh⁵², Tania Gardner⁵³, Stéphane Genevay⁵⁴, Tiffany K. Gill⁵⁵, Rosemary Giuriano²⁵, Christine Goertz⁵⁶, Luis Antunes Gomes^{57,58}, Rebecca Grainger⁵⁹, Fredrik Granviken^{60,61}, Jim Greenwood⁶², Søren Grøn⁶³, Douglas P. Gross⁶⁴, Amitabh Gupta²², Travis Haber⁶⁵, Melanie Hamilton^{66,67}, Mark J. Hancock⁶⁸, Ian A. Harris⁶⁹, Jan Hartvigsen^{63,70}, A.T.M. Tanveer Hasan⁷¹, Jill A. Hayden⁷², Jeffrey J. Hebert^{73,74}, Bibi Dige Heiberg⁷⁵, Lise Hestbaek^{70,63}, Cesar A. Hincapié^{76,77,78,79}, Johan Holte⁸⁰, Ben Hudson⁸¹, Aminu Alhassan Ibrahim⁸², Wilhelmina IJzelenberg⁸³, Hazel J. Jenkins²⁵, Fabianna Resende de Jesus-Moraleida⁸⁴, Renea V. Johnston¹, Pascale Jonckheer⁸⁵, Matthew D. Jones^{33,32}, Steven J. Kamper^{86,87}, Steve Karas⁸⁸, Jaro Karppinen^{89,90}, Gerard Keown, Eugene Khoo⁹¹, Bart Koes⁹², P. Paul F.M. Kuijer⁹³, Richard Kwiatek⁹⁴, Andrew Lawson^{33,95}, Italo Ribeiro Lemes^{96,97}, Trevor A. Lentz^{56,98}, Chung-Wei Christine Lin^{66,67}, Esther T. Maas⁸³, Rajiv Maharjan⁹⁹, Genevieve Mahieu¹⁰⁰, Peter Malliaras¹⁰¹, Ramakrishnan Mani^{102,103}, Paul W. Marshall^{104,105}, Denis Martin^{106,107}, Stephanie Mathieson^{67,108}, Suzanne McIlroy^{109,110}, Annie (Annette) McPherson^{29,111}, Rebecca Mellor^{112,113}, Ney Meziat-Filho^{114,115}, Sarah Mingels¹¹⁶, Gisela Cristiane Miyamoto¹¹⁷, Allan Richard Molloy^{118,119}, Laura R.C. Montgomery⁵³, Javier Muñoz Laguna^{76,77,78}, Leigh E. Murrant⁶⁷, Charbel Najem¹²⁰, Josh Naunton^{42,121}, Tania Inés Nava-Bringas¹²², Christopher Neason²⁷, Dennis R. Neuen^{123,124,125}, Emmanuel Jacob Ng'wiza¹²⁶, Casper Nim^{127,128,63}, Tim Noblet^{129,130,131}, Edel O'Hagan¹³², Mary O'Keefe¹³³, Crystian Bitencourt Oliveira^{66,67}, Cecilie K. Øverås^{134,135}, Patrick J. Owen^{26,27}, Feng Pan¹³⁶, Evangelos Pappas^{137,53}, Anupa Pathak⁶⁷, Kerry Peek⁵³, Romain S. Perera^{138,139}, Wajida Perveen¹⁴⁰, Andrés Pierobon¹⁴¹, Rafael Z. Pinto^{142,143}, Lisa G. Pont¹⁴⁴, Susanna Proudman^{145,146}, Firmansyah Purwanto¹⁴⁷, Jonathan G. Quicke^{148,149}, Helen E. Ramsay¹, Filippo Recenti^{150,151}, Michael Reddington¹⁵², Roland R. Reezigt^{37,153}, M.F. Reneman¹⁵⁴, Daniel Cury Ribeiro¹⁹, Bethan Richards^{66,123}, Raju Rijal⁹⁹, Ebonie Rio^{155,156,157}, Claire L. Samanna²⁷, Veronica Souza Santos¹¹⁴, Bruno T. Saragiotto^{143,158}, Michael Schneider¹⁵⁹, Dorte Schoeler Ziegler^{160,128}, Gwendolyn G.M. Scholten-Peeters³⁷, E. Michael Shanahan¹⁶¹, Saurab Sharma^{33,32,119,162,163}, Sweekriti Sharma^{66,67}, Stephen A. Sharp^{68,25}, Juscelio Pereira da Silva^{142,164}, Helen Slater¹⁹, Rob J.E.M. Smeets^{165,166,167}, Robyn Speerin¹⁶⁸, Geoffrey C. Speldewinde¹⁶⁹, Michael Steimling¹⁷⁰, Meenakshi Sundaram¹⁷¹, Michael S. Swain²⁵, Martin Underwood^{172,173}, Marianne Larsen van Gastel¹⁷⁴, Dina C. Janse van Rensburg¹⁷⁵, R. Vanderstraeten¹⁷⁶,

Lenny Vasanthan¹⁷⁷, Simon P. Vella^{66,67}, Carel Viljoen¹⁷⁸, Nina K. Vøllestad¹⁷⁹, Tom Vredevelde^{180,37}, Jason A. Wallis¹, Yuanyuan Wang⁴⁵, Ishanka Weerasekara¹⁸¹, Andrew D.H. Weiss^{118,119}, Peter D. Werth, Evert Onno Wesselink^{37,182}, Samuel L. Whittle^{1,183}, Louise K. Wiles¹⁸⁴, Hannah G. Withers^{25,68}, Lianne Wood¹⁸⁵, Anika Young²⁵, Joshua R. Zadro^{66,67}

1. Musculoskeletal Health and Wiser Health Care Units, School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia
2. Faculty of Medicine and Health, The University of Sydney
3. ZHAW Zurich University of Applied Sciences, School of Health Sciences, Winterthur, Switzerland
4. University of Lucerne, University Research Centre Health and Society, Lucerne, Switzerland
5. Department of Anesthesiology and Pain Management, Mount Sinai Hospital, Toronto, Canada
6. Department of Surgery, Faculty of Medicine, University of Toronto, Toronto, Canada
7. Physiotherapy Department, Wollongong Hospital, Illawarra Shoalhaven Local Health District, NSW, Australia
8. School of Medical Sciences, UNSW Medical School, University of New South Wales, Sydney, Australia
9. Karolinska Institutet, Institute of Environmental Medicine, Unit of Intervention and Implementation Research for Worker Health, Stockholm, Sweden
10. Bayero University Kano, Nigeria
11. Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Pharmacologie et évaluation des soins, AP-HP, Hôpital Pitié Salpêtrière, Centre de la douleur, Paris, France
12. Redcliffe Hospital, Redcliffe, Queensland, Australia
13. Faculty of Medicine, The University of Queensland, Brisbane, Australia
14. Arnold School of Public Health, University of South Carolina, Columbia, USA
15. Universidade cidade de São Paulo, UNICID, São Paulo, Brasil
16. School of Allied Health, Sport and Social Work, Griffith University, Brisbane and Gold Coast, Australia
17. The Bone Clinic, Brisbane, Australia
18. Department of Physiotherapy, Ariel University, Ariel, Israel
19. Curtin School of Allied Health, Faculty of Health Sciences, Curtin University, Perth, Australia
20. Center for Pain Research and Behavioral Health, University of Florida, Gainesville, Florida, USA
21. Department of Research and Innovation, Division of Clinical Neuroscience, Oslo University Hospital HF, Oslo, Norway
22. School of Health Sciences, Western Sydney University, Sydney, NSW, Australia
23. Translational Health Research Institute, Western Sydney University, Sydney, Australia
24. Diabetes Obesity Metabolism Translational Research Unit, Western Sydney University, Sydney, Australia
25. Department of Chiropractic, Faculty of Medicine, Health and Human Sciences, Macquarie University, Sydney, Australia
26. Eastern Health Emergency Medicine Program, Melbourne, Australia
27. Eastern Health Clinical School, Monash University, Melbourne, Australia
28. Department of Anesthesiology, Universidad del Cauca, Colombia; Department of Anesthesiology, Erasmus MC Rotterdam, The Netherlands
29. Consumer Advisory Group (CAG), Australia and New Zealand Musculoskeletal (ANZMUSC) Clinical Trials Network
30. Department of Physical Medicine and Rehabilitation, School of Medicine, University of Colorado Anschutz Medical Campus, Aurora CO, USA
31. School of Chiropractic, Universidad Central del Caribe, Bayamón, Puerto Rico
32. Centre for Pain IMPACT, Neuroscience Research Australia, Sydney, Australia
33. School of Health Sciences, Faculty of Medicine and Health, University of New South Wales, Sydney, Australia
34. Médecine physique et de réadaptation, Centre Hospitalier de Laval, Laval, France
35. Genesis Physical Therapy, Taipei City, Taiwan
36. HESAV School of Health Sciences - Vaud, HES-SO University of Applied Sciences and Arts Western Switzerland
37. Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam, Amsterdam Movement Sciences - Program Musculoskeletal Health, Amsterdam, The Netherlands
38. The University of Queensland's Clinical Trials Capability (ULTRA) Team, Centre for Clinical Research, Faculty of Health, Medicine and Behavioural Sciences, The University of Queensland, Brisbane, Australia
39. Centre for Medicine Use and Safety, Faculty of Pharmacy and Pharmaceutical Sciences, Monash University, Parkville, Australia
40. Department of Primary Care and General Practice, University of Otago, Wellington, New Zealand
41. School of Health, Medical and Applied Sciences, CQUniversity, Brisbane, Australia
42. La Trobe University, La Trobe Rural Health School, Department of Rural Allied Health, Australia
43. Practice Consultant for Physiotherapy, Department for Quality and Medicine, Region Centre, Denmark
44. Falls, Balance and Injury Research Centre, Neuroscience Research Australia, Sydney, Australia

45. School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia
46. Department of Integrative Medicine, Chris O'Brien Lifehouse Hospital, Sydney, Australia
47. Department of Social and Preventive Medicine, Faculty of Medicine, Université Laval, Québec City (QC), Canada
48. School of Medicine and Psychology, Australian National University, Canberra, Australia
49. Discipline of Physiotherapy, College of Healthcare Sciences, James Cook University, Townsville, Australia
50. Department of Physical Therapy & Athletic Training, University of Utah, Salt Lake City, Utah, USA
51. Surgical Specialities and Anesthesiology Department, Botucatu Medical School, São Paulo State University (UNESP), Botucatu, Brazil
52. Composite Rehabilitation Centre for Skill Development, Rehabilitation, and Empowerment of Persons with Disabilities, Lucknow, India
53. Sydney School of Health Sciences, Faculty of Medicine and Health, The University of Sydney, Sydney, Australia
54. Division of Rheumatology, Department of Medicine, University Hospitals of Geneva, Switzerland
55. Adelaide Medical School, The University of Adelaide, Adelaide, Australia
56. Department of Orthopaedic Surgery, School of Medicine, Duke University, Durham, North Carolina, USA
57. CHRC, Comprehensive Health Research Centre, NOVA Medical School, Universidade NOVA de Lisboa, Lisbon, Portugal
58. Departamento de Fisioterapia, Escola Superior de Saúde, Instituto Politécnico de Setúbal, Setúbal, Portugal
59. Department of Medicine, University of Otago Wellington, Wellington, New Zealand
60. Department of Public Health and Nursing, Norwegian University of Science and Technology (NTNU), Trondheim, Norway
61. Clinic of Rehabilitation, St Olav's Hospital, Trondheim University Hospital, Trondheim, Norway
62. Victor Horsley Department of Neurosurgery, University College London Hospital, Queen Square, London, UK
63. Center for Muscle and Joint Health, Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark
64. Department of Physical Therapy, University of Alberta, Edmonton, Canada
65. Centre for Health, Exercise and Sports Medicine, Department of Physiotherapy, School of Health Sciences, The University of Melbourne, Melbourne, Australia
66. Institute for Musculoskeletal Health, Sydney Local Health District, Sydney, NSW, Australia
67. Sydney School of Public Health, Faculty of Medicine and Health, The University of Sydney, Sydney, Australia
68. Department of Health Sciences and Spinal Pain Research Centre, Macquarie University, Sydney, Australia
69. School of Clinical Medicine, UNSW Medicine & Health, UNSW Sydney, Australia
70. Chiropractic Knowledge Hub, Odense, Denmark
71. Department of Rheumatology, Green Life Medical College, Dhaka, Bangladesh
72. Department of Community Health & Epidemiology, Dalhousie University, Halifax, Nova Scotia, Canada
73. Faculty of Kinesiology, University of New Brunswick, Fredericton, New Brunswick, Canada
74. School of Allied Health, Murdoch University, Murdoch, Western Australia, Australia
75. Research Unit of Clinical Biomechanics, Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark
76. Musculoskeletal Epidemiology Research Group, University of Zurich and Balgrist University Hospital, Zurich, Switzerland
77. Epidemiology, Biostatistics and Prevention Institute (EBPI), University of Zurich, Zurich, Switzerland
78. University Spine Centre Zurich (UWZH), Balgrist University Hospital, University of Zurich, Zurich, Switzerland
79. Dalla Lana School of Public Health, University of Toronto, Toronto, Canada
80. Epsom and St Helier University Hospitals NHS Trust, London, UK
81. Department of Primary Care and Clinical Simulation, University of Otago, Christchurch, New Zealand
82. Physiotherapy Department, Tishk International University, Erbil, Iraq
83. Department of Health Sciences, Faculty of Science and Amsterdam Movement Sciences Research Institute, Vrije Universiteit, Amsterdam, The Netherlands
84. Master Program in Physiotherapy and Functioning, Department of Physiotherapy, Universidade Federal do Ceará, Fortaleza, Brazil
85. Belgian Health Care Knowledge Centre, Brussels, Belgium
86. School of Health Sciences, The University of Sydney, Sydney, Australia
87. Nepean Blue Mountains Local Health District, Sydney, Australia
88. Chatham University, Pittsburgh, USA
89. Research Unit of Health Sciences and Technology, University of Oulu, Oulu, Finland
90. Rehabilitation Services of Wellbeing Services County of South Karelia, Lappeenranta, Finland

91. Department of Occupational Health, Injury Care Australia, Epping, New South Wales, Australia
92. Department of General Practice, Erasmus MC, Rotterdam, The Netherlands
93. Department of Public and Occupational Health, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands
94. University of the Sunshine Coast, Birtinya, Queensland, Australia
95. Whitlam Orthopaedic Research Centre, Ingham Institute for Applied Medical Research, Sydney, Australia
96. Department of Physical Therapy, São Paulo State University (UNESP), Presidente Prudente, Brazil
97. Department of Physical Therapy, Faculdade Israelita de Ciências da Saúde Albert Einstein (FICSAE), São Paulo, Brazil
98. Duke Clinical Research Institute, Duke University, Durham, USA
99. BP Koirala Institute of Health Sciences, Dharan, Nepal
100. CHU Ucl Namur, Dinant, Belgium
101. Physiotherapy Department, School of Primary and Allied Health Care, Monash University, Melbourne, Australia
102. Centre for Health, Activity and Rehabilitation Research, School of Physiotherapy, University of Otago, Dunedin, New Zealand
103. Pain@Otago Research Theme, University of Otago, Dunedin, New Zealand
104. Centre for Health and Rehabilitation Research, University of Auckland, New Zealand
105. Department of Exercise Sciences, University of Auckland, New Zealand
106. School of Health and Life Sciences, Teesside University, UK
107. NIHR Applied Research Collaboration (ARC) North East and North Cumbria, UK
108. Sydney Pharmacy School, Faculty of Medicine and Health, The University of Sydney, Sydney, Australia
109. King's College Hospital NHS Foundation Trust, London, UK
110. King's College London, London, UK
111. Ankylosing Spondylitis Victoria Inc., Victoria, Australia
112. The University of Queensland, St Lucia, Queensland, Australia
113. Gallipoli Medical Research, Greenslopes, Queensland, Australia
114. School of Rehabilitation Science, Faculty of Health Sciences, McMaster University, Hamilton, Ontario, Canada
115. Postgraduate Program in Rehabilitation Sciences, Augusto Motta University Centre - UNISUAM, Rio de Janeiro, Brazil
116. Musculoskeletal Research Unit, Department of Rehabilitation Sciences, Faculty of Movement and Rehabilitation Sciences, Leuven University, Belgium
117. Department of Health Sciences, Ribeirão Preto Medical School, Universidade de São Paulo, Ribeirão Preto, São Paulo, Brazil
118. Northern Clinical School, Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia
119. Pain Management Research Institute, Kolling Institute, Faculty of Medicine and Health, The University of Sydney and Northern Sydney Local Health District, Sydney, NSW, Australia
120. Physical Therapy Department, Faculty of Public Health, Antonine University, Lebanon
121. Physiotherapy Department, Bendigo Health, Bendigo, Australia
122. National Institute of Rehabilitation "Luis Guillermo Ibarra Ibarra", Mexico
123. Department of Rheumatology, Royal Prince Alfred Hospital, Sydney, Australia
124. University of New South Wales, Sydney, Australia
125. University of Notre Dame, Sydney, Australia
126. Muhimbili Orthopedic Institute, Tanzania
127. Spine Centre of Southern Denmark, University Hospital of Southern Denmark, Kolding, Denmark
128. Department of Regional Health Research, University of Southern Denmark, Odense, Denmark
129. St George's University Hospitals NHS Foundation Trust, London, UK
130. School of Physical Therapy, Western University, Canada
131. Faculty of Medicine, Health and Human Sciences, Macquarie University, Sydney, Australia
132. Westmead Applied Research Centre, Faculty of Medicine and Health, The University of Sydney, Western Sydney (Baludarri) Precinct, Westmead, Australia
133. School of Public Health, Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland
134. Department of Public Health and Nursing, Norwegian University of Science and Technology, Trondheim, Norway
135. The Norwegian Chiropractors' Research Foundation - Et Liv i Bevegelse (ELiB), Oslo, Norway
136. Menzies Institute for Medical Research, University of Tasmania
137. School of Health and Biomedical Sciences, Royal Melbourne Institute of Technology (RMIT) University, Melbourne, Australia
138. Department of Cell and Developmental Biology, University College London, London, UK
139. Center for Sport and Exercise Medicine, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka

140. Department of Physical Therapy, School of Allied Health Sciences, CMH Lahore Medical College & Institute of Dentistry (NUMS) Lahore, Pakistan
141. Department of Primary Health Care and General Practice, University of Otago, Wellington, New Zealand
142. Department of Physical Therapy, Universidade Federal de Minas Gerais (UFMG), Belo Horizonte, Minas Gerais, Brazil
143. Discipline of Physiotherapy, Graduate School of Health, Faculty of Health, University of Technology Sydney, Sydney, Australia
144. Graduate School of Health, Faculty of Health, University of Technology Sydney, Sydney, Australia
145. Rheumatology Unit, Royal Adelaide Hospital, Adelaide, Australia
146. Discipline of Medicine, The University of Adelaide, Adelaide, Australia
147. Department of Physiotherapy, Dustira Hospital Institute of Health Sciences, West Java, Indonesia
148. Surgical Treatment and Rehabilitation Service (STARS) Education and Research Alliance, The University of Queensland and Metro North Health, Herston, Brisbane, Queensland, Australia
149. Centre for Innovation in Pain and Health Research, School of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Australia
150. Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health, University of Genova, Genoa, Italy
151. Department of Clinical Sciences Lund, Orthopedics, Clinical Epidemiology Unit, Lund University, Lund, Sweden
152. Sheffield Teaching Hospitals NHS Trust, Sheffield, UK
153. Academy of Health, Department of Physical Therapy, Hanze University of Applied Sciences, Groningen, The Netherlands
154. University of Groningen, University Medical Centre Groningen, Department of Rehabilitation Medicine, Groningen, The Netherlands
155. La Trobe Sport and Exercise Medicine Research Centre, La Trobe University, Bundoora, Australia
156. The Victorian Institute of Sport, Melbourne, Australia
157. The Australian Ballet, Melbourne, Australia
158. Masters and Doctoral Programs in Physical Therapy, Universidade Cidade de Sao Paulo, Brazil
159. Doctor of Chiropractic Program, University of Pittsburgh, Pittsburgh, USA
160. Medical Spinal Research Unit, University Hospital of Southern Denmark, Kolding, Denmark
161. Flinders University, Bedford Park, South Australia, Australia
162. Pain Management and Research Centre, Royal North Shore Hospital, Northern Sydney Local Health District, Sydney, Australia
163. Department of Physiotherapy, Manipal Academy of Higher Education, Manipal University, Manipal, India
164. Graduate Program in Rehabilitation Sciences, Universidade Federal de Alfenas, Alfenas, Brazil
165. Department of Rehabilitation Medicine, Care and Public Health Research Institute (CAPHRI), Faculty of Health, Life Sciences and Medicine, Maastricht University, Maastricht, The Netherlands
166. Clinics in Revalidatie (CIR), The Netherlands
167. Pain in Motion International Research Group (PiM)
168. The University of Sydney, Sydney, Australia
169. Capital Pain and Rehabilitation Clinic, Canberra, Australia
170. School of Rehabilitation Sciences, Moravian University, Pennsylvania, USA
171. Department of Physical Therapy, University of Pittsburgh, Pittsburgh, USA
172. Warwick Clinical Trials Unit, Coventry, UK
173. University Hospitals Coventry and Warwickshire, Coventry, UK
174. Department of Physical Therapy, Ziekenhuis Amstelland, Amstelveen, The Netherlands
175. Section Sports Medicine and SEMLI, Faculty of Health Science, University of Pretoria, Pretoria, South Africa
176. University of Antwerp, Faculty of Medicine and Health Sciences (MOVANT), Antwerp, Belgium
177. Physiotherapy, Department of Physical Medicine and Rehabilitation, Christian Medical College, Vellore, India
178. Department of Physiotherapy, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa
179. Department of Public Health Science and Interdisciplinary Science, Institute of Health and Society, University of Oslo, Norway
180. Centre of Expertise Urban Vitality, Faculty of Health, Sport and Physical Activity, Amsterdam University of Applied Sciences, The Netherlands
181. Institute of Health and Wellbeing, Federation University Australia, Melbourne, Australia
182. Division of Pain Medicine, Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University School of Medicine, Palo Alto, CA, USA
183. Rheumatology Unit, The Queen Elizabeth Hospital, Woodville South, Australia
184. IIMPACT in Health, Allied Health and Human Performance, University of South Australia, Kaurna Country, Adelaide, Australia
185. Department of Public Health and Sports Science, Faculty of Health and Life Sciences, University of Exeter, Exeter, UK

Funding

Bayden McKenzie is supported by a PhD scholarship from the Chiropractic Australia Research Foundation and a top-up scholarship from the National Health and Medical Research Council (NHMRC) Australia & New Zealand Low Back Pain Research Network Centre of Research Excellence (ANZBACK CRE; [APP1171459](#)). Giovanni Ferreira is supported by an NHMRC Early Leadership Fellowship ([APP2009808](#)), Chris Maher and Rachelle Buchbinder are supported by NHMRC Leadership Grants ([APP1194283](#) and [APP1194483](#) respectively). The funding bodies were not involved in any aspect of the study design, collection, analysis or interpretation of the data or writing of the manuscript.

Data availability

All data are available upon reasonable request to the corresponding author.

Declaration of competing interest

One or more of the authors declare financial or professional relationships on ICMJE-TSJ disclosure forms.

CRediT authorship contribution statement

Bayden J. McKenzie: Writing – original draft, Visualization, Project administration, Methodology, Formal analysis, Data curation, Conceptualization. **Giovanni E. Ferreira:** Writing – review & editing, Supervision, Methodology, Investigation, Formal analysis, Conceptualization. **Romi Haas:** Writing – review & editing, Supervision, Methodology, Investigation, Formal analysis, Conceptualization. **Alexandra Gorelik:** Writing – review & editing, Visualization, Validation, Formal analysis, Data curation. **Chris G. Maher:** Writing – review & editing, Supervision, Methodology, Formal analysis, Conceptualization. **Rachelle Buchbinder:** Writing – review & editing, Visualization, Supervision, Methodology, Investigation, Formal analysis, Conceptualization.

Acknowledgments

We acknowledge and thank all Delphi participants for their contribution to this work. A special thanks to the consumer advocacy groups of ANZMUSC and ANZBACK, Arthritis Australia, Arthritis New Zealand, and the Consortium for Low Back Pain in Low- and Middle-Income Countries for promoting our study across your networks.

Supplementary materials

Supplementary material associated with this article can be found in the online version at <https://doi.org/10.1016/j.spinee.2025.10.003>.

References

- [1] Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F, et al. A systematic review of the global prevalence of low back pain. *Arthritis Rheum* 2012;64(6):2028–37.
- [2] Ferreira ML, De Luca K, Haile LM, Steinmetz JD, Culbreth GT, Cross M, et al. Global, regional, and national burden of low back pain, 1990–2020, its attributable risk factors, and projections to 2050: a systematic analysis of the Global Burden of Disease Study 2021. *Lancet Rheumatol* 2023;5(6):e316–e29.
- [3] Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, et al. What low back pain is and why we need to pay attention. *Lancet* 2018;391(10137):2356–67.
- [4] Foster NE, Anema JR, Cherkin D, Chou R, Cohen SP, Gross DP, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *Lancet* 2018;391(10137):2368–83.
- [5] Buchbinder R, van Tulder M, Öberg B, Costa LM, Woolf A, Schoene M, et al. Low back pain: a call for action. *Lancet* 2018;391(10137):2384–8.
- [6] Australian Institute of Health and Welfare [Internet]. Health labour force national minimum data set: National Health Data Dictionary version 12. Canberra ACT: AIHW; 2008. Available at: <https://www.aihw.gov.au/reports/workforce/health-labour-force-nmds/contents/table-of-contents>. Accessed November 13, 2024.
- [7] Australian Institute of Health and Welfare [Internet]. Data set specifications and national minimum data sets. Canberra ACT: Meteor Metadata Online Registry; 2024. Available at: <https://meteor.aihw.gov.au/content/344846>. Accessed December 2, 2024.
- [8] Australian Institute of Health and Welfare [Internet]. Admitted patient care national minimum data set: national health data dictionary version 12. Canberra ACT: AIHW; 2003. Available at: <https://www.aihw.gov.au/reports/hospitals/admitted-patient-care-nmds/summary>. Updated 2024. Accessed November 13, 2024.
- [9] Health New Zealand Te Whatu Ora [Internet]. National Minimum Dataset (hospital events). Wellington: Te Kāwanatanga o Aotearoa New Zealand Government; 1999. Available at: <https://www.aihw.gov.au/reports/hospitals/admitted-patient-care-nmds/summary>. Updated 2024. Accessed November 13, 2024.
- [10] NHS England [Internet]. Data sets. Leeds, West Yorkshire: NHS England; 2024. Available at: <https://digital.nhs.uk/data-and-information/data-collections-and-data-sets/data-sets>. Accessed November 13, 2024.
- [11] Freguia F, Danielis M, Moreale R, Palese A. Nursing minimum data sets: Findings from an umbrella review. *J Health Informatics* 2022;28(2):14604582221099826.
- [12] Sermeus W, van den Heede K, Michiels D, Delesie L, Thonon O, Van Boven C, et al. Revising the Belgian Nursing Minimum Dataset: from concept to implementation. *Int J Med Inform* 2005;74(11–12):946–51.
- [13] Bryant E, Broomfield C, Burrows J, McLean S, Marks P, Maloney D, et al. Gaining consensus on clinical quality outcomes for eating disorders: Framework for the development of an Australian national minimum dataset. *BMJ Open* 2023;13(4):e071150.
- [14] Safdari R, Saeedi MG, Masoumi-Asl H, Rezaei-Hachesu P, Mirmia K, Mohammadzadeh N, et al. National minimum data set for antimicrobial resistance management: toward global surveillance system. *Iran J Med Sci* 2018;43(5):494.
- [15] Christey G, Warren J, Palmer CS, Burrell M, Vallmuur K. Development of a standardized minimum dataset for including low-severity trauma patients in trauma registry collections in Australia and Aotearoa New Zealand. *ANZ J Surg* 2023;93(3):572–6.
- [16] Australian Commission for Safety and Quality in Health Care. Low back pain clinical care standard. Sydney: ACSQHC; 2022. Available at: <https://www.safetyandquality.gov.au/standards/clinical-care-standards/low-back-pain-clinical-care-standard>. Accessed November 28, 2024.

- [17] Health Quality Ontario. Low back pain care for adults with acute low back pain Available at: <https://www.ontariohealth.ca/providing-health-care/clinical-quality-improvement>; 2019. Accessed January 18, 2025.
- [18] National Institute for Health and Care Excellence. Low back pain and sciatica in over 16s (Quality standard [QS155]) Available at: <https://www.nice.org.uk/guidance/qs155>; 2017. Accessed January 14, 2025.
- [19] Kelley E., Hurst J. OECD health working papers No. 23. Health care quality indicators project: conceptual framework paper Available at: https://www.oecd.org/content/dam/oecd/en/publications/reports/2006/03/health-care-quality-indicators-project_g17a17b5/440134737301.pdf; 2006. Accessed September 7, 2025.
- [20] Dos Santos VL, Sato KS, Maher CG, Vidal RV, Grande GH, Costa LOP, et al. Clinical indicators to monitor health care in low back pain: a scoping review. *Int J Qual Health Care* 2024;36(2):mzae044. <https://doi.org/10.1093/intqhc/mzae044>.
- [21] McKenzie B.J., Ferreira G.E., Haas R., Di Donato M., Maher C.G., Buchbinder R. Routinely collected indicators to monitor low back pain burden and its care in Australia: A scoping review. [Manuscript under review].
- [22] Barrett D, Heale R. What are Delphi studies? *Evid Based Nurs* 2020;23(3):68–9.
- [23] Nasa P, Jain R, Juneja D. Delphi methodology in healthcare research: how to decide its appropriateness. *World J Methodol* 2021;11(4):116.
- [24] McKenzie BJ, Haas R, Ferreira GE, Maher CG, Buchbinder R. The environmental impact of health care for musculoskeletal conditions: a scoping review. *PLoS One* 2022;17(11):e0276685. <https://doi.org/10.1371/journal.pone.0276685>.
- [25] Okoli C, Pawlowski SD. The Delphi method as a research tool: an example, design considerations and applications. *Inf Manage J* 2004;42(1):15–29.
- [26] Manyara AM, Purvis A, Ciani O, Collins GS, Taylor RS. Sample size in multistakeholder Delphi surveys: at what minimum sample size do replicability of results stabilize? *J Clin Epidemiol* 2024;174:111485.
- [27] McKenzie BJ, Haas R, Ferreira GE, Gorelik A, Cyril S, Han JX, Gilbert S, Grobler L, Maher CG, Buchbinder R. Agreement between high-quality clinical practice guidelines in their treatment recommendations for low back pain: A systematic review. *Spine J* 2025. <https://doi.org/10.1016/j.spinee.2025.07.015>.
- [28] Australia and New Zealand Musculoskeletal (ANZMUSC). Clinical Trials Network [Internet]. Melbourne, Australia: Optimising Musculoskeletal Health; 2024. Available at: <https://anzmusc.org>. Accessed November 28, 2024.
- [29] Sharma S, McAuley JH. Low back pain in low-and middle-income countries, part 1: the problem. *J Orthop Sports Phys Ther* 2022;52(5):233–5.
- [30] National Spine Network [Internet]. Holborn, London. Available at: <https://www.nationalspinenetwork.co.uk>; 2024. Accessed November 28, 2024.
- [31] Mauck MC, Lotz J, Psioda MA, Carey TS, Clauw DJ, Majumdar S, et al. The back pain consortium (BACPAC) research program: structure, research priorities, and methods. *Pain Medicine* 2023;24(Supplement 1):S3–12. <https://doi.org/10.1093/pm/pnac202>.
- [32] Australia and New Zealand Low Back Pain Research Network (ANZ-BACK). Melbourne, Australia. Available at: <https://www.anzback.net/>. 2024. Accessed November 28, 2024.
- [33] Arthritis Australia [Internet]. Glebe, NSW, Australia Available at: <https://arthritisaustralia.com.au>; 2024. Accessed November 28, 2024.
- [34] Arthritis New Zealand [Internet]. Wellington, New Zealand Available at: <https://www.arthritis.org.nz/>; 2024. Accessed November 28, 2024.
- [35] Musculoskeletal Health Australia [Internet]. Elsternwick, Victoria, Australia Available at: <https://muscha.org/>; 2024. Accessed November 28, 2024.
- [36] Cochrane Musculoskeletal [Internet] Available at: <https://musculoskeletal.cochrane.org>; 2024. Accessed November 28, 2024.
- [37] Keeney E, Thom H, Turner E, Martin RM, Sanghera S. Using a modified Delphi approach to gain consensus on relevant comparators in a cost-effectiveness model: application to prostate cancer screening. *Pharmacoeconomics* 2021;39(5):589–600.
- [38] Williams MT, Lewthwaite H, Brooks D, Jensen D, Abdallah SJ, Johnston KN. Chronic breathlessness explanations and research priorities: findings from an international Delphi survey. *J Pain Symp Manage* 2020;59(2):310–319.e12.
- [39] Slade SC, Dionne CE, Underwood M, Buchbinder R. Standardised method for reporting exercise programmes: protocol for a modified Delphi study. *BMJ Open* 2014;4(12):e006682.
- [40] Hepworth LR, Rowe FJ. Using Delphi methodology in the development of a new patient-reported outcome measure for stroke survivors with visual impairment. *Brain Behav* 2018;8(2):e00898.
- [41] Diamond IR, Grant RC, Feldman BM, Pencharz PB, Ling SC, Moore AM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67(4):401–9.
- [42] Schifano J, Niederberger M. How Delphi studies in the health sciences find consensus: a scoping review. *Syst Rev* 2025;14(1):14.
- [43] Jayawardana S, Forman R, Johnston-Webber C, Campbell A, Berterame S, de Joncheere C, et al. Global consumption of prescription opioid analgesics between 2009 and 2019: a country-level observational study. *EclinicalMedicine* 2021;42:101198. <https://doi.org/10.1016/j.eclinm.2021.101198>.
- [44] Laur C, Kithulegoda N, McCleary N, Nicholas Angl E, Strange M, Sklar B, et al. Changing or validating physician opioid prescribing behaviors through audit and feedback and academic detailing interventions in primary care. *Implement Res Pract* 2025;6:26334895241307638.
- [45] Barnett MJ, Clubbs B, Woodyard A, Wasem V, Prinze J, Tirupasur A, et al. Barriers and motivational factors for engaging in novel opioid and medication disposal-related services in community pharmacies. *Res Social Adm Pharm* 2024;20(8):713–22.
- [46] Canadian Institute of Health Information (CIHI) [Internet]. National Ambulatory Care Reporting System (NACRS) metadata. Ottawa, Ontario: CIHI; 2025. Available at: <https://www.cihi.ca/en/national-ambulatory-care-reporting-system-nacrs-metadata>. Accessed January 18, 2025.
- [47] Ontario Health [Internet]. Data Catalogue: Datasets. Toronto, Ontario: Ministry of Health; 2025. Available at: https://data.ontario.ca/dataset/?keywords_en=Health. Updated 2024; Accessed January 18, 2025.
- [48] Ontario Health [Internet]. Narcotics monitoring system. Toronto, Ontario: Ministry of Health; 2025. Available at: <https://www.ontario.ca/page/narcotics-monitoring-system>. Updated October 31, 2024; Accessed January 18, 2025.
- [49] Paridaens R, Vaes B, Van den Bulck S, Soetaert J. Benchmarks for low back pain in general practice in Flanders: electronic audit of INTEGEO. *BMC Prim Care* 2024;25(1):431.
- [50] Deyo RA, Dworkin SF, Amtmann D, Andersson G, Borenstein D, Carragee E, et al. Report of the NIH Task Force on research standards for chronic low back pain. *Phys Ther* 2015;95(2):e1–e18.
- [51] Verburg A, Van Dulmen S, Kiers H, Nijhuis-Van Der Sanden M, Van Der Wees P. Development of a standard set of outcome measures for non-specific low back pain in Dutch primary care physiotherapy practices: a Delphi study. *Eur Spine J* 2019;28(7):1550–64.
- [52] Clement RC, Welander A, Stowell C, Cha TD, Chen JL, Davies M, et al. A proposed set of metrics for standardized outcome reporting in the management of low back pain. *Acta Orthop* 2015;86(5):523–33.
- [53] Hyatt A, Gough K, Chung H, Wood W, Aston R, Cockwill J, et al. Development of consensus quality indicators for cancer supportive care: a Delphi study and pilot testing. *BMC Health Serv Res* 2024;24(1):377.
- [54] Aktaa S, Batra G, Wallentin L, Baigent C, Erlinge D, James S, et al. European Society of Cardiology methodology for the development of quality indicators for the quantification of cardiovascular care and outcomes. *EHJ-QCCO* 2022;8(1):4–13.